



HAL
open science

Del Pezzo quintics as equivariant compactifications of vector groups

Adrien Dubouloz, Takashi Kishimoto, Pedro Montero

► **To cite this version:**

Adrien Dubouloz, Takashi Kishimoto, Pedro Montero. Del Pezzo quintics as equivariant compactifications of vector groups. 2022. <hal-03772297>

HAL Id: hal-03772297

<https://hal.science/hal-03772297v1>

Preprint submitted on 8 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

DEL PEZZO QUINTICS AS EQUIVARIANT COMPACTIFICATIONS OF VECTOR GROUPS

ADRIEN DUBOULOZ, TAKASHI KISHIMOTO, AND PEDRO MONTERO

ABSTRACT. We study faithful actions with a dense orbit of abelian unipotent groups on quintic del Pezzo varieties over a field of characteristic zero. Such varieties are forms of linear sections of the Grassmannian of planes in a 5-dimensional vector space. We characterize which smooth forms admit these types of actions and show that in case of existence, the action is unique up to equivalence by automorphisms. We also give a similar classification for mildly singular quintic del Pezzo threefolds and surfaces.

INTRODUCTION

Vector group varieties are defined by analogy to toric varieties as being varieties X endowed with an effective action of an abelian unipotent group $\mathbb{U} \cong \mathbb{G}_a^n$ with a Zariski dense open orbit. For varieties defined over a field of characteristic zero, the group \mathbb{U} then embeds equivariantly as the open orbit, making X into a partial equivariant completion of \mathbb{U} . The study of such equivariant completions which are Fano varieties was initiated by Hassett and Tschinkel in [21] with views towards Manin's conjecture on the asymptotic distribution of rational points of bounded height over number fields (see e.g. [9, 10]). Over algebraically closed fields of characteristic zero, besides projective spaces of every dimension, many families of Fano varieties and other Mori fibers spaces including for instance smooth projective quadrics, Grassmannians and flag varieties are known to be vector group varieties, see e.g. [1, 3, 17, 18, 22, 31, 37].

Of particular interest in this context is the question of classification of possible equivalence classes of structures of vector group variety up to isomorphisms on a given variety. Indeed, it was observed by Hassett and Tschinkel [21] that in contrast to toric structures, vector group variety structures on $\mathbb{P}_{\mathbb{C}}^n$, $n \geq 2$, are not unique and that for $n \geq 6$ there are even infinitely many equivalence classes of such structures. In contrast, it is known that over algebraically closed fields of characteristic zero, Grassmannians other than projective spaces [3] and smooth quadrics [37] admit a unique equivalence class of structure of vector group variety. For Fano varieties of Picard rank one over the field of complex numbers, Fu and Hwang [15] gave a uniform characterization of uniqueness of vector group variety structures in terms in the smoothness of the Variety of Minimal Rational Tangents (VMRT) of a dominating family of minimal rational curves.

In this article, we consider the problem of existence and uniqueness of vector group variety structures on smooth and mildly singular quintic del Pezzo varieties over an arbitrary field k of characteristic zero. By definition, a *smooth quintic del Pezzo variety* is a smooth geometrically connected projective k -variety X whose base extension $X_{\bar{k}} = X \times_{\mathrm{Spec}(k)} \mathrm{Spec}(\bar{k})$ to an algebraic closure \bar{k} of k has an ample invertible sheaf \mathcal{L} of degree 5 such that $\omega_{X_{\bar{k}}}^{\vee} \cong \mathcal{L}^{\otimes(n-1)}$. If $n = 2$ then $X_{\bar{k}}$ is a del Pezzo surface of degree 5. On the other hand, by [19], a smooth quintic del Pezzo k -variety X exists if and only if $n \leq 6$ and furthermore, for each $n = 3, 4, 5, 6$, $X_{\bar{k}}$ is unique up to isomorphism, isomorphic to any smooth linear section of the Grassmannian $G(2, 5) \subset \mathbb{P}_{\bar{k}}^9$ of 2-dimensional vector subspaces of $\bar{k}^{\oplus 5}$. For $n \leq 3$, the automorphism group of $X_{\bar{k}}$ is either finite if $n = 2$ or isomorphic to $\mathrm{PGL}_2(\bar{k})$ if $n = 3$ (see e.g. [7, Proposition 7.1.10]). In particular, it cannot contain any abelian unipotent subgroup defining a vector group variety structure on X . Over the field of complex numbers, existence and uniqueness of vector group variety structures on the remaining varieties, which are all Fano of Picard rank 1, has been settled affirmatively by Fu and Hwang as a consequence of a series of articles [16, 17] devoted to the broader study using VMRT techniques of so-called Euler-symmetric projective varieties.

Here, since we work over arbitrary fields k of characteristic zero, possibly non-closed, the actual question becomes to determine and classify k -forms of vector group variety structures on del Pezzo quintics. Of course, the non-existence of such structures after base extension to an algebraic closure is a clear obstruction for existence of these structures over the given base field, But on the other hand, neither the existence nor its combination with uniqueness up to equivalence after base extension is enough in general to conclude, say by Galois descent arguments, the existence of such structures defined over the base field. For instance, a smooth n -dimensional quadric in $\mathbb{P}_{\mathbb{Q}}^{n+1}$ without \mathbb{Q} -rational point does not admit any vector group variety structure defined over \mathbb{Q} , even though its base extension to $\bar{\mathbb{Q}}$ admits infinitely many such structures, which, as a consequence of [37], are all equivalent. Our approach is thus by necessity different from that using

VMRT techniques in [16, 17, 18], which, in particular, depend on the Cartan-Fubini analytic extension theorem [24] that has no arguably straightforward counterpart over arbitrary fields. Instead we build on elementary birational geometry of the Grassmannian $G(2, 5)$, its linear Schubert sub-varieties and their associated rational projections, a material which, over algebraically closed fields, goes back to a classical article of Todd [41]. Our first main result is a classification of k -forms of vector group variety structures on smooth del Pezzo quintics of dimension ≥ 4 which can be summarized as follows:

Theorem 1. *For a smooth quintic del Pezzo k -variety X_n of dimension $n \in \{4, 5, 6\}$, the following hold:*

- 1) *If $n = 6$ then X_6 admits a vector group variety structure if and only if it has a k -rational point. If so, X_6 is isomorphic to $G(2, 5) \subset \mathbb{P}_k^9$ and it admits a unique vector group variety structure up to equivalence.*
- 2) *If $n = 4, 5$ then X_n is unique up to isomorphism, isomorphic to any smooth section of $G(2, 5) \subset \mathbb{P}_k^9$ by a linear subspace of codimension $6 - n$. Furthermore, it admits precisely one vector group variety structure up to equivalence.*

In a second step, we apply the same methods to k -forms of vector group variety structures on mildly singular del Pezzo quintic threefolds and surfaces. The case of quintic del Pezzo surfaces with canonical singularities has already been fully settled by Derenthal and Loughran [12] who studied vector group variety structures on del Pezzo surfaces with canonical singularities over arbitrary fields of characteristic zero. We therefore mainly focus on the case of quintic del Pezzo threefolds with terminal singularities. We obtain the following characterization, which says in particular that in contrast to smooth del Pezzo quintics of dimension 4 and 5 and canonical del Pezzo quintic surfaces which have no non-trivial k -forms for any field k , trinodal del Pezzo quintics threefolds do in general have non-trivial k -forms:

Theorem 2. *A quintic del Pezzo threefold X_3 with terminal singularities admits a vector group variety structure if and only if its base extension to \bar{k} has precisely three ordinary double points. In this case, the vector group structure is unique up to isomorphism.*

Furthermore, isomorphism classes of such threefolds are in one-to-one correspondence with $\mathrm{PGL}_2(k)$ -orbits of smooth 0-dimensional sub-schemes of \mathbb{P}_k^1 of length three.

The article is organized as follows. In the first section we collect standard facts on Grassmannians and basic properties of vector groups and their actions on varieties. Section two is devoted to a review of certain classes of linear Schubert sub-varieties of the Grassmannian $G(2, 5)$ and its smooth linear sections, and of their associated rational linear projections. These preliminary results are then applied in the third and fourth sections to derive the proofs of Theorem 1 and Theorem 2, respectively.

Acknowledgements. This research was initiated during the stay of the first and the third authors at Saitama University in March 2020 at the occasion of the last in-person workshop “Affine and Birational Geometry” held before the Covid-19 pandemic.

The first author was partially supported by the French ANR project "FIBALGA" ANR-18-CE40-0003 and acknowledge the support of the EIPHI Graduate School ANR-17-EURE-0002 to the Institute of Mathematics of Burgundy. The second author was partially funded by JSPS KAKENHI Grant Number 19K03395. The third author was partially supported by Fondecyt ANID projects 11190323 and 1200502.

1. PRELIMINARIES

We work over a field k of characteristic zero, with a fixed algebraic closure \bar{k} and associated Galois group $\Gamma = \mathrm{Gal}(\bar{k}/k)$. We consider Γ as the profinite group $\varprojlim \mathrm{Gal}(k'/k)$ endowed with the profinite topology, the limit being taken over the directed set of finite Galois extension $k \subset k'$ of k , each group $\mathrm{Gal}(k'/k)$ being endowed with the discrete topology. A k -variety is an integral k -scheme of finite type.

1.1. Notation and conventions. For vector bundles and projective bundles, we follow [20]. Namely, given a quasi-coherent sheaf \mathcal{F} on a scheme X , we let $\mathrm{Sym}\mathcal{F}$ be the symmetric algebra of \mathcal{F} and we denote by $p : \mathbb{V}_X(\mathcal{F}) = \mathrm{Spec}_X(\mathrm{Sym}\mathcal{F}) \rightarrow X$ the “vector bundle” over X associated to \mathcal{F} and by

$$\pi : \mathbb{P}_X(\mathcal{F}) = \mathrm{Proj}_X(\mathrm{Sym}\mathcal{F}) \cong (\mathbb{V}_X(\mathcal{F}) \setminus 0_X) / \mathbb{G}_{m,X} \rightarrow X,$$

where 0_X is the zero section of p , its associated “projective bundle”. We denote by $\mathcal{O}_{\mathbb{P}_X(\mathcal{F})}(1)$ the canonical coherent invertible sheaf of $\mathcal{O}_{\mathbb{P}_X(\mathcal{F})}$ -modules associated to $\mathrm{Sym}\mathcal{F}$ viewed as graded sheaf of modules over itself with the degree shifted by 1. When \mathcal{E} is a coherent locally free sheaf, $\mathbb{V}_X(\mathcal{E})$ is a usual Zariski locally trivial vector bundle of finite rank and $\mathbb{P}_X(\mathcal{E})$ is its associated projective bundle of lines.

Given a quasi-coherent sheaf \mathcal{F} on a scheme X and an integer $d \geq 0$, we denote by $\rho : \mathbb{G}_X(\mathcal{F}, d) \rightarrow X$ the Grassmann bundle whose T -points, where $f : T \rightarrow X$ is any X -scheme, are equivalence classes of

coherent locally free quotients \mathcal{E} of $f^*\mathcal{F}$ of constant rank d , two such quotients being called equivalent if the corresponding surjections $q : f^*\mathcal{F} \rightarrow \mathcal{E}$ and $q' : f^*\mathcal{F} \rightarrow \mathcal{E}'$ have the same kernel, see e.g. [26]. We denote by $\rho^*\mathcal{F} \rightarrow \mathcal{Q}$ the universal coherent locally free quotient of constant rank d and refer the kernel \mathcal{S} of this surjection to as the universal subsheaf of $\rho^*\mathcal{F}$ so that we have the universal exact sequence $0 \rightarrow \mathcal{S} \rightarrow \rho^*\mathcal{F} \rightarrow \mathcal{Q} \rightarrow 0$. For a coherent locally free sheaf \mathcal{F} of constant rank $n \geq d + 1$, there is a canonical isomorphism $\mathbb{G}_X(\mathcal{F}, d) \rightarrow \mathbb{G}_X(\mathcal{F}^\vee, n - d)$ given on T -points by mapping a quotient $q : f^*\mathcal{F} \rightarrow \mathcal{E}$ with kernel \mathcal{K} to the quotient $f^*\mathcal{F}^\vee \rightarrow \mathcal{K}^\vee$. For $d = 1$, $\mathbb{G}_X(\mathcal{F}, d) \cong \mathbb{P}_X(\mathcal{F})$ and the universal exact sequence coincides under this isomorphism with the relative Euler exact sequence of $\mathbb{P}_X(\mathcal{F})$

$$0 \rightarrow \Omega_{\mathbb{P}_X(\mathcal{F})}^1(1) := \Omega_{\mathbb{P}_X(\mathcal{F})}^1 \otimes \mathcal{O}_{\mathbb{P}_X(\mathcal{F})}(1) \rightarrow \pi^*\mathcal{F} \rightarrow \mathcal{O}_{\mathbb{P}_X(\mathcal{F})}(1) \rightarrow 0.$$

1.2. Grassmannians. We summarize basic properties of Grassmannian varieties, see e.g. [28] and [43, Chapters 3 and 4] for the details. For a k -vector space V of dimension $n \geq 2$ and an integer $1 \leq d \leq n - 1$, the Grassmann bundle $\rho : \mathbb{G}_k(V^\vee, d) \rightarrow \text{Spec}(k)$ is the $d(n - d)$ -dimensional Grassmannian whose geometric points correspond to equivalence classes of d -dimensional quotients of V_k^\vee , equivalently to d -dimensional \bar{k} -vector subspaces E of $V_{\bar{k}}$.

1.2.1. Tautological sheaves. We put $V_{\mathbb{G}_k(V^\vee, d)}^\vee = V^\vee \otimes_k \mathcal{O}_{\mathbb{G}_k(V^\vee, d)}$ and we write

$$0 \rightarrow \mathcal{S} = \mathcal{S}_{\mathbb{G}_k(V^\vee, d)} \rightarrow V_{\mathbb{G}_k(V^\vee, d)}^\vee \rightarrow \mathcal{Q} = \mathcal{Q}_{\mathbb{G}_k(V^\vee, d)} \rightarrow 0$$

for the universal sequence of coherent locally free sheaves on $\mathbb{G}_k(V^\vee, d)$. The sheaf of Kähler differential $\Omega_{\mathbb{G}_k(V^\vee, d)/k}^1$ is canonically isomorphic to $\mathcal{H}om(\mathcal{Q}, \mathcal{S}) \cong \mathcal{S} \otimes \mathcal{Q}^\vee$ and its determinant $\omega_{\mathbb{G}_k(V^\vee, d)}$ is canonically isomorphic to $(\det \mathcal{S})^{\otimes n-d} \otimes (\det \mathcal{Q}^\vee)^{\otimes d} \cong (\det \mathcal{Q}^\vee)^{\otimes n}$. The k -vector spaces $H^0(\mathbb{G}_k(V^\vee, d), \mathcal{Q})$ and $H^0(\mathbb{G}_k(V^\vee, d), \mathcal{S}^\vee)$ are canonically isomorphic to V^\vee and V respectively, $H^1(\mathbb{G}_k(V^\vee, d), \Omega_{\mathbb{G}_k(V^\vee, d)/k}^1) \cong k$ and all other cohomology spaces of \mathcal{Q} , \mathcal{Q}^\vee , \mathcal{S} , \mathcal{S}^\vee and $\Omega_{\mathbb{G}_k(V^\vee, d)/k}^1$ are zero.

1.2.2. Plücker embedding and automorphisms. We denote by $j_P : \mathbb{G}_k(V^\vee, d) \rightarrow \mathbb{P}_k(\Lambda^d V^\vee)$ the *Plücker embedding*, that is, the closed immersion determined by the surjection $\Lambda^d V_{\mathbb{G}_k(V^\vee, d)}^\vee \rightarrow \det \mathcal{Q}$ induced by the universal quotient homomorphism. Letting $\text{Aut}_k(\mathbb{P}_k(\Lambda^d V^\vee), \mathbb{G}_k(V^\vee, d))$ be the stabilizer of $j_P(\mathbb{G}_k(V^\vee, d))$ in $\text{Aut}_k(\mathbb{P}_k(\Lambda^d V^\vee))$, it follows from [8] that the composition of the homomorphism of k -group schemes

$$(1.1) \quad \text{PGL}_k(V^\vee) = \text{Aut}_k(\mathbb{P}_k(V^\vee)) \rightarrow \text{Aut}_k(\mathbb{P}_k(\Lambda^d V^\vee), \mathbb{G}_k(V^\vee, d)), \varphi \mapsto \Lambda^d \varphi$$

with the restriction homomorphism $\text{Aut}_k(\mathbb{P}_k(\Lambda^d V^\vee), \mathbb{G}_k(V^\vee, d)) \rightarrow \text{Aut}_k(\mathbb{G}_k(V^\vee, d))$ is a closed immersion, which is an isomorphism when $n \neq 2d$ (otherwise, its image is a k -subgroup scheme of index 2).

1.3. Vector groups and vector group varieties.

1.3.1. Vector groups. A *vector k -group* is an abelian unipotent algebraic k -group scheme. By [11, IV.2.4], there is an equivalence between the category of finite dimensional k -vector spaces and the category of vector k -groups, given by the map associating to a finite dimensional k -vector space V the k -group scheme $(\mathbb{V}_k(V^\vee), +)$, where the comorphism of the k -group scheme structure $+$ is induced by the diagonal homomorphism $V^\vee \rightarrow V^\vee \oplus V^\vee$, and to a k -linear homomorphism $f : W \rightarrow V$ the k -group homomorphism $(\mathbb{V}_k(W^\vee), +) \rightarrow (\mathbb{V}_k(V^\vee), +)$ induced by ${}^t f : V^\vee \rightarrow W^\vee$. The choice of a k -basis of V determines an isomorphism of k -group schemes $(\mathbb{V}_k(V^\vee), +) \cong \mathbb{G}_{a,k}^n$. We will repeatedly use the following elementary results.

Lemma 3. *With the notation above, the following hold:*

- (1) *Every k -subgroup and quotient k -group of a vector k -group is a vector k -group.*
- (2) *Every extension $0 \rightarrow \mathbb{U}' \rightarrow \mathbb{U} \rightarrow \mathbb{U}'' \rightarrow 0$ of vector k -groups has a splitting $h : \mathbb{U} \xrightarrow{\cong} \mathbb{U}' \times \mathbb{U}''$.*
- (3) *The set $M^\mathbb{U}$ of \mathbb{U} -invariants of a rational \mathbb{U} -module M of finite positive dimension is nonzero.*

Recall [29, I.3] (see also [23, 4.2]) that a *quasi-coherent G -sheaf* of \mathcal{O}_X -modules on a k -scheme X with an action $\mu : G \times X \rightarrow X$ of an algebraic k -group G is a pair (\mathcal{F}, θ) consisting of a coherent sheaf of \mathcal{O}_X -modules \mathcal{F} and an isomorphism $\theta : \mu^*\mathcal{F} \xrightarrow{\cong} p_2^*\mathcal{F}$ of coherent sheaves of $\mathcal{O}_{G \times X}$ -modules, called a *G -linearization* of \mathcal{F} , that satisfies the cocycle relation $(m_G \times \text{id}_X)^*\theta = p_{23}^*\theta \circ (\text{id}_G \times \mu)^*\theta$ on $G \times G \times X$, where $m_G : G \times G \rightarrow G$ is the group law on G and where $p_{23} : G \times G \times X \rightarrow G \times X$ is the projection onto the last two factors. In particular, when G acts trivially on X , a G -linearization of \mathcal{F} is the same as a homomorphism of G into the group $\text{Aut}_X(\mathcal{F})$ of \mathcal{O}_X -module automorphisms of \mathcal{F} . Two G -linearizations θ and θ' of \mathcal{F} are called *equivalent* if there exists an \mathcal{O}_X -module automorphism φ of \mathcal{F} such that $p_2^*\varphi \circ \theta' = \theta \circ \mu^*\varphi$.

Lemma 4. *Let X be a normal k -variety endowed with an action of a vector k -group \mathbb{U} . Then every coherent invertible sheaf of \mathcal{O}_X -modules \mathcal{L} admits a \mathbb{U} -linearization $\theta_{\mathcal{L}}$ unique up to equivalence.*

Proof. This follows from [5, Lemma 2.13] and the fact that since $\mathbb{U} \cong \mathbb{A}_k^n$, for every normal k -variety X , the pullback homomorphisms $p_X^* : H^i(X, \mathcal{O}_X^*) \rightarrow H^i(X \times \mathbb{U}, \mathcal{O}_{X \times \mathbb{U}}^*)$, $i = 0, 1$, are isomorphisms. \square

1.3.2. Vector group structures and vector group varieties.

Definition 5. A *vector group variety* is a k -variety X endowed with an effective action $\mu : \mathbb{U} \times X \rightarrow X$ of a vector k -group \mathbb{U} which has a Zariski dense open orbit U_X . The action μ is said to define a *vector group structure* on X . Two vector group structures $\mu : \mathbb{U} \times X \rightarrow X$ and $\mu' : \mathbb{U}' \times X \rightarrow X$ on X are said to have the same equivalence class if there exists an isomorphism of k -groups $\alpha : \mathbb{U}' \rightarrow \mathbb{U}$ and a k -automorphism φ of X such that $\varphi \circ \mu' = \mu \circ (\alpha \times \varphi)$.

Lemma 6. *Let X be a vector group variety with open orbit U_X . Then U_X is a trivial \mathbb{U} -torsor. In particular, U_X contains a k -point of X .*

Proof. Since the action of \mathbb{U} is effective and U_X is Zariski dense, the morphism $(\mu, p_2) : \mathbb{U} \times_k U_X \rightarrow U_X \times U_X$ is an isomorphism, i.e., U_X endowed with the induced action of \mathbb{U} is a \mathbb{U} -torsor. The conclusion then follows from the additive form of Hilbert Theorem 90 which asserts that every such torsor is trivial. \square

Example 7. Since all orbits of unipotent group actions on a quasi-affine k -variety are closed [36, Theorem 2], Lemma 6 implies that a quasi-affine vector group variety X is a trivial \mathbb{U} -torsor. In particular, \mathbb{A}_k^n is the unique quasi-affine k -variety with a $\mathbb{G}_{a,k}^n$ -structure and this structure is unique up to isomorphism. On the other hand, by Sumihiro's equivariant completion [38, Theorem 3], every normal vector group k -variety X admits a \mathbb{U} -equivariant open immersion $j : X \hookrightarrow \bar{X}$ into a complete vector group k -variety \bar{X} . When X is smooth, the existence of equivariant resolution of singularities [27, Theorem 3.36 and Proposition 3.9.1] implies in addition that \bar{X} can be chosen to be smooth and such that $\bar{X} \setminus j(X)$ is the support of a \mathbb{U} -stable smooth normal crossing divisor on \bar{X} .

Proposition 8. *Let X be a k -variety endowed with a vector group structure $\mu : \mathbb{U} \times X \rightarrow X$, let $f : X \rightarrow Y$ be a proper morphism to a k -variety Y such that $f_* \mathcal{O}_X = \mathcal{O}_Y$ and let $i : F \hookrightarrow X$ be the scheme-theoretic fiber of f over a k -point y_0 of Y in the image by f of the open \mathbb{U} -orbit U_X . Then there exists an extension of vector k -groups $0 \rightarrow \mathbb{U}' \xrightarrow{a} \mathbb{U} \xrightarrow{b} \bar{\mathbb{U}} \rightarrow 0$ such that the following hold:*

- (1) *The variety Y is endowed with a vector group structure $\mu_Y : \bar{\mathbb{U}} \times_k Y \rightarrow Y$ with open $\bar{\mathbb{U}}$ -orbit U_Y and $f : X \rightarrow Y$ is equivariant with respect to the homomorphism $b : \mathbb{U} \rightarrow \bar{\mathbb{U}}$.*
- (2) *The scheme F is a k -variety endowed with a vector group structure $\mu_F : \mathbb{U}' \times_k F \rightarrow F$ and the closed immersion $i : F \hookrightarrow X$ is equivariant with respect to the homomorphism $a : \mathbb{U}' \rightarrow \mathbb{U}$.*
- (3) *Given any section $c : \bar{\mathbb{U}} \rightarrow \mathbb{U}$ of $b : \mathbb{U} \rightarrow \bar{\mathbb{U}}$, the morphism*

$$j : \mu \circ (c \times i) \circ (\mu_Y^{-1}(\cdot, y_0) \times \text{id}_F) : U_Y \times F \xrightarrow{\cong} \bar{\mathbb{U}} \times F \rightarrow \mathbb{U} \times X \rightarrow X$$

is a $\mathbb{U}' \times \bar{\mathbb{U}}$ -equivariant open immersion with image $f^{-1}(U_Y)$.

Proof. By Blanchard's lemma [6, Theorem 7.2.1], there exists a unique action $\nu : \mathbb{U} \times Y \rightarrow Y$ such that f is \mathbb{U} -equivariant. Let $\mathbb{U}' \subset \mathbb{U}$ be the stabilizer of y_0 and let $\bar{\mathbb{U}} = \mathbb{U}/\mathbb{U}'$. Since y_0 belongs to $f(U_X)$, the \mathbb{U} -orbit of y_0 is a constructible set which is not contained in any proper closed subset of Y . It thus contains a Zariski dense open subset of Y , hence, being homogeneous under the action of \mathbb{U} , is a Zariski dense open subset of Y . This implies that \mathbb{U}' acts trivially on Y and that the induced action $\mu_Y : \bar{\mathbb{U}} \times Y \rightarrow Y$ of $\bar{\mathbb{U}}$ is a vector group structure on Y with the property that $f : X \rightarrow Y$ is equivariant with respect to the k -group homomorphism $b : \mathbb{U} \rightarrow \bar{\mathbb{U}}$. This proves (1). The closed subscheme F is \mathbb{U}' -stable, with \mathbb{U}' -action $\mu_F : \mathbb{U}' \times F \rightarrow F$ induced by μ . Given a section $c : \bar{\mathbb{U}} \rightarrow \mathbb{U}$ of b , we have the following cartesian square of $\bar{\mathbb{U}}$ -equivariant morphisms

$$\begin{array}{ccc} \bar{\mathbb{U}} \times F & \xrightarrow{\mu \circ (c \times i)} & X \\ \text{id}_{\bar{\mathbb{U}}} \times f \downarrow & & \downarrow f \\ \bar{\mathbb{U}} \times \{y_0\} & \xrightarrow{\mu_Y(\cdot, y_0)} & Y \end{array}$$

where $\bar{\mathbb{U}}$ acts on $\bar{\mathbb{U}} \times F$ and $\bar{\mathbb{U}} \times \{y_0\}$ by translations on the first factor and on X by $\mu \circ (c \times \text{id}_X)$. For every $v \in \bar{\mathbb{U}}(k)$, $\mu \circ (c(v) \times \text{id}_X)$ is an automorphism of X which maps F isomorphically onto the scheme-theoretic fiber of f over the point $\mu_Y(v, y_0)$. Since $\mu_Y(\cdot, y_0) : \bar{\mathbb{U}} \times \{y_0\} \rightarrow Y$ is an open immersion with image U_Y , it follows that $\mu \circ (c \times i)$ is an open immersion with image $f^{-1}(U_Y)$. Furthermore, $\mu \circ (c \times i)$ is equivariant for the isomorphism $h = (c, a) : \bar{\mathbb{U}} \times \mathbb{U}' \rightarrow \mathbb{U}$ with respect to the product action of $\bar{\mathbb{U}} \times \mathbb{U}'$ on $\bar{\mathbb{U}} \times F$ and the \mathbb{U} -action on X . Assertion (3) follows. Finally, assertion (2) follows from the observation that the intersection of the inverse image of $U_X \subset f^{-1}(U_Y)$ by $\mu \circ (c \times i)$ with $\{0\} \times F$ is a Zariski dense \mathbb{U}' -stable open subset U_F of F which is a principal homogeneous space of \mathbb{U}' . \square

Recall that a coherent locally free sheaf \mathcal{E} on a k -variety X is called *simple* if its only endomorphisms are scalar homotheties. Lemma 4 and Proposition 8 imply the following result.

Corollary 9. *For a simple coherent locally free sheaf \mathcal{E} of rank $r \geq 2$ on a normal k -variety X , the total space of the projective bundle $\pi : \mathbb{P}_X(\mathcal{E}) \rightarrow X$ does not admit a vector group structure.*

Proof. Assume that $\mathbb{P}_X(\mathcal{E})$ admits a vector group structure given by the action of vector group \mathbb{U} . By Lemma 4, the invertible sheaf $\mathcal{O}_{\mathbb{P}_X(\mathcal{E})}(1)$ is canonically \mathbb{U} -linearized. By Proposition 8, $\pi : \mathbb{P}_X(\mathcal{E}) \rightarrow X$ is \mathbb{U} -equivariant for a uniquely determined \mathbb{U} -action on X , which factors through an effective action of a non-trivial quotient $\bar{\mathbb{U}} = \mathbb{U}/\mathbb{U}'$ defining a vector group structure on X . Since π is \mathbb{U} -equivariant, $\mathcal{E} \cong \pi_* \mathcal{O}_{\mathbb{P}_X(\mathcal{E})}(1)$ is endowed with an induced \mathbb{U} -linearization for the action on X , hence with a linearization for the trivial action of the positive dimensional vector group \mathbb{U}' . The latter is determined by some group homomorphism $\mathbb{U}' \rightarrow \text{Aut}_X(\mathcal{E})$ which is injective by Proposition 8. But this is impossible since $\text{Aut}_X(\mathcal{E}) \cong \mathbb{G}_{m,X}$ by hypothesis. \square

2. LINEAR SECTIONS OF $G(2, 5)$ AND THEIR RATIONAL LINEAR PROJECTIONS

We collect results concerning linear Schubert sub-varieties of dimension ≥ 2 of the Grassmannian $G(2, 5)$ of 2-dimensional k -vector subspaces of $k^{\oplus 5}$ and of its smooth linear sections in the Plücker embedding. We then review the description of the rational maps given by projections with respect to these linear Schubert sub-varieties. Over algebraically closed fields, all this material is classical, see e.g. [13, 19, 33, 41].

2.1. Linear Schubert subvarieties of $G(2, 5)$ and its hyperplane sections. We put $G = \mathbb{G}_k(V^\vee, 2) \cong G(2, 5)$ for some fixed 5-dimensional k -vector space V . We denote by $0 \rightarrow \mathcal{S} \rightarrow V_G^\vee \rightarrow \mathcal{Q} \rightarrow 0$ the universal sequence on G and by $j_P : G \hookrightarrow \mathbb{P}_k(\Lambda^2 V^\vee)$ the Plücker embedding. For any algebraic extension k' of k , we interpret k' -points of $G_{k'}$ either as 2-dimensional k' -vector subspaces $E \subset V_{k'}$ or as their corresponding lines $\mathbb{P}_{k'}(E^\vee)$ in $\mathbb{P}_{k'}(V_{k'}^\vee)$. For concrete examples, we fix the following coordinate convention:

Notation 10. For a chosen basis e_1, \dots, e_5 of V with dual basis $e_1^\vee, \dots, e_5^\vee$, we identify $\mathbb{P}_k(V^\vee)$ with \mathbb{P}_k^4 and $G(2, 5)$ with the closed subvariety of $\mathbb{P}_k(\Lambda^2 V^\vee) = \mathbb{P}_k^9$ endowed with the Plücker coordinates $w_{ij} = e_i^\vee \wedge e_j^\vee$, $1 \leq i < j \leq 5$, defined by the equations

$$\begin{cases} w_{12}w_{34} - w_{13}w_{24} + w_{14}w_{23} = 0 \\ w_{12}w_{35} - w_{13}w_{25} + w_{15}w_{23} = 0 \\ w_{12}w_{45} - w_{14}w_{25} + w_{15}w_{24} = 0 \\ w_{13}w_{45} - w_{14}w_{35} + w_{15}w_{34} = 0 \\ w_{23}w_{45} - w_{24}w_{35} + w_{25}w_{34} = 0 \end{cases}$$

2.1.1. Solids and planes in $G(2, 5)$. We consider the following linear Schubert sub-varieties of G :

Definition 11. Let $\{V_1 \subset V_3 \subset V_4\}$ be a partial flag of k -vector subspaces of V , with $\dim_k V_i = i$:

- The $\sigma_{3,0}$ -solid $\sigma_{3,0}(V_1) \cong \mathbb{P}_k((V/V_1)^\vee)$ associated to V_1 is the zero scheme of the homomorphism $V_{1,G} \hookrightarrow V_G \rightarrow \mathcal{S}^\vee$. Its intersection $\sigma_{3,1}(V_1 \subset V_4) \cong \mathbb{P}_k((V_4/V_1)^\vee)$ with the zero scheme of the homomorphism $(V/V_4)_G^\vee \hookrightarrow V_G^\vee \rightarrow \mathcal{Q}$ is called the $\sigma_{3,1}$ -plane associated to $\{V_1 \subset V_4\}$.
- The $\sigma_{2,2}$ -plane $\sigma_{2,2}(V_3) = \mathbb{G}_k(V_3^\vee, 2) \cong \mathbb{P}_k(V_3)$ associated to V_3 is the zero scheme of the homomorphism $(V/V_3)_G^\vee \hookrightarrow V_G^\vee \rightarrow \mathcal{Q}$.

The above sub-schemes are linear subspaces of G in the Plücker embedding $G \subset \mathbb{P}_k(\Lambda^2 V^\vee)$, given respectively as the intersections $G \cap \mathbb{P}_k(\Lambda^2 V^\vee / \Lambda^2(V/V_1)^\vee)$, $\sigma_{3,0}(V_1) \cap \mathbb{P}_k(\Lambda^2 V_4^\vee)$ and $G \cap \mathbb{P}_k(\Lambda^2 V_3^\vee)$. It follows from [41] that they are the only linear k -subspaces of dimension ≥ 2 contained in G .¹ Geometrically, the closed points of $\sigma_{3,0}(V_1)_{\bar{k}}$ and $\sigma_{3,1}(V_1 \subset V_4)_{\bar{k}}$ correspond respectively to lines in $\mathbb{P}_{\bar{k}}(V_{\bar{k}}^\vee)$ passing through the point $\mathbb{P}_{\bar{k}}(V_{1,\bar{k}}^\vee)$ and its subset of those which are contained in the subspace $\mathbb{P}_{\bar{k}}(V_{4,\bar{k}}^\vee)$. The closed points of $\sigma_{2,2}(V_3)_{\bar{k}}$ correspond to lines contained in the subspace $\mathbb{P}_{\bar{k}}(V_{3,\bar{k}}^\vee)$ of $\mathbb{P}_{\bar{k}}(V_{\bar{k}}^\vee)$.

Remark 12. The conormal sheaves in G of a solid $\Pi = \sigma_{3,0}(V_1)$ and of a plane $\Xi = \sigma_{2,2}(V_3)$ are canonically isomorphic to $\Omega_{\Pi}^1(1) \otimes V_{1,\Pi}$ and to $\Omega_{\Xi}^1(1) \otimes (V/V_3)_{\Xi}^\vee$, respectively. Indeed, since Π is the zero scheme of $V_{1,G} \rightarrow \mathcal{S}^\vee$, its conormal sheaf $\mathcal{C}_{\Pi/G}$ is isomorphic to $(\mathcal{S} \otimes V_{1,G})|_{\Pi} \cong \mathcal{S}|_{\Pi} \otimes V_{1,\Pi}$ and the canonical

¹The two types of planes in G are called planes of the first and second system in [41], and in [19], planes of vertex type and of non-vertex type, respectively.

isomorphism $\mathcal{S}|_{\Pi} \cong \Omega_{\Pi}^1(1)$ is given by the following commutative diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \rightarrow & \Omega_{\Pi}^1(1) & \rightarrow & (V/V_1)_{\Pi}^{\vee} & \rightarrow & \mathcal{O}_{\Pi}(1) \rightarrow 0 \\
 & & \parallel & & \downarrow & & \downarrow \\
 0 & \rightarrow & \mathcal{S}|_{\Pi} & \rightarrow & V_{\Pi}^{\vee} & \rightarrow & \mathcal{Q}|_{\Pi} \rightarrow 0 \\
 & & \searrow^0 & & \downarrow & & \downarrow \\
 & & & & V_{1,\Pi}^{\vee} & \xlongequal{\quad} & V_{1,\Pi}^{\vee} \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

Similarly, since Ξ is a codimension 4 local complete intersection equal to the zero-scheme of the homomorphism $(V/V_3)_G^{\vee} \rightarrow \mathcal{Q}$, the conormal sheaf $\mathcal{C}_{\Xi/G}$ is canonically isomorphic to $(\mathcal{Q}^{\vee} \otimes (V/V_3)_G^{\vee})|_{\Xi} \cong (\mathcal{Q}^{\vee}|_{\Xi}) \otimes (V/V_3)_{\Xi}^{\vee}$, and since $\mathcal{Q}|_{\Xi}$ equals the universal quotient sheaf on $\Xi \cong \mathbb{G}_k(V_3^{\vee}, 2)$, the canonical isomorphism $\mathbb{G}_k(V_3^{\vee}, 2) \cong \mathbb{P}_k(V_3)$ gives the identification $\mathcal{Q}^{\vee}|_{\Xi} \cong \Omega_{\Xi}^1(1)$.

Notation 13. Given a d -dimensional k -vector subspace $V_d \subset V$, $1 \leq d \leq 4$, we denote by Δ_{V_d} the stabilizer in $\mathrm{GL}_k(V^{\vee})$ of the subspace $(V/V_d)^{\vee} \subset V^{\vee}$. The choice of a splitting $V \cong V_d \oplus (V/V_d)$ identifies this k -group scheme with that consisting of block-matrices of the form

$$(2.1) \quad M(A_{5-d}, A_d, U) = \begin{pmatrix} A_{5-d} & U \\ 0 & A_d \end{pmatrix} \in \mathrm{GL}_k((V/V_d)^{\vee} \oplus V_d^{\vee})$$

with $A_{5-d} \in \mathrm{GL}_k((V/V_d)^{\vee})$, $A_d \in \mathrm{GL}_k(V_d^{\vee})$ and $U \in \mathrm{Hom}_k(V_d^{\vee}, (V/V_d)^{\vee})$. The associated subgroups $\mathrm{P}\Delta_{V_d}$ of $\mathrm{PGL}_k(V^{\vee})$ correspond under the isomorphism $\mathrm{PGL}_k(V^{\vee}) \cong \mathrm{Aut}_k(G)$ of (1.1) to the stabilizer subgroups of the solid $\sigma_{3,0}(V_1)$ if $d = 1$ and of the plane $\sigma_{2,2}(V_3)$ if $d = 3$.

2.1.2. Smooth hyperplane sections. By a hyperplane section of G , we mean the zero scheme $Z_{\langle s \rangle}$ of a non-zero global section $s \in H^0(G, \Lambda^2 \mathcal{Q}) = \Lambda^2 V^{\vee}$, equivalently the intersection of G in the Plücker embedding with the hyperplane $\mathbb{P}_k(\Lambda^2 V^{\vee} / \langle s \rangle)$ of $\mathbb{P}_k(\Lambda^2 V^{\vee})$. Denote by $\tilde{s} : V \rightarrow V^{\vee}$ the k -linear homomorphism corresponding to the form s under the canonical isomorphism $\mathrm{Hom}_k(V \otimes V, k) \cong \mathrm{Hom}_k(V, V^{\vee})$. The fivefold $Z_{\langle s \rangle}$ is the *isotropic Grassmannian* $\mathrm{I}_s \mathbb{G}_k(V^{\vee}, 2)$: a closed point $E \subset V_{\bar{k}}$ of $G_{\bar{k}}$ belongs to $(Z_{\langle s \rangle})_{\bar{k}}$ if and only if the homomorphism $\tilde{s}_{\bar{k}}|_E : E \rightarrow V_{\bar{k}} \rightarrow V_{\bar{k}}^{\vee}$ has image contained in $(V_{\bar{k}}/E)^{\vee}$, hence if and only if E is $s_{\bar{k}}$ -isotropic. Considering the conormal sequence

$$0 \rightarrow \mathcal{C}_{Z_{\langle s \rangle}/G} = \Lambda^2 \mathcal{Q}^{\vee}|_{Z_{\langle s \rangle}} \otimes \mathcal{O}_{Z_{\langle s \rangle}} \langle s \rangle_{Z_{\langle s \rangle}} \xrightarrow{d} \Omega_{G/k}^1|_{Z_{\langle s \rangle}} = \mathcal{H}om(\mathcal{Q}|_{Z_{\langle s \rangle}}, \mathcal{S}|_{Z_{\langle s \rangle}}) \rightarrow \Omega_{Z_{\langle s \rangle}/k}^1 \rightarrow 0$$

we see that $Z_{\langle s \rangle, \bar{k}}$ is smooth at $E \subset V_{\bar{k}}$ if and only if the map

$$d|_E : \Lambda^2 E \subset \mathrm{Hom}_k(E^{\vee} \otimes E^{\vee}, k) \cong \mathrm{Hom}_k(E^{\vee}, E) \rightarrow \mathrm{Hom}_{\bar{k}}(E^{\vee}, (V_{\bar{k}}/E)^{\vee}), \quad f \mapsto \tilde{s}_{\bar{k}}|_E \circ f$$

is nonzero, hence if and only if $E \not\subset \mathrm{Ker} \tilde{s}_{\bar{k}}$. In particular, $Z_{\langle s \rangle}$ is smooth if and only if \tilde{s} has rank 4. This yields a functorial correspondence between k -points $\langle s \rangle^{\vee}$ of $\mathbb{P}_k(\Lambda^2 V) \setminus \mathbb{G}_k(V, 2)$ and smooth hyperplane sections $Z_{\langle s \rangle}$ of G from which we get in particular that the action of $\mathrm{Aut}_k(G)(k) \cong \mathrm{PGL}_5(k)$ on the set of smooth hyperplane sections $Z_{\langle s \rangle}$ of G is transitive.

Given a smooth hyperplane section $Z_{\langle s \rangle}$, the s -orthogonal $V^{\perp} = \mathrm{Ker} \tilde{s}$ of V has dimension 1. We put $\bar{V} = V/V^{\perp}$ and denote by $\bar{s} \in \Lambda^2 \bar{V}^{\vee}$ the symplectic form on \bar{V} induced by s . Letting $\bar{G} = \mathbb{G}_k(\bar{V}^{\vee}, 2)$ with universal sequence $0 \rightarrow \mathcal{S}_{\bar{G}} \rightarrow \bar{V}_{\bar{G}}^{\vee} \rightarrow \mathcal{Q}_{\bar{G}} \rightarrow 0$, the zero scheme $Q_{\langle \bar{s} \rangle} \subset \bar{G}$ of \bar{s} is the *Lagrangian Grassmannian* $\mathrm{L}_{\bar{s}} \mathbb{G}_k(\bar{V}^{\vee}, 2)$ whose k -points are the maximal \bar{s} -isotropic subspaces of \bar{V} . The Plücker embedding $\bar{G} \hookrightarrow \mathbb{P}_k(\Lambda^2 \bar{V}^{\vee})$ induces a closed immersion of $Q_{\langle \bar{s} \rangle}$ in $\mathbb{P}_k(\Lambda^2 \bar{V}^{\vee} / \langle \bar{s} \rangle)$ as the zero scheme of the non-degenerate quadratic form \bar{q} associated to the symmetric bi-linear form $\bar{b} \in \mathrm{Sym}^2(\Lambda^2 \bar{V}^{\vee} / \langle \bar{s} \rangle)$ induced by the bi-linear form $b : \Lambda^2 \bar{V} \otimes \Lambda^2 \bar{V} \rightarrow k$, $\bar{u}_1 \wedge \bar{v}_1 \otimes \bar{u}_2 \wedge \bar{v}_2 \mapsto (\bar{s} \wedge \bar{s})(\bar{u}_1 \wedge \bar{v}_1 \wedge \bar{u}_2 \wedge \bar{v}_2)$.

Lemma 14. *With the notation above, the following hold:*

- (1) *Every smooth hyperplane section $Z_{\langle s \rangle}$ of G contains a unique solid $\Pi_{\langle s \rangle} = \sigma_{3,0}(V^{\perp})$ and conversely every solid of G is contained in a (non-unique) smooth hyperplane section of G .*
- (2) *A plane $\sigma_{2,2}(V_3)$ of G is contained in a smooth hyperplane section $Z_{\langle s \rangle}$ if and only if $V^{\perp} \subset V_3$ and V_3/V^{\perp} is \bar{s} -isotropic. In other words, the $\sigma_{2,2}$ -planes in $Z_{\langle s \rangle}$ are in one-to-one correspondence with the k -points of $Q_{\langle \bar{s} \rangle}$ and they intersect the unique solid $\Pi_{\langle s \rangle}$ of $Z_{\langle s \rangle}$ along lines.*

Proof. A solid $\sigma_{3,0}(V_1)$ of G is contained in $Z_{\langle s \rangle}$ if and only if $V = V_1^{\perp}$ hence, since by hypothesis V^{\perp} is 1-dimensional, if and only if $V_1 = V^{\perp}$. Conversely, for every 1-dimensional k -vector subspace V_1 of V ,

any choice of symplectic form $\bar{s} \in \Lambda^2(V/V_1)^\vee$ on the 4-dimensional k -space V/V_1 determines through the inclusion $\Lambda^2(V/V_1)^\vee \subset \Lambda^2V^\vee$ a skew-symmetric form $s \in \Lambda^2V^\vee$ with $\text{Ker } \bar{s} = V_1$, hence a corresponding smooth hyperplane section $Z_{(s)}$ of G containing $\sigma_{3,0}(V_1)$. A plane $\sigma_{2,2}(V_3)$ of G is contained in $Z_{(s)}$ if and only if every 2-dimensional k -vector subspace $E \subset V_3$ is s -isotropic, which holds if and only if V_3 is s -isotropic. This property is equivalent to the fact that V_3 contains V^\perp and $\bar{V}_3 = V_3/V^\perp$ is an \bar{s} -isotropic subspace of \bar{V} . Every $\sigma_{2,2}$ -plane with this property intersects $\Pi_{(s)} \cong \mathbb{P}_k(\bar{V}^\vee)$ along the line $\mathbb{P}_k(\bar{V}_3^\vee)$. \square

Remark 15. For a smooth hyperplane section $Z = Z_{(s)}$ of G , the exact sequence

$$0 \rightarrow \mathcal{C}_{Z/G}|_\Pi \cong \Lambda^2\mathcal{Q}^\vee|_\Pi \xrightarrow{\bar{s}} \mathcal{C}_{\Pi/G} \cong \Omega_\Pi^1(1) \rightarrow \mathcal{C}_{\Pi/Z} \rightarrow 0$$

identifies the conormal sheaf $\mathcal{C}_{\Pi/Z}$ of the unique solid $\Pi = \Pi_{(s)} \cong \mathbb{P}_k(\bar{V}^\vee)$ of $Z_{(s)}$ with the null-correlation rank 2 locally sheaf $\mathcal{N}_{(\bar{s})}$ associated to the symplectic form $\bar{s} \in \Lambda^2\bar{V}^\vee$, that is, the sheaf whose fiber over a closed point $\ell \subset \bar{V}_k$ of $\mathbb{P}_k(\bar{V}_k^\vee)$ is the quotient ℓ^\perp/ℓ , where ℓ^\perp is the \bar{s}_k -orthogonal of ℓ .

2.1.3. Smooth linear sections of codimension 2. A codimension 2 linear section of G is the zero scheme W_L of a homomorphism $L_G \rightarrow \Lambda^2\mathcal{Q}$ determined by a 2-dimensional k -vector subspace $L \subset H^0(G, \Lambda^2\mathcal{Q}) = \Lambda^2V^\vee$, equivalently the intersection of G in the Plücker embedding with the codimension 2 linear subspace $\mathbb{P}_k(\Lambda^2V^\vee/L)$ of $\mathbb{P}_k(\Lambda^2V^\vee)$. A closed point $E \subset V_k$ of G_k belongs to W_L if and only if $s_k|_E = 0$ for every skew-symmetric form s in L and, arguing as in § 2.1.2, we see from the conormal sequence

$$\mathcal{C}_{W_L/G} = \Lambda^2\mathcal{Q}^\vee|_{W_L} \otimes L_{W_L} \xrightarrow{d} \Omega_{G/k}^1|_{W_L} = \mathcal{H}om(\mathcal{Q}|_{W_L}, \mathcal{S}|_{W_L}) \rightarrow \Omega_{W_L/k}^1 \rightarrow 0$$

that $W_{L,\bar{k}}$ is smooth at a closed point $E \subset V_k$ if and only if $E \not\subset \text{Ker } \bar{s}_k$ for all $s \in L \setminus \{0\}$. In particular, W_L is smooth if and only if the line $\mathbb{P}_k(L^\vee) \subset \mathbb{P}_k(\Lambda^2V)$ is contained in $\mathbb{P}_k(\Lambda^2V) \setminus \mathbb{G}_k(V, 2)$. This yields a natural correspondence between smooth codimension 2 linear sections W_L of G and k -points of the open subset of $\mathbb{G}_k(\Lambda^2V, 2)$ parametrizing such lines, which is a homogeneous space under the action of $\text{PGL}_5(k)$.

Lemma 16. *A smooth linear section W_L of G does not contain any $\sigma_{3,0}$ -solid of G . It contains a unique $\sigma_{2,2}$ -plane $\Xi_L = \sigma_{2,2}(V_{3,L})$, where $V_{3,L} \subset V$ is the linear span of the kernels of the linear maps \tilde{s} , $s \in L \setminus \{0\}$.*

Proof. By adjunction formula, a solid Π contained in W_L would have normal sheaf isomorphic to $\mathcal{O}_\Pi(-1)$, and hence $\mathcal{O}_{W_L}(\Pi)$ would be an invertible sheaf of degree 1 on W_L . This is impossible since $\text{Pic}(W_L) \cong \mathbb{Z}$ is generated by an ample invertible sheaf of degree 5. A plane $\sigma_{2,2}(V_3)$ of G is contained in W_L if and only if V_3 contains the linear span of the kernels of the linear maps \tilde{s} corresponding to the forms $s \in L \setminus \{0\}$. The image of $\mathbb{P}_k(L^\vee) \subset \mathbb{P}_k(\Lambda^2V) \setminus \mathbb{G}_k(V, 2)$ by the morphism $\mathbb{P}_k(\Lambda^2V) \setminus \mathbb{G}_k(V, 2) \rightarrow \mathbb{G}_k(V, 4)$ given by the complete linear system of quadrics containing $\mathbb{G}_k(V, 2)$ is a smooth conic C_L in $\mathbb{G}_k(V, 4) \cong \mathbb{P}_k(V^\vee)$ whose k -points are the kernels $\text{Ker } \tilde{s} \subset V$ of the elements $s \in L \setminus \{0\}$. Letting $V_{3,L} \subset V$ be the unique 3-dimensional k -vector subspace such that $\mathbb{P}_k(V_{3,L}^\vee)$ contains C_L , we conclude that $\sigma_{2,2}(V_{3,L})$ is the unique $\sigma_{2,2}$ -plane in W_L . \square

2.2. Projections from $\sigma_{3,0}$ -solids. Let $V_1 \subset V$ be a 1-dimensional k -vector subspace, let $p : V \rightarrow \bar{V} = V/V_1$ be the quotient morphism and let $\bar{G} = \mathbb{G}_k(\bar{V}^\vee, 2)$ with universal sequence $0 \rightarrow \mathcal{S}_{\bar{G}} \rightarrow \bar{V}_G^\vee \rightarrow \mathcal{Q}_{\bar{G}} \rightarrow 0$. Let $Z_{(s)} \subset G$ be a smooth hyperplane section determined by the image $s \in \Lambda^2V^\vee$ of a symplectic form $\bar{s} \in \Lambda^2\bar{V}^\vee$ and let $Q_{(\bar{s})} \subset \bar{G}$ be the zero scheme of \bar{s} . Let $\Pi = \sigma_{3,0}(V_1) = G \cap \mathbb{P}_k(\Lambda^2V^\vee/\Lambda^2(\bar{V}^\vee))$ be the solid of G contained in $Z_{(s)}$ determined by V_1 . The *projection of G from the solid Π* is the dominant rational map

$$(2.2) \quad \pi_\Pi : G = \mathbb{G}_k(V^\vee, 2) \dashrightarrow \bar{G} = \mathbb{G}_k(\bar{V}^\vee, 2)$$

given by the restriction to G of the linear projection $\mathbb{P}_k(\Lambda^2V^\vee) \dashrightarrow \mathbb{P}_k(\Lambda^2\bar{V}^\vee)$. The morphism $\pi_\Pi|_{G \setminus \Pi}$ maps a k -point $E \subset V$ of G not containing V_1 to the k -point $p(E)$ of \bar{G} and conversely, the closure in G of a fiber of π_Π over a k -point $\bar{E} \subset \bar{V}$ of \bar{G} is the plane $\sigma_{2,2}(p^{-1}(\bar{E}))$ of G .

The restriction $\pi_\Pi : Z_{(s)} \dashrightarrow Q_{(\bar{s})}$ of π_Π to $Z_{(s)}$ is called the *projection of $Z_{(s)}$ from its solid $\Pi = \Pi_{(s)}$* . The restriction $\pi_\Pi|_{Z_{(s)} \setminus \Pi}$ maps a k -point of $Z_{(s)}$ represented by an s -isotropic k -point $E \subset V$ of G not containing V_1 to the k -point of $Q_{(\bar{s})}$ represented by the \bar{s} -isotropic k -point $p(E)$ of \bar{G} .

To state the next result, we put $(X_6, \mathbf{Q}_4, \mathcal{E}_6) = (G, \bar{G}, \mathcal{Q}_{\bar{G}}^\vee)$ and $(X_5, \mathbf{Q}_3, \mathcal{E}_5) = (Z_{(s)}, Q_{(\bar{s})}, \mathcal{S})$, where $\mathcal{S} = \mathcal{Q}_{\bar{G}}^\vee|_{Q_{(\bar{s})}}$ is the spinor locally free sheaf of rank 2 on the quadric threefold $Q_{(\bar{s})} \subset \bar{G}$, see e.g. [32].

Proposition 17. *For $i = 5, 6$, let $Y_\Pi \subset X_i \times \mathbf{Q}_{i-2}$ be the graph of π_Π with projections $p_{X_i} : Y_\Pi \rightarrow X_i$ and let $p_{\mathbf{Q}_{i-2}} : Y_\Pi \rightarrow \mathbf{Q}_{i-2}$. Then we have the following Sarkisov link*

$$(2.3) \quad \begin{array}{ccccc} \mathbb{P}_\Pi(\mathcal{C}_{\Pi/X_i}) \cong \mathbb{P}_{\mathbf{Q}_{i-2}}(\mathcal{E}_i) & \xrightarrow{\quad} & Y_\Pi & \xrightarrow{\cong} & \mathbb{P}_{\mathbf{Q}_{i-2}}(\mathcal{E}_i \oplus \mathcal{O}_{\mathbf{Q}_{i-2}}) \\ \downarrow & & \swarrow p_{X_i} & & \downarrow p_{\mathbf{Q}_{i-2}} \\ \Pi & \xrightarrow{\quad} & X_i & \xrightarrow{\pi_\Pi} & \mathbf{Q}_{i-2} \end{array}$$

where $p_{X_i} : Y_{\Pi} \rightarrow X_i$ is the contraction of the sub-bundle $\mathbb{P}_{\mathbf{Q}_{i-2}}(\mathcal{E}_i) \subset \mathbb{P}_{\mathbf{Q}_{i-2}}(\mathcal{E}_i \oplus \mathcal{O}_{\mathbf{Q}_{i-2}})$ onto Π .

Proof. The projection $p_{\bar{G}} : Y_{\Pi} \rightarrow \bar{G}$ is isomorphic to the projective bundle $\mathbb{G}_{\bar{G}}(\mathcal{F}, 2) \cong \mathbb{P}_{\bar{G}}(\mathcal{F}^{\vee}) \rightarrow \bar{G}$, where \mathcal{F} is the cokernel of the injective homomorphism $\mathcal{S}_{\bar{G}} \rightarrow \bar{V}_{\bar{G}}^{\vee} \rightarrow V_{\bar{G}}^{\vee}$. The latter is locally free of rank 3, isomorphic to an extension of $V_{1, \bar{G}}^{\vee}$ by $\mathcal{Q}_{\bar{G}}$, hence to $\mathcal{Q}_{\bar{G}} \oplus \mathcal{O}_{\bar{G}}$ due to the vanishing of $H^1(\bar{G}, \mathcal{Q}_{\bar{G}})$, and the projection $p_G : Y_{\Pi} \rightarrow G$ contracts the projective sub-bundle $\mathbb{P}_{\bar{G}}(\mathcal{Q}_{\bar{G}}^{\vee}) \subset \mathbb{P}_{\bar{G}}(\mathcal{Q}_{\bar{G}}^{\vee} \oplus \mathcal{O}_{\bar{G}})$ to Π . This identifies in particular $\mathbb{P}_{\bar{G}}(\mathcal{Q}_{\bar{G}}^{\vee})$ with the exceptional divisor $\mathbb{P}_{\Pi}(\mathcal{C}_{\Pi/\bar{G}})$ of the blow-up of Π in G . The corresponding diagram for the smooth hyperplane section $Z_{\langle s \rangle}$ follows immediately by restriction. \square

Example 18. With Notation 10, the kernel V^{\perp} of the linear map \bar{s} associated to the skew-symmetric form $s = e_1^{\vee} \wedge e_3^{\vee} - e_2^{\vee} \wedge e_4^{\vee}$ equals $\langle e_5 \rangle$. The associated hyperplane section $Z_{\langle s \rangle} = G \cap \{w_{13} - w_{24} = 0\}$ is the smooth fivefold in $\mathbb{P}_k^8 \subset \mathbb{P}_k^9$ with coordinates w_{ij} , $(i, j) \neq (2, 4)$, defined by the equations

$$\begin{cases} w_{12}w_{34} - w_{13}^2 + w_{14}w_{23} = 0 \\ w_{12}w_{35} - w_{13}w_{25} + w_{15}w_{23} = 0 \\ w_{12}w_{45} - w_{14}w_{25} + w_{13}w_{15} = 0 \\ w_{13}w_{45} - w_{14}w_{35} + w_{15}w_{34} = 0 \\ w_{23}w_{45} - w_{13}w_{35} + w_{25}w_{34} = 0 \end{cases}$$

Putting $\bar{V} = V/\langle e_5 \rangle$, the image of G by the projection

$$\mathbb{P}_k(\Lambda^2 V^{\vee}) = \mathbb{P}_k^9 \dashrightarrow \mathbb{P}_k^5 = \mathbb{P}_k(\Lambda^2 \bar{V}^{\vee}), [w_{ij}]_{1 \leq i < j \leq 5} \mapsto [w_{12} : w_{13} : w_{14} : w_{23} : w_{24} : w_{34}]$$

from the solid $\Pi = \sigma_{3,0}(V^{\perp}) = \{w_{12} = w_{13} = w_{14} = w_{23} = w_{24} = w_{34} = 0\}$ is the smooth quadric fourfold $\bar{G} = \mathbb{G}_k(\bar{V}^{\vee}, 2) = \{\bar{w}_{12}\bar{w}_{34} - \bar{w}_{13}\bar{w}_{24} + \bar{w}_{14}\bar{w}_{23} = 0\}$ in \mathbb{P}_k^5 with Plücker coordinates $\bar{w}_{ij} = \bar{e}_i^{\vee} \wedge \bar{e}_j^{\vee}$, $1 \leq i < j \leq 4$, where \bar{e}_i denotes the image of e_i in \bar{V} . The image of $Z_{\langle s \rangle}$ by this projection is the smooth quadric threefold $Q_{\langle \bar{s} \rangle} = \{\bar{w}_{12}\bar{w}_{34} - \bar{w}_{13}^2 + \bar{w}_{14}\bar{w}_{23} = 0\}$ in $\mathbb{P}_k^4 \subset \mathbb{P}_k^5$ with coordinates \bar{w}_{ij} , $(i, j) \neq (2, 4)$.

With the notation above, let $\mathrm{Sp}_k(\bar{V}^{\vee}, \bar{s})$ be the symplectic group of the symplectic form $\bar{s} \in \Lambda^2 \bar{V}^{\vee}$ and let $\mathrm{PSp}_k(\bar{V}^{\vee}, \bar{s})$ be its image in $\mathrm{PGL}_k(\bar{V}^{\vee})$. We infer the following description:

Corollary 19. *There exists a split exact sequence of k -group schemes*

$$0 \rightarrow \mathrm{Aut}_k(Z_{\langle s \rangle}, \Pi_{\langle s \rangle})_0 \cong \mathbb{G}_{a,k}^4 \rtimes \mathbb{G}_{m,k} \rightarrow \mathrm{Aut}_k(Z_{\langle s \rangle}, \Pi_{\langle s \rangle}) = \mathrm{Aut}_k(Z_{\langle s \rangle}) \rightarrow \mathrm{Aut}_k(Q_{\langle s \rangle}) \cong \mathrm{PSp}_k(\bar{V}^{\vee}, \bar{s}) \rightarrow 0,$$

where $\mathrm{Aut}_k(Z_{\langle s \rangle}, \Pi_{\langle s \rangle})_0$ is the kernel of the restriction homomorphism $\mathrm{Aut}_k(Z_{\langle s \rangle}, \Pi_{\langle s \rangle}) \rightarrow \mathrm{Aut}_k(\Pi_{\langle s \rangle})$. Moreover, up to the choice of a splitting $V \cong \bar{V} \oplus V^{\perp}$, $\mathrm{Aut}_k(Z_{\langle s \rangle})$ is the image under the restriction homomorphism $\mathrm{Aut}_k(G, Z_{\langle s \rangle}) \rightarrow \mathrm{Aut}_k(Z_{\langle s \rangle})$ of the subgroup

$$\left\{ \begin{pmatrix} \mathrm{Sp}_k(\bar{V}^{\vee}, \bar{s}) & \mathrm{Hom}_k((V^{\perp})^{\vee}, \bar{V}^{\vee}) \\ 0 & \mathbb{G}_{m,k} \end{pmatrix} \right\} / \{\pm \mathrm{Id}\}$$

of $\mathrm{PGL}_k(V^{\vee})$ under the isomorphism $\Phi : \mathrm{PGL}_k(V^{\vee}) \rightarrow \mathrm{Aut}_k(G)$ of (1.1).

Proof. Since, by Lemma 14, $\Pi = \Pi_{\langle s \rangle}$ is the unique solid contained in $Z = Z_{\langle s \rangle}$, we have $\mathrm{Aut}_k(Z) = \mathrm{Aut}_k(Z, \Pi)$. The action of $\mathrm{Aut}_k(Z, \Pi)$ lifts to the blow-up $p_Z : Y \rightarrow Z$ of Π and since the fibers of the projection $\pi_{\Pi} : Z \dashrightarrow Q_{\langle \bar{s} \rangle}$ of Z from Π over k -points of $Q_{\langle \bar{s} \rangle}$ meet Π along lines, $\mathrm{Aut}_k(Z, \Pi)_0$ equals the kernel of the homomorphism $\mathrm{Aut}_k(Z, \Pi) \rightarrow \mathrm{Aut}_k(Q_{\langle \bar{s} \rangle})$ induced by $p_{Q_{\langle \bar{s} \rangle}} : Y \rightarrow Q_{\langle \bar{s} \rangle}$. Let $p : \mathrm{Aut}_k(Z, \Pi) \rightarrow B := \mathrm{Aut}_k(Z, \Pi) / \mathrm{Aut}_k(Z, \Pi)_0$ be the quotient morphism and let $\gamma : B \rightarrow \mathrm{Aut}_k(Q_{\langle \bar{s} \rangle})$ be the induced injective homomorphism. Let $\Delta_{V^{\perp}} \subset \mathrm{GL}_k(V^{\vee})$ be the stabilizer of $\bar{V}^{\vee} \subset V^{\vee}$, see Notation 13. Let $\Delta_{V^{\perp}, 0} \cong \mathbb{G}_{a,k}^4 \rtimes \mathbb{G}_{m,k}$ be its normal subgroup consisting of matrices $M(\mathrm{id}_4, \lambda, U)$ and let $S_{V^{\perp}}$ be its subgroup consisting of matrices $M(A_4, \pm 1, 0)$ with $A_4 \in \mathrm{Sp}_k(\bar{V}^{\vee}, \bar{s})$. The image of $\mathrm{P}\Delta_{V^{\perp}} \subset \mathrm{PGL}_k(V^{\vee})$ by $\Phi : \mathrm{PGL}_k(V^{\vee}) \rightarrow \mathrm{Aut}_k(G)$ is contained in the stabilizer $\mathrm{Aut}_k(G, (Z, \Pi))$ of the pair (Z, Π) . A direct verification shows that the homomorphism $j : \mathrm{P}\Delta_{V^{\perp}} \rightarrow \mathrm{Aut}_k(Z)$ obtained by composing with the restriction homomorphism $\mathrm{Aut}_k(G, Z) \rightarrow \mathrm{Aut}_k(Z)$ is injective and maps $\mathrm{P}\Delta_{V^{\perp}, 0} \cong \Delta_{V^{\perp}, 0}$ isomorphically onto $\mathrm{Aut}_k(Z, \Pi)_0$. Letting $\bar{q} \in \mathrm{Sym}^2(\Lambda^2 \bar{V}^{\vee} / \langle \bar{s} \rangle)$ be the quadratic form associated to the symplectic form \bar{s} (see § 2.1.2), the conclusion then follows from the fact that the restriction of the composition

$$\mathrm{P}\Delta_{V^{\perp}} \xrightarrow{j} \mathrm{Aut}_k(Z, \Pi) \xrightarrow{p} B \xrightarrow{\gamma} \mathrm{Aut}_k(Q_{\langle \bar{s} \rangle}) = \mathrm{PO}_k(\Lambda^2 \bar{V}^{\vee} / \langle \bar{s} \rangle, \bar{q}) = \mathrm{SO}_k(\Lambda^2 \bar{V}^{\vee} / \langle \bar{s} \rangle, \bar{q})$$

to the subgroup $\mathrm{PS}_{V^{\perp}} \cong \mathrm{PSp}_k(\bar{V}^{\vee}, \bar{s})$ is an isomorphism onto its image $\mathrm{SO}_k(\Lambda^2 \bar{V}^{\vee} / \langle \bar{s} \rangle, \bar{q})$. \square

2.3. Projections from $\sigma_{2,2}$ -planes. Let $V_3 \subset V$ be a 3-dimensional k -vector subspace, let $K = \Lambda^2 V / \Lambda^2 V_3$ and let $\langle s \rangle \subset K^\vee$ and $L \subset K^\vee$ be respectively a 1-dimensional and a 2-dimensional linear subspace of skew-symmetric bilinear forms on V whose non-zero elements all have maximal rank. These data determine respectively a plane $\Xi = \sigma_{2,2}(V_3) = G \cap \mathbb{P}_k(\Lambda^2 V_3^\vee)$ of G , a smooth hyperplane section $Z_{\langle s \rangle}$ of G and a smooth codimension 2 linear section W_L of G which both contain Ξ .

- The projection of G from the plane Ξ is the birational map

$$(2.4) \quad \pi_\Xi : G = \mathbb{G}_k(V^\vee, 2) \dashrightarrow \mathbb{P}_k(K^\vee)$$

induced by the restriction to G of the linear projection $\mathbb{P}_k(\Lambda^2 V^\vee) \dashrightarrow \mathbb{P}_k(K^\vee)$. The morphism $\pi_\Xi|_{G \setminus \Xi}$ maps a k -point $E \subset V$ of G not contained in V_3 to the image of $\Lambda^2 E \subset \Lambda^2 V$ by the quotient homomorphism $\Lambda^2 V \rightarrow K$. Let $Z_\Xi = G \cap \mathbb{P}_k(\Lambda^2 V^\vee / \Lambda^2(V/V_3)^\vee)$ and $H_G = \mathbb{P}_k(K^\vee / \Lambda^2(V/V_3)^\vee)$ be the hyperplane sections of G and $\mathbb{P}_k(K^\vee)$ determined by the 1-dimensional k -vector subspace $\Lambda^2(V/V_3)^\vee$ of $K^\vee \subset \Lambda^2 V^\vee$. Then the image S_G of the rational map $\pi_\Xi|_{Z_\Xi} : Z_\Xi \dashrightarrow H_G$ equals that of the Segre embedding

$$s_{1,1} : \mathbb{P}_k(V_3^\vee) \times \mathbb{P}_k((V/V_3)^\vee) \hookrightarrow \mathbb{P}_k(V_3^\vee \otimes_k (V/V_3)^\vee) \cong \mathbb{P}_k(K^\vee / \Lambda^2(V/V_3)^\vee).$$

A k -point $E \subset V$ of G belongs to Z_Ξ if and only if $E \cap V_3 \neq \{0\}$, and the closure in G of the fiber of $\pi_\Xi|_{Z_\Xi \setminus \Xi}$ over the image by $s_{1,1}$ of a k -point $(V_1 \subset V_3, V_4/V_3 \subset V/V_3)$ of $\mathbb{P}_k(V_3^\vee) \times \mathbb{P}_k((V/V_3)^\vee)$ is the $\sigma_{3,1}$ -plane $\sigma_{3,1}(V_1 \subset V_4)$.

- Since $\langle s \rangle \subset K^\vee$, the projection of G from Ξ restricts on $Z_{\langle s \rangle}$ to the birational map

$$(2.5) \quad \pi_\Xi : Z_{\langle s \rangle} \dashrightarrow \mathbb{P}_k(K^\vee / \langle s \rangle)$$

defined by the complete linear system of hyperplane sections of $Z_{\langle s \rangle}$ containing Ξ , called the *projection of $Z_{\langle s \rangle}$ from the plane Ξ* . Letting $H_{Z_{\langle s \rangle}} = H_G \cap \mathbb{P}_k(K^\vee / \langle s \rangle)$, the image of the induced rational map $\pi_\Xi|_{Z_\Xi \cap Z_{\langle s \rangle}} : Z_\Xi \cap Z_{\langle s \rangle} \dashrightarrow H_{Z_{\langle s \rangle}}$ is the smooth cubic scroll $S_{Z_{\langle s \rangle}} = S_G \cap \mathbb{P}_k(K^\vee / \langle s \rangle)$ in $\mathbb{P}_k(K^\vee / \langle s \rangle)$.

- Since $L \subset K^\vee$, the projection of G from Ξ restricts on W_L to the birational map

$$(2.6) \quad \pi_\Xi : W_L \dashrightarrow \mathbb{P}_k(K^\vee / L)$$

defined by the complete linear system of hyperplane sections of W_L containing Ξ , called the *projection of W_L from the plane Ξ* . Putting $H_{W_L} = H_G \cap \mathbb{P}_k(K^\vee / L)$, the image of the induced rational map $\pi_\Xi|_{Z_\Xi \cap W_L} : Z_\Xi \cap W_L \dashrightarrow H_{W_L}$ is the smooth rational cubic curve $S_{W_L} = S_G \cap \mathbb{P}_k(K^\vee / L)$ in $\mathbb{P}_k(K^\vee / L)$.

To state the next result, we put $(X_6, \mathbf{P}_6) = (G, \mathbb{P}_k(K^\vee))$, $(X_5, \mathbf{P}_5) = (Z_{\langle s \rangle}, \mathbb{P}_k(K^\vee / \langle s \rangle))$ and $(X_4, \mathbf{P}_4) = (W_L, \mathbb{P}_k(K^\vee / L))$. We denote by $Y_{\Xi, i} \subset X_i \times \mathbf{P}_i$, the graph of $\pi_{\Xi, i} = \pi_\Xi : X_i \dashrightarrow \mathbf{P}_i$, $i = 4, 5, 6$.

Proposition 20. *With the notation above, for $i = 4, 5, 6$ we have the following Sarkisov link*

$$(2.7) \quad \begin{array}{ccccc} \text{Bl}_{S_{X_i}} H_{X_i} \cong \mathbb{P}_\Xi(\mathcal{C}_{\Xi/X_i}) & \xrightarrow{\quad} & Y_{\Xi, i} & \xleftarrow{\quad} & \text{Bl}_\Xi(Z_\Xi \cap X_i) \\ \downarrow & & \swarrow \text{p}_{X_i} & & \searrow \text{p}_2 \\ \Xi & \hookrightarrow & Z_\Xi \cap X_i & \xrightarrow{\quad} & X_i & \xrightarrow{\quad} & \mathbf{P}_i & \xleftarrow{\quad} & H_{X_i} & \xleftarrow{\quad} & S_{X_i} \\ & & & & \swarrow \Phi_i & & & & & & \downarrow \end{array}$$

where $\text{p}_{X_i} : Y_{\Xi, i} \rightarrow X_i$ is the blow-up of Ξ , $\text{Bl}_\Xi(Z_\Xi \cap X_i)$ is the proper transform of $Z_\Xi \cap X_i$, $\text{p}_2 : Y_{\Xi, i} \rightarrow \mathbf{P}_i$ is the blow-up of S_{X_i} and $\text{Bl}_{S_{X_i}} H_{X_i}$ is the proper transform of $H_{X_i} \subset \mathbf{P}_i$. The birational inverse Φ_i of $\pi_{\Xi, i}$ is given by the complete linear system of quadrics in \mathbf{P}_i containing S_{X_i} .

Proof. All the properties follow from [41] and the description above. \square

Example 21. With Notation 10, let $V_3 = \langle e_3, e_4, e_5 \rangle$, let

$$\Xi = \sigma_{2,2}(V_3) = \{w_{12} = w_{13} = w_{14} = w_{15} = w_{23} = w_{24} = w_{25} = 0\}$$

be the associated plane of G and let

$$\mathbb{P}_k(\Lambda^2 V^\vee) = \mathbb{P}_k^9 \dashrightarrow \mathbb{P}_k^6 = \mathbb{P}_k(K^\vee), [w_{ij}]_{1 \leq i < j \leq 5} \mapsto [w_{12} : w_{13} : w_{14} : w_{15} : w_{23} : w_{24} : w_{25}]$$

be the associated linear projection. The skew-symmetric forms $s = e_1^\vee \wedge e_3^\vee - e_2^\vee \wedge e_4^\vee$ and $s' = e_1^\vee \wedge e_4^\vee - e_2^\vee \wedge e_5^\vee$ generate a subspace $L \subset \Lambda^2 V^\vee$ whose non-zero elements all have rank 4. The associated smooth linear section $W_L = G \cap \{w_{13} - w_{24} = 0\} \cap \{w_{14} - w_{25} = 0\}$ is the smooth fourfold in $\mathbb{P}_k^7 \subset \mathbb{P}_k^9$ with coordinates w_{ij} , $(i, j) \neq (2, 4), (2, 5)$ defined by the equations

$$\begin{cases} w_{12}w_{34} - w_{13}^2 + w_{14}w_{23} = 0 \\ w_{12}w_{35} - w_{13}w_{14} + w_{15}w_{23} = 0 \\ w_{12}w_{45} - w_{14}^2 + w_{13}w_{15} = 0 \\ w_{13}w_{45} - w_{14}w_{35} + w_{15}w_{34} = 0 \\ w_{23}w_{45} - w_{13}w_{35} + w_{14}w_{34} = 0 \end{cases}$$

The smooth hyperplane section $Z_{\langle s \rangle} = G \cap \{w_{13} - w_{24} = 0\}$ of Example 18 and the above smooth linear section W_L both contain Ξ . The images of $Z_\Xi = G \cap \{w_{12} = 0\}$, $Z_\Xi \cap Z_{\langle s \rangle}$ and $Z_\Xi \cap W_L$ by the projection π_Ξ and its successive restrictions are the smooth threefold $S_G \cong \mathbb{P}_k^1 \times \mathbb{P}_k^2$, the smooth rational cubic surface $S_{Z_{\langle s \rangle}} \cong \mathbb{P}_{\mathbb{P}_k^1}(\mathcal{O}_{\mathbb{P}_k^1}(1) \oplus \mathcal{O}_{\mathbb{P}_k^1}(2)) \cong \mathbb{F}_1$ and the smooth rational normal curve $S_{W_L} \cong \mathbb{P}_k^1$ with equations

$$\begin{cases} -w_{13}w_{24} + w_{14}w_{23} = 0 \\ -w_{13}w_{25} + w_{15}w_{23} = 0 \\ -w_{14}w_{25} + w_{15}w_{24} = 0 \end{cases} \quad \begin{cases} -w_{13}^2 + w_{14}w_{23} = 0 \\ -w_{13}w_{25} + w_{15}w_{23} = 0 \\ -w_{14}w_{25} + w_{15}w_{13} = 0 \end{cases} \quad \text{and} \quad \begin{cases} -w_{13}^2 + w_{14}w_{23} = 0 \\ -w_{13}w_{14} + w_{15}w_{23} = 0 \\ -w_{14}^2 + w_{15}w_{13} = 0 \end{cases}$$

in \mathbb{P}_k^5 , \mathbb{P}_k^4 and \mathbb{P}_k^3 respectively.

Remark 22. By a result attributed to Weil [42], the smooth varieties $\mathbb{P}_k^1 \times \mathbb{P}_k^2 \subset \mathbb{P}_k^5$, $\mathbb{F}_1 \subset \mathbb{P}_k^4$ and the rational normal curve $\mathbb{P}_k^1 \subset \mathbb{P}_k^3$ are the only smooth cubics which are not hypersurfaces.

For a smooth codimension 2 linear section W_L of G with unique $\sigma_{2,2}$ -plane $\Xi_L = \sigma_{2,2}(V_{3,L})$, we infer the following description of $\text{Aut}_k(W_L)$:

Corollary 23. *There exists a split exact sequence of k -group schemes*

$$0 \rightarrow \text{Aut}_k(W_L, \Xi_L)_0 \cong \mathbb{G}_{a,k}^4 \rtimes \mathbb{G}_{m,k} \rightarrow \text{Aut}_k(W_L, \Xi_L) = \text{Aut}_k(W_L) \rightarrow \text{Aut}_k(S_{W_L}) \cong \text{PGL}_k(L) \rightarrow 0,$$

where $\text{Aut}_k(W_L, \Xi_L)_0$ is the kernel of the restriction homomorphism $\text{Aut}_k(W_L, \Xi_L) \rightarrow \text{Aut}_k(\Xi_L)$.

Proof. Since, by Lemma 16, $\Xi = \Xi_L$ is the unique $\sigma_{2,2}$ -plane contained in W_L and since the intersection $Z_\Xi \cap W_L$ is the union of all the $\sigma_{3,1}$ -planes contained in W_L , we have $\text{Aut}_k(W_L) = \text{Aut}_k(W_L, (Z_\Xi, \Xi))$. Since the $\sigma_{3,1}$ -planes of W_L intersect $\Xi \cong \mathbb{P}_k(V_{3,L})$ along the smooth conic C_{L^\vee} dual to $C_L \cong \mathbb{P}_k(L^\vee) \hookrightarrow \mathbb{P}_k(V_{3,L}^\vee)$ (see the proof of Lemma 16), the image of the restriction homomorphism $\text{Aut}_k(W_L, \Xi_L) \rightarrow \text{Aut}_k(\Xi_L)$ is contained in the subgroup $\text{Aut}(\Xi_L, C_{L^\vee}) \cong \text{Aut}_k(C_{L^\vee}) \cong \text{PGL}_k(L)$. Since on the other hand the $\sigma_{3,1}$ -planes of W_L are the closures of the fibers of $\pi_\Xi : W_L \setminus \Xi \rightarrow \mathbb{P}_k(K^\vee/L)$ over the k -points of S_{W_L} , the projection $\pi_\Xi : W_L \dashrightarrow \mathbb{P}_k(K^\vee/L)$ induces an isomorphism of k -group schemes $\text{Aut}_k(W_L, (Z_\Xi, \Xi)) \cong \text{Aut}_k(\mathbb{P}_k(K^\vee/L), (H_{W_L}, S_{W_L}))$ which maps $\text{Aut}_k(W_L, \Xi_L)_0$ isomorphically onto the kernel $\mathbb{G}_{a,k}^4 \rtimes \mathbb{G}_{m,k}$ of the restriction homomorphism $\text{Aut}_k(\mathbb{P}_k(K^\vee/L), (H_{W_L}, S_{W_L})) \rightarrow \text{Aut}_k(S_{W_L})$. The latter homomorphism is a split surjection, which identifies $\text{Aut}_k(S_{W_L})$ with $\text{Aut}_k(H_{W_L}, S_{W_L})$. \square

3. SMOOTH QUINTIC DEL PEZZO VARIETIES WITH VECTOR GROUP STRUCTURES

Recall that a smooth quintic del Pezzo k -variety of dimension $n \in \{2, \dots, 6\}$ is a k -form X of a smooth section of the Grassmannian $G(2, 5) \subset \mathbb{P}_k^9$ by a linear subspace of dimension $6 - n$. For $n \leq 3$, the automorphism group of $X_{\bar{k}}$ is too small to allow the existence of a vector subgroup structure on $X_{\bar{k}}$. In this section, we consider the case of smooth quintic del Pezzo k -varieties of dimension 4, 5 and 6.

3.1. Toric vector groups structures on linear sections of $G(2, 5)$. Let V be a k -vector space of dimension 5 and, with the notation introduced in § 2.3, let $X_6 = \mathbb{G}_k(V^\vee, 2)$, let $V_3 \subset V$ be a 3-dimensional k -vector subspace with associated plane $\Xi = \sigma_{2,2}(V_3)$ of X_6 and let $K_6 = \Lambda^2 V / \Lambda^2 V_3$. Let $\langle s \rangle \subset K_6^\vee$ and $L \subset K_6^\vee$ be respectively a 1-dimensional and a 2-dimensional linear subspace of skew-symmetric bilinear forms on V whose non-zero elements all have maximal rank. Put $K_5^\vee = K_6^\vee / \langle s \rangle$, $K_4^\vee = K_6^\vee / L$ and let $X_5 = Z_{\langle s \rangle}$ and $X_4 = W_L$ be the smooth linear sections of X_6 containing Ξ defined by $\langle s \rangle$ and L respectively. Let $F_i = \text{Hom}_k(K_i^\vee / \Lambda^2(V/V_3)^\vee, \Lambda^2(V/V_3)^\vee) \cong k^{\oplus i}$, $i = 4, 5, 6$, and let $\mathbb{V}_k(F_i^\vee)$ be the associated vector group. We derive from the exact sequence

$$0 \rightarrow \Lambda^2(V/V_3)^\vee \xrightarrow{a} K_i^\vee \xrightarrow{b} K_i^\vee / \Lambda^2(V/V_3)^\vee \rightarrow 0, \quad i = 4, 5, 6$$

a faithful homomorphism of k -group schemes $\mathbb{V}_k(F_i^\vee) \rightarrow \text{GL}_k(K_i^\vee)$, $f \mapsto \text{id}_{K_i^\vee} + a \circ f \circ b$ corresponding to a $\mathbb{V}_k(F_i^\vee)$ -action on $\mathbb{P}_k(K_i^\vee)$ restricting to the trivial action on the invariant hyperplane $H_{X_i} = \mathbb{P}_k(K_i^\vee / \Lambda^2(V/V_3)^\vee)$ and having $\mathbb{P}_k(K_i^\vee) \setminus H_{X_i}$ as an open orbit. By Proposition 20, this action lifts through the birational projection $\pi_\Xi : X_i \dashrightarrow \mathbb{P}_k(K_i^\vee)$ from the plane Ξ to a $\mathbb{V}_k(F_i^\vee)$ -action on X_i with open orbit $X_i \setminus (Z_\Xi \cap X_i)$ and whose restriction to Ξ is trivial. As a consequence, we obtain the following:

Proposition 24. *Every smooth section of $G(2, 5) \subset \mathbb{P}_k^9$ by a linear subspace of codimension ≤ 3 admits a vector group structure.*

Example 25. With the notation of Example 21, let $V_3 = \langle e_3, e_4, e_5 \rangle$ with associated linear projection

$$\mathbb{P}_k^9 \dashrightarrow \mathbb{P}_k^6, [w_{ij}]_{1 \leq i < j \leq 5} \mapsto [w_{12} : w_{13} : w_{14} : w_{15} : w_{23} : w_{24} : w_{25}],$$

let $s = e_1^\vee \wedge e_3^\vee - e_2^\vee \wedge e_4^\vee \in K_6^\vee$ and $L \subset K_6^\vee$ be the subspace generated by s and $s' = e_1^\vee \wedge e_4^\vee - e_2^\vee \wedge e_5^\vee$.

• For the basis $t_{ij} = (e_i^\vee \wedge e_j^\vee) \otimes (\bar{e}_1 \wedge \bar{e}_2)$, $i = 1, 2$, $j = 3, 4, 5$ of F_6^\vee , the corresponding action of $\mathbb{V}_k(F_6^\vee) \cong \text{Spec}(k[t_{ij}])$ on $\mathbb{P}_k(K_6^\vee) = \mathbb{P}_k^6$ with open orbit $\mathbb{P}_k^6 \setminus \{w_{12} = 0\}$ is the “toric” \mathbb{G}_a^6 -structure on \mathbb{P}_k^6 given by $w_{12} \mapsto w_{12}$ and $w_{ij} \mapsto w_{ij} + t_{ij}w_{12}$ for $(i, j) \neq (1, 2)$.² Its lift to $X_6 = G$ is the restriction of the $\mathbb{V}_k(F_6^\vee)$ -action on $\mathbb{P}_k(\Lambda^2 V^\vee)$ induced by the second exterior power of the representation

$$(3.1) \quad \rho_6 : \mathbb{V}_k(F_6^\vee) \rightarrow \text{GL}_k(V^\vee), \quad (t_{13}, t_{23}, t_{14}, t_{24}, t_{15}, t_{25}) \mapsto \begin{pmatrix} 1 & 0 & -t_{23} & -t_{24} & -t_{25} \\ 0 & 1 & t_{13} & t_{14} & t_{15} \\ \vdots & \ddots & 1 & 0 & 0 \\ \vdots & & \ddots & 1 & 0 \\ 0 & \dots & \dots & 0 & 1 \end{pmatrix}.$$

Comparing with the classification in [39, Chapter 3, § 3, IV], the image of ρ_6 is one of the two maximal abelian unipotent subgroups of GL_5 corresponding to cases N_2 and N_3 . The other one, given by the representation dual to ρ_6 , corresponds to a vector group structure on $\mathbb{G}_k(V, 2) \cong \mathbb{G}_k(V^\vee, 3)$.

• The corresponding action of $\mathbb{V}_k(F_5^\vee) \cong \text{Spec}(k[t_{13}, t_{23}, t_{14}, t_{15}, t_{25}])$ on $\mathbb{P}_k(K_5^\vee) = \mathbb{P}_k^5$ with open orbit $\mathbb{P}_k^5 \setminus \{w_{12} = 0\}$ is the toric \mathbb{G}_a^5 -structure on \mathbb{P}_k^5 given by $w_{12} \mapsto w_{12}$ and $w_{ij} \mapsto w_{ij} + t_{ij}w_{12}$ for $(i, j) \neq (1, 2), (2, 4)$. Its lift to $X_5 = Z_{(s)}$ is the restriction of the $\mathbb{V}_k(F_5^\vee)$ -action on $\mathbb{P}_k(\Lambda^2 V^\vee)$ preserving $Z_{(s)} = G \cap \{w_{13} - w_{24} = 0\}$ induced by the second exterior power of the representation

$$(3.2) \quad \rho_5 : \mathbb{V}_k(F_5^\vee) \rightarrow \text{GL}_k(V^\vee), \quad (t_{13}, t_{23}, t_{14}, t_{15}, t_{25}) \mapsto \begin{pmatrix} 1 & 0 & -t_{23} & -t_{13} & -t_{25} \\ 0 & 1 & t_{13} & t_{14} & t_{15} \\ \vdots & \ddots & 1 & 0 & 0 \\ \vdots & & \ddots & 1 & 0 \\ 0 & \dots & \dots & 0 & 1 \end{pmatrix}.$$

The stabilizer $\text{Stab}(\langle e_5^\vee \rangle)$ of the subspace $\langle e_5^\vee \rangle$ is the subgroup $\text{Spec}(k[t_{15}, t_{25}])$ of $\mathbb{V}_k(F_5^\vee)$ and the induced action of the vector group $\mathbb{V}_k(F_5^\vee)/\text{Stab}(\langle e_5^\vee \rangle) \cong \text{Spec}(k[\bar{t}_{13}, \bar{t}_{23}, \bar{t}_{14}])$ on $\bar{V}^\vee = (V/\langle e_5^\vee \rangle)^\vee$ endowed with the basis dual to that determined by the images \bar{e}_i of the e_i , $i = 1, \dots, 4$, is given by the representation

$$(3.3) \quad \bar{\rho}_5 : \mathbb{V}_k(F_5^\vee)/\text{Stab}(\langle e_5^\vee \rangle) \rightarrow \text{Sp}_4(\bar{V}^\vee, \bar{s}), \quad (\bar{t}_{13}, \bar{t}_{23}, \bar{t}_{14}) \mapsto \begin{pmatrix} 1 & 0 & -\bar{t}_{23} & -\bar{t}_{13} \\ 0 & 1 & \bar{t}_{13} & \bar{t}_{14} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

where $\bar{s} = \bar{e}_1^\vee \wedge \bar{e}_3^\vee - \bar{e}_2^\vee \wedge \bar{e}_4^\vee$ is the symplectic form on \bar{V} induced by s .

• The corresponding action of $\mathbb{V}_k(F_4^\vee) \cong \text{Spec}(k[t_{13}, t_{23}, t_{14}, t_{15}, t_{25}])$ on $\mathbb{P}_k(K_4^\vee) = \mathbb{P}_k^4$ with open orbit $\mathbb{P}_k^4 \setminus \{w_{12} = 0\}$ is the toric \mathbb{G}_a^4 -structure on \mathbb{P}_k^4 given by $w_{12} \mapsto w_{12}$ and $w_{ij} \mapsto w_{ij} + t_{ij}w_{12}$ for $(i, j) \neq (1, 2), (2, 4), (2, 5)$. Its lift to $X_4 = W_L$ is the restriction of the $\mathbb{V}_k(F_4^\vee)$ -action on $\mathbb{P}_k(\Lambda^2 V^\vee)$ preserving $W_L = G \cap \{w_{13} - w_{24} = w_{14} - w_{25} = 0\}$ induced by the second exterior power of the representation

$$(3.4) \quad \rho_4 : \mathbb{V}_k(F_4^\vee) \rightarrow \text{GL}_k(V^\vee), \quad (t_{13}, t_{23}, t_{14}, t_{15}) \mapsto \begin{pmatrix} 1 & 0 & -t_{23} & -t_{13} & -t_{14} \\ 0 & 1 & t_{13} & t_{14} & t_{15} \\ \vdots & \ddots & 1 & 0 & 0 \\ \vdots & & \ddots & 1 & 0 \\ 0 & \dots & \dots & 0 & 1 \end{pmatrix}.$$

3.2. Proof of Theorem 1. We now proceed to the proof of Theorem 1, each case $n \in \{4, 5, 6\}$ is treated separately in the next subsections.

3.2.1. Proof of Theorem 1 for sixfolds. A smooth quintic del Pezzo sixfold X is a k -form of the Grassmannian $\text{G}(2, 5)$. Recall [4] that isomorphism classes of k -forms of a projective k -variety X are in one-to-one correspondence with the elements of the Galois cohomology set $H^1(\Gamma, \text{Aut}_{\bar{k}}(X_{\bar{k}}))$ of continuous Galois 1-cocycles $\gamma : \Gamma = \text{Gal}(\bar{k}/k) \rightarrow \text{Aut}_{\bar{k}}(X_{\bar{k}})$, where $\text{Aut}_{\bar{k}}(X_{\bar{k}})$ is endowed with the discrete topology and the natural action of Γ by conjugation. Since $\text{Aut}_{\bar{k}}(\text{G}(2, 5)_{\bar{k}}) \cong \text{PGL}_5(\bar{k}) = \text{Aut}_{\bar{k}}(\mathbb{P}_{\bar{k}}^4)$ (see § 1.2), isomorphism classes of k -forms of $\text{G}(2, 5)$ are in one-to-one correspondence with isomorphism classes of k -forms P of \mathbb{P}_k^4 . In view of Lemma 6, the existence of a k -rational point is a necessary condition for the existence of a vector group structure on X . Conversely, for a k -form X of $\text{G}(2, 5)$ containing a k -rational point, the corresponding k -form P of \mathbb{P}_k^4 contains a closed sub-variety C defined over k whose base extension $C_{\bar{k}}$ is a line in $P_{\bar{k}} \cong \mathbb{P}_{\bar{k}}^4$.

²By [2], every complete toric variety admitting a vector group structure has such a structure which is normalized by the torus. We used the term “toric” here to indicate the fact that the given \mathbb{G}_a^6 -action is normalized by the toric structure on \mathbb{P}_k^6 .

Since the restriction of a canonical divisor K_P of P to C has odd degree, C is the trivial k -form of \mathbb{P}_k^1 and hence, P is the trivial k -form of \mathbb{P}_k^4 . This implies in turn that X is isomorphic to $G(2, 5)$. By Proposition 24, $G(2, 5)$ admits at least one vector group structure.

The following proposition completes the proof of Theorem 1 in the case $n = 6$.

Proposition 26. *The Grassmannian $G(2, 5)$ has a unique class of vector group structure.*

Proof. Write $G(2, 5) = \mathbb{G}_k(V^\vee, 2) = G$ for some 5-dimensional k -vector space V . Through the isomorphism $\mathrm{PGL}_k(V^\vee) \cong \mathrm{Aut}_k(G)$ of (1.1), a vector group structure on G is given by the projective representation associated to a faithful representation $\rho : \mathbb{U} \rightarrow \mathrm{GL}_k(V^\vee)$ of a vector group \mathbb{U} . Let $V_1 \subset V$ be a 1-dimensional k -vector subspace of V that is invariant for the representation dual to ρ , its existence being guaranteed by Lemma 3(3). Since the action of $\mathrm{GL}_k(V)$ on such 1-dimensional subspaces is transitive, up to the choice of a basis of V as in Notation 10 and up to changing ρ by its conjugate by a suitable automorphism of G we henceforth assume without loss of generality that $V_1 = \langle e_5 \rangle$ and put $\bar{V} = V/V_1 \cong \langle e_1, \dots, e_4 \rangle$. The image of ρ is then contained in the stabilizer Δ_{V_1} of the subspace $\bar{V}^\vee \subset V^\vee$, see Notation 13. Let \mathbb{U}' be the kernel of the induced representation $\mathbb{U} \rightarrow \mathrm{GL}_k(\bar{V}^\vee)$ and let $\bar{\rho} : \bar{\mathbb{U}} = \mathbb{U}/\mathbb{U}' \rightarrow \mathrm{GL}_k(\bar{V}^\vee)$ be the induced faithful representation. With the notation of § 2.2 and Example 18, the projection

$$\pi_\Pi : G \dashrightarrow \bar{G} = \{\bar{w}_{12}\bar{w}_{34} - \bar{w}_{13}\bar{w}_{24} + \bar{w}_{14}\bar{w}_{23} = 0\} \subset \mathbb{P}_k^5$$

from the solid $\Pi = \sigma_{3,0}(V_1)$ is then \mathbb{U} -equivariant for the action of \mathbb{U} on \bar{G} factoring through the action of $\bar{\mathbb{U}}$ determined under the isomorphism $\mathrm{Aut}_k(\bar{G}) \cong \mathrm{PGL}_k(\bar{V}^\vee)$ by the projective representation induced by $\bar{\rho}$. The \mathbb{U} -action on G lifts to a vector group structure on the blow-up $Y_\Pi \rightarrow G$ of Π and, by Proposition 8 applied to the induced morphism $p_{\bar{G}} : Y_\Pi \rightarrow \bar{G}$, the $\bar{\mathbb{U}}$ -action on \bar{G} defines a vector group structure on it. In particular, $\bar{\mathbb{U}}$ is 4-dimensional, say $\bar{\mathbb{U}} = \mathrm{Spec}(k[\bar{t}_{13}, \bar{t}_{23}, \bar{t}_{14}, \bar{t}_{24}])$. By [37], \bar{G} admits a unique vector group structure given up to isomorphism by the projective representation associated to the representation

$$\bar{\mathbb{U}} \rightarrow \mathrm{GL}_k(\bar{V}^\vee), \bar{t} = (\bar{t}_{ij})_{i=1,2,j=3,4} \mapsto A(\bar{t}) = \begin{pmatrix} 1 & 0 & -\bar{t}_{23} & -\bar{t}_{24} \\ 0 & 1 & \bar{t}_{13} & \bar{t}_{14} \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Write $\mathbb{U}' = \mathrm{Spec}(k[s_1, s_2])$, $\mathbb{U} \cong \mathrm{Spec}(k[\bar{t}][s_1, s_2]) \cong \bar{\mathbb{U}} \times \mathbb{U}'$ and put $u = (\bar{t}, s_1, s_2)$. Then, with Notation 13, the representation $\rho : \mathbb{U} \rightarrow \Delta_{V_1}$ lifting $\bar{\rho} : \bar{\mathbb{U}} \rightarrow \mathrm{GL}_k(\bar{V}^\vee)$ has the form $u \mapsto M(u) = M(A(\bar{t}), 1, {}^t L(u))$ for some row matrix $L(u) = (f_i(u))_{i=1,\dots,4}$ of elements of $k[u]$ such that $M(u+u') = M(u)M(u') = M(u)M(u') = M(u'+u)$. By direct computation, this identity implies that $f_3 = f_4 = 0$ and that f_1 and f_2 are linear polynomials. Moreover, since ρ is injective, we must have $k[\bar{t}, f_1, f_2] = k[\bar{t}, s_1, s_2]$. Denoting \bar{t}_{ij} , f_1 and f_2 anew by t_{ij} , $-t_{25}$ and t_{15} then identifies ρ with the representation (3.1) in Example 25. \square

3.2.2. Proof of Theorem 1 for fivefolds. The following proposition establishes the first part of the assertion of Theorem 1 for $n = 5$.

Proposition 27. *A smooth quintic del Pezzo fivefold is isomorphic to a smooth hyperplane section of $G(2, 5)$ and all these sections are isomorphic.*

Proof. Let X be smooth quintic del Pezzo fivefold. By [19], $X_{\bar{k}}$ is isomorphic to a smooth hyperplane section $Z_{(s)}$ of $G = \mathbb{G}_{\bar{k}}(V^\vee, 2)$ for some 5-dimensional \bar{k} -vector space V . By Lemma 14, X contains a unique 3-dimensional sub-scheme Π whose base extension $\Pi_{\bar{k}}$ to \bar{k} is a $\sigma_{3,0}$ -solid of $X_{\bar{k}}$. Let $\mathcal{I}_{\Pi_{\bar{k}}} \subset \mathcal{O}_{X_{\bar{k}}}$ be the ideal sheaf of $\Pi_{\bar{k}}$, let $\mathcal{O}_{X_{\bar{k}}}(1) = \Lambda^2 \mathcal{Q}|_{X_{\bar{k}}}$ and consider the projection

$$\pi_{\Pi_{\bar{k}}} : X_{\bar{k}} = Z_{(s)} \dashrightarrow Q_{(\bar{s})} \subset \mathbb{P}_{\bar{k}}(H^0(X_{\bar{k}}, \mathcal{I}_{\Pi_{\bar{k}}}(1)) \cong \mathbb{P}_{\bar{k}}^4$$

from $\Pi_{\bar{k}}$. Since the action of the Galois group $\Gamma = \mathrm{Gal}(\bar{k}/k)$ on $X_{\bar{k}}$ maps smooth hyperplane sections of $X_{\bar{k}}$ to smooth hyperplane sections, the projective space $\mathbb{P}_{\bar{k}}(H^0(X_{\bar{k}}, \mathcal{I}_{\Pi_{\bar{k}}}(1)))$ inherits a natural continuous linear Galois action of Γ . By Galois descent for quasi-projective varieties and rational maps between these, the map $\pi_{\Pi_{\bar{k}}}$ thus descends to a rational map $\pi_\Pi : X \dashrightarrow Q \subset P_4$ whose image is a k -form Q of $Q_{(\bar{s})}$ in a k -form P_4 of \mathbb{P}_k^4 . The divisor $-K_{P_4} - 2Q$ being defined over k and of degree 1, P_4 is the trivial form of \mathbb{P}_k^4 . Let $p_X : Y \rightarrow X$ be the blow-up of Π . Then, by Proposition 17, the induced morphism $p_Q : Y \rightarrow Q$ is an étale locally trivial \mathbb{P}^2 -bundle whose base extension to \bar{k} is isomorphic to $\mathbb{P}_{Q_{(s)}}(\mathcal{S} \oplus \mathcal{O}_{Q_{(s)}})$ where \mathcal{S} denotes the spinor sheaf on $Q_{(\bar{s})}$. Furthermore, the restriction $p_Q : E \rightarrow Q$ of p_Q to the exceptional divisor of p_X is an étale locally trivial \mathbb{P}^1 -sub-bundle of $p_Q : Y \rightarrow Q$, whose base extension to \bar{k} is isomorphic to the sub-bundle $\mathbb{P}_{Q_{(\bar{s})}}(\mathcal{S})$ of $\mathbb{P}_{Q_{(\bar{s})}}(\mathcal{S} \oplus \mathcal{O}_{Q_{(\bar{s})}})$. Thus, considering the direct image by p_Q of the exact sequence $0 \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_Y(E) \rightarrow \mathcal{O}_E(E) \rightarrow 0$ on Y , we conclude that $p_Q : Y \rightarrow Q$ is isomorphic to the \mathbb{P}^2 -bundle $\mathbb{P}_Q(\mathcal{E})$ where $\mathcal{E} = (p_Q)_* \mathcal{O}_Y(E)$ is a locally free sheaf of rank 3 on Q . Furthermore, $\mathcal{S}_Q = (p_Q)_* \mathcal{O}_E(E)$ is a locally

free sheaf of rank 2 whose base extension to \bar{k} is isomorphic to the spinor sheaf \mathcal{S} on $Q_{\bar{k}} \cong Q_{\langle \bar{s} \rangle}$. Since $H^0(Q_{\langle \bar{s} \rangle}, \mathcal{S}^\vee) \cong H^0(Q, \mathcal{S}_Q^\vee) \otimes_k \bar{k}$ by flat base change and since \mathcal{S}^\vee is globally generated by sections whose zero schemes are lines of $Q_{\langle \bar{s} \rangle} \subset \mathbb{P}_{\bar{k}}^4$ (see e.g. [32]), we infer that Q contains a line of $P_4 \cong \mathbb{P}_k^4$. A smooth quadric of \mathbb{P}_k^4 containing a line being unique up to isomorphism and equal to a hyperplane section of the smooth quadric $G(2, 4) \subset \mathbb{P}_k^5$, we conclude by reading the Sarkisov link of Proposition 20 backwards that X is isomorphic to a smooth hyperplane section of $G(2, 5)$ over k . The transitivity of the action of $\text{Aut}_k(G)(k)$ on the set of such smooth sections, see § 2.1.2, implies that they are all isomorphic. \square

Since by Proposition 24 every smooth hyperplane section of $G(2, 5) \subset \mathbb{P}_k^9$ admits a vector group structure, the following proposition completes the proof of Theorem 1 in the case $n = 5$.

Proposition 28. *A smooth hyperplane section of $G(2, 5)$ has a unique class of vector group structure.*

Proof. Write $G(2, 5) = \mathbb{G}_k(V^\vee, 2) = G$ for some 5-dimensional k -vector space V . The action of $\text{Aut}_k(G)$ on the set of smooth hyperplane sections being transitive, we are reduced without loss of generality to prove that the smooth hyperplane section $Z_{\langle s \rangle}$ associated to the skew-symmetric form $s = e_1^\vee \wedge e_3^\vee - e_2^\vee \wedge e_4^\vee$ with $V^\perp = \langle e_5 \rangle$ considered in Example 25 admits a unique class of vector group structure. Under the isomorphism of Corollary 19, a vector group structure on $Z_{\langle s \rangle}$ is given by a certain faithful representation $\rho : \mathbb{U} \rightarrow \text{GL}_k(V^\vee)$ of a vector group \mathbb{U} with image contained in the subgroup of the stabilizer Δ_{V^\perp} of the subspace $\bar{V}^\vee \subset V^\vee$ consisting of matrices of the form

$$M(A_4, \lambda, U) = \begin{pmatrix} A_4 & U \\ 0 & \lambda \end{pmatrix} \text{ with } A_4 \in \text{Sp}_4(\bar{V}^\vee, \bar{s}), U \in \text{Hom}_k((V^\perp)^\vee, \bar{V}^\vee), \lambda \in \text{GL}_k((V^\perp)^\vee) = \mathbb{G}_{m,k}.$$

Let \mathbb{U}' be the kernel of the induced representation $\mathbb{U} \rightarrow \text{Sp}_4(\bar{V}^\vee, \bar{s})$ and let $\bar{\rho} : \bar{\mathbb{U}} = \mathbb{U}/\mathbb{U}' \rightarrow \text{Sp}_4(\bar{V}^\vee, \bar{s})$ be the induced injective homomorphism. With the notation of § 2.2 and Example 18, the projection

$$\pi_\Pi : Z_{\langle s \rangle} \dashrightarrow Q_{\langle \bar{s} \rangle} = \{\bar{w}_{12}\bar{w}_{34} - \bar{w}_{13}^2 + \bar{w}_{14}\bar{w}_{23} = 0\}$$

from the solid $\Pi = \sigma_{3,0}(V^\perp)$ is then \mathbb{U} -equivariant for the action of \mathbb{U} on $Q_{\langle \bar{s} \rangle}$ factoring through the action of $\bar{\mathbb{U}}$ determined under the isomorphism $\text{Aut}_k(Q_{\langle \bar{s} \rangle}) \cong \text{PSp}_k(\bar{V}^\vee, \bar{s})$ by the projective representation induced by $\bar{\rho}$. The \mathbb{U} -action on $Z_{\langle s \rangle}$ lifts to a vector group structure on the blow-up $Y_\Pi \rightarrow Z_{\langle s \rangle}$ of Π and, by Proposition 8 applied to the induced morphism $\text{p}_{Q_{\langle \bar{s} \rangle}} : Y_\Pi \rightarrow Q_{\langle \bar{s} \rangle}$, the $\bar{\mathbb{U}}$ -action on $Q_{\langle \bar{s} \rangle}$ defines a vector group structure on it. In particular, $\bar{\mathbb{U}}$ is 3-dimensional, say $\bar{\mathbb{U}} = \text{Spec}(k[\bar{t}_{13}, \bar{t}_{23}, \bar{t}_{24}])$. By [37], $Q_{\langle \bar{s} \rangle}$ admits a unique vector group structure up to isomorphism, which, with our choice of coordinates, is induced by the representation $\bar{\rho}_5 : \mathbb{V}_k(F_5^\vee)/\text{Stab}(\langle e_5^\vee \rangle) \rightarrow \text{Sp}_4(\bar{V}^\vee, \bar{s})$ of (3.3) in Example 25. Writing $\mathbb{U}' = \text{Spec}(k[s_1, s_2])$, $\mathbb{U} \cong \text{Spec}(k[\bar{t}][s_1, s_2]) \cong \bar{\mathbb{U}} \times \mathbb{U}'$ and $u = (\bar{t}, s_1, s_2)$, the same argument as in the proof of Proposition 26 implies that the lift $\rho : \mathbb{U} \rightarrow \Delta_{V^\perp} \subset \text{GL}_k(V^\vee)$ of $\bar{\rho} : \bar{\mathbb{U}} \rightarrow \text{Sp}_4(\bar{V}^\vee, \bar{s})$ has the form $u \mapsto M(u) = M(A(\bar{t}), 1, {}^t L(s))$ where $L(s) = (f_1(s), f_2(s), 0, 0)$ is a row matrix of linear elements of $k[s]$ with the property that $k[\bar{t}, f_1, f_2] = k[\bar{t}, s_1, s_2]$. Denoting the \bar{t}_{ij} , f_1 and f_2 anew by t_{ij} , $-t_{25}$ and t_{15} respectively, we get that $\rho : \mathbb{U} \rightarrow \text{GL}_k(V^\vee)$ is the representation (3.2) of Example 25. \square

3.2.3. *Proof of Theorem 1 for fourfolds.* The following proposition completes the proof of Theorem 1.

Proposition 29. *A smooth quintic del Pezzo fourfold is isomorphic to a smooth section of $G(2, 5) \subset \mathbb{P}_k^9$ by a linear subspace of codimension 2. Furthermore, all such sections are isomorphic and admit exactly one class of vector group structure.*

Proof. Let X be smooth quintic del Pezzo fourfold. By [19], we can assume that $X_{\bar{k}}$ is a smooth section W_L of $G(2, 5)_{\bar{k}}$ by a linear subspace of codimension 2. By Lemma 16, X contains a unique 2-dimensional sub-scheme Ξ whose base extension $\Xi_{\bar{k}}$ to \bar{k} is a $\sigma_{2,2}$ -plane of W_L . Let $\mathcal{I}_{\Xi_{\bar{k}}} \subset \mathcal{O}_{X_{\bar{k}}}$ be the ideal sheaf of $\Xi_{\bar{k}}$, let $\mathcal{O}_{X_{\bar{k}}}(1) = \Lambda^2 \mathcal{Q}|_{X_{\bar{k}}}$ and consider the projection

$$\pi_{\Xi_{\bar{k}}} : X_{\bar{k}} = W_L \dashrightarrow \mathbb{P}_{\bar{k}}(H^0(X_{\bar{k}}, \mathcal{I}_{\Xi_{\bar{k}}}(1)) \cong \mathbb{P}_{\bar{k}}^4)$$

from $\Xi_{\bar{k}}$ as in § 2.3. The projective space $\mathbb{P}_{\bar{k}}(H^0(X_{\bar{k}}, \mathcal{I}_{\Xi_{\bar{k}}}(1)))$ inherits a natural continuous linear action of the Galois group $\Gamma = \text{Gal}(\bar{k}/k)$, which stabilizes the hyperplane corresponding to the unique section of $\mathcal{I}_{\Xi_{\bar{k}}}(1)$ whose zero scheme is the special hyperplane section $Z_{\Xi_{\bar{k}}} \cap X_{\bar{k}}$ of $X_{\bar{k}}$ spanned by the $\sigma_{3,1}$ -planes of $X_{\bar{k}}$. In particular, the divisor $Z_{\Xi_{\bar{k}}} \cap X_{\bar{k}}$ is defined over k , say $Z_{\Xi_{\bar{k}}} \cap X_{\bar{k}} = (Z_{\Xi})_{\bar{k}}$ for some geometrically irreducible divisor $Z_{\Xi} \subset X$ containing Ξ . In view of Proposition 20, the projection $\pi_{\Xi_{\bar{k}}}$ thus descends to a birational map $\pi_{\Xi} : X \dashrightarrow P_4$, with image equal to a k -form P_4 of \mathbb{P}_k^4 , which contract Z_{Ξ} onto a smooth curve C contained in a hypersurface $P_3 \subset P_4$ such that the triple (P_4, P_3, C) is a k -form of the triple $(\mathbb{P}_k^4, \mathbb{P}_k^3, C_3)$, where \mathbb{P}_k^3 is a hyperplane of \mathbb{P}_k^4 and $C_3 \subset \mathbb{P}_k^3$ is a smooth rational cubic curve. Thus, P_4 and P_3 are trivial forms of \mathbb{P}_k^4 and \mathbb{P}_k^3 respectively, and, since the intersection of C with a hyperplane of \mathbb{P}_k^3 is a divisor of degree

3 on C , it follows that $C \cong \mathbb{P}_k^1$. Reading the Sarkisov link of Proposition 20 backwards, we conclude that X is isomorphic to a smooth section of $G(2, 5)$ by a linear subspace of codimension 2, and the uniqueness up to isomorphism follows from the transitivity of the action $\mathrm{PGL}_5(k)$ on triples $(\mathbb{P}_k^4, \mathbb{P}_k^3, C_3)$ where $C_3 \subset \mathbb{P}_k^3$ is a smooth rational cubic. Proposition 24 implies that X admits a vector group structure. To show the uniqueness, we observe that being unique, the $\sigma_{2,2}$ -plane Ξ is stable under any faithful action of a unipotent abelian group \mathbb{U} on X defining a vector group structure on X . This vector group structure lifts to the blow-up $p_\Xi : Y \rightarrow X$ of Ξ and then, by Proposition 8, descends via the contraction $p_2 : Y \rightarrow \mathbb{P}_k^4$ of the proper transform of Z_Ξ to a faithful \mathbb{U} -action on \mathbb{P}_k^4 defining a vector group structure, for which the pair (\mathbb{P}_k^3, C) is globally \mathbb{U} -stable. By the classification of vector group structures on \mathbb{P}_k^4 , see [21] or [22, Corollary 3.6], the unique class of vector group structure with this property is the toric \mathbb{G}_a^4 -structure on \mathbb{P}_k^4 described in Example 25. \square

Remark 30. A by-product of Proposition 29 is that among the four non-isomorphic compactifications of $\mathbb{A}_{\mathbb{C}}^4$ into the smooth quintic del Pezzo fourfold classified by Prokhorov [34, Theorem 3.1], only the case (i) can be endowed with a vector group structure making it an *equivariant* compactification of $\mathbb{G}_{a,\mathbb{C}}^4$.

Remark 31. By [14, § 2.2], smooth sections of $G(2, 5) \subset \mathbb{P}_k^9$ by linear subspaces of codimension 3 have in general non-trivial k -forms, whose isomorphism classes are parametrized by equivalence classes of non-degenerate ternary quadratic forms over k . In contrast, Proposition 27 and Proposition 29 imply that smooth del Pezzo quintics of dimension $n = 4, 5$ do not have non-trivial k -forms. Since these are compactifications of \mathbb{A}_k^n , one deduces from the techniques in *loc. cit.* that a proper morphism $f : X \rightarrow Y$ between normal varieties over an algebraically closed field of characteristic zero whose general closed fibers are smooth quintic del Pezzo varieties of dimension $n = 4, 5$ contains a vertical \mathbb{A}^n -cylinder in the sense of [14]. Similarly, a proper morphism $f : X \rightarrow Y$ whose general closed fibers are isomorphic to $G(2, 5)$ and which has a rational section contains a vertical \mathbb{A}^6 -cylinder.

4. VECTOR GROUP STRUCTURES ON TERMINAL QUINTIC DEL PEZZO THREEFOLDS AND CANONICAL QUINTIC DEL PEZZO SURFACES

Classifying vector group structures on all del Pezzo quintics, including singular ones, is a challenging problem. For instance, many families of singular del Pezzo quintics of dimension $n = 3, 4, 5$ endowed with a vector group structure can be constructed as hyperplanes sections of $G = G(2, 5)$ with respect to the Plücker embedding containing a $\sigma_{2,2}$ -plane Ξ of G (see § 2.3 for the notation) which is fixed by the $\mathbb{G}_{a,k}^6$ -structure on G corresponding under the birational projection $\pi_\Xi : G = G(2, 5) \dashrightarrow \mathbb{P}_k^6 = \mathbb{P}(H^0(G, \mathcal{I}_\Xi(1)))$ to the toric $\mathbb{G}_{a,k}^6$ -structure on \mathbb{P}_k^6 described in Example 25. Namely, every linear subspace L of codimension $m = 1, 2, 3$ of \mathbb{P}_k^6 which intersects the complement $H_\infty = \{w_{12} = 0\}$ of the open orbit transversely has stabilizer isomorphic to $\mathbb{G}_{a,k}^m$ and the induced effective action of the quotient group $\mathbb{G}_{a,k}^{6-m}$ on L defines a vector group structure on it. The proper transform of L by π_Ξ is then a linear section X_L of G of dimension $6 - m$, singular in general, endowed with a vector structure for which the inclusion $X_L \hookrightarrow G$ is equivariant for the induced action of a subgroup $\mathbb{G}_{a,k}^{6-m}$ of the group $\mathbb{G}_{a,k}^6$ which acts with an open orbit on G .

Here, we consider mildly singular del Pezzo quintics of dimension 3 and 2, a case of special interest due to the fact that no smooth del Pezzo quintics in these dimensions admit a vector group structure.

4.1. Vector group structures on terminal quintic del Pezzo threefolds. Over algebraically closed fields of characteristic zero, non-smooth terminal quintic del Pezzo threefolds are classified in [35]. By Corollary 5.3 in *loc. cit.*, the main invariant of such a threefold X is the rank $r(X) = \mathrm{rk}_{\mathbb{Z}} \mathrm{Cl}(X)$ of its divisor class group, which is either 2, 3 or 4. Furthermore, all the singularities of X are ordinary double points and, by [35, Corollary 8.3.1], the number of such nodes equals $r(X) - 1$.

Lemma 32. *A terminal quintic del Pezzo threefold X over k whose base extension $X_{\bar{k}}$ to \bar{k} has one or two nodes does not admit a vector group structure.*

Proof. Since it is enough to show that $X_{\bar{k}}$ does not admit any vector group structure, we henceforth assume that $k = \bar{k}$. Assume that X has a vector group structure. Then the latter lifts to a vector group structure on a small \mathbb{Q} -factorialization $\xi : \hat{X} \rightarrow X$ of X . If X has a unique node then, by [25, Theorem 3.6] or [35, Theorem 5.2], \hat{X} is isomorphic to the total space of a \mathbb{P}^1 -bundle $\pi : \hat{X} \cong \mathbb{P}_k^2(\mathcal{E}) \rightarrow \mathbb{P}_k^2$ for some stable, hence simple, locally free sheaf \mathcal{E} of rank 2 on \mathbb{P}_k^2 . We would thus obtain a vector group structure on $\mathbb{P}_{\mathbb{P}_k^2}(\mathcal{E})$ which is impossible by Corollary 9. If X has two nodes then, by [35, Theorem 8.1 and Corollary 8.1.1], \hat{X} is isomorphic to the blow-up $\sigma : \hat{X} \rightarrow \tilde{X}$ of the total space of the projective bundle $\pi : \tilde{X} = \mathbb{P}_{\mathbb{P}_k^2}(\mathcal{T}_{\mathbb{P}_k^2}(-1)) \rightarrow \mathbb{P}_k^2$ at a point. By Proposition 8, the vector group structure on \hat{X} descends to a vector group structure on \tilde{X} . But again, this is impossible since $\mathcal{T}_{\mathbb{P}_k^2}(-1)$ is a simple sheaf. \square

We now consider the case of terminal quintic del Pezzo threefolds whose base extensions to \bar{k} possess exactly three nodes, which we henceforth call for short *trinodal quintic del Pezzo threefolds*.

Proposition 33. *A trinodal quintic del Pezzo threefold X_3 is isomorphic to a section of $G(2, 5) \subset \mathbb{P}_k^9$ by a linear subspace of codimension 3. It contains a unique $\sigma_{2,2}$ -plane Ξ of $G(2, 5)$ and the projection $\pi_\Xi : G(2, 5) \dashrightarrow \mathbb{P}_k^6$ from Ξ induces the following Sarkisov link*

$$(4.1) \quad \begin{array}{ccccc} \text{Bl}_{S_{X_3}} H_{X_3} & \hookrightarrow & \text{Bl}_\Xi X_3 \cong \text{Bl}_{S_{X_3}} \mathbb{P}_k^3 & & \\ \downarrow & & \swarrow \text{p}_{X_3} & & \searrow \text{p}_2 \\ \Xi & \hookrightarrow & X_3 & \xleftarrow{\text{---} \pi_\Xi \text{---}} & \mathbb{P}_k^3 \xleftarrow{\text{---} \Phi \text{---}} H_{X_3} \xleftarrow{\text{---} S_{X_3}} \end{array}$$

where $\text{p}_{X_3} : \text{Bl}_\Xi X_3 \rightarrow X_3$ is the blow-up of Ξ , $\text{p}_2 : \text{Bl}_\Xi X_3 \rightarrow \mathbb{P}_k^3$ is the blow-up of a smooth 0-dimensional sub-scheme S_{X_3} of length 3 of \mathbb{P}_k^3 not contained in a line, H_{X_3} is the unique hyperplane of \mathbb{P}_k^3 containing S_{X_3} and $\text{Bl}_{S_{X_3}} H_{X_3}$ is its proper transform. Furthermore, the birational inverse Φ of π_Ξ is given by the complete linear system of quadrics of \mathbb{P}_k^3 containing S_{X_3} .

Proof. By [35, Theorem 7.1], $X_{3,\bar{k}}$ is isomorphic to a threefold obtained from $\mathbb{P}_{\bar{k}}^3$ as the blow-up $\sigma : Y \rightarrow \mathbb{P}_{\bar{k}}^3$ of three non-colinear closed points, say p_1, p_2, p_3 , followed by the contraction $\xi : Y \rightarrow X_{3,\bar{k}}$ of the proper transforms by σ of the lines in $\mathbb{P}_{\bar{k}}^3$ passing through p_i and p_j , $1 \leq i < j \leq 3$ to the nodes of $X_{3,\bar{k}}$. The class group of $X_{3,\bar{k}}$ is freely generated by the classes of the proper transform \bar{H} of the unique hyperplane $H \subset \mathbb{P}_{\bar{k}}^3$ containing the points p_i and of the images F_i , $1 \leq i \leq 3$, of the exceptional divisors of σ . By [35, Theorem 7.2], P and the F_i are the only planes contained in $X_{3,\bar{k}}$. Their union is thus defined over k and since P is the only plane among these which fully contains the singular locus of $X_{3,\bar{k}}$ it is defined over k as well, say $P = \Xi_{\bar{k}}$ for some closed sub-scheme Ξ of X_3 . This implies in turn that the union of the F_i is defined over k , say $\bigcup F_i = F_{\bar{k}}$ for some closed sub-scheme F of X_3 . Since the divisor $(\Xi + F)_{\bar{k}} = P + \sum_{i=1}^3 F_i$ is linearly equivalent to the proper transform in $X_{3,\bar{k}}$ of any hyperplane $H' \subset \mathbb{P}_{\bar{k}}^3$ not passing through the points blown-up, it is Cartier. The invertible sheaf $\mathcal{O}_{X_{3,\bar{k}}}(P + \sum_{i=1}^3 F_i)$ is then the base extension to \bar{k} of the invertible sheaf $\mathcal{O}_{X_3}(1) := \mathcal{O}_{X_3}(\Xi + F)$ and, letting $\mathcal{I} \subset \mathcal{O}_{X_3}$ be the ideal sheaf of the singular locus of X_3 , the rational map $\sigma \circ \xi^{-1} : X_{3,\bar{k}} \dashrightarrow \mathbb{P}_{\bar{k}}^3$ is the base extension of the birational map $\pi : X_3 \dashrightarrow \mathbb{P}(H^0(X_3, \mathcal{I}(1))) \cong \mathbb{P}_k^3$. The latter maps Ξ to a hyperplane $H_{X_3} \subset \mathbb{P}_k^3$ and contracts F to a smooth closed sub-scheme $S_{X_3} \subset H_{X_3}$ of length 3 whose base extension to \bar{k} equals the union of the points p_i . Since S_{X_3} is not contained in a line, there exists a smooth rational cubic curve $S_{X_4} \subset \mathbb{P}_k^3$ whose scheme-theoretic intersection with H_{X_3} equals S_{X_3} . Considering \mathbb{P}_k^3 as a hyperplane H_{X_4} of \mathbb{P}_k^4 , it follows from Proposition 20 that the image of the rational map $\mathbb{P}_k^4 \dashrightarrow \mathbb{P}_k^7$ given by the complete linear system of quadrics containing $S_{X_4} \subset H_{X_4}$ is a smooth quintic del Pezzo fourfold X_4 containing X_3 as a hyperplane section and which has the proper transform Ξ of H_{X_4} as its unique $\sigma_{2,2}$ -plane. By construction, the birational map $\pi : X_3 \dashrightarrow \mathbb{P}_k^3$ then coincides with the restriction to X_3 of the projection $\pi_\Xi : X_4 \dashrightarrow \mathbb{P}_k^4$ from Ξ , which completes the proof. \square

Corollary 34. *Isomorphism classes of trinodal quintic del Pezzo threefolds are in one-to-one correspondence with $\text{PGL}_2(k)$ -orbits of smooth 0-dimensional sub-schemes of \mathbb{P}_k^3 of length three. Furthermore, every such threefold admits a unique class of vector group structure.*

Proof. By Proposition 33, two trinodal quintic del Pezzo threefolds X_3 and X'_3 are isomorphic if and only if there exists an automorphism of \mathbb{P}_k^3 which maps the pair (H_{X_3}, S_{X_3}) onto the pair $(H_{X'_3}, S_{X'_3})$. Being of length 3 and not contained in a line, the schemes S_{X_3} and $S_{X'_3}$ are contained in smooth k -rational conics C_{X_3} and $C_{X'_3}$ of H_{X_3} and $H_{X'_3}$, respectively. Since $\text{Aut}_k(\mathbb{P}_k^3)$ acts transitively on pairs (H, C) consisting of a hyperplane H of \mathbb{P}_k^3 and a smooth k -rational conic $C \cong \mathbb{P}_k^1$ in it and since for such pairs the restriction homomorphism $\text{Aut}_k(H, C) \rightarrow \text{Aut}_k(C)$ is an isomorphism, we conclude that X_3 and X'_3 are isomorphic if and only if there exists an isomorphism $\varphi : C_{X_3} \rightarrow C_{X'_3}$ which maps S_{X_3} onto $S_{X'_3}$. This holds if and only if S_{X_3} and $\varphi^{-1}(S_{X'_3})$ belong to the same orbit of the action of $\text{Aut}_k(C_{X_3})(k) \cong \text{PGL}_2(k)$.

For the second assertion, since the singular locus of X_3 and the unique $\sigma_{2,2}$ -plane Ξ of X_3 are stable under any vector group action on X_3 , it follows from Proposition 8 applied to the birational morphism $\text{Bl}_\Xi X \rightarrow \mathbb{P}_k^3$ that the Sarkisov link of Proposition 33 is equivariant for any vector group structure on X_3 and that the corresponding vector group structure on \mathbb{P}_k^3 stabilizes the non-linear closed sub-scheme S_{X_3} of length 3. By the classification [21] of vector group structures on $\mathbb{P}_k^3 = \text{Proj}_k(k[x_0, x_1, x_2, x_3])$, the unique class of vector group structure with this property is that of the toric \mathbb{G}_a^3 -structure defined by $x_0 \mapsto x_0$ and $x_i \mapsto x_i + t_i x_0$, $1 \leq i \leq 3$. Conversely, this structure lifts to a vector group structure on the blow-up of S_{X_3} , which, by Proposition 8 again, descends in turn to a vector group structure on X_3 . \square

Remark 35. By Proposition 33 and Corollary 34, every trinodal quintic del Pezzo threefold contains the affine 3-space \mathbb{A}_k^3 as a Zariski open subset. In contrast, there exist in general k -forms of smooth quintic del Pezzo threefolds which do not contain \mathbb{A}_k^3 , see [14, Theorem 12].

Example 36. With Notation 10, let $V_3 = \langle e_3, e_4, e_5 \rangle$ and let $\Xi = \sigma_{2,2}(V_3)$ be the associated plane of $G = G(2, 5) \subset \mathbb{P}_k^9$ as in Example 21. For every $\beta \in k^*$, the linear section

$$X_3(\beta) = G \cap \{w_{13} - w_{24} = 0\} \cap \{\beta w_{14} - w_{25} = 0\} \cap \{\beta w_{15} + w_{23} = 0\}$$

is a trinodal quintic del Pezzo threefold containing Ξ , isomorphic to the sub-variety in \mathbb{P}_k^6 with coordinates w_{ij} , $(i, j) \neq (2, 3), (2, 4), (2, 5)$ defined by the equations

$$\begin{cases} w_{12}w_{34} - w_{13}^2 - \beta w_{14}w_{15} = 0 \\ w_{12}w_{35} - \beta w_{13}w_{14} - \beta w_{15}^2 = 0 \\ w_{12}w_{45} - \beta w_{14}^2 + w_{13}w_{15} = 0 \\ w_{13}w_{45} - w_{14}w_{35} + w_{15}w_{34} = 0 \\ -\beta w_{15}w_{45} - w_{13}w_{35} + \beta w_{14}w_{34} = 0 \end{cases}$$

Its singular locus $\text{Sing}(X_3(\beta))$ is the closed sub-scheme of $\Xi \cong \text{Proj}_k(k[w_{34}, w_{35}, w_{45}])$ with equations

$$\beta w_{34}w_{45} - w_{35}^2 = w_{34}w_{35} - \beta w_{45}^2 = w_{35}w_{45} - w_{34}^2 = 0.$$

Letting $\lambda, \epsilon \in \bar{k}$ be respectively a third root of β and a primitive third root of unity, the singular locus of $X_3(\beta)_{\bar{k}}$ is the union of the three closed points $[\lambda\epsilon^m : (\lambda\epsilon^m)^2 : 1]$, $0 \leq m \leq 2$, of Ξ . Thus, according to whether β is cube in k^* or not and k^* contains a primitive primitive third root of unity or not, $\text{Sing}(X_3(\beta))$ consists either of a single closed point, or the union of a k -point and a single other closed point, or the union of three k -points. The image of Ξ by the restriction $X_3(\beta) \dashrightarrow \mathbb{P}_k^3 = \text{Proj}_k(k[w_{12}, w_{13}, w_{14}, w_{15}])$ of the projection from Ξ is the hyperplane $H_{X_3(\beta)} = \{w_{12} = 0\}$. The associated smooth 0-dimensional sub-scheme $S_{X_3(\beta)}$ of length 3 is the closed sub-scheme of $H_{X_3(\beta)}$ defined by the equations

$$\beta w_{14}w_{15} + w_{13}^2 = w_{13}w_{14} + w_{15}^2 = w_{13}w_{15} - \beta w_{14}^2 = 0.$$

The restriction to $X_3(\beta)$ of the action of the sub-group $\mathbb{U} \cong \mathbb{G}_{a,k}^3$ of the vector group $\mathbb{V}_k(F_6^\vee)$ of (3.1) in Example 25 defined by $t_{24} = t_{13}$, $t_{25} = \beta t_{14}$ and $t_{23} = -\beta t_{15}$ induces a vector group structure

$$\begin{cases} w_{12} \mapsto w_{12} \\ w_{13} \mapsto w_{13} + t_{13}w_{12} \\ w_{14} \mapsto w_{14} + t_{14}w_{12} \\ w_{15} \mapsto w_{15} + t_{15}w_{12} \\ w_{34} \mapsto w_{34} + 2t_{13}w_{13} + \beta(t_{15}w_{14} + t_{14}w_{15}) + (t_{13}^2 + \beta t_{14}t_{15})w_{12} \\ w_{35} \mapsto w_{35} + \beta(t_{14}w_{13} + t_{13}w_{14} + 2t_{15}w_{15}) + \beta(t_{13}t_{14} + t_{15}^2)w_{12} \\ w_{45} \mapsto w_{45} + (2\beta t_{14}w_{14} - t_{15}w_{13} - t_{13}w_{15}) + (\beta t_{14}^2 - t_{15}t_{13})w_{12} \end{cases}$$

on $X_3(\beta)$ with open orbit $X_3(\beta) \setminus \{w_{12} = 0\}$.

4.2. Vector group structures on canonical quintic del Pezzo surfaces. Del Pezzo surfaces with canonical singularities admitting a vector group structure are classified in [12], see also [30]. For the sake of completeness, we record the following consequence of the classification in the quintic case:

Proposition 37. *Up to isomorphism, there exist two quintic del Pezzo surfaces with canonical singularities which admit a vector group structure:*

a) *A surface S with an A_3 -singularity, whose neutral component $\text{Aut}^0(S)$ of the automorphism group is isomorphic to $\mathbb{G}_{a,k}^2 \rtimes \mathbb{G}_{m,k}$ and which admits a unique class of vector group structure.*

b) *A surface S' with an A_4 -singularity, whose neutral component $\text{Aut}^0(S')$ of the automorphism group is isomorphic to $\mathbb{U}_3 \rtimes \mathbb{G}_{m,k}$, where \mathbb{U}_3 is a maximal unipotent subgroup of $\text{PGL}_3(k)$, and which admits exactly two classes of vector group structures.*

Proof. All the properties but those concerning the actual number of equivalence classes of vector group structures are established in [12] and for the description of the automorphism groups in [30, Table 1], cases 5E and 5F for S and S' respectively. We briefly recall the principle of the argument in *loc. cit.* and explain how derive from it the equivalence classes of vector group structures. Equivalence classes of vector group structures on a del Pezzo surface S with canonical singularities are in one-to-one correspondence with those on its minimal desingularization $\tilde{S} \rightarrow S$, which is obtained from S by performing a finite sequence of successive blow-ups of singular loci of intermediate surfaces and normalizations. Indeed, a vector group structure stabilizes singular loci, hence canonically lifts to their blow-ups and, by universal property, canonically lifts as well to normalizations. Conversely, Proposition 8 ensures that every vector group structure on \tilde{S}

10. A. Chambert-Loir and Y. Tschinkel, *Integral points of bounded height on partial equivariant compactifications of vector groups*, Duke Math. J., 161(15):2799-2836, 2012.
11. M. Demazure, P. Gabriel, *Groupes algébriques*, Masson, Paris, 1970.
12. U. Derenthal and D. Loughran, *Singular del Pezzo surfaces that are equivariant compactifications*, J. Math. Sci. (N.Y.) **171** (2010), no. 6, 714–724.
13. R. Donagi, *On the geometry of Grassmannians*, Duke Math. J. 44 (1977), no. 4, 795-837.
14. A. Dubouloz and T. Kishimoto, *Cylindres dans les fibrations de Mori: formes du volume quintique de del Pezzo*, Annales de L'institut Fourier, **69** (2019), 2377-2393.
15. B. Fu and J.-M. Hwang, *Uniqueness of equivariant compactifications of \mathbb{C}^n by a Fano manifold of Picard number 1*, Math. Res. Lett., **21** (2014), no. 1, 121-125.
16. B. Fu and J. M. Hwang, *Special birational transformations of type (2, 1)*, J. Algebraic Geom., **27** (2018), no. 1, 55-89.
17. B. Fu and J. M. Hwang *Euler-symmetric projective varieties*, Algebr. Geom. 7 (2020), no. 3, 377-389.
18. B. Fu and P. Montero, *Equivariant compactifications of vector groups with high index*, C. R. Acad. Sci. Paris, Ser. I, **357** (2019), 455-461.
19. T. Fujita, *On the structure of polarized manifolds with total deficiency one, II*, J. Math. Soc. Japan, **33** (1981), 415-434.
20. A. Grothendieck, *Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes*, Inst. Hautes Études Sci. Publ. Math. No. 8 (1961).
21. B. Hassett and Y. Tschinkel, *Geometry of Equivariant Compactifications of \mathbb{G}_a^n* , International Mathematics Research Notices, **22** (1999), 1211-1230.
22. Z. Huang and P. Montero, *Fano threefolds as equivariant compactifications of the vector group*, Michigan Math. J., **69** (2020), 341-368.
23. D. Huybrechts and M. Lehn, *The geometry of moduli spaces of sheaves*, Second edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2010.
24. J.M. Hwang and N. Mok, *Cartan-Fubini type extension of holomorphic maps for Fano manifolds of Picard number 1*, J. Math. Pures Appl. 80 (2001), no. 6, 563-575.
25. P. Jahnke and T. Peternell, *Almost del Pezzo manifolds*, Adv. in Geom., **8** (2008), 387-411.
26. S. L. Kleiman, *Geometry on Grassmannians and applications to splitting bundles and smoothing cycles*, Inst. Hautes Études Sci. Publ. Math. No. 36 (1969), 281-297.
27. J. Kollár, *Lectures on Resolution of Singularities*, Annals of Mathematics Studies 166, Princeton University Press, 2007.
28. V. Lakshmibai and J. Brown, *The Grassmannian variety. Geometric and representation-theoretic aspects*, Developments in Mathematics, 42. Springer, New York, 2015.
29. D. Mumford, J. Fogarty and F. Kirwan, *Geometric invariant theory*, Third edition. Ergebnisse der Mathematik und ihrer Grenzgebiete (2), 34. Springer-Verlag, Berlin, 1994.
30. G. Martin and C. Stadlmayr, *Weak del Pezzo surfaces with global vector fields*, arXiv:2007.03665.
31. M. Nagaoka, *\mathbb{G}_a^3 -structures on del Pezzo fibrations*, Michigan Math. J., 71 (2022), 601-610.
32. G. Ottaviani, *Spinor bundles on quadrics*, Trans. Amer. Math. Soc. **307** (1988), no. 1, 301-316.
33. J. Piontkowski and A. Van de Ven, *The automorphism group of linear sections of the Grassmannians $\mathbf{G}(1, N)$* , Doc. Math. 4 (1999), 623-664.
34. Yu. Prokhorov, *Compactifications of \mathbb{C}^4 of index 3*, Algebraic Geometry and its Applications. Aspects of Mathematics, Vieweg, Braunschweig, vol. E25 (1994), 159-169.
35. Yu. Prokhorov, *G-Fano threefolds, I*, Adv. in Geom., **13** (2013), 389-418.
36. M. Rosenlicht, *On quotient varieties and the affine embedding of certain homogeneous spaces*, Trans. Amer. Math. Soc. 101(1961) 211-223.
37. E.V. Sharoyko, *Hassett-Tschinkel correspondence and automorphisms of a quadric*, Sb. Math., **200** (2009), 145-160.
38. H. Sumihiro, *Equivariant completion*, J. Math. Kyoto Univ. 14(1): 1-28 (1974).
39. D. A. Suprunenko and R.Y Tylčevič, *Commutative matrices*, Nauka i Tekhnika, Minsk, 1966 104 pp.
40. M. Szurek and J. Wiśniewski, *On Fano manifolds, which are \mathbf{P}^k -bundles over \mathbf{P}^2* , Nagoya Math. J. 120 (1990), 89-101.
41. J. A. Todd, *The Locus Representing the Lines of Four-Dimensional Space and its Application to Linear Complexes in Four Dimensions*, Proc. London Math. Soc. (2) 30 (1930), no. 7, 513–550.
42. *Correspondence*, Attributed to A. Weil, Amer. J. Math. 79 (1957), 951–952.
43. J. Weyman, *Cohomology of vector bundles and syzygies*, Cambridge Tracts in Mathematics, 149. Cambridge University Press, Cambridge, 2003.

IMB UMR5584, CNRS, UNIV. BOURGOGNE FRANCHE-COMTÉ, F-21000 DIJON, FRANCE.
 Email address: adrien.dubouloz@u-bourgogne.fr

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, SAITAMA UNIVERSITY, SAITAMA 338-8570, JAPAN
 Email address: tkishimo@rimath.saitama-u.ac.jp

DEPARTAMENTO DE MATEMÁTICA UNIVERSIDAD TÉCNICA FEDERICO SANTA MARÍA
 AVENIDA ESPAÑA 1680, VALPARAÍSO, CHILE
 Email address: pedro.montero@usm.cl