



Gas-Phase Ozone Reaction Kinetics of C-5-C-8 Unsaturated Alcohols of Biogenic Interest

Asma Grira, Cornelia Amarandei, Claudiu Roman, Oumaya Bejaoui, Nouha Aloui, Gisele El Dib, Cecilia Arsene, Iustinian G. Bejan, Romeo I. Olariu, André Canosa, et al.

► To cite this version:

Asma Grira, Cornelia Amarandei, Claudiu Roman, Oumaya Bejaoui, Nouha Aloui, et al.. Gas-Phase Ozone Reaction Kinetics of C-5-C-8 Unsaturated Alcohols of Biogenic Interest. *Journal of Physical Chemistry A*, 2022, 126 (27), pp.4413-4423. 10.1021/acs.jpca.2c02805 . hal-03772466

HAL Id: hal-03772466

<https://hal.science/hal-03772466>

Submitted on 17 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Gas-phase ozone reaction kinetics of C₅-C₈ unsaturated alcohols of biogenic interest

Asma Grira^{1,2#}, Cornelia Amarandei^{3,4#}, Claudiu Roman^{3,4#}, Oumaya Bejaoui¹, Nouha Aloui¹, Gisèle El Dib², Cecilia Arsene^{3,4}, Iustinian G. Bejan^{3,4}, Romeo I. Olariu^{3,4}, André Canosa², Alexandre Tomas^{1,*}*

¹IMT Nord Europe, Institut Mines-Télécom, Univ. Lille, Center for Energy and Environment, 59000 Lille, France

²Université de Rennes, CNRS, IPR (Institut de Physique de Rennes), UMR 6251, Rennes, F-35000, France

³Faculty of Chemistry, Department of Chemistry, 11 Carol I, “Alexandru Ioan Cuza” University of Iasi, 700506 Iasi, Romania

⁴Integrated Centre of Environmental Science Studies in the North Eastern Region, 11 Carol I, “Alexandru Ioan Cuza” University of Iasi, 700506 Iasi, Romania.

first three authors have equal contributions

*Corresponding authors: alexandre.tomas@imt-nord-europe.fr, +33 327 712 651; oromeo@uaic.ro, +40232201354

ABSTRACT

Unsaturated alcohols are volatile organic compounds (VOCs) characterizing the emissions of plants. Changes in climate together with related increases of biotic and abiotic stresses are expected to strengthen these emissions in the future. Ozonolysis of unsaturated alcohols is one of the oxidation pathways that control their fate in the atmosphere. The rate coefficients of gas-phase O_3 reaction with seven C_5 to C_8 unsaturated alcohols have been determined at 296 K using both absolute and relative kinetic methods. The following rate coefficients have been obtained (in $cm^3 molecule^{-1} s^{-1}$) from the absolute method: $(1.1 \pm 0.2) \times 10^{-16}$ for *cis*-2-penten-1-ol, $(1.2 \pm 0.2) \times 10^{-16}$ for *trans*-2-hexen-1-ol, $(6.4 \pm 1.0) \times 10^{-17}$ for *trans*-3-hexen-1-ol, $(5.8 \pm 0.9) \times 10^{-17}$ for *cis*-3-hexen-1-ol, $(2.0 \pm 0.3) \times 10^{-17}$ for 1-octen-3-ol and $(8.4 \pm 1.3) \times 10^{-17}$ for *trans*-2-octen-1-ol and from the relative method: $(1.27 \pm 0.11) \times 10^{-16}$ for *trans*-2-hexen-1-ol, $(5.01 \pm 0.30) \times 10^{-17}$ for *trans*-3-hexen-1-ol, $(4.13 \pm 0.34) \times 10^{-17}$ for *cis*-3-hexen-1-ol, $(1.40 \pm 0.12) \times 10^{-16}$ for *trans*-4-hexen-1-ol. Alkenols display high reactivity with ozone with lifetimes in the hour range. Rate coefficients show a strong and complex dependence on the structure of the alkenol, particularly the relative position of the OH group towards the C=C double bond. The results are discussed and compared to the available literature data and four Structure Activity Relationship (SAR) methods.

KEYWORDS: unsaturated alcohols, Green Leaf Volatile (GLV), ozonolysis, absolute kinetic method, relative kinetic method, Structure Activity Relationship (SAR) approaches





Introduction



It is widely accepted that ozone plays a key role in controlling both atmospheric chemistry and climate change. This is due to its oxidative and light-absorbing properties which present a threat for all the ecosystem elements^{1,2} and to its role in producing hydroxyl (OH) radical which initiates the majority of oxidation processes in the atmosphere³. Therefore, ozone reactivity has been a subject of intensive studies and continues to be of significant concern worldwide. Background ozone concentrations in remote areas are generally within a few $\mu\text{g m}^{-3}$, yet, in many locations, they have been shown to increase steadily in the past decades partly due to increased global emissions of nitrogen oxides (NO_x) and volatile organic compounds (VOCs)⁴⁻⁷.

Unsaturated alcohols or alkenols are VOCs emitted by numerous plants due to biological activity (photosynthesis, transpiration...) and cell decomposition⁸. More specifically, C_6 and C_9 compounds have been shown to be ubiquitous species characterizing the emissions of plants, with stronger releases under biotic and abiotic stresses with transient emission rates up to 2 orders of magnitude larger than monoterpenes⁹. Once in the atmosphere, the double bond present in the carbon skeleton makes such alcohols quite reactive towards the atmospheric oxidants, thus contributing to the oxidative capacity of the atmosphere. While reactions of unsaturated alcohols with OH radical are relatively well known¹⁰, reactions with ozone have been less investigated and uncertainties remain on their reactivity and atmospheric impact. These latter reactions make unsaturated alcohols relatively short-lived (atmospheric lifetimes around minutes to hours)¹¹ and usually associated with the formation of organic acids¹², organic hydroperoxides¹³, and secondary organic aerosols¹⁴ (SOA) which often appear as a haze over forests¹⁵.

Regarding O₃ + alkenols literature kinetics data, only ambient temperature rate coefficients have been reported. The kinetics of *cis*-2-penten-1-ol was first investigated by Grosjean and Grosjean¹⁶, but the obtained value for the rate coefficient $(1.69 \pm 0.25) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ is inconsistent with two more recent studies^{14,17} which reported $(1.15 \pm 0.07) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ and $(1.05 \pm 0.20) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, respectively. The O₃ kinetics of *trans*-2-hexen-1-ol was recently studied by Gibilisco et al.¹⁸ and Lin et al.¹⁹. The obtained rate coefficients differ by almost a factor of 3, with values of $(0.60 \pm 0.07) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ and $(1.66 \pm 0.22) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, respectively. The rate coefficients for the reaction of O₃ with *trans*-3-hexen-1-ol measured by Gibilisco et al.¹⁸, Lin et al.¹⁹ and Kalalian et al.¹⁷ are $(5.83 \pm 0.86) \times 10^{-17}$, $(6.19 \pm 0.72) \times 10^{-17}$ and $(6.24 \pm 1.37) \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, respectively. For *cis*-3-hexen-1-ol, the study of Grosjean et al.²⁰ is about twice higher than those of Atkinson et al.²¹, Gibilisco et al.¹⁸, Lin et al.¹⁹, and Chen et al.²², raising the concern about the kinetic rate coefficients obtained for *cis*-2-penten-1-ol by Grosjean and Grosjean¹⁶. The rate coefficient for the reaction of O₃ with *trans*-4-hexen-1-ol was measured only by Lin et al.¹⁹, who reported $(1.05 \pm 0.21) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. For 1-octen-3-ol ozonolysis, to the best of our knowledge, only Li et al.²³ investigated the kinetics. Table S1 summarizes the above available literature data.

The objective of this work was thus to investigate the ozone kinetics of a series of seven unsaturated alcohols from C₅ to C₈:





Two complementary experimental setups were implemented using absolute and relative kinetics methods.

The results will help resolving the inconsistencies noted above on the rate coefficients and better understand the atmospheric impacts of such compounds in the context of increasing O_3 background concentrations and climate change. Note that this work represents the first investigation of *trans*-2-octen-1-ol + O_3 kinetics. Further, developing structure-activity relationships (SAR) for atmospheric chemistry models needs consistent and extended kinetic databases to which the present work aims to contribute, especially the effect of the position of the double bond with respect to the $-OH$ functional group. Four different SAR approaches proposed in the literature^{24,25,26,27} were tested against alkenols ozonolysis kinetics, including the present results.

Experimental part

Pseudo first order kinetic set-up (Douai Teflon chamber)

A series of kinetic experiments for *cis*-2-penten-1-ol, *trans*-2-hexen-1-ol, *trans*-3-hexen-1-ol, *cis*-3-hexen-1-ol, 1-octen-3-ol and *trans*-2-octen-1-ol were performed in a Teflon chamber of about 300 L at a temperature of 296 ± 2 K and atmospheric pressure of purified air^{28,29}. This chamber is equipped with inlet and outlet ports used for introducing reactants and taking samples from the reaction mixture. All the connections between the reaction chamber and the instruments were operated via Teflon tubes. Liquid aliquots of alcohols were first vaporized in the chamber under a stream of pure air and the mixture was left to stabilize for ~1h. Preliminary tests were

dedicated to the study of alcohol stability in the reactor. Wall losses of alcohols appeared to be negligible in the present conditions (loss rate < 3% per hour for all alcohols). Ozone was produced separately by an O₃ generator (C-Lasky C-010-DTI) based on electrical discharge on pure zero air flow. After stabilization of the alcohol inside the chamber (1σ of three measurements by Fourier Transform Infrared (FTIR) spectroscopy below 3%), a few tens of mL of O₃ were sampled with a gas syringe and added to the reaction mixture under a high flow of air (about 20 L min⁻¹), thus enhancing the prompt mixing of the reactants and starting the reaction.

Pseudo-first order conditions were set up with alcohol initial concentrations $[alc]_{t_0}$ ((0.34-3.6)×10¹⁴ molecules cm⁻³) in excess (at least 10 times higher) over that of ozone $[O_3]_{t_0}$ ((1.4-17)×10¹² molecules cm⁻³). The alcohol concentration can thus be assumed to be constant during the experiment. Eq. 1 describes the ozone concentration decay taking into account the ozone wall loss rate (k_w in s⁻¹).

$$-\frac{d[O_3]_t}{dt} = k \times [alc]_{t_0} \times [O_3]_t + k_w \times [O_3]_t \quad (\text{Eq. 1})$$

with k the rate coefficient for the reaction between the studied alcohol and O₃ and $[O_3]_t$ is the concentration of ozone at time t . Eq. 1 can be simplified into:

$$-\frac{d[O_3]_t}{dt} = k_{obs} \times [O_3]_t \quad (\text{Eq. 2})$$

where $k_{obs} = k \times [alc]_{t_0} + k_w$ is the pseudo-first-order rate constant. Integrating Eq. 2 yields:

$$\ln\left(\frac{[O_3]_{t_0}}{[O_3]_t}\right) = k_{obs} \times t \quad (\text{Eq. 3})$$

Plotting $\ln\left(\frac{[O_3]_{t_0}}{[O_3]_t}\right)$ vs t leads to a straight line with zero intercept and a regression coefficient $k_{obs} = k \times [alc]_{t_0} + k_w$. Plotting k_{obs} vs $[alc]_{t_0}$ finally gives the rate coefficient k as the regression coefficient and k_w as the intercept.

The ozone decay was monitored by a UV-absorption O₃ analyzer (Environment SA 42M) with a resolution time of 10 s. Preliminary tests showed that the investigated alcohols were not absorbing at the wavelength of the ozone instrument (253.7 nm). Alcohol calibration curves (Figures S1-S6) were carefully determined through FTIR spectroscopy using a 10 m optical length White cell³⁰.


Formation of OH radical through the ozonolysis of hexenols was reported by Atkinson et al.²¹ (OH yield of about 26%). No OH scavenger was used in the Douai Teflon chamber, except for 1-octen-3-ol where two experiments were performed with excess of cyclohexane (200 and 1000 ppm). In these two experiments, about 90% of the OH were scavenged by cyclohexane. It should be stressed, however, that any OH radical formed will react with the alcohol but would not affect the ozone decay in a significant manner at the high initial alcohol/ozone concentration ratios employed. The two experiments noted above were carried out to confirm the absence of effect on the O₃ decay.

Primary products from reactions of unsaturated compounds with O₃ are mainly carbonyls³¹. Potential interferences from these carbonyls in the O₃ analyzer are expected to be negligible since the absorption cross section ratios between ozone and carbonyls at 253.7 nm (detection wavelength of the O₃ analyzer) are in the range of 400 to 4000³². Hence, carbonyl absorption will be negligible compared to that of ozone. Consistently, Peters et al.³³ found no interferences for commercial ozone monitor for a series of VOCs including alcohols and carbonyls.

Concentrations of *cis*-2-penten-1-ol, *trans*-2-hexen-1-ol, *trans*-3-hexen-1-ol, *cis*-3-hexen-1-ol, 1-octen-3-ol and *trans*-2-octen-1-ol were measured in the following IR ranges, respectively: 960-1081 cm⁻¹, 956-1032 cm⁻¹, 934-1000 cm⁻¹, 997-1096 cm⁻¹, 913-940 cm⁻¹, and 900-1050 cm⁻¹.

Relative rate kinetic set-up (Iasi ESC-Q-UAIC reaction chamber)

A series of kinetic experiments for *trans*-2-hexene-1-ol, *trans*-3-hexen-1-ol, *cis*-3-hexen-1-ol, and *trans*-4-hexene-1-ol were performed at 298 K and 1 atm of air pressure, using the facilities of the Environmental Simulation Chamber made out of Quartz from “Alexandru Ioan Cuza” University of Iasi (ESC-Q-UAIC), Romania. The reactor consists of three quartz tubes with a total length of 4.1 m and a total volume of 760 L. The homogeneous mixture of compounds inside the reactor is achieved by two Teflon fans. The reactor can be evacuated down to 10^{-2} mbar. Both ends are closed with anodized aluminum flanges and are provided with multiple inlet and outlet ports. A Bruker Vertex 80 FT-IR spectrometer was used for collecting IR spectra over a White type mirror system with an optical path length of 492 m at a resolution of 1 cm^{-1} . A more detailed description of the reactor can be found in literature³⁴. The relative kinetics method was employed for rate coefficient measurements using *trans*-2-butene and cyclohexene as reference compounds with $k(\text{trans-2-butene} + \text{O}_3) = (2.00 \pm 0.50) \times 10^{-17}\text{ cm}^3\text{ molecule}^{-1}\text{ s}^{-1}$ ³⁵ and $k(\text{cyclohexene} + \text{O}_3) = (7.44 \pm 1.86) \times 10^{-17}\text{ cm}^3\text{ molecule}^{-1}\text{ s}^{-1}$ ³⁶. 1,3,5-Trimethylbenzene was used as OH scavenger in concentrations such that more than 90% of any OH radicals produced was suppressed during the ozonolysis of unsaturated compounds. For *trans*-3-hexene-1-ol only, two other additional scavengers were employed: cyclohexane and di-butyl ether. The reason was to check whether or not the nature of the scavengers plays any role in this study. Preliminary tests to measure wall depositions of *trans*-2-hexene-1-ol, *trans*-3-hexen-1-ol, *cis*-3-hexen-1-ol, and *trans*-4-hexene-1-ol revealed the following wall loss rates (k_{WL}): $(4.93 \pm 0.45) \times 10^{-5}\text{ s}^{-1}$, $(4.35 \pm 0.50) \times 10^{-5}\text{ s}^{-1}$, $(5.08 \pm 0.51) \times 10^{-5}\text{ s}^{-1}$ and $(6.04 \pm 0.53) \times 10^{-5}\text{ s}^{-1}$, respectively. No wall deposition was observed for the reference compounds. In the gas-phase system, besides the reaction of alkenols with ozone, reference compounds would also react with ozone (reactions (8) and (9)):





For the unsaturated alcohols and reference compounds the following kinetic equations are valid in the relative method approach:

$$-\frac{d[alc]_t}{dt} = k \times [alc]_t \times [O_3]_t + k_{WL} \times [alc]_t \quad (\text{Eq. 4})$$

$$-\frac{d[reference]_t}{dt} = k_{ref} \times [reference]_t \times [O_3]_t \quad (\text{Eq. 5})$$

where $[alc]_t$ and $[reference]_t$ are the concentrations of the alcohols and reference compounds at time t . Integrating and combining these two equations yields the following relation between the alcohols and reference concentrations and the O_3 reaction rate coefficients:

$$\ln \frac{[alc]_{t_0}}{[alc]_t} - k_{WL} \times (t - t_0) = \frac{k}{k_{ref}} \ln \frac{[reference]_{t_0}}{[reference]_t} \quad (\text{Eq. 6})$$

where $[alc]_{t_0}$ and $[reference]_{t_0}$ are the initial concentrations of unsaturated alcohols and reference compounds, k and k_{ref} are the rate coefficients of the reaction of alcohols and reference compounds with ozone.

The initial concentrations were estimated based on the weighted amounts injected into the reaction chamber, as the difference of the syringe weights with the liquid compound and without liquid compound measured right after the injection, leading to the following average values (in molecules cm^{-3} , where the uncertainties are two standard deviations): $(7.37 \pm 0.39) \times 10^{13}$ for alcohols, $(8.12 \pm 1.62) \times 10^{13}$ for *trans*-2-butene, $(4.69 \pm 0.78) \times 10^{13}$ for cyclohexene. Regarding OH scavengers, their concentrations (in molecules cm^{-3}) were calculated based on the volume injected: 2.30×10^{15} for 1,3,5-trimethylbenzene, 3.00×10^{16} for cyclohexane and 5.60×10^{15} for di-butyl ether. The OH scavenger concentration was estimated to suppress up to 90% of OH radicals considering

the initial concentrations of the alcohols and reference compounds and their reactivity toward OH radicals³⁷.

The target alcohols, references and OH scavengers were obtained commercially with the following purities: *cis*-2-penten-1-ol (Aldrich, > 96%), *trans*-2-hexen-1-ol (Aldrich, ≥ 95%), *trans*-3-hexen-1-ol (Aldrich, 97%), *cis*-3-hexen-1-ol (Fluka, > 98%), 1-octen-3-ol (Aldrich, ≥ 98%), *trans*-2-octen-1-ol (Aldrich, 97%), *trans*-4-hexen-1-ol (Aldrich, 97%), *trans*-2-butene (Aldrich, 99%), cyclohexene (Aldrich, 99%), 1,3,5-trimethylbenzene (Aldrich, 98%), cyclohexane (Carl-Roth, > 99.5%), di-butyl ether (Aldrich, 99.3%). They were used as received without further purification. At Iasi ESC-Q-UAIC, synthetic air (Messer-Griesheim, 99.999%) and oxygen (Messer-Griesheim, 99.999%) were used while ozone was produced photochemically by passing a flow of O₂ over a VUV-Hg lamp mounted inside a tube and connected to the reactor. Extra pure zero air was produced in Douai with pure air generator (Claind AZ 2020) with relative humidity RH < 2 ppm, CO and CO₂ < 80 ppb.

Results and discussion

Rate coefficients

An example of pseudo-first order kinetic plots representing k_{obs} as a function of $[alc]_{t_0}$ is displayed in Figure 1 for *trans*-2-octen-1-ol. The kinetic plots for the other 5 alkenols are provided in Supporting Information (Figures S7-S11). All plots show very good linearity with close to zero intercepts. The ozone rate coefficients obtained from the slope of these plots are (in 10⁻¹⁷ cm³ molecule⁻¹ s⁻¹): 11.0±2.0 for *cis*-2-penten-1-ol, 12.0±2.0 for *trans*-2-hexen-1-ol, 6.4±1.0 for *trans*-3-hexen-1-ol, 5.8±0.9 for *cis*-3-hexen-1-ol, 2.0±0.3 for 1-octen-3-ol and 8.4±1.3 for *trans*-2-octen-1-ol. Quoted uncertainties are two standard deviations and represent the statistical error on the ozone decays (about 5%) combined with the uncertainty coming from alcohol concentrations (of

about 15%, both statistical and systematic errors). The final uncertainty on k , $\Delta(k)$, was calculated using the error propagation method through Eq. 7 and is about 16% for each rate coefficient.

$$\frac{\Delta(k)}{k} = \sqrt{\left(\frac{\Delta k_{obs}}{k_{obs}}\right)^2 + \left(\frac{\Delta[alc]_{t_0}}{[alc]_{t_0}}\right)^2} \quad (\text{Eq. 7})$$

Studies performed in Iasi by using the relative kinetic method measured simultaneously the reference decay and the degradation of hexenols during the ozonolysis. The total measured decay of hexenols during the experiments was counted from 20% to 35% (about 48% for O_3) from which (91.0 \pm 3.3)% of *trans*-2-hexen-1-ol, (91.5 \pm 7.8)% of *trans*-3-hexen-1-ol, (83.4 \pm 4.6)% of *cis*-3-hexen-1-ol and (91.4 \pm 1.8)% of *trans*-4-hexen-1-ol reacted with ozone. Fairly linear distributions of the data points on the kinetic slopes can be observed in Figure 2 (*trans*-2-hexen-1-ol) and Figures S12-S14 (*trans*-3-hexen-1-ol, *cis*-3-hexen-1-ol, and *trans*-4-hexen-1-ol) in Supplementary Information. Table 1 reports the experimental results obtained from the relative kinetic investigations. The ratios between rate coefficients of hexenols and reference compounds (k/k_{ref}) and the attributed uncertainties (2σ) were retrieved from linear regression analysis. Uncertainties for k values were obtained by propagating 10% error from the k_{ref} values and 2σ from the slope of our experimental data. The value of k (average) and its associated uncertainty is derived from the weighted average for k values. The control test experiment (relative kinetics between *trans*-2-butene and cyclohexene) presented in Supplementary Information (Figure S15) shows a maximum of 5% difference between the measured and literature rate coefficients ratios ($k_{cyclohexene+O_3}/k_{trans-2-butene+O_3}$), this slight difference being covered by the proposed uncertainties. One could notice the very good agreement between the O_3 rate coefficients obtained for *trans*-3-hexen-1-ol when three different OH radical scavengers were used. These results proved that the effect of OH radical could be suppressed by using any of the scavengers.

Table 2 compares the rate coefficients obtained in the present work (absolute and relative methods) with literature data, where available. The rate coefficient obtained for *cis*-2-penten-1-ol ($(1.1 \pm 0.2) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) is in excellent agreement with the determinations of O'Dwyer et al.¹⁴ and Kalalian et al.¹⁷ who found $(1.15 \pm 0.07) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ and $(1.05 \pm 0.20) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, respectively, while about 33% lower than the study from Grosjean and Grosjean¹⁶ of $(1.69 \pm 0.25) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. All previous literature studies used the absolute kinetics method except Kalalian et al.¹⁷, who used the relative method. Grosjean and Grosjean¹⁶ carried out only three experiments under 55% relative humidity using cyclohexane as scavenger with no variation of the initial alkenol concentration, leading to probably less reliable results. A higher rate coefficient value may also point to unknown secondary losses such as impurities in *cis*-2-penten-1-ol. Yet, with a stated purity of 99%,¹⁶ the 1% impurity should have reacted with O₃ at a fairly high rate coefficient of about $5 \times 10^{-15} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ in order to yield the 30% higher *cis*-2-penten-1-ol + O₃ rate coefficient measured by Grosjean and Grosjean¹⁶. Reactions of Criegee Intermediates (CI) with alkenols may also be evoked as potential additional loss of alkenols. In the absence of kinetics data, an alkenol + CI rate coefficient of about $1 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ can be estimated based on the reactivity of CI with alkenes³⁸ and alcohols³⁹. The comparison of the rate of alkenol + O₃ ($r_{\text{O}_3} = k \times [\text{O}_3] \times [\text{alc}]$, with $k = 1.1 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ in the case of *cis*-2-penten-1-ol and $[\text{O}_3] = 1 \times 10^{12} \text{ molecules cm}^{-3}$) with the estimated rate of alkenol + CI ($r_{\text{CI}} = k_{\text{CI}} \times [\text{CI}] \times [\text{alc}]$ with $k_{\text{CI}} = 1 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ and $[\text{CI}]$ estimated at $1 \times 10^6 \text{ molecules cm}^{-3}$) leads to a ratio of about 1000. This value indicates that alkenol reactions with CI are probably not significant in the present chamber experiments. Hence, the possible discrepancies in the rate constants between the present work and literature values cannot be explained by CI reactions with alkenols.

For *trans*-2-hexen-1-ol, a rate coefficient of $(1.2 \pm 0.2) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ was obtained by the absolute kinetic method and $(1.27 \pm 0.11) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ by the relative rate kinetic approach. Both investigations are virtually similar and are in good agreement with the last study by Lin et al.¹⁹ $((1.66 \pm 0.22) \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1})$ while significantly larger (about a factor of two) than Gibilisco et al.¹⁸ who found $(5.98 \pm 0.73) \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. However, in this latter study, it was mentioned that OH radicals were not sufficiently scavenged and the results should be considered with caution. Gibilisco et al.¹⁸ corrected the obtained rate coefficients by inclusion of self-canceling terms of $-k_{\text{hexenol}}[\text{OH}]dt$ and $-k_{\text{reference}}[\text{OH}]dt$ based on the assumption that references have identical rate coefficients against ozone and similar OH radical yields of 26% as reported for isoprene and *trans*-3-hexen-1-ol.

For *trans*-3-hexen-1-ol and *cis*-3-hexen-1-ol where the double bond is further from the -OH group compared to *trans*-2-hexen-1-ol, lower rate coefficients are obtained in the present work, $(6.4 \pm 1.0) \times 10^{-17}$ and $(5.8 \pm 0.9) \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, respectively, when employing the pseudo-first order kinetic method and $(5.01 \pm 0.30) \times 10^{-17}$ and $(4.13 \pm 0.34) \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ when using relative rate kinetics. Experimental evidence for other unsaturated alcohols^{14,40} as well as quantum chemistry calculations¹⁹ supports a higher rate coefficient when the double bond is in α position with respect to the -OH functional group, as is the case with *trans*-2-hexen-1-ol. It should be noted that both 3-hexen-1-ol isomers (*cis*- and *trans*-) display a relatively similar reactivity towards O₃, while literature reviews on alkene + O₃ kinetics indicate that *trans*- isomers are generally more reactive than *cis*- due to a higher pre-exponential factor^{41,42}. Consistently, a slightly higher rate coefficient value can be observed for the *trans*- isomer in comparison with the *cis*- isomer, though this conclusion could be encompassed if the uncertainties level is considered. It is important to mention that rate coefficient values obtained for 3-hexen-1-ol isomers agree within

30% with the literature values obtained by using different methods and simulation chambers as those used in present investigations (see Table 2). However, most of the difference could be explained considering the assumed errors of about 20% of the reference rate coefficients used in the relative rate method. As for *cis*-2-penten-1-ol, only three experiments were performed by Grosjean et al.²⁰ for *cis*-3-hexen-1-ol with no variation of the initial alkenol concentration. This may explain the discrepancies observed for $k_{cis-3-hexen-1-ol+O_3}$ between Grosjean et al.²⁰, on the one hand, and the other studies (including the present work), on the other hand.

The kinetic values for *trans*-4-hexen-1-ol (in $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) are in good agreement for both reference compounds ($(14.1 \pm 1.7) \times 10^{-17}$ with *trans*-2-butene and $(13.9 \pm 1.8) \times 10^{-17}$ with cyclohexene) and with Lin et al.¹⁹ ($(10.5 \pm 1.4) \times 10^{-17}$) considering the uncertainties. In an attempt to compare the rate coefficient of *trans*-2-hexen-1-ol, *trans*-3-hexen-1-ol and *trans*-4-hexen-1-ol, a virtually similar reactivity for *trans*-2-hexen-1-ol and *trans*-4-hexen-1-ol is observed while the rate coefficient for *trans*-3-hexen-1-ol is about three times lower.

For 1-octen-3-ol, our value is in excellent agreement with the only literature data from Li et al.²³. Finally, the present work reports the first O_3 rate coefficient for *trans*-2-octen-1-ol.

As observed in many studies on alkenes + O_3 reaction kinetics^{43,44}, a higher degree of alkyl substitution of the double bond generally leads to a higher rate coefficient. Considering all rate coefficients known for alkenols + O_3 (Table S1) suggests that it is also the case for alkenols. Note that the highest k value is for 3-methyl-2-buten-1-ol ($> 4.39 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$) which is a tri-substituted compound. This phenomenon has been explained by an inductive effect produced via the enhancement of the π -orbital electron density for higher substituted compounds, leading to a lowering of the activation energy.

In the case of unsaturated compounds, the hydroxyl function group is expected to induce an electron-donating effect on the adjacent bonds, thus enhancing the electronic density of the double bond and increasing the reactivity of the unsaturated compound. Comparing the O₃ reactivity of alkenols with that of the corresponding alkenes as a function of the distance between –OH and C=C displays a puzzling picture, as shown in Figure 3. While for 1-alken-3-ols, the rate coefficient ratio $k(\text{O}_3+\text{alkenol})/k(\text{O}_3+\text{alkene})$ is higher than 1, as expected, the reverse is observed for all other alkenols. The highest ratio of about 2 is calculated for 1-octen-3-ol/1-octene and 1-nonen-3-ol/1-nonene. Mason et al. showed that C₃ to C₁₀ 1-alkenes all react with O₃ with rate coefficients around $1 \times 10^{-17} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ without any effect of the carbon chain length⁴⁵. Interestingly, the present results indicate that the substitution by OH at the α -position to the C=C bond in 1-alkenes increases the rate coefficient up to a factor two. In addition to the electron-donating effect of the –OH group, this may be due to the formation of a strong pre-reactive complex through hydrogen bond interaction between O₃ and –OH¹⁹. Noteworthy, 2-alken-1-ols like 2-penten-1-ol (which are α -alkenols) do not show such reactivity increase between alkenes and alkenols, probably due to steric effects. Further experimental and theoretical investigations of reaction mechanisms of α -alkenols + O₃ may help understanding their reactivity.

Structure-Activity Relationships (SAR)

Rate coefficients can be estimated using different methods involving structure reactivity relationships. Atkinson and Carter²⁴ applied a methodology used by the US Environmental Protection Agency under the AOPWIN (Atmospheric Oxidation Program for Windows) to estimate ozone rate coefficients. Calvert et al.²⁵ developed SAR methods using the group-additivity approach and factors contributions. Pfrang et al.²⁶ used correlations and rate coefficients for basic structures multiplied by group reactivity factors for corresponding substituents. More

recently, a new SAR method was developed implying the contribution of inductive effect of heteroatoms and total steric effects based on the earlier work of McGillen et al.⁴⁶ on alkenes and extended by McGillen et al.²⁷ to heteroatomic unsaturated VOCs.

The different SAR approaches cited above were applied to evaluate their suitability as alternative tools for rate coefficient estimation for the reactions of unsaturated alcohols with ozone. Where applicable, the experimental values were calculated by averaging rate coefficients obtained in the present work by the pseudo-first order approach and the relative rate kinetic method. In Table 3, the experimental k values (k_{exp}) can be compared with the calculated values obtained using the different SAR approaches (k_{SAR}) on the investigated compounds. The earliest methodologies proposed in the literature on SAR estimations of rate coefficients show large discrepancies with the experimental results. Poor agreement was also noticed when using the SAR approach from Pfrang et al.²⁶ (except for 3-hexen-1-ol isomers) despite the more complex methodology involved. Using McGillen et al.²⁷ methodology (details are provided in the Supplementary Information), a very satisfactory estimate is obtained for *cis*-2-penten-1-ol, *trans*-2-hexen-1-ol and *trans*-2-octen-1-ol, with agreement with our results between 2% and 20%. For the others alkenols, the ratios $k_{\text{SAR}}/k_{\text{exp}}$ vary between 0.29 and 0.59. The strongest disagreement is obtained for *trans*-4-hexen-1-ol ($k_{\text{SAR}}/k_{\text{exp}} = 0.29$) for which the experimental finding is three times larger than the calculated rate coefficient. Because of the distance between –OH and C=C in *trans*-4-hexen-1-ol (–OH is in δ position towards C=C), no inductive effect is considered in the SAR calculation. Consistently, the experimental rate coefficient of the corresponding alkene (*trans*-2-hexene) with O₃ ($1.53 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)³⁷ is close to that obtained in the present work for *trans*-4-hexen-1-ol ($1.40 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$). Therefore, the low $k_{\text{SAR}}/k_{\text{exp}}$ ratio reflects a weakness in the SAR method, probably regarding a change in the reaction mechanisms as suggested by McGillen et al.²⁷. Based

on the overall agreement, it is suggested that the SAR method from McGillen et al.²⁷ estimates the rate coefficients from ozonolysis of unsaturated alcohols much closer to the experimental data than the other above-mentioned approaches.

Ozonolysis rate coefficients were also computed for a range of other alkenols using the SAR method from McGillen et al.²⁷. Values are reported in Figure 4 as a function of experimental ones (see Table S1). Besides the compounds already included in McGillen et al.²⁷ analysis (with updated values from Kalalian et al.¹⁷ and the present work), fourteen new compounds were added (see Table S1, compounds in italic). It can be seen that calculated rate coefficients are within a factor of 2 of the experimental rate coefficients. A linear regression (forced through zero) on the data yields a slope coefficient of 0.64 ± 0.11 (2σ uncertainty), indicating that SAR calculated rate coefficients are mostly lower than experimental ones. As stressed above, this may suggest unknown changes in reaction mechanisms of alkenol ozonolysis. Additional experiments are clearly needed to better characterize these mechanisms and improve structure activity relationships.

Atmospheric implications

The rate coefficients determined in the present work allow calculating the lifetime τ of alkenols in the atmosphere when reacting with O_3 , using $\tau = 1/(k \times [O_3])$. Assuming an average ozone concentration of 1.6×10^{12} molecules cm^{-3} (about 65 ppb at 295 K and 1 atm)⁴⁷ leads to lifetimes of a few hours (Table 4). The corresponding lifetimes when reacting with OH, NO_3 and Cl can be similarly calculated and are reported also in Table 4. Comparing with OH, NO_3 and Cl reactions indicates that ozonolysis is a competitive oxidation process, especially during the night and at low NO_x environments where OH and NO_3 concentrations are low, respectively. In addition, in the

perspective of increasing ozone concentrations in the future⁴⁸, an increasing impact of ozonolysis may be expected in the degradation of alkenols.

Conclusions

The rate coefficients of a series of seven unsaturated alcohols with ozone have been determined using two complementary setups, allowing both absolute and relative kinetics methods to be employed. Good agreement has been obtained between the two methods. Alkenols appear to have high reactivity with ozone with lifetime in the hour range. Rate coefficients show a strong and complex dependence on the structure of the alkenol, particularly the relative position of the OH group towards the C=C double bond. Four SAR approaches were tested against the investigated compounds, the most recent from McGillen et al.²⁷ displaying the best results. Yet, efforts are still needed to better understand the reaction mechanisms and unravel the links between the structure and the reactivity of alkenols with O₃.

Supporting Information

Details for evaluating SAR rate coefficients using McGillen et al.²⁷ methodology, ambient temperature ozonolysis rate coefficients of alkenols (Table S1), IR calibration curves for unsaturated alcohols with the corresponding integrated band intensities (IBI, Figures S1-S6), kinetic plots from the absolute and relative rate methods (Figures S7-S14), relative kinetic plot of ozone reaction with *trans*-2-butene vs cyclohexene, (data obtained from control experiments in ESC-Q-UAIC chamber; 1,3,5-trimethylbenzene was used as OH scavenger) (Figure S15).

Acknowledgements

The Center for Energy and Environment acknowledges funding by the French ANR Agency under contract No. ANR-11-LabX-0005-01 CaPPA (Chemical and Physical Properties of the Atmosphere), the Région Hauts-de-France, the Ministère de l'Enseignement Supérieur et de la

Recherche (CPER Climibio), and the European Fund for Regional Economic Development. AG is grateful for a PhD grant from Brittany Region and IMT Nord Europe. AG, GED, AC and AT are grateful to the INSU-LEFE-CHAT program for funding this research. CAr, IGB, RIO and AT acknowledge the financial support from European Union's Horizon 2020 Research and Innovation Framework Program, through the EUROCHAMP-2020 Infrastructure Activity Grant (grant agreement No 730997). CAr, IGB, RIO, CAm and CR and acknowledge the Operational Program Competitiveness 2014-2020, Axis 1, under POC / 448 / 1 / 1 Research infrastructure projects for public R&D institutions / Sections F 2018, through the Research Center with Integrated Techniques for Atmospheric Aerosol Investigation in Romania (RECENT AIR) project, under grant agreement MySMIS no. 127324 for its co-support. All the authors are thankful to the PHC Brancusi program for funding the OzOA project (38385UD).

References

- (1) Reich, P. B. Quantifying Plant Response to Ozone: A Unifying Theory. *Tree Physiol.* **1987**, 3 (1), 63–91.
- (2) Scebba, F.; Giuntini, D.; Castagna, A.; Soldatini, G.; Ranieri, A. Analysing the Impact of Ozone on Biochemical and Physiological Variables in Plant Species Belonging to Natural Ecosystems. *Environ. Exp. Bot.* **2006**, 57 (1–2), 89–97.
- (3) Finlayson-Pitts, B. J.; Pitts, J. N. Atmospheric Chemistry of Tropospheric Ozone Formation: Scientific and Regulatory Implications. *Air Waste* **1993**, 43 (8), 1091–1100.
- (4) Bortz, S. E.; Prather, M. J.; Cammas, J.-P.; Thouret, V.; Smit, H. Ozone, Water Vapor, and Temperature in the Upper Tropical Troposphere: Variations over a Decade of MOZAIC Measurements. *J. Geophys. Res.* **2006**, 111 (D5), D05305. .
- (5) Sicard, P.; Serra, R.; Rossello, P. Spatiotemporal Trends in Ground-Level Ozone Concentrations and Metrics in France over the Time Period 1999–2012. *Environ. Res.* **2016**, 149, 122–144.
- (6) Jenkin, M. E. Trends in Ozone Concentration Distributions in the UK since 1990: Local, Regional and Global Influences. *Atmos. Environ.* **2008**, 42 (21), 5434–5445.
- (7) Cooper, O. R.; Schultz, M. G.; Schröder, S.; Chang, K.-L.; Gaudel, A.; Benítez, G. C.; Cuevas, E.; Fröhlich, M.; Galbally, I. E.; Molloy, S. et al., Multi-Decadal Surface Ozone Trends at Globally Distributed Remote Locations. *Elem. Sci. Anthr.* **2020**, 8, 23.
- (8) Kleist, E.; Mentel, T. F.; Andres, S.; Bohne, A.; Folkers, A.; Kiendler-Scharr, A.; Rudich, Y.; Springer, M.; Tillmann, R.; Wildt, J. Irreversible Impacts of Heat on the Emissions of Monoterpenes, Sesquiterpenes, Phenolic BVOC and Green Leaf Volatiles from Several Tree Species. *Biogeosciences* **2012**, 9 (12), 5111–5123.

- (9) Heiden, A. C.; Kobel, K.; Langebartels, C.; Schuh-Thomas, G.; Wildt, J. Emissions of Oxygenated Volatile Organic Compounds from Plants Part I: Emissions from Lipoxygenase Activity. *J. Atmospheric Chem.* **2003**, *45* (2), 143–172.
- (10) Calvert, J. G.; Mellouki, A.; Orlando, J. J.; Pilling, M. J.; Wallington, T. J. *Mechanisms of Atmospheric Oxidation of the Oxygenates*; Oxford University Press, 2011.
- (11) Atkinson, R. Gas-Phase Tropospheric Chemistry of Volatile Organic Compounds: 1. Alkanes and Alkenes. *J. Phys. Chem. Ref. Data* **1997**, *26* (2), 215–290.
- (12) Horie, O.; Moortgat, G. K. Gas-Phase Ozonolysis of Alkenes. Recent Advances in Mechanistic Investigations. *Acc. Chem. Res.* **1998**, *31* (7), 387–396.
- (13) Hellpointner, E.; Gäb, S. Detection of Methyl, Hydroxymethyl and Hydroxyethyl Hydroperoxides in Air and Precipitation. *Nature* **1989**, *337* (6208), 631–634.
- (14) O Dwyer, M. A.; Carey, T. J.; Healy, R. M.; Wenger, J. C.; Picquet-Varrault, B.; Doussin, J. F. The Gas-Phase Ozonolysis of 1-Penten-3-Ol, (Z)-2-Penten-1-Ol and 1-Penten-3-One: Kinetics, Products and Secondary Organic Aerosol Formation. *Z. Für Phys. Chem.* **2010**, *224* (7–8), 1059–1080.
- (15) Graedel, T. E. Terpenoids in the Atmosphere. *Rev. Geophys.* **1979**, *17* (5), 937.
- (16) Grosjean, E.; Grosjean, D. Rate Constants for the Gas-Phase Reactions of Ozone with Unsaturated Aliphatic Alcohols. *Int. J. Chem. Kinet.* **1994**, *26* (12), 1185–1191.
- (17) Kalalian, C.; El Dib, G.; Singh, H. J.; Rao, P. K.; Roth, E.; Chakir, A. Temperature Dependent Kinetic Study of the Gas Phase Reaction of Ozone with 1-Penten-3-Ol, Cis-2-Penten-1-Ol and Trans-3-Hexen-1-Ol: Experimental and Theoretical Data. *Atmos. Environ.* **2020**, *223* (October 2019), 117306.
- (18) Gibilisco, R. G.; Bejan, I.; Barnes, I.; Wiesen, P.; Teruel, M. A. FTIR Gas Kinetic Study of the Reactions of Ozone with a Series of Hexenols at Atmospheric Pressure and 298K. *Chem. Phys. Lett.* **2015**, *618*, 114–118.
- (19) Lin, X.; Ma, Q.; Yang, C.; Tang, X.; Zhao, W.; Hu, C.; Gu, X.; Fang, B.; Gai, Y.; Zhang, W. Kinetics and Mechanisms of Gas Phase Reactions of Hexenols with Ozone. *RSC Adv.* **2016**, *6* (87), 83573–83580.
- (20) Grosjean, D.; Grosjean, E.; Williams, E. L. Rate Constants for the Gas-Phase Reactions of Ozone with Unsaturated Alcohols, Esters, and Carbonyls. *Int. J. Chem. Kinet.* **1993**, *25* (9), 783–794.
- (21) Atkinson, R.; Arey, J.; Aschmann, S. M.; Corchnoy, S. B.; Shu, Y. Rate Constants for the Gas-Phase Reactions of Cis-3-Hexen-1-Ol, Cis-3-Hexenylacetate, Trans-2-Hexenal, and Linalool with OH and NO₃ Radicals and O₃ at 296 ± 2 K, and OH Radical Formation Yields from the O₃ Reactions. *Int. J. Chem. Kinet.* **1995**, *27* (10), 941–955.
- (22) Chen, Y.; Wang, J.; Zhao, S.; Tong, S.; Ge, M. An Experimental Kinetic Study and Products Research of the Reactions of O₃ with a Series of Unsaturated Alcohols. *Atmos. Environ.* **2016**, *145*, 455–467.
- (23) Li, W.; Chen, Y.; Tong, S.; Guo, Y.; Zhang, Y.; Ge, M. Kinetic Study of the Gas-Phase Reaction of O₃ with Three Unsaturated Alcohols. *J. Environ. Sci.* **2018**, *71*, 292–299.
- (24) Atkinson, R.; Carter, W. P. L. Kinetics and Mechanisms of the Gas-Phase Reactions of Ozone with Organic Compounds under Atmospheric Conditions. *Chem. Rev.* **1984**, *84* (5), 437–470.
- (25) Calvert, J. G.; Atkinson, R.; Kerr, J. A.; Madronich, S.; Moortgat, G. K.; Wallington, T. J.; Yarwood, G. *The Mechanisms of Atmospheric Oxidation of the Alkenes*; Oxford University Press, 2000.

- (26) Pfrang, C.; King, M. D.; Braeckvelt, M.; Canosa-Mas, C. E.; Wayne, R. P. Gas-Phase Rate Coefficients for Reactions of NO₃, OH, O₃ and O(³P) with Unsaturated Alcohols and Ethers: Correlations and Structure-Activity Relations (SARs). *Atmos. Environ.* **2008**, *42* (13), 3018–3034.
- (27) McGillen, M. R.; Archibald, A. T.; Carey, T.; Leather, K. E.; Shallcross, D. E.; Wenger, J. C.; Percival, C. J. Structure–Activity Relationship (SAR) for the Prediction of Gas-Phase Ozonolysis Rate Coefficients: An Extension towards Heteroatomic Unsaturated Species. *Phys Chem Chem Phys* **2011**, *13* (7), 2842–2849.
- (28) Szabo, E.; Tarmoul, J.; Tomas, A.; Fittschen, C.; Dobe, S.; Coddeville, P. Kinetics of The •OH-Radical Initiated Reactions of Acetic Acid and Its Deuterated Isomers. **2009**, *96* (2), 299–309.
- (29) Crunaire, S.; Tarmoul, J.; Fittschen, C.; Tomas, A.; Lemoine, B.; Coddeville, P. Use of Cw-CRDS for Studying the Atmospheric Oxidation of Acetic Acid in a Simulation Chamber. *Appl. Phys. B* **2006**, *85* (2–3), 467–476.
- (30) Bouzidi, H.; Djehiche, M.; Coddeville, P.; Fittschen, C.; Tomas, A. Atmospheric Chemistry of α-Diketones: Kinetics of C5 and C6 Compounds with Cl Atoms and OH Radicals. *Int. J. Chem. Kinet.* **2017**, *49* (2), 112–118.
- (31) Johnson, D.; Marston, G. The Gas-Phase Ozonolysis of Unsaturated Volatile Organic Compounds in the Troposphere. *Chem. Soc. Rev.* **2008**, *37* (4), 699.
- (32) Turnipseed, A. A.; Andersen, P. C.; Williford, C. J.; Ennis, C. A.; Birks, J. W. Use of a Heated Graphite Scrubber as a Means of Reducing Interferences in UV-Absorbance Measurements of Atmospheric Ozone. *Atmospheric Meas. Tech.* **2017**, *10* (6), 2253–2269.
- (33) Peters, S.; Bejan, I.; Kurtenbach, R.; Liedtke, S.; Villena, G.; Wiesen, P.; Kleffmann, J. Development of a New LOPAP Instrument for the Detection of O₃ in the Atmosphere. *Atmos. Environ.* **2013**, *67*, 112–119.
- (34) Roman, C.; Arsene, C.; Bejan, I. G.; Olariu, R. I. Investigations into the Gas-Phase Photolysis and OH Radical Kinetics of Nitrocatechols: Implications of Intramolecular Interactions on Their Atmospheric Behaviour. *Atmospheric Chem. Phys.* **2022**, *22* (4), 2203–2219.
- (35) McGillen, M. R.; Carter, W. P. L.; Mellouki, A.; Orlando, J. J.; Picquet-Varrault, B.; Wallington, T. J. Database for the Kinetics of the Gas-Phase Atmospheric Reactions of Organic Compounds. *Earth Syst. Sci. Data* **2020**, *12* (2), 1203–1216.
- (36) Treacy, J.; Curley, M.; Wenger, J.; Sidebottom, H. Determination of Arrhenius Parameters for the Reactions of Ozone with Cycloalkenes. *J. Chem. Soc. - Faraday Trans.* **1997**, *93* (16), 2877–2881.
- (37) Calvert, J. G.; Orlando, J. J.; Stockwell, W. R.; Wallington, T. J. *The Mechanisms of Reactions Influencing Atmospheric Ozone*; Oxford University Press: Oxford, UK, 2015.
- (38) Buras, Z. J.; Elsamra, R. M. I.; Jalan, A.; Middaugh, J. E.; Green, W. H. Direct Kinetic Measurements of Reactions between the Simplest Criegee Intermediate CH₂OO and Alkenes. *J. Phys. Chem. A* **2014**, *118*, 1997–2006.
- (39) McGillen, M. R.; Curchod, B. F. E.; Chhantyal-Pun, R.; Beames, J. M.; Watson, N.; Khan, M. A. H.; McMahon, L.; Shallcross, D. E.; Orr-Ewing, A. J. Criegee Intermediate–Alcohol Reactions, A Potential Source of Functionalized Hydroperoxides in the Atmosphere. *ACS Earth Space Chem.* **2017**, *1* (10), 664–672.
- (40) Gai, Y.; Ge, M.; Wang, W. Kinetics of the Gas-Phase Reactions of Some Unsaturated Alcohols with Cl Atoms and O₃. *Atmos. Environ.* **2011**, *45* (1), 53–59.

- (41) Treacy, J.; Hag, M. El; O'Farrell, D.; Sidebottom, H. Reactions of Ozone with Unsaturated Organic Compounds. *Berichte Bunsenges. Für Phys. Chem.* **1992**, *96* (3), 422–427.
- (42) Grosjean, E.; Grosjean, D. Rate Constants for the Gas-Phase Reaction of Ozone with 1,2-Disubstituted Alkenes. *Int. J. Chem. Kinet.* **1996**, *28* (6), 461–466.
- (43) Avzianova, E. V.; Ariya, P. A. Temperature-Dependent Kinetic Study for Ozonolysis of Selected Tropospheric Alkenes. *Int. J. Chem. Kinet.* **2002**, *34* (12), 678–684.
- (44) Leather, K. E.; McGillen, M. R.; Percival, C. J. Temperature-Dependent Ozonolysis Kinetics of Selected Alkenes in the Gas Phase: An Experimental and Structure–Activity Relationship (SAR) Study. *Phys. Chem. Chem. Phys.* **2010**, *12* (12), 2935.
- (45) Mason, S. A.; Arey, J.; Atkinson, R. Rate Constants for the Gas-Phase Reactions of NO₃ Radicals and O₃ with C₆–C₁₄ 1-Alkenes and 2-Methyl-1-Alkenes at 296 ± 2 K. *J. Phys. Chem. A* **2009**, *113* (19), 5649–5656.
- (46) McGillen, M. R.; Carey, T. J.; Archibald, A. T.; Wenger, J. C.; Shallcross, D. E.; Percival, C. J. Structure-Activity Relationship (SAR) for the Gas-Phase Ozonolysis of Aliphatic Alkenes and Dialkenes. *Phys. Chem. Chem. Phys.* **2008**, *10* (13), 1757–1768.
- (47) Dayan, U.; Ricaud, P.; Zbinden, R.; Dulac, F. Atmospheric Pollution over the Eastern Mediterranean during Summer – a Review. *Atmospheric Chem. Phys.* **2017**, *17* (21), 13233–13263.
- (48) Lin, M.; Horowitz, L. W.; Payton, R.; Fiore, A. M.; Tonnesen, G. US Surface Ozone Trends and Extremes from 1980 to 2014: Quantifying the Roles of Rising Asian Emissions, Domestic Controls, Wildfires, and Climate. *Atmospheric Chem. Phys.* **2017**, *17*, 2943–2970.
- (49) Pfrang, C.; King, M. D.; Canosa-Mas, C. E.; Wayne, R. P. Structure–Activity Relations (SARs) for Gas-Phase Reactions of NO₃, OH and O₃ with Alkenes: An Update. *Atmos. Environ.* **2006**, *40* (6), 1180–1186. <https://doi.org/10.1016/j.atmosenv.2005.09.080>.
- (50) Hein, R.; Crutzen, P. J.; Heimann, M. An Inverse Modeling Approach to Investigate the Global Atmospheric Methane Cycle. *Glob. Biogeochem. Cycles* **1997**, *11* (1), 43–76.
- (51) Shu, Y.; Atkinson, R. Atmospheric Lifetimes and Fates of a Series of Sesquiterpenes. *J. Geophys. Res. Atmospheres* **1995**, *100* (D4), 7275–7281.
- (52) Wingenter, O. W.; Kubo, M. K.; Blake, N. J.; Smith Jr., T. W.; Blake, D. R.; Rowland, F. S. Hydrocarbon and Halocarbon Measurements as Photochemical and Dynamical Indicators of Atmospheric Hydroxyl, Atomic Chlorine, and Vertical Mixing Obtained during Lagrangian Flights. *J. Geophys. Res. Atmospheres* **1996**, *101* (D2), 4331–4340.
- (53) Orlando, J. J.; Tyndall, G. S.; Ceazan, N. Rate Coefficients and Product Yields from Reaction of OH with 1-Penten-3-Ol, (Z)-2-Penten-1-Ol, and Allyl Alcohol (2-Propen-1-Ol). *J. Phys. Chem. A* **2001**, *105* (14), 3564–3569.
- (54) Pfrang, C.; Baeza Romero, M. T.; Cabanas, B.; Canosa-Mas, C. E.; Villanueva, F.; Wayne, R. P. Night-Time Tropospheric Chemistry of the Unsaturated Alcohols (Z)-Pent-2-En-1-Ol and Pent-1-En-3-Ol: Kinetic Studies of Reactions of NO₃ and N₂O₅ with Stress-Induced Plant Emissions. *Atmos. Environ.* **2007**, *41* (8), 1652–1662.
- (55) Zhao, Z.; Husainy, S.; Smith, G. D. Kinetics Studies of the Gas-Phase Reactions of NO₃ Radicals with Series of 1-Alkenes, Dienes, Cycloalkenes, Alkenols, and Alkenals. *J. Phys. Chem. A* **2011**, *115* (44), 12161–12172.
- (56) Grira, A.; Amarandei, C.; Romanias, M. N.; El Dib, G.; Canosa, A.; Arsene, C.; Bejan, I. G.; Olariu, R. I.; Coddeville, P.; Tomas, A. Kinetic Measurements of Cl Atom Reactions with C₅–C₈ Unsaturated Alcohols. *Atmosphere* **2020**, *11* (3), 256.

- (57) Rodríguez, A.; Rodríguez, D.; Garzón, A.; Soto, A.; Aranda, A.; Notario, A. Kinetics and Mechanism of the Atmospheric Reactions of Atomic Chlorine with 1-Penten-3-Ol and (Z)-2-Penten-1-Ol: An Experimental and Theoretical Study. *Phys. Chem. Chem. Phys.* **2010**, *12* (38), 12245–12258.
- (58) Gibilisco, R. G.; Santiago, A. N.; Teruel, M. A. OH-Initiated Degradation of a Series of Hexenols in the Troposphere. Rate Coefficients at 298 K and 1 Atm. *Atmos. Environ.* **2013**, *77*, 358–364.
- (59) Davis, M. E.; Burkholder, J. B. Rate Coefficients for the Gas-Phase Reaction of OH with (Z)-3-Hexen-1-Ol, 1-Penten-3-Ol, (E)-2-Penten-1-Ol, and (E)-2-Hexen-1-Ol between 243 and 404 K. *Atmospheric Chem. Phys.* **2011**, *11* (7), 3347–3358.
- (60) Gibilisco, R. G.; Bejan, I.; Barnes, I.; Wiesen, P.; Teruel, M. A. Rate Coefficients at 298 K and 1 Atm for the Tropospheric Degradation of a Series of C6, C7 and C8 Biogenic Unsaturated Alcohols Initiated by Cl Atoms. *Atmos. Environ.* **2014**, *94*, 564–572.
- (61) Gibilisco, R. G.; Blanco, M. B.; Bejan, I.; Barnes, I.; Wiesen, P.; Teruel, M. A. Atmospheric Sink of (E)-3-Hexen-1-Ol, (Z)-3-Hepten-1-Ol, and (Z)-3-Octen-1-Ol: Rate Coefficients and Mechanisms of the OH-Radical Initiated Degradation. *Environ. Sci. Technol.* **2015**, *49* (13), 7717–7725.
- (62) Peirone, S. A.; Barrera, J. A.; Taccone, R. A.; Cometto, P. M.; Lane, S. I. Relative Rate Coefficient Measurements of OH Radical Reactions with (Z)-2-Hexen-1-Ol and (E)-3-Hexen-1-Ol under Simulated Atmospheric Conditions. *Atmos. Environ.* **2014**, *85*, 92–98.
- (63) Jiménez, E.; Lanza, B.; Antiñolo, M.; Albaladejo, J. Photooxidation of Leaf-Wound Oxygenated Compounds, 1-Penten-3-Ol, (Z)-3-Hexen-1-Ol, and 1-Penten-3-One, Initiated by OH Radicals and Sunlight. *Environ. Sci. Technol.* **2009**, *43* (6), 1831–1837.

Table 1: Experimental results from the relative kinetic measurements employed in the present study for the ozonolysis of a series of alkenols.








Alcohol	OH scavenger	Reference	k/k_{ref}	$k \times 10^{17}$ ($\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	$k \text{ (average)} \times 10^{17}$ ($\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)
<i>trans</i> -2-hexen-1-ol	1,3,5-TMB	<i>trans</i> -2-butene	0.66±0.04	13.1±1.5	12.7±1.1
		cyclohexene	1.65±0.12	12.3±1.5	
<i>trans</i> -3-hexen-1-ol	1,3,5-TMB	<i>trans</i> -2-butene	0.24±0.02	4.84±0.62	5.01±0.30
		cyclohexene	0.74±0.04	5.49±0.62	
	cyclohexane	<i>trans</i> -2-butene	0.24±0.03	4.77±0.70	
		cyclohexene	0.65±0.08	4.86±0.74	
	di-butyl ether	cyclohexene	0.67±0.08	4.98±0.75	
<i>cis</i> -3-hexen-1-ol	1,3,5-TMB	<i>trans</i> -2-butene	0.20±0.01	4.07±0.47	4.13±0.34
		cyclohexene	0.57±0.04	4.21±0.50	
<i>trans</i> -4-hexen-1-ol	1,3,5-TMB	<i>trans</i> -2-butene	0.71±0.05	14.1±1.7	14.0±1.2
		cyclohexene	1.87±0.16	13.9±1.8	

Table 2: Rate coefficients obtained in the present work using the absolute rate (pseudo first order) method (AR) and relative rate method (RR) compared with literature data.

Alcohol	$k \times 10^{17} \text{ (cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}\text{)}^a$	Reference
<i>cis</i> -2-penten-1-ol	11.0±2.0 (AR)	This work
	16.9±2.5 (AR)	Grosjean and Grosjean ¹⁶
	11.5±0.7 (AR)	O'Dwyer et al. ¹⁴
	10.5±2.0 (RR)	Kalalian et al. ¹⁷
<i>trans</i> -2-hexen-1-ol	12.0±2.0 (AR)	This work
	12.7±1.1 (RR)	
	6.0±0.7 (RR)	Gibilisco et al. ¹⁸
	16.6±2.2 (AR)	Lin et al. ¹⁹
<i>trans</i> -3-hexen-1-ol	6.4±1.0 (AR)	This work
	5.01±0.30 (RR)	
	5.8±0.9 (RR)	Gibilisco et al. ¹⁸
	6.2±0.7 (AR)	Lin et al. ¹⁹
	6.2±1.4 (RR)	Kalalian et al. ¹⁷
<i>cis</i> -3-hexen-1-ol	5.8±0.9 (AR)	This work
	4.13±0.34 (RR)	
	10.5±0.7 (AR)	Grosjean et al. ²⁰
	6.4±1.7 (RR)	Atkinson et al. ²¹
	6.0±0.9 (RR)	Gibilisco et al. ¹⁸
	5.5±0.7 (AR)	Lin et al. ¹⁹
	6.05±0.38 (RR)	Chen et al. ²²
	5.78±0.65 (AR)	
<i>trans</i> -4-hexen-1-ol	14.0±1.2 (RR)	This work
	10.5±1.4 (AR)	Lin et al. ¹⁹
1-octen-3-ol	2.0±0.3 (AR)	This work
	1.9±0.2 (AR)	Li et al. ²³

<i>trans</i> -2-octen-1-ol	8.4±1.3 (AR)	This work
a: quoted uncertainties from the present work correspond to 2σ		

Table 3: The comparison of different SAR methods and the ratio of $k_{\text{exp}}/k_{\text{SAR}}$. k_{exp} are the average of the absolute and relative rate coefficients obtained in the present work, where applicable.

Alcohol	k_{exp} (this work) ($\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	k_{SAR} ($\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$)	$k_{\text{SAR}}/k_{\text{exp}}$
 <i>cis</i> -2-penten-1-ol	1.10×10^{-16}	$1.30 \times 10^{-16} \text{ a}$	1.18
		$1.20 \times 10^{-16} \text{ b}$	1.09
		$1.38 \times 10^{-17} \text{ c}$	0.13
		$1.08 \times 10^{-16} \text{ d}$	0.98
 <i>trans</i> -2-hexen-1-ol	1.23×10^{-16}	$2.00 \times 10^{-16} \text{ a}$	1.61
		$1.90 \times 10^{-16} \text{ b}$	1.54
		$1.38 \times 10^{-17} \text{ c}$	0.11
		$0.99 \times 10^{-16} \text{ d}$	0.80
 <i>trans</i> -3-hexen-1-ol	5.71×10^{-17}	$2.00 \times 10^{-16} \text{ a}$	3.57
		$1.90 \times 10^{-16} \text{ b}$	3.33
		$3.65 \times 10^{-17} \text{ c}$	0.64
		$2.80 \times 10^{-17} \text{ d}$	0.49
 <i>cis</i> -3-hexen-1-ol	4.97×10^{-17}	$1.30 \times 10^{-16} \text{ a}$	2.63
		$1.20 \times 10^{-16} \text{ b}$	2.44
		$3.68 \times 10^{-17} \text{ c}$	0.74
		$2.80 \times 10^{-17} \text{ d}$	0.56
 <i>trans</i> -4-hexen-1-ol	1.40×10^{-16}	$2.00 \times 10^{-16} \text{ a}$	1.43
		$1.90 \times 10^{-16} \text{ b}$	1.35
		$5.66 \times 10^{-17} \text{ e}$	0.40
		$4.11 \times 10^{-17} \text{ d}$	0.29
 1-octen-3-ol	2.00×10^{-17}	$1.20 \times 10^{-17} \text{ a}$	0.60
		$1.00 \times 10^{-17} \text{ b}$	0.50
		$3.72 \times 10^{-18} \text{ c}$	0.19
		$1.18 \times 10^{-17} \text{ d}$	0.59
 <i>trans</i> -2-octen-1-ol	8.40×10^{-17}	$2.00 \times 10^{-16} \text{ a}$	2.38
		$1.90 \times 10^{-16} \text{ b}$	2.27

	1.38×10^{-17} ^c	0.16
	0.99×10^{-16} ^d	1.18

^a Atkinson and Carter²⁴, ^b Calvert et al.²⁵, ^c Pfrang et al.²⁶, ^d McGillen et al.²⁷, ^e Pfrang et al.⁴⁹ (for *trans*-4-hexen-1-ol, the OH group being far from the C=C double bond, the SAR from Pfrang et al.²⁶ for alkenols was not used but that developed for alkenes by Pfrang et al.⁴⁹).

Table 4: Comparison of atmospheric lifetimes of unsaturated alcohols with respect to removal by O₃, OH radicals, NO₃ radicals and Cl atoms. Average concentrations used: [O₃] = 1.6×10¹² molecules cm⁻³ ⁴⁷, [OH] = 2×10⁶ radicals cm⁻³ ⁵⁰, [NO₃] = 5×10⁸ radicals cm⁻³ ⁵¹, [Cl] = 10⁴ atoms cm⁻³ ⁵². Values in italic correspond to SAR calculations.

Alcohol	k _{O3} (cm ³ molecule ⁻¹ s ⁻¹) ^a	τ _{O3} (h)	k _{OH} (10 ⁻¹⁰ cm ³ molecule ⁻¹ s ⁻¹)	τ _{OH} (h)	k _{NO3} (10 ⁻¹³ cm ³ molecule ⁻¹ s ⁻¹)	τ _{NO3} (h)	k _{Cl} (10 ⁻¹⁰ cm ³ molecule ⁻¹ s ⁻¹)	τ _{Cl} (h)
<i>cis</i> -2-penten-1-ol	11.0	1.5	1.17 ^b	1.2	2.34 ^c	2.4	3.00 ^d	93
<i>trans</i> -2-hexen-1-ol	12.3	1.4	0.81 ^e	1.7	1.56 ^f	3.6	3.45 ^g	81
<i>trans</i> -3-hexen-1-ol	5.71	3.0	1.05 ^h	1.3	2.67 ^f	2.1	3.24 ^g	86
<i>cis</i> -3-hexen-1-ol	4.97	3.4	1.12 ⁱ	1.2	3.58 ^j	1.6	3.04 ^g	91
<i>trans</i> -4-hexen-1-ol	14.0	1.2	0.534 ^m	2.6	2.93 ^f	1.9	-	-
1-octen-3-ol	2.00	8.6	0.16 ⁿ	8.7	0.0099 ⁿ	560	4.03 ^k	69
<i>trans</i> -2-octen-1-ol	8.40	2.0	0.28 ⁿ	5.0	0.20 ⁿ	28	-	-

^a this work (average of absolute and relative determinations); ^b Orlando et al. ⁵³; ^c Average of Pfrang et al. ⁵⁴ and Zhao et al. ⁵⁵; ^d Average of Grira et al. ⁵⁶ and Rodriguez et al. ⁵⁷; ^e Average of Gibilisco et al. ⁵⁸ and Davis and Burkholder ⁵⁹; ^f Pfrang et al. ²⁶; ^g Average of Grira et al. ⁵⁶ and Gibilisco et al. ⁶⁰; ^h Average of Gibilisco et al. ⁶¹; Peirone et al. ⁶² and Gibilisco et al. ⁵⁸; ⁱ Average of Gibilisco et al. ⁵⁸, Jimenez et al. ⁶³, Atkinson et al. ²¹ and Davis and Burkholder ⁵⁹; ^j Average of Pfrang et al. ²⁶ and Atkinson et al. ²¹; ^k Grira et al. ⁵⁶; ^m Pfrang et al. ⁴⁹; ⁿ Pfrang et al. ²⁶; -: no data

Figures

Figure 1: Kinetic plot of *trans*-2-octen-1-ol ozonolysis.

Figure 2: Relative kinetic plot of ozone reaction with *trans*-2-hexen-1-ol versus (●) cyclohexene and (○) *trans*-2-butene, using 1,3,5-trimethylbenzene as OH scavenger.

Figure 3: Relationship between O₃ rate coefficient ratio and number of carbon atoms between -OH and C=C. For example, *cis*-2-penten-1-ol (OHCH₂-CH=CH-CH₂-CH₃) has one carbon atom between -OH and C=C. Stars (*) represent data for 1-alken-3-ols (7 compounds) while circles (●) represent other alkenols (14 compounds). Rate coefficients for alkenes are from Calvert et al.²⁵ while those for alkenols are reported in Table S1.

Figure 4: Ozonolysis of alkenols: rate coefficients calculated by SAR (see McGillen et al.²⁷ and Supplementary Information) vs those determined experimentally at 298 K (see **Table S1**). The solid line is a linear regression over all the data (26 data) going through zero. The uncertainty on the slope corresponds to 2 standard deviations (2σ). One compound, 3-methyl-2-buten-1-ol, was excluded from the linear regression (see main text).




Figure 1: Kinetic plot of *trans*-2-octen-1-ol ozonolysis.




Figure 2: Relative kinetic plot of ozone reaction with *trans*-2-hexen-1-ol versus (●) cyclohexene and (○) *trans*-2-butene, using 1,3,5-trimethylbenzene as OH scavenger.




Figure 3: Relationship between O₃ rate coefficient ratio and number of carbon atoms between -OH and C=C. For example, cis-2-penten-1-ol (OHCH₂-CH=CH-CH₂-CH₃) has one carbon atom between -OH and C=C. Stars (*) represent data for 1-alken-3-ols (7 compounds) while circles (●) represent other alkenols (14 compounds). Rate coefficients for alkenes are from Calvert et al.²⁵ while those for alkenols are reported in Table S1.




Figure 4: Ozonolysis of alkenols: rate coefficients calculated by SAR (see McGillen et al.²⁷ and Supplementary Information) vs those determined experimentally at 298 K (see **Table S1**). The solid line is a linear regression over all the data (26 data) going through zero. The uncertainty on the slope corresponds to 2 standard deviations (2σ). One compound, 3-methyl-2-buten-1-ol, was excluded from the linear regression (see main text).

TOC Graphic

