HAL
open science

New insights into the Van Krevelen diagram: Automated molecular formula determination from HRMS for a large chemical profiling of lichen extracts

Simon Ollivier, Philippe Jehan, Damien Olivier-Jimenez, Fabian Lambert, Joël Boustie, Françoise Lohezic-Le Devehat, Nicolas Le Yondre

To cite this version:

Simon Ollivier, Philippe Jehan, Damien Olivier-Jimenez, Fabian Lambert, Joël Boustie, et al.. New insights into the Van Krevelen diagram: Automated molecular formula determination from HRMS for a large chemical profiling of lichen extracts. Phytochemical Analysis, 2022, 33 (7), pp.1111-1120. 10.1002/pca.3163 . hal-03772625

HAL Id: hal-03772625

https://hal.science/hal-03772625v1

Submitted on 9 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



New Insights into the Van Krevelen Diagram: Automated Molecular Formula Determination from HR-MS for a Large Chemical Profiling of Lichen Extracts

Journal:	*Phytochemical Analysis*
Manuscript ID	PCA-22-0063.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Ollivier, Simon; Univ Rennes, UAR 2025 - ScanMAT - CRMPO (Centre Régional de Mesures Physiques de l'Ouest); Univ Rennes, CNRS - ISCR (Institut des Sciences Chimiques de Rennes) - UMR 6226 Jéhan, Phillipe; Univ Rennes, UAR 2025 - ScanMAT - CRMPO (Centre Régional de Mesures Physiques de l'Ouest) Olivier-Jimenez, Damien; Univ Rennes, CNRS - ISCR (Institut des Sciences Chimiques de Rennes) - UMR 6226 Lambert, Fabian; Univ Rennes, UAR 2025 - ScanMAT - CRMPO (Centre Régional de Mesures Physiques de l'Ouest) BOUSTIE, Joël; Univ Rennes, CNRS - ISCR (Institut des Sciences Chimiques de Rennes) - UMR 6226 LE DEVEHAT, Françoise; Univ Rennes, CNRS - ISCR (Institut des Sciences Chimiques de Rennes) - UMR 6226 LE YONDRE, Nicolas; Univ Rennes, UAR 2025 - ScanMAT - CRMPO (Centre Régional de Mesures Physiques de l'Ouest)
Keywords:	Lichens, Metabolomics, Structural Classification, Extract Comparison

SCHOLARONE™
Manuscripts

New Insights into the Van Krevelen Diagram: Automated Molecular Formula Determination from HR-MS for a Large Chemical Profiling of Lichen Extracts

Simon OLLIVIER[1,2,3], Philippe JÉHAN[1], Damien OLIVIER-JIMENEZ[2], Fabian LAMBERT[1], Joël BOUSTIE[2], Françoise LOHÉZIC – LE DÉVÉHAT[2]*, Nicolas LE YONDRE[1]*

[1] Univ Rennes, CNRS, ScanMAT UAR 2025, CRMPO (Centre Régional de Mesures Physiques de l'Ouest), F-35000 Rennes, France

[2] Univ Rennes, CNRS, ISCR (Institut des Sciences Chimiques de Rennes) - UMR 6226, F-35000 Rennes, France

[3] *INRAE, UR BIA, F-44300 Nantes, France and INRAE, PROBE research infrastructure, BIBS facility, F-44300 Nantes, France*

***Correspondence**

Dr. Nicolas Le Yondre

Phone: +33 (0)2 23 23 73 01

Email: nicolas.leyondre@univ-rennes1.fr

Fax: +33 (0)2 23 23 67 43

Dr. Françoise Lohézic-Le Dévéhat

Phone: +33 (0)2 23 23 48 16

Email: francoise.le-devehat@univ-rennes1.fr

For Peer Review

Short title (< 70 char): **Chemical Profiling of Lichen Extracts using the Van Krevelen Diagram**

Short abstract (< 80 words):

Because most metabolomics tools are designed for LC-MS, it is often difficult to obtain structural information when straying from this gold standard. The Van Krevelen diagram is adapted for this purpose. Therefore, a new classification method is benchmarked against the classification tool ClassyFire, and various extracts from four lichens are analyzed with high resolution mass spectrometry (HRMS). Lichen metabolites are efficiently classified and the obtained results are consistent with the extraction protocols. This approach provides structural insight in lichen metabolites from HRMS.

For Peer Review

ABSTRACT

Introduction: In recent years, LC-MS has become the gold standard for metabolomic studies. Indeed, liquid chromatography is relatively easy to couple with the soft electrospray ionization. As a consequence, many tools have been developed for the structural annotation of tandem mass spectra. However, it is sometimes difficult to do Data Dependent Acquisition (DDA), especially when developing new methods that stray from the classical LC-MS workflow.

Objective: An old tool from petroleomics that has recently gained popularity in metabolomics, the Van Krevelen (VK) diagram, is adapted for an overview of the molecular diversity profile in lichens through HR-MS.

Methods: A new method is benchmarked against the state-of-the-art classification tool ClassyFire using a database containing most known lichen metabolites ($n \approx 2{,}000$). Four lichens known for their contrasted chemical composition were selected, and extractions with apolar, aprotic polar and protic polar solvents were performed to cover a wide range of polarities. Extracts were analyzed with Direct Infusion ElectroSpray Ionization Mass Spectrometry (DI-ESI-MS) and Atmospheric Solids Analysis Probe Mass Spectrometry (ASAP-MS) techniques to be compared with the chemical composition described in the literature.

Results: The most common lichen metabolites were efficiently classified, with more than 90% of the molecules in some classes being matched with ClassyFire. Results from this method are consistent with the various extraction protocols in the present case study.

Conclusion: This approach is a rapid and efficient tool to gain structural insight regarding lichen metabolites analyzed by high resolution mass spectrometry without relying on DDA by LC-MS/MS analysis. It may notably be of use during the development phase of novel MS-based

For Peer Review

metabolomic approaches.

Keywords: Lichens, Metabolomics, Structural Classification, Extract Comparison

For Peer Review

INTRODUCTION

In recent years, significant advances in mass spectrometry have led to an increase in the measurement precision, sensitivity and frequency of acquisition of tandem mass spectra. These different parameters, combined with the extensive separation capabilities of liquid chromatography, have made LC-MS the gold standard in metabolomics.[1,2] It is indeed a preferred analytical technique because it provides an insight into the composition of an extract in a form that could then be used by the industry.[3] However, this also implies (i) that a time consuming extraction step must first be carried out and (ii) that a certain number of compounds will not be detected because they will be retained on the column during separation. The extraction step determines, through the choice of solvent, the fraction of the metabolome that can be studied.[4] To obtain a more comprehensive insight into the chemical composition of an organism, it becomes necessary to perform various extraction steps, which further increases the workload. Therefore, it is interesting to develop new tools for a comprehensive understanding of this composition by directly analyzing the biological material. Natural products (NPs), which are the first historical source of drugs and remain an important subject of study for the pharmaceutical industry today,[5] are of special concern.

With the establishment of LC-MS as a standard for metabolomics, many biocomputing tools have been developed and combined to facilitate the processing and interpretation of the massive amounts of data generated.[6] Most, if not all, of these tools have been created to process data containing tandem mass spectra, which are commonly acquired in a Data Dependent Acquisition (DDA).[7] It is possible to isolate features characterized by a dataset each comprising one m/z value, one retention time, and one tandem mass spectrum, which can be linked together by scores or shared patterns, notably on the GNPS platform.[8–10] Finally, these tandem mass spectra allow to propose identifications of a substantial number of molecules for each sample by comparing

For Peer Review

them with experimental[11,12] or *in silico* databases.[13] This structural annotation can allow a hierarchical classification of the molecules composing an extract, in particular thanks to the ClassyFire tool.[14]

The use of tandem mass spectra, while being the strength of these tools, is also their weakness. For instance, when developing new mass spectrometry tools, it is not always possible to proceed *ab initio* to such an acquisition associating mass and tandem mass spectra. Data Independent Acquisition (DIA) is also possible and fragments all ions present in a wide range of *m/z* regardless of intensity. However, it then becomes difficult to interpret the resulting multiplexed spectra, and the *ad hoc* algorithms require substantial computation time, which is not suitable for a rapid screening.[15] As an alternative, the option was to develop a tool providing structural insight directly from the molecular formula (MF), which can readily be computed from high resolution MS data.[16] Indeed, an old tool from petroleomics – the Van Krevelen (VK) diagram[17] – was previously used to estimate the structural class associated with a MF through element ratios, notably for studying the composition of some alcohols.[18,19] To avoid the computational burden of MF determination, a local database search can be performed to determine element ratios for Van Krevelen analysis,[20] however in this approach the analysis ignores masses absent from the database, which can be problematic in a compound discovery context. Furthermore, a database search can sometimes generate mismatches because of cation adducts or the presence of 'atypical' atoms – e.g. chlorinated metabolites, often described in lichens[21] – so, direct MF computation was chosen. For users wishing to try this kind of Van Krevelen-based approach, a Python package (PyKrev) has recently been published and offers a number of processing options, including a classification function – but not tailored for phytochemicals.[22]

Lichens, as organisms constituted of a fungus living in a symbiotic relationship with an alga

For Peer Review

and/or a cyanobacterium,[23] possess an original chemistry comprising varied and unique compounds[21] and are thus well suited for the development and assessment of such a tool. They are also an interesting source of bioactive compounds,[24,25] and have been shown to be the ancestors of major fungal lineages (e.g. *Penicillium spp., Aspergillus spp.*).[26]

This paper is the first part of a study aiming to establish a novel concept using Ambient Mass Spectrometry for NP research, focused on four lichen species: *Evernia prunastri* (L.) Ach., *Lichina pygmaea* (Lightf.) C. Agardh., *Parmelia saxatilis* (L.) Ach. and *Roccella fuciformis* (L.) DC. The lichens were selected for their chemical diversity and an extensive list of the metabolites described in these species is provided as **Table S1**. In this paper a method capable of obtaining large structural insights into the composition of lichen extracts is presented without being reliant on tandem mass spectra.

EXPERIMENTAL

General Procedures.

All solvents and reagents used in this study were HPLC grade (Sigma-Aldrich) and were used without further purification. In order to avoid any contamination by plastic compounds, the use of glass containers was preferred and the use of colored plastics was strictly prohibited. Eppendorf micropipetting systems were used for metering the extract solutions. For experiments requiring the thermal desorption of extracts, samples were deposited on Marienfeld melting point capillaries.

Lichen Material.

This study was conducted on herbarium specimens. They were collected in: Cressensac, France (N 45°0’19321” O 1°30’51.191”), August 2006 for *E. prunastri*; Roscoff, France (N 48°43'31.458" O 3°58'8.651"), June 2016 for *L. pygmaea*; Girona, Spain (N 42°27’ 53.399” O 1°47’23.816”), July

For Peer Review

126 2017 for *P. saxatilis*; and Saint-Coulomb, France (N 48°41'29.299" O 1°56'45.855"), May 2003 for
127 *R. fuciformis*. Voucher specimens are kept in the herbarium of the University of Rennes, France
128 under the respective reference codes: JB/06/51, JB/16/204, JB/17/213 and JB/03/02.

129 **Extraction Protocol.**

130 The dried weights were chosen for each extraction so as to extract a sufficient quantity of material
131 for all analyses (**Figure S1**). For each lichen, 2 extracts were obtained with methanol and acetone,
132 and 2 extracts were from a successive extraction using cyclohexane, acetone then methanol (CAM)
133 and using cyclohexane, acetone then water (CAW). CAM and CAW extracts were recombined *pro*
134 *rata* to the extracted quantities. The selected quantities were 2 g for both the single-step 'Acetone'
135 and the multiple-step 'CAM' extracts, 500 mg for the 'Methanol' extract and 4 g for the 'CAW'
136 extract. All extractions were performed using an Accelerated Speed Extractor Büchi
137 SpeedExtractor E-914 after dispersion of ground lichens in celite (1:1 ratio, m/m). A 2 g sand bed
138 made of Büchi quartz sand of 0.3-0.9 mm granulometry was used to avoid clogging of the sintered
139 part of the extraction cartridges. Each extraction step consisted of 3 maceration cycles of 10 min
140 in 40 °C solvent, with a discharge time of 2 min and a degassing between every cycle.

141 **HR-MS Analyses.**

142 Direct infusion electrospray mass spectrometry (DI-ESI-MS) analyses were acquired on a Q-
143 Exactive mass spectrometer (Thermo Scientific, Bremen, Germany) equipped with a Thermo
144 HESI-II ion source. DI-ESI-MS analyses were performed on 15 µg/mL extract solutions using two
145 ionization solvents, acetone and methanol. Acquisition time was set to 2 min (230 scans).
146 Optimized parameters included a sheath gas set at 20 a.u., a transfer capillary voltage of 2800 V
147 and an S-lens RF level of 50 a.u., the other parameters were set to default. The Thermo Scientific

For Peer Review

148 Q-Exactive™ was calibrated with Pierce LTQ Velos ESI Positive and Negative Ion calibration
149 solutions and lock-mass calibration was applied on palmitic acid during acquisition (deprotonated
150 molecule $[M–H]^–$ *m/z* 255.2330 and protonated molecule $[M+H]^+$ *m/z* 257.2475). Experiments
151 were performed in a single batch with analytical triplicates. The system was rinsed with the
152 working solvent up to noise level between each sample. No loss of sensitivity was observed when
153 rerunning samples as control.

154 Atmospheric Solids Analysis Probe Mass Spectrometry (ASAP-MS) analyses were
155 acquired on a Maxis 4G spectrometer (Bruker Daltonics, Bremen, Germany) equipped with an
156 APCI ion source and a direct insertion probe (DIP). The extracts were deposited on Marienfeld
157 melting point capillaries as homogeneous solutions and dried with nitrogen before insertion in the
158 APCI source. Preliminary tests having shown a major thermolability of some compounds, analyses
159 were performed with a temperature ramp-up (50 °C every 15 sec) over 2 min. The temperature
160 range covered was therefore 100-400 °C with 120 scans in the range *m/z* 100-2000. Source
161 parameters were optimized as following with a nebulization gas (nitrogen) pressure of 2.5 bars and
162 a dry gas at 2.5 $L.min^{-1}$. Voltages were set to: 500 V for the transfer capillary, -500 V for the end
163 plate offset, with a 2000 V charging voltage and a 4000 V working voltage. An ion-cooler of 175
164 V_{p-p} was applied on the pusher. The Corona discharge was set to 500 nA for positive ionization and
165 2000 nA for negative ionization. External calibration of the spectrometer was done with PEG 600
166 in positive mode and PEG diacid 600 in negative mode. Lock-mass calibration was applied on
167 palmitic acid by post-processing with R-script.[27] During the batch two replicates were performed
168 to obtain a total of 8 acquisitions on the merged data for interpretation. Experiments were
169 performed as a single batch with a thermal desorption to noise level between each run, no loss of
170 sensitivity was observed on controls.

For Peer Review

Bioinformatic Data Processing.

Thermo Scientific *.RAW files were converted to *.mzXML using readw 4.3.1 and Bruker *.d folders were converted to *.mzXML using Bruker CompassXport 3.0.6.9 from the command line. The obtained files were processed with R-script and Microsoft Visual Basic, following the Seven Golden Rules of MF determination,[16] every step was controlled manually while encoding the scripts, and exported in the form of *.csv files.

Peak lists were extracted from the *.mzXML files using R packages `xcms`,[28–30] `MSnbase`,[31] and `CAMERA`,[32] and automated instrument detection was performed with the `readMzXmlData`[33] package. The peak picking parameters were optimized for each instrument. Isotopic contributions calculation and clustering followed by deisotoping were performed with a 3 ppm tolerance using element restrictions depending on *m/z* as described in the Seven Golden Rules. For this purpose the mass ranges were refined according to the compounds referenced in the online Dictionary of Natural Products (DNP) v26.2[34] (**Table 1**). After an in-depth study of the mass spectra, it appeared that isotopic contributions could not be seen for monoisotopic ions of less than 0.01 % relative intensity, so this value was set as a threshold for the calculation of isotopic clusters.

The molecular formulae were then calculated by querying the online platform ChemCalc[35] with R-script, taking into account the calculated isotopic patterns, allowing an error of 4 carbons and 2 nitrogens. $[M+Na]^+$ adducts were further accounted for, when generating the formulae for ESI-MS. When no isotope pattern was detected, **Table 1** features were considered. However, sulfur and chlorine were not considered unless detected. The absolute mass range error was set to 1.0 milli-unified atomic mass (mu) for orbitrap data (DI-ESI-MS) and 1.7 mu for Q-TOF data (ASAP-MS). Based on the Double Bond Equivalent (DBE) the script filtered out the formulae corresponding to radical ions for ESI-MS. The results were coherent with manual MF

For Peer Review

194 determination of known compounds and yielded most often a single formula, though the script was
195 able to calculate the five best formulae.

196 The different elemental ratios were then calculated with Visual Basic in order to filter the
197 formulae according to the Seven Golden Rules to remove the potentially incoherent ones. An
198 adduct correction was applied according to the formulae and DBE with the following considered
199 adducts: ASAP-MS: $M^{+\bullet}$, $[M+H]^+$, $M^{-\bullet}$, $[M–H]^-$ and ESI-MS: $[M+H]^+$, $[M+Na]^+$, $[M–H]^-$,
200 $[M+Cl]^-$. The corresponding exact molecular masses were subsequently calculated.

201 Chemical classes associated with the Van Krevelen diagram were retrieved from the
202 literature and double-checked against an in-house database of lichen compounds, the complete
203 projet is available[12] (see **Technical validation**). The *Van Krevelen coordinates i.e. the H/C and*
204 *O/C ratios* were then imported in R-script and merged in a single matrix that can be filtered by all
205 experimental variables. A rectangle approximation of the chemical groups presented **Figure 1** was
206 applied to tag the detected compounds, which were filtered for plotting in the form of a histogram
207 (relative proportion within a method) stacked with a heatmap (absolute count of detected features).
208 The plots were created with R package `plot3D`[36] and were made interactive with `plot3Drgl`.
209 [37] Although the presented plots were constructed with the merged data of all lichens, the provided
210 scripts allow for species filtering (e.g. **Figure S2**). The Van Krevelen coordinates were also used
211 to plot the Van Krevelen diagrams using R-script.

212 The entire processed dataset is available at https://osf.io/6pyuq/
213 (dataset_solvents_comp_publi1.tsv) along with sample files for Q-Orbitrap and Q-TOF data in the
214 *.mzXML format. The scripts are deposited at
215 https://github.com/siollivier/directacquisitionproject.

For Peer Review

Technical validation.

Database establishment. An in-house database of lichen compounds was compiled from the literature. The total number of structures contained in this compiled database is about 2,000 entries[12]. The database will be provided by the authors upon reasonable request.

The structural information was translated into InChIKey, a particularly popular format in chemoinformatics, because it is standardized and of fixed length, allowing easy data processing (example: mycosporine serinol, VVTDHOIRNPCGTH-NSHDSACASA-N).

A detailed classification of the lichen compounds was then obtained by submitting these InChIKeys to ClassyFire,[14] a tool to obtain a hierarchical classification of compounds, through the `classyfireR` package.[38] This structure-based bioinformatics approach allows for a comprehensive result through the establishment of a chemical taxonomy—in hierarchical order: chemical kingdom, superclass, class, subclass, and additional levels 5-9.

Evaluation of the method against a state-of-the-art classification tool. The accuracy of the classification established by **Figure 1** was examined in comparison to this database of lichen compounds. Within the database, compounds were selected according to different levels of their ClassyFire classification, which was compared to the classification obtained with Van Krevelen.

The same level of precision cannot be expected from a classification based only on the MF. However, depending on the structure of the compounds, it is possible to expect a result in one or more classes (more or less specific) of this diagram. Thus, some areas of the diagram (i.e. benzenoids or prenyl derivatives) can be considered as areas that extend more specific ones. During the evaluation of the protocol, the compounds were considered as correctly classified if their Van Krevelen annotation was in accordance with the expected matches between ClassyFire and Van

For Peer Review

238 Krevelen that are shown in **Table 2**.

239 **RESULTS AND DISCUSSION**

240 **Determination and validation of the *Van Krevelen chemical groups*.**

241 First, a search was conducted in the literature to identify the positioning of the defined chemical
242 groups on the Van Krevelen diagram.[18,19,39–41] These bibliographic results were then refined
243 according to a database organized with ClassyFire and the final ranges are specified in **Figure 1**.
244 The database used to validate these annotations mostly contains compounds of MW < 500 Da,
245 including mainly, in terms of Classyfire classes, depsides and depsidones (> 20 %), as illustrated
246 in **Figure 2**.

247 In order to guarantee the quality of the following results, it is first necessary to evaluate the
248 accuracy of the classification on datasets not acquired in DDA. The proportion of correctly
249 classified molecules in each class is an effective way to perform this validation.

250 However, two biases to this estimation should be considered: (i) the classification being
251 based on H/C and O/C ratios, classes containing a large number of heteroatoms (N, S, *etc*.) such as
252 amino acids are more likely to be incorrectly classified. Others have addressed this issue by
253 including various elemental ratios, but this promising approach has yet to achieve the same level
254 of specificity for classification;[42] (ii) some ClassyFire classifications include very structurally
255 diverse derivatives within the same class, particularly for compounds with a lipid or carbohydrate
256 function.

257 The proportions of molecules correctly classified in each of the classes considered are
258 presented in **Figure 3**. As expected, amino acids and analogues, fatty acyls and carbohydrates have
259 a lesser overlap between the two classifications, with respectively 56%, 59% and 68% of

For Peer Review

compounds being correctly classified.

On the other hand, all the other structural classes considered have an overlap of more than 70%. For 4 of the 10 structural types considered, the similarity between the two classifications is even higher than 90%. These are benzenoid derivatives, chromones, depsides and depsidones as well as prenol lipids, which include terpenes. These chemical families are among the most frequently reported in lichens, as illustrated in the recently published tandem mass spectra database of the most common lichen compounds.[12]

The encouraging results of the method provided by processing non-experimental data gave us the incentive to proceed to a practical evaluation. In the following case study, this method is used to obtain a rapid evaluation of the diversity of compounds detected by multiple ionization techniques, presenting a comparison of results from two DI-ESI-MS conditions vs ASAP-MS.

Case Study: Evaluation of the Metabolic Diversity of Lichen Extracts.

In order to cover a large panel of structures from lichen extracts, four lichen samples known to have very distinct metabolite profiles were chosen (the number of compound in the ClassyFire classes are given in **Table S2** for each lichen). Protocols used for each lichen species (two single-step and two three-step ones) were established to have a comparison with a simple extraction generally used in chemotaxonomic studies in lichenology and more exhaustive extractions (pooling successive extracts from polar to apolar ones) (**Figure S1**). These extracts were analyzed with DI-ESI-MS and ASAP-MS to compare the performance of these techniques to reveal lichen compounds and evaluate their distribution in structural families (**Figure S2**).

The DI-ESI-MS experiments were conducted with two ionization solvents, acetone and methanol since this aspect of the experiment heavily impacts the compound detection. Indeed, with

For Peer Review

282 positive ion mode data, Principal Component Analysis (PCA) shows that the two ionization
283 solvents form distinct groups on the PC2 component (**Figure S3**). PC1 clearly differentiates DI-
284 ESI-MS from ASAP-MS analysis and post-treated data with an adduct correction resulted in a
285 slightly better clustering of samples with regard to raw data. While the positive ion mode detects a
286 huge number of compounds when compared to the negative ion mode, the Van Krevelen diagram
287 provides a broad and clear overview of the chemical diversity of the diverse chemical families in
288 lichens. The complementarity of these three methods is also revealed through the distinct zones
289 mostly covered by these 3 MS conditions on the Van Krevelen diagram with all discriminant
290 coordinates (**Figure S4**).

291 In other words, a single detection mode is not sufficient to cover the metabolic diversity of
292 lichen compounds and each technique is more or less suitable for a class of compounds to be
293 revealed. This assumption was verified using a bioinformatic quantification of the number of
294 molecules ionized in various samples obtained with different extraction protocols: depending on
295 their MF, a structural class was assigned to the compounds according to the cartography presented
296 **Figure 1**.

297 With regard to the relative performance of the extraction protocols compared in this study
298 the quantitative and the qualitative aspects are considered. Comparing extractions with a single
299 solvent, yields are higher when a polar protic solvent (i.e. methanol) is used instead of a polar
300 aprotic solvent (i.e. acetone). The Cyclohexane-Acetone-Methanol (CAM) and Cyclohexane-
301 Acetone-Water (CAW) extractions gave yields in the same range compared to a single methanol
302 extraction (**Figure S1**).

303 Considering the performance of the MS techniques used to reveal the composition of
304 extracts, the results obtained with the two multi-step solvents were analyzed, which are considered

For Peer Review

305 to be the more exhaustive. In **Figure 4** ASAP-MS allowed the detection of more compounds than
306 ESI-MS in negative mode (average number of ions: 424/run for ASAP-MS vs. 116/run for ESI-
307 MS in acetone), but fewer in positive mode (average: 482/run for ASAP-MS vs. 1434/run for ESI-
308 MS in acetone). Furthermore, these results are coherent with what could be expected when
309 considering the protic solvent selected, and the most appropriate extraction and ionization methods
310 for each class of compounds are summarized in **Table 3**. In addition, the comparison with the in-
311 house database[12] exhibits differences for Fatty acyls or Amino-acids, as examples, with higher
312 frequencies in the experimental data than in the database.

313 *Regarding the extraction protocols*.

314 The successive extraction steps method appears to be better suited than the methanol extraction to
315 gather terpenes and prenol derivatives (respectively 2.0 and 1.5 times more, in positive ASAP-
316 MS), while 1.3 times more carbohydrates were extracted with water (negative ESI-MS in
317 methanol).

318 A comparison with the single-step extractions (**Figure S5**) shows that an acetone extraction
319 would appear as a good choice for studies targeting polyphenols & derivatives or condensed
320 aromatic structures as well as terpenes. On the other hand, as methanol has the closest extraction
321 profile to that of the successive solvent extraction and comparable yields, it seems to be a good
322 compromise. However as expected with a protic solvent fewer terpenes are extracted with
323 methanol.

324 *Regarding the MS methods*.

325 Compared to ESI-MS, ASAP-MS favors the ionization of terpenes (relative frequency within an
326 ionization method up to r.f. 20 %) and unsaturated hydrocarbons (r.f. 10 %) in positive mode, and

For Peer Review

327 benzenoids (r.f. 35 %) in negative mode. However, ASAP-MS appears less favorable for
328 polyphenols & derivatives but, interestingly, quite favorable for condensed aromatic compounds
329 (up to r.f. 20% in some conditions, see **Figure 4d**).

330 This piece of data also shows differences between the two DI-ESI-MS ionization solvents
331 that may be of use for targeted studies: in negative mode, carbohydrates and fatty acyls' ionization
332 is superior in methanol (15 vs. 10 % and 20 vs. 10 % r.f. respectively) whereas that of polyphenols
333 & derivatives and condensed aromatic compounds is better in acetone (40 vs. 25 % and 15 vs. 10
334 % respectively). Differences are less marked in positive ionization, except for the condensed
335 aromatic structures that are better ionized in methanol (> 15 vs. 10 %). Such compounds are
336 common occurrences in lichens (e.g. depsides for polyphenols & derivatives and xanthones for
337 condensed aromatic compounds).[21,43]

338 Furthermore, in positive mode there appears to be a similar ionization profile for benzenoids
339 in all ionization methods (**Figure 4**, **Figure S5**), questioning the complementarity of the methods
340 in this aspect. A referral to the Van Krevelen diagram however reveals that the ionized compounds
341 have some discriminant elemental ratios with each method (Figure 5). VK discriminant coordinates
342 are clearly found in low H/C ratio for ASAP-MS and in high H/C ratio for DI-ESI-MS in Methanol,
343 whereas DI-ESI-MS in Acetone as few discriminant coordinates, showing a complementarity
344 within the ionization modes. A referral to the Van Krevelen diagram however reveals that some
345 elemental ratios are common to 2 or 3 ionization modes (**Figure 6**). Moreover, an overall higher
346 number of occurrences in acetone extract is clearly observed, with a specificity in high H/C ratio–
347 once again showing the usefulness of the Van Krevelen diagram for natural products metabolomics.
348 To illustrate the potential of the use of Ambient MS for validation or new identification of Natural
349 Product, in comparison with the in-house database, classical expected compounds but also new

For Peer Review

coordinates are found in DI-ESI-MS and ASAP-MS as proofs of principle (**Figures S6-S10**).

Overall, the interest of the Van Krevelen diagram was demonstrated for the structural classification of metabolites (without the need to acquire tandem mass spectra data), and its applicability was shown with real-life cases by studying the chemical composition of lichens.

REFERENCES

1. Wolfender J-L, Marti G, Thomas A, Bertrand S. Current approaches and challenges for the metabolite profiling of complex natural extracts. *J Chromatogr A*. 2015;1382:136-164. doi:10.1016/j.chroma.2014.10.091

2. Wolfender JL, Nuzillard JM, Van J der H, Renault JH, Bertrand S. Accelerating metabolite identification in natural product research: toward an ideal combination of LC-HRMS/MS and NMR profiling, in silico databases and chemometrics. *Anal Chem*. Published online November 2018. doi:10.1021/acs.analchem.8b05112

3. Sarker SD, Latif Z, Gray AI. Natural Product Isolation. In: Sarker SD, Latif Z, Gray AI, eds. *Natural Products Isolation*. Methods in Biotechnology. Humana Press; 2005:1-25. doi:10.1385/1-59259-955-9:1

4. Choi YH, Verpoorte R. Metabolomics: What You See is What You Extract. *Phytochem Anal*. 2014;25(4):289-290. doi:10.1002/pca.2513

5. Basmadjian C, Zhao Q, Bentouhami E, et al. Cancer wars: natural products strike back. *Front Chem*. 2014;2. doi:10.3389/fchem.2014.00020

6. Ernst M, Kang KB, Caraballo-Rodríguez AM, et al. MolNetEnhancer: Enhanced Molecular Networks by Integrating Metabolome Mining and Annotation Tools. *Metabolites*. 2019;9(7):144. doi:10.3390/metabo9070144

7. Allard P-M, Genta-Jouve G, Wolfender J-L. Deep metabolome annotation in natural products research: towards a virtuous cycle in metabolite identification. *Curr Opin Chem Biol*. 2017;36:40-49. doi:10.1016/j.cbpa.2016.12.022

8. Wang M, Carver JJ, Phelan VV, et al. Sharing and community curation of mass spectrometry data with Global Natural Products Social Molecular Networking. *Nat Biotechnol*. 2016;34:828-837. doi:10.1038/nbt.3597

9. Wandy J, Zhu Y, van der Hooft JJJ, Daly R, Barrett MP, Rogers S. Ms2lda.org: web-based topic modelling for substructure discovery in mass spectrometry. *Bioinformatics*. 2018;34(2):317-318. doi:10.1093/bioinformatics/btx582

10. Hooft JJJ van der, Wandy J, Barrett MP, Burgess KEV, Rogers S. Topic modeling for untargeted substructure exploration in metabolomics. *Proc Natl Acad Sci*.

For Peer Review

2016;113(48):13738-13743. doi:10.1073/pnas.1608041113

11. Ramos AEF, Le Pogam P, Alcover CF, et al. Collected mass spectrometry data on monoterpene indole alkaloids from natural product chemistry research. *Sci Data*. 2019;6(1):1-6. doi:10.1038/s41597-019-0028-3

12. Olivier-Jimenez D, Chollet-Krugler M, Rondeau D, et al. A database of high-resolution MS/MS spectra for lichen metabolites. *Sci Data*. 2019;6(1):1-11. doi:10.1038/s41597-019-0305-1

13. Allard P-M, Péresse T, Bisson J, et al. Integration of Molecular Networking and In-Silico MS/MS Fragmentation for Natural Products Dereplication. *Anal Chem*. 2016;88:3317-3323. doi:10.1021/acs.analchem.5b04804

14. Djoumbou Feunang Y, Eisner R, Knox C, et al. ClassyFire: automated chemical classification with a comprehensive, computable taxonomy. *J Cheminformatics*. 2016;8(1):61. doi:10.1186/s13321-016-0174-y

15. Bilbao A, Varesio E, Luban J, et al. Processing strategies and software solutions for data-independent acquisition in mass spectrometry. *Proteomics*. 2015;15(5-6):964-980. doi:10.1002/pmic.201400323

16. Kind T, Fiehn O. Seven Golden Rules for heuristic filtering of molecular formulas obtained by accurate mass spectrometry. *BMC Bioinformatics*. 2007;8:105. doi:10.1186/1471-2105-8-105

17. Van Krevelen DW. Graphical-statistical method for the study of structure and reaction processes of coal. *Fuel*. 1950;29:269-284.

18. Roullier-Gall C, Witting M, Gougeon RD, Schmitt-Kopplin P. High precision mass measurements for wine metabolomics. *Front Chem*. 2014;2. doi:10.3389/fchem.2014.00102

19. Roullier-Gall C, Lucio M, Noret L, Schmitt-Kopplin P, Gougeon R. How Subtle Is the "Terroir" Effect? Chemistry-Related Signatures of Two "Climats de Bourgogne." *PloS One*. 2014;9:e97615. doi:10.1371/journal.pone.0097615

20. Brockman SA, Roden EV, Hegeman AD. Van Krevelen diagram visualization of high resolution-mass spectrometry metabolomics data with OpenVanKrevelen. *Metabolomics*. 2018;14(4):48. doi:10.1007/s11306-018-1343-y

21. Huneck S, Yoshimura I. *Identification of Lichen Substances*. Springer Berlin Heidelberg; 1996. Accessed February 17, 2019. http://public.eblib.com/choice/publicfullrecord.aspx?p=3097560

22. Kitson E, Kew W, Ding W, Bell NGA. PyKrev: A Python Library for the Analysis of Complex Mixture FT-MS Data. *J Am Soc Mass Spectrom*. 2021;32(5):1263-1267. doi:10.1021/jasms.1c00064

For Peer Review

23. Honegger R. The Lichen Symbiosis—What is so Spectacular about it?*. *The Lichenologist*. 1998;30(3):193-212. doi:10.1006/lich.1998.0140

24. Boustie J, Grube M. Lichens—a promising source of bioactive secondary metabolites. *Plant Genet Resour*. 2005;3(2):273-287. doi:10.1079/PGR200572

25. Ranković B, Kosanić M. Lichens as a Potential Source of Bioactive Secondary Metabolites. In: Ranković B, ed. *Lichen Secondary Metabolites: Bioactive Properties and Pharmaceutical Potential*. Springer International Publishing; 2015:1-26. doi:10.1007/978-3-319-13374-4_1

26. Lutzoni F, Pagel M, Reeb V. Major fungal lineages are derived from lichen symbiotic ancestors. *Nature*. 2001;411(6840):937-940. doi:10.1038/35082053

27. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria; 2018. https://www.R-project.org/

28. Smith CA, Want EJ, O'Maille G, Abagyan R, Siuzdak G. XCMS: processing mass spectrometry data for metabolite profiling using nonlinear peak alignment, matching, and identification. *Anal Chem*. 2006;78(3):779-787. doi:10.1021/ac051437y

29. Tautenhahn R, Böttcher C, Neumann S. Highly sensitive feature detection for high resolution LC/MS. *BMC Bioinformatics*. 2008;9(1):504. doi:10.1186/1471-2105-9-504

30. Benton HP, Want EJ, Ebbels TMD. Correction of mass calibration gaps in liquid chromatography–mass spectrometry metabolomics data. *Bioinformatics*. 2010;26(19):2488-2489. doi:10.1093/bioinformatics/btq441

31. Gatto L, Lilley KS. MSnbase-an R/Bioconductor package for isobaric tagged mass spectrometry data visualization, processing and quantitation. *Bioinformatics*. 2012;28(2):288-289. doi:10.1093/bioinformatics/btr645

32. Kuhl C, Tautenhahn R, Böttcher C, Larson TR, Neumann S. CAMERA: an integrated strategy for compound spectra extraction and annotation of liquid chromatography/mass spectrometry data sets. *Anal Chem*. 2012;84(1):283-289. doi:10.1021/ac202450g

33. Gibb S. *ReadMzXmlData: Reads Mass Spectrometry Data in MzXML Format*.; 2015. Accessed February 21, 2019. https://CRAN.R-project.org/package=readMzXmlData

34. Dictionary of Natural Products 26.2. Published April 13, 2018. Accessed April 13, 2018. http://dnp.chemnetbase.com/faces/chemical/ChemicalSearch.xhtml

35. Patiny L, Borel A. ChemCalc: A Building Block for Tomorrow's Chemical Infrastructure. *J Chem Inf Model*. 2013;53(5):1223-1228. doi:10.1021/ci300563h

36. Soetaert K. *Plot3D: Plotting Multi-Dimensional Data*.; 2017. Accessed February 21, 2019. https://CRAN.R-project.org/package=plot3D

For Peer Review

37. Soetaert K. *Plot3Drgl: Plotting Multi-Dimensional Data - Using "Rgl."*; 2016. Accessed February 21, 2019. https://CRAN.R-project.org/package=plot3Drgl

38. Wilson T, Finch J. *ClassyfireR: R Interface to the ClassyFire RESTful API*.; 2019. Accessed August 16, 2019. https://CRAN.R-project.org/package=classyfireR

39. D'Andrilli J, Cooper W, Foreman C, G. Marshall A. An ultrahigh-resolution mass spectrometry index to estimate natural organic matter lability. *Rapid Commun Mass Spectrom*. 2015;29:2385-2401. doi:10.1002/rcm.7400

40. Kuhnert N, Dairpoosh F, Yassin G, Golon A, Jaiswal R. *What Is under the Hump? Mass Spectrometry Based Analysis of Complex Mixtures in Processed Food – Lessons from the Characterisation of Black Tea Thearubigins, Coffee Melanoidines and Caramel*. Vol 4.; 2013. doi:10.1039/c3fo30385c

41. Lu Y, Li X, Mesfioui R, et al. Use of ESI-FTICR-MS to Characterize Dissolved Organic Matter in Headwater Streams Draining Forest-Dominated and Pasture-Dominated Watersheds. *PloS One*. 2015;10:e0145639. doi:10.1371/journal.pone.0145639

42. Rivas-Ubach A, Liu Y, Bianchi TS, Tolić N, Jansson C, Paša-Tolić L. Moving beyond the van Krevelen Diagram: A New Stoichiometric Approach for Compound Classification in Organisms. *Anal Chem*. 2018;90(10):6152-6160. doi:10.1021/acs.analchem.8b00529

43. Le Pogam P, Boustie J. Xanthones of Lichen Source: A 2016 Update. *Molecules*. 2016;21(3):294. doi:10.3390/molecules21030294

TABLES

Table 1. Element restriction ranges established from the online Dictionary of Natural Products v26.2.

Mass range [Da]	C max	H max	N max	O max	S max	Cl max
< 200	15	30	8	7	6	4
< 400	30	58	10	14	12	7
< 600	42	86	13	21	12	8
< 800	56	108	16	25	20	10
< 1000	66	126	25	37	20	11
< 1500	100	182	26	44	20	11

For Peer Review

476 **Table 2.** Matches between ClassyFire and the Van Krevelen diagram for the most common lichen
477 metabolites. The matches were determined to fit properly with the usual classification of lichen
478 compounds by Huneck and Yoshimura.[21]

ClassyFire Classification (level)	Expected Area(s) in the Van Krevelen Diagram
Amino Acids and analogues (subclass)	Amino acids; Nucleic acids
Anthraquinones (subclass)	Condensed aromatic compounds
Benzene derivatives (class)	***Polyphenols and derivatives***; Benzenoids
Carbohydrates and conjugates (subclass)	Carbohydrates
Chromones (level 5)	***Polyphenols and derivatives***; Benzenoids
Depsides and depsidones (class)	***Polyphenols and derivatives***; Benzenoids
Dibenzofurans (subclass)	Condensed aromatic compounds; Benzenoids
Fatty acyls (class)	Fatty acyls; Prenol derivatives
Prenol lipids (class)	Terpenes; Prenol derivatives; Unsaturated hydrocarbons
Xanthones (level 7)	Condensed aromatic compounds

479

480 **Table 3.** An overview of the most efficient parameters in this study for the extraction and MS
481 analysis of lichen metabolites, with regards to the number of compounds ionized and the amount
482 of other metabolites that may hinder the interpretation for each structural class.

Class	Extraction	Ionization	MS Polarity
Carbohydrates	CAW	ESI in methanol	NEG
Amino Acids	Methanol	ESI in acetone	POS
Nucleic Acids	CAW	ESI in acetone	NEG
Prenol Derivatives	CAM	ASAP (APCI)	POS
Fatty Acyls	Methanol	ESI in methanol	NEG
Unsaturated Hydrocarbons	CAM	ASAP (APCI)	POS

For Peer Review

Terpenes	CAM or Acetone	ASAP (APCI)	POS
Benzenoids	Acetone	ESI in acetone	POS
Polyphenols and derivatives	Acetone	ESI in acetone	NEG
Condensed Aromatic Compounds	Acetone	ESI in methanol	POS

483

484 **FIGURE LEGENDS**

485 **Figure 1.** Identification of the chemical groups on the Van Krevelen diagram.

486 **Figure 2.** (a) The distribution of the molecular weights of the compounds contained in the
487 validation database (LDB-Lit, solid line) shows a good overlap with the experimental data (dotted
488 line); (b) The Classyfire classes of the compounds in the LDB-Lit illustrate the diversity of the
489 molecules used for technical validation.

490 **Figure 3.** Proportions of molecules in the database accurately classified by the Van Krevelen
491 diagram (in accordance with **Table 2**).

492 **Figure 4.** Evaluation of the chemical diversity for the merged data of the three-step extractions:
493 'Cyclohexane-Acetone-Methanol' in positive mode (a) and negative mode (b), and 'Cyclohexane-
494 Acetone-Water' in positive mode (c) and negative mode (d). Both the relative frequency for each
495 technique (histogram, vertical axis) and the absolute number of detected ions (heatmap) are
496 presented. The relative frequency represents the proportion of a chemical class within a given
497 ionization method. The database content is proposed in comparison to show the existing potential
498 in new identifications.

499 **Figure 5.** Extracted Van Krevelen diagram for benzenoids i.e. in the (O/C 0.2-0.4; H/C 0.8-1.8)
500 range, in all three ionization methods for the acetone (a) and Cyclohexane-Acetone-Methanol
501 extracts (b). The size of the circle, represented by a coefficient of expansion (cex), is proportional
502 to the number of occurrences of the (O/C, H/C) coordinates without correction factor (100 gives a
503 cex of 100). The colors (green, blue and red, alpha of 0.4) indicate respectively discriminant VK
504 coordinates for DI-ESI-MS in Methanol, ASAP-MS and DI-ESI-MS in Acetone ionization mode.

505 **Figure 6.** Extracted Van Krevelen diagram for benzenoids i.e. in the (O/C 0.2-0.4; H/C 0.8-1.8)
506 range, in all three ionization methods for the acetone (a) and Cyclohexane-Acetone-Methanol
507 extracts (b). The size of the circle is proportional to the number of occurrences of the (O/C, H/C)
508 coordinates with a correction factor of 0.1 (100 gives a cex of 10). Non-discriminant VK
509 coordinates for a single ionization mode are plotted in violet.

510

511 **Acknowledgements**

For Peer Review

The authors thank Isabelle Rouaud (ISCR, CorInt, Rennes) for her technical assistance regarding the extraction protocols.

Author Notes

SO, PJ, FLD, JB and NLY designed the experiments. SO and FLD selected the lichen material, which was identified by JB. DO optimized the ASE protocol; SO, PJ and FL carried out the experiments. SO and NLY encoded the R-scripts and Visual Basic macro used for data processing. DO compiled the database for validation, DO and SO curated the database. All authors contributed to the redaction of the manuscript. None of the authors declare any competing financial interest.

For Peer Review



Figure 1. Identification of the chemical groups on the Van Krevelen diagram.

35x23mm (300 x 300 DPI)



Figure 2. (a) The distribution of the molecular weights of the compounds contained in the validation database (LDB-Lit, solid line) shows a good overlap with the experimental data (dotted line); (b) The Classyfire classes of the compounds in the LDB-Lit illustrate the diversity of the molecules used for technical validation.

950x328mm (200 x 200 DPI)



Figure 3. Proportions of molecules in the database accurately classified by the Van Krevelen diagram (in accordance with Table 2).

3102x1281mm (72 x 72 DPI)



Figure 4. Evaluation of the chemical diversity for the merged data of the three-step extractions: 'Cyclohexane-Acetone-Methanol' in positive mode (a) and negative mode (b), and 'Cyclohexane-Acetone-Water' in positive mode (c) and negative mode (d). Both the relative frequency for each technique (histogram, vertical axis) and the absolute number of detected ions (heatmap) are presented. The relative frequency represents the proportion of a chemical class within a given ionization method. The database content is proposed in comparison to show the existing potential in new identifications.

1420x969mm (96 x 96 DPI)



Figure 5. Extracted Van Krevelen diagram for benzenoids i.e. in the (O/C 0.2-0.4; H/C 0.8-1.8) range, in all three ionization methods for the acetone (a) and Cyclohexane-Acetone-Methanol extracts (b). The size of the circle, represented by a coefficient of expansion (cex), is proportional to the number of occurrences of the (O/C, H/C) coordinates without correction factor (100 gives a cex of 100). The colors (green, blue and red, alpha of 0.4) indicate respectively discriminant VK coordinates for DI-ESI-MS in Methanol, ASAP-MS and DI-ESI-MS in Acetone ionization mode.

950x328mm (200 x 200 DPI)



Figure 6. Extracted Van Krevelen diagram for benzenoids i.e. in the (O/C 0.2-0.4; H/C 0.8-1.8) range, in all three ionization methods for the acetone (a) and Cyclohexane-Acetone-Methanol extracts (b). The size of the circle is proportional to the number of occurrences of the (O/C, H/C) coordinates with a correction factor of 0.1 (100 gives a cex of 10). Non-discriminant VK coordinates for a single ionization mode are plotted in violet.

950x328mm (200 x 200 DPI)