



**HAL**  
open science

## **Are nanomaterials leading to more efficient agriculture? Outputs from 2009 to 2022 research metadata analysis**

E. Santos, G. Montanha, M. F. Gomes, N. Duran, C. Corrêa, S. L. Z. Romeu,  
A. E.S. Pereira, J. Oliveira, E. Almeida, A. Pérez-De-Luque, et al.

### ► **To cite this version:**

E. Santos, G. Montanha, M. F. Gomes, N. Duran, C. Corrêa, et al.. Are nanomaterials leading to more efficient agriculture? Outputs from 2009 to 2022 research metadata analysis. *Environmental science.Nano*, 2022, 2022, 10.1039/D1EN01078F . hal-03774435

**HAL Id: hal-03774435**

**<https://hal.science/hal-03774435>**

Submitted on 10 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Environmental  
Science**  
Nano

**Are nanomaterials leading to more efficient agriculture?  
Outputs from 2009-2022 research metadata analysis**

Journal:	<i>Environmental Science: Nano</i>
Manuscript ID	EN-CRV-11-2021-001078.R2
Article Type:	Critical Review

**SCHOLARONE™**  
Manuscripts

**Environmental Significance Statement**

The unique properties of materials at the nanoscale has raised expectations for new products and solutions for agriculture, such as enhanced fertilizers, biostimulants, and pesticides. Aiming at gathering and summarizing the collective knowledge so far produced, this metadata review assembled and organized quantitative information focused on the scientific literature of the last decade. The present study represents a source of comprehensive data that can help scientists to clearly grasp what has been done and how much is still to be done.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## ARTICLE

## Are nanomaterials leading to more efficient agriculture? Outputs from 2009-2022 research metadata analysis

Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Eduardo Santos,<sup>a</sup> Gabriel Sgarbiero Montanha,<sup>a,b</sup> Marcos Henrique Feresin Gomes,<sup>a</sup> Nádia Marion Duran,<sup>a</sup> Camila Grazielle Corrêa,<sup>a</sup> Sara Luiza Zachy Romeu,<sup>a</sup> Anderson Espirito Santo Pereira,<sup>c</sup> Jhones Luiz de Oliveira,<sup>c</sup> Eduardo de Almeida,<sup>a</sup> Alejandro Pérez-de-Luque,<sup>d</sup> Subhasis Ghoshal,<sup>e</sup> Catherine Santaella,<sup>f</sup> Renata de Lima,<sup>g</sup> Leonardo Fernandes Fraceto,<sup>b</sup> Hudson Wallace Pereira de Carvalho<sup>a,\*</sup>

Agriculture is responsible for supplying food, feed, fibres, and an increasing fraction of fuel and raw chemicals for industry. Fulfilling such demands sustainably is one of the major challenges of our time. In this metadata analysis, we offer a quantitative overview of how scientists have been addressing the effects of nanomaterials on plants between 2009 and 2022. The analysis showed that cultivated crops (55%) and plant nutrients (52%) are mostly employed in the studies, pointing to the relevance of these aspects to agriculture. Nevertheless, it also revealed that the concentration of elements as nanomaterials is, generally, more than 2-fold higher than the elemental concentration applied as traditionally formulated fertilisers or those naturally found in soil. Furthermore, the median time span of most studies, *i.e.*, 49 days for plants cultivated in soil, is still quite short compared to annual crop life cycles (90-120 days), and little attention (19% of treatments) has been devoted to soil microorganisms. Also, only a small fraction of experiments (6%) has been carried out under field conditions. Therefore, the data did not allow establishing correlations between effects and experimental parameters, such as concentration range, soil pH, or time of exposure. These observations point to the intricate relationship between our ability to infer conclusions and the experimental design employed. Finally, this comprehensive and up-to-date overview of the effects of nanomaterials in plant systems raises the question of whether nanomaterials will lead to incremental yield gains by replacing current inputs with nanotechnology-based ones, such as the controlled release of fertilizers and pesticides, or it will disrupt agriculture by attacking problems so far not practically addressed, such as hacking plant stress and defence mechanisms or modulating metabolism and photosystems.

### Agriculture under pressure

Agriculture is currently under a huge strain as the demand for food, feed, fibre, fuel, and biomass-based molecules rise.<sup>1,2</sup> It is expected that agricultural outputs must be increased by 70% in the next few decades,<sup>3</sup> yet shrinking farmlands make it difficult to support the demands on agriculture.<sup>4</sup>

Meeting higher agricultural yields implies overcoming many challenges, such as promoting nutrient use efficiency, soil protection, drought, salt and high-temperature tolerance, and

the management of several harms induced by nematodes, arthropods, and pathogenic fungi, bacteria, and viruses, as well as parasitic weeds.<sup>5-9</sup>

In this context, how could nanotechnology contribute to developing more sustainable and productive agriculture? Although nanosized particles (NPs) do naturally occur in the environment,<sup>10,11</sup> their many inherent properties, *e.g.*, larger contact surface, size and shape-dependent solubility, and surface charge alterability<sup>12,13</sup> are often claimed as beneficial to plant-based systems as it might play crucial roles in plant nutrition, disease management, and crop production.<sup>14</sup> NPs are also credited for driving better protection against biotic and abiotic stresses,<sup>15</sup> as well as enhanced photosynthetic efficiency,<sup>16,17</sup> controlled fertiliser's release,<sup>18-20</sup> and even the synthesis of doped-nanodevices for 'smart'-delivery of bioactive compounds and minerals to target tissues.<sup>21,22</sup>

In order to establish the current stage of nanomaterials-based research-driven toward plant systems, the present study brings a quantitative overview of previously published studies involving inorganic nanomaterials and plants. It includes 1154 peer-reviewed publications found in the Web of Science database between 2009 and 2022. The following keywords and booleans were employed: "*nanomaterial* OR *nanoparticle* OR

<sup>a</sup> University of São Paulo (USP), Centre of Nuclear Energy in Agriculture (CENA), Laboratory of Nuclear Instrumentation, Piracicaba, 13416-000, São Paulo, Brazil.

<sup>b</sup> Biology and Biotechnology Department "Charles Darwin" (DBBCD), Sapienza University of Rome, Rome, 00185, Italy.

<sup>c</sup> São Paulo State University (UNESP), Institute of Science and Technology, Sorocaba, 18087-180, São Paulo, Brazil.

<sup>d</sup> IFAPA Alameda del Obispo, Area of Genomic and Biotechnology, Avenida Menéndez Pidal, S/N, 14004, Córdoba, Spain.

<sup>e</sup> McGill University, Department of Civil Engineering, Montreal, Quebec, Canada H3A 0C3

<sup>f</sup> Aix Marseille Univ, CEA, CNRS, BIAM, Laboratory of Microbial Ecology of the Rhizosphere, F-13008 Saint-Paul-lès-Durance, France

<sup>g</sup> University of Sorocaba (Uniso), Bioactivity Assessment and Toxicology of Nanomaterials Lab (Labiton), Sorocaba, 18023-000, São Paulo, Brazil.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

nanotechnology" and "plant OR seed OR agriculture". Papers approaching nanoencapsulation of chemicals, e.g., pesticides and sensors, were not considered, since unbundling the reported effects as either a function of NPs or the loaded chemical is not straightforward (for a comprehensive review of this topic, please read Wang *et al.*<sup>14</sup>). Therefore, this systematic metadata survey aimed at answering nine broad questions:

1) Which nanomaterials/nanoparticles (NPs) have been tested?; 2) Which plant species have been exposed to NPs?; 3) What is the size range of the NPs?; 4) Did the studies with plants employ any kind of positive control?; 5) Where have the experiments been carried out?; 6) Which plant parts or organs have been exposed to NPs?; 7) How long plants have been exposed to NPs?; 8) What was the concentration range of NPs?; 9) What were the effect of nanomaterials on soil microorganisms?

Replying to these questions may give a glimpse into how nanomaterials-plant interactions have been addressed. Additionally, it might point out whether the knowledge generated allows us to conclude the real contribution of nanomaterials for optimized and sustainable agriculture.

Due to the variability of experimental strategies and the lack of a standardized procedure employed by the different authors, the survey presented here does not aim at making statistical analyses or extrapolations. We included papers strictly following the keywords and booleans criteria mentioned above and no intentional bias was exerted in the selection. Further details on the criteria used to select the manuscripts are provided in the electronic supplementary information (ESI). The raw data for the figures, the strategy for data mining, and the error/uncertainty estimation method are also provided in the ESI.

## Experimental design

Fig. 1 shows that most studies employed NPs as metal (25%) or metal oxides (54%), whereas 11% of the studies explored carbon-based NPs such as fullerenes, graphene, and carbon nanotubes. Among the metal oxides, ZnO, TiO<sub>2</sub>, CuO, and CeO<sub>2</sub> account for more than a third of all studied NPs. Moreover, Ag appears in 12% of them, outstanding as the most frequently employed metallic NP.

The analysed articles cluster NPs into four main branches: 1) Plant nutrient supply; 2) Stimulation of plant metabolism; 3) Mechanisms of NP uptake, transport, storage, and metabolism in plants; 4) Impacts and toxicity on plants. The survey identified the leading topic of interest of the scientific community, which is increasing plant productivity through nano-based fertilizers since 52% of the data entries employed plant nutrient NPs or also evaluated NP-driven crop protection.

However, a noteworthy fraction of those studies were either related to toxicity or the evaluation of NPs uptake. Although it encompasses crucial steps for determining the behaviour of the NPs, one must keep in mind that most of these studies do not relate to the practical application of NPs in cropping systems. In those studies, both NPs concentration and

experimental conditions explored were usually quite different from practical




Fig. 1 Most frequently employed NPs ( $n = 1456$  NPs) were found in 1154 original research articles published between 2009 and 2022. Each bar colour encompasses an NP group, namely, metal oxides, elemental ones, carbon-based, sulphides and sulphates, core-shells, and phosphate. Particles whose frequency was smaller than 1%, *i.e.*, nano-hydroxides, and mixed oxides, among others are not shown, thereby the summarisation might not reach 100%. The complete data and references are presented as a worksheet in Table S1 of the Electronic Supplementary Information, as well as in a data repository.<sup>23</sup> The estimated data uncertainty is *ca.* 4 %.

scenarios, and thus the obtained results might be difficult to be transposed to real field applications.


Another approach concerns the use of NPs made of beneficial or non-nutrient materials with stimulatory effects, *e.g.*, enhancing germination, plant development, nutrient translocation, and antioxidant responses.<sup>24-28</sup> ZnO, TiO<sub>2</sub>, CeO<sub>2</sub>, and Ag NPs are produced in large quantities by industry,<sup>29</sup> hence several studies aimed at evaluating whether these NPs contained in commercial goods, will eventually end up in the environment, thereby affecting plants.<sup>30</sup> One can highlight that ZnO and TiO<sub>2</sub> are present in sunscreen products and cosmetics, while Ag presents antimicrobial properties and is used in antimicrobial coatings, and many textiles, keyboards, wound dressings, and biomedical devices.<sup>31-37</sup>

Plants are fundamental components of ecosystems, and due to the characteristics of absorption and bioaccumulation, they are widely used as models for toxicity tests.<sup>38</sup> Furthermore, understanding the toxicity mechanisms of these NPs is an important point to better design them, either through component changes or even synthesis routes, to obtain systems with the least environmental side effects. It is worth noting that, for a redesign of these systems, an important aspect is that in addition to reducing the toxicity to non-target organisms and relevant environmental compartments, the nanomaterials must still have appropriate properties for their different applications.<sup>39</sup>

Since Ag, Au, TiO<sub>2</sub>, CeO<sub>2</sub>, and Al<sub>2</sub>O<sub>3</sub> are not found in plants at appreciable concentrations, their background signals are very low, easing their detection and making them suitable for mechanistic studies investigating the uptake, and transport, storage, and metabolism of NPs.

Most studies employed crop species responsible for producing staple food. Fig. 2 reveals that species from *Poaceae*,

*Fabaceae*, and *Solanaceae* families encompass *ca.* 55% of all plant species studied. According to the Food and Agriculture Organization of the United Nations, these families address *ca.*



**Fig. 2** Frequency of plant species and families found in 1154 original research articles published between 2009 and 2022, summing up an amount of 1374 plants from 248 species and 74 families. Those species and families encompassing frequencies smaller than 1% were not reported. The complete data and references are presented as a worksheet in a data repository.<sup>23</sup> The estimated data uncertainty is *ca.* 4%.

60% of all harvested areas in 2019.<sup>40</sup> Besides, a considerable fraction of the studies raise concerns about the NP toxic effects on crop plants and the introduction of NPs in the food chain. Here one may perceive an opportunity to carry out studies using perennial and arboreal plant species such as those producing coffee, orange, olives, apples, pears, and vines among others.


Fig. 3 reveals some aspects regarding the experimental design explored by the studies, *e.g.*, particle size, use of positive experimental controls (*i.e.*, the insoluble form of the same NP composition at micrometre range or a soluble compound containing the element of interest), and conditions of plant growth.

Since NP properties are intrinsically related to particle size, this is a piece of crucial information to be considered. Thereby, Fig. 3(a) shows the size range of individual NPs, usually determined by scanning or transmission electron microscopy, and the hydrodynamic diameter of NPs dispersed in liquid media, measured by light scattering techniques. The NPs size ranged from a few nanometres up to hundreds of nm, with an average of 45 nm, and a median size of 28 nm. It is worth mentioning that Auffan *et al.* argue that non-bulk properties arise mainly for diameters below 30 nm,<sup>41</sup> which calls into question the account of a real nano effect for many of these studies.

NPs tend to agglomerate depending on physical-chemical interactions, and the primary particle size assessed by electron microscopy does not necessarily reflect the actual size in which they are going to reach plant tissues. The hydrodynamic diameter median size was 174 nm, *ca.* 6-fold higher than the NP primary median size. The attempt of establishing a mathematical relationship between each size analysis did not return a clear correlation.

From an agronomic standpoint, either for nutritional or pest management, the NP size is a key parameter. It is well-known that the release of ions can be controlled as a function of

particle size since its dissolution rate is inversely proportional to the particle size.<sup>42-44</sup> Ideally, one could develop nanoparticles for which the dissolution rate could be controlled to match plant demand avoiding losses by volatilization, leaching, runoff,



**Fig. 3** (a) Boxplots of primary particle diameter and hydrodynamic diameter of NPs. The central line represents the median, whereas the black dots indicate the mean value. The estimated data uncertainty is *ca.* 5%. Noteworthy highlights are that from all the articles consulted, 20% did not provide any, or supplied incomplete information about NP primary size, while 63% did not report NPs hydrodynamic diameter. (b) Frequency of studies that reported the use of positive control, *i.e.*, either bulk or soluble compounds, at the same experiment as NPs, the estimated data uncertainty is *ca.* 3%. (c) Frequency of the environment in which the experiments were carried out, namely, ambient condition (without control of light and temperature factors), field, greenhouse, and growth chamber. The estimated data uncertainty is *ca.* 2%. In Fig. 3(a), *n* represents the number of treatments, whereas in Fig. 3(b) *n* indicates the number of papers, and in Fig. 3(c) *n* represents the number of experimental conditions. The complete data and references are presented as a worksheet public available in a data repository.<sup>23</sup>

or strong adsorption in soil colloids.

Size also determines whether entire particles will enter plant tissues, several size exclusions limits have been reported

1  
2  
3 varying from *ca.* 5 nm up to 50 nm.<sup>45-48</sup> However, some studies  
4 have demonstrated the uptake and translocation of  
5 nanomaterials with diameters greater than the size range  
6 stated above,<sup>47,49</sup> which suggests that the uptake of intact  
7 particles depends also on NP composition, surface charge,<sup>50</sup> and  
8 plant species. The ions released from NPs may be taken up by  
9 roots or leaves, depending on the dosing method, and then  
10 these ions might eventually, be precipitated within plants.  
11 Additionally, size and chemical reactivity can also be modified  
12 through the interaction of these nanomaterials with  
13 components of the environment, such as microorganisms and  
14 soil organic matter.<sup>51</sup> In this context, dealing with plants is even  
15 more challenging because roots can exude both protons,  
16 organic acids and siderophores that act as chelating agents,<sup>52</sup> as  
17 well as several secondary metabolites that might interact with  
18 the NPs.<sup>53</sup>

19 Fig. 3(b) shows that *ca.* 71% of the articles do not employ  
20 bulk or soluble ionic positive controls, whereas 15.1 % used  
21 soluble controls, 7.7% bulk control, and only 6.3% considered  
22 both soluble ionic and bulk controls. Although establishing  
23 soluble controls are not possible for several compounds, such  
24 as TiO<sub>2</sub> and carbon-based NPs, one should consider that these  
25 figures often challenge attributing the observed results to  
26 properties arising from the nano dimension of particles. For  
27 example, it is known that metallic nanomaterials can be  
28 transformed producing reactive ions able to bind into functional  
29 molecules in living organisms.<sup>54</sup> This leads to a myriad of  
30 harmful effects, such as destabilization of cell membrane  
31 integrity, alteration of cell osmotic balance, deactivation of  
32 enzymes, and reactive oxygen species (ROS) production, usually  
33 resulting in altered metabolism and cell damage.<sup>54,55</sup> Because of  
34 that, providing this kind of control could help interpret the  
35 effects of nanomaterials.<sup>54</sup> Nevertheless, several papers aimed  
36 at comparing the effects of surface charge, type of coating, or  
37 particle size on uptake properties, and in such cases positive  
38 controls might not be useful or necessary.

39 Fig. 3(c) shows the environment in which the experiments  
40 were carried out, revealing that 59% of them were performed  
41 in growth chamber conditions, whereas *ca.* 27% took place in  
42 greenhouses, and only a 6% fraction was conducted under field  
43 conditions. Experiments that could not be grouped in these  
44 categories were named "ambient condition"; they correspond  
45 to 8% of reports. As researchers working on this topic, we  
46 acknowledge the difficulties involved in setting up a safe  
47 experiment with nanomaterials in the field. On the other hand,  
48 to facilitate real-world agricultural applications, this barrier  
49 must be overcome by organizing controlled field trials, because  
50 those are equivalent to clinical trials in medicine for practical  
51 applications.<sup>56</sup>

52 Growth chambers offer controlled temperature, air  
53 moisture, light, and water irrigation, while greenhouses usually  
54 grant controlled temperature and irrigation. Most of the  
55 experiments carried out either in growth chambers or  
56 greenhouse are performed in pots or trays. Nevertheless, it may  
57 subject plants to root space confinement and increase the  
58 probability of interaction between plant tissues and NPs or their  
59 released components leading to results that would likely differ  
60

from those in field trials. A compromise between field and pot  
can be met by a mesocosms environment, which usually  
remains within a greenhouse with controlled external  
parameters, but at the same time, provides enough space for  
roots and allows NPs to diffuse across distinct environments.  
Prominent examples can be found in the studies by the groups  
of Lowry,<sup>57,58</sup> Wiesner<sup>59,60</sup> and Colman.<sup>61</sup>

Fig. 4(a), shows that 65% of the studies exposed plant root  
tissues to NPs, whereas only 17% and 14% exposed leaves and  
seeds, respectively. Within root exposure, 54% of them were  
carried out using growth solutions or agar-like media, which  
allow a great level of experimental control, but the results are  
hardly representative of soil. The preference for root exposure  
might arise from the fact that roots are the plant's organs  
designed for water and nutrient absorption, and hence it may  
unintentionally take up those NPs released into the  
environment. Nevertheless, although leaves are mainly  
responsible for transpiration and gas exchange, they can also  
absorb chemicals, including fertilizers and NPs.<sup>62-64</sup>

If NPs are to be employed as fertilizers, stimulants, or  
pesticides, their broadcast in the soil makes them more subject  
to losses. Additionally, the chemical interactions between NPs  
and soil colloids, such as clay and organic matter, and the effect  
of pH, can modify the surface properties which will influence  
the bioavailability and mobility of NPs.<sup>65</sup>

From a practical point of view, the application of NPs in  
agriculture through seed and foliar treatment would lead to the  
direct deposition of NPs on plants increasing the likelihood of  
assimilation. In this scenario, ZnO NPs applied to seeds enhance  
germination and increase root and shoot tissue length in maize,  
rice, and soybean plants.<sup>24,66,67</sup> Hence, the low number of  
studies evaluating this way of exposure points to an opportunity  
that can be seized in future studies. For example, coffee plants  
that received foliar sprays of *ca.* 68 nm ZnO NPs at 10 mg L<sup>-1</sup>  
and, after 40 days of treatment, presented 55% higher  
photosynthetic activity and 90% higher stomatal conductance  
compared to ZnSO<sub>4</sub> and control treatments.<sup>68</sup> Also, foliar  
application of Cu NPs at 250 mg L<sup>-1</sup> mitigated salt stress in  
tomatoes by improving the Na<sup>+</sup>/K<sup>+</sup> ratio. Besides, fruits of plants  
under salinity had their content of Cu, vitamin C, glutathione,  
and phenols significantly increased compared to controls  
through foliar exposure of Cu NPs.<sup>69</sup>

Fig. 4(b) shows that three-quarters of the studies carried out  
under hydroponics medium, and soil span up between 21 and  
90 days, respectively, with a 10- and 49-days median duration.  
The most frequently investigated crops shown in Fig. 2, require  
around three to five months to complete their cycles, *i.e.*, from  
sowing to harvesting. Here one could point to a gap, and hence  
an opportunity for long-term studies on the uptake of  
biotransformation of NPs. Considering the putative potential of  
nanotechnology to increase agricultural productivity, we should  
also focus on evaluating the impact of NPs on crop yield. Last,  
but not least, in the middle of a debate on whether the nutrient  
content of agricultural outputs has been declining compared to  
the past because of higher yields,<sup>70,71</sup> we should also consider  
evaluating more frequently the effect of NPs on the quality of  
vegetal products.<sup>72</sup>

Another parameter that might influence the results is the concentration range of NPs applied, as shown in Fig. 4(c). The median concentration of NPs dispersed in the liquid was 50 mg L<sup>-1</sup> while the average concentration was 644 mg L<sup>-1</sup>. In a general-

values presented in Fig. 4(c) correspond to those NPs in which most of the elements are not supposed to be dissolved. It is noteworthy that the median concentration range of NPs dispersed in liquid media exceeds more than 10-fold that of essential micronutrients.

In regular hydroponics, the nutrients are depleted from the solution as they are absorbed by plants. Conversely, in a colloidal NP dispersion, one could expect a more constant concentration level due to equilibrium between solid and aqueous phases. Several studies indicate the fraction of dissolved ions in ZnO dispersions may vary from *ca.* 4 up to 9%,<sup>44,74</sup> while for CuO these values reach *ca.* 0.1 up to 1.1%,<sup>75,76</sup> and for CeO<sub>2</sub> they range from <0.1% up to 0.9%.<sup>77,78</sup> Nevertheless, the fraction of dissolved ions depends not only on the chemical composition of the NPs but also on the pH and chelating molecules present in the solution.<sup>79</sup> Among the surveyed papers, 103 out of 740 reported NPs tested at concentrations below 1 mg L<sup>-1</sup>, *ca.* 27% of these studies investigated treatments based on silver or gold, while *ca.* 40% employed Mn, Cu, Zn, and Fe plant micronutrients based nanomaterials.


The range employed in soil was higher than in hydroponics, displaying median and average NPs of 125 and 989 mg kg<sup>-1</sup>, respectively. The availability of nutrients depends on soil features, *i.e.*, clay fraction content, pH, organic matter content, and cation exchange capacity. Therefore, the adequate available concentration varies on soil texture, for phosphorus are slightly above 10 mg kg<sup>-1</sup>, and higher than 50, 600, 2.5-5, 0.8-3, and 0.5-2 mg kg<sup>-1</sup> for K, Ca, Fe, Zn, and Cu, respectively.<sup>80</sup>

From an environmental safety standing point, Reimann *et al.*<sup>81</sup> determined the HNO<sub>3</sub>/HCl/H<sub>2</sub>O extractable concentration of 53 elements in 2108 samples collected in agricultural and grazing lands of 33 European countries. They found that geochemical threshold values, *i.e.*, concentrations unusually high based on the value of the 98<sup>th</sup> percentiles of the sampling, for Zn, Ag, Cu, and Ce were 129, 0.19, 78, and 81 mg kg<sup>-1</sup>, respectively. These elements comprise *ca.* 57% of the treatments in which NPs were dispersed into soils, sand or substrate, the median concentration of Zn, Ag, Cu, and Ce in these studies were 200 (*n* = 264), 31 (*n* = 99), 100 (*n* = 106), and 250 (*n* = 140) mg kg<sup>-1</sup>, respectively (*n* corresponds to the number of treatments).

The amount of zinc employed in conventional fertilization depends on the crop and soil features, but it usually remains between a few to up to a dozen kilograms per hectare.<sup>82,83</sup> Liu *et al.*<sup>83</sup> obtained high maize yield by applying Zn at a rate of 11.3 kg ha<sup>-1</sup> (3.83 mg kg<sup>-1</sup> of Zn in soil, considering a typical 0.2 m deep soil layer and 1.5 kg dm<sup>-3</sup> soil density).

To evaluate the potential gains offered by nanotechnology for fertilization, experiments aiming at replacing conventional bulk particulate or soluble fertilizers with nanomaterials should employ realistic field concentrations such as those above reported.

Thus, one should keep in mind that the nanosized properties of particles might have a crucial role in understanding the harmful effects of NPs. As well, the effects of high exposure concentrations should be accounted for. This puzzle could be



**Fig. 4** (a) Plant part or plant organ exposed to NPs, and the media employed in case of root exposure, the estimated data uncertainty is *ca.* 4%. (b) Boxplots presenting the exposure time of plants to NPs either dispersed in liquid solutions or soil. Herein, the estimated data uncertainty is *ca.* 5%. (c) Boxplots of NPs concentration range either dispersed in liquid or solid, the estimated data uncertainty is *ca.* 4%. Centrelines represent the median, whereas the dots indicate the mean values in b and c. In Fig. 4(a) *n* represents the number of experiments, in Fig. 4(b) *n* indicates the number of exposure times described by the studies, and Fig. 4(c) *n* represents the number of treatments reported by the manuscripts. The complete data and references are presented as a worksheet in a data repository.<sup>23</sup>

purpose solution for plant growth, essential macroelements such as potassium, calcium, and phosphorus are found at 235, 160, and 62 mg L<sup>-1</sup>, respectively. On the other hand, micronutrients such as zinc, copper, and iron are present at 0.13, 0.03, and 1-3 mg L<sup>-1</sup>, respectively.<sup>73</sup> These concentrations correspond to the number of available nutrients, while the



more assertively solved if the results are backed up by adequate negative and positive experimental controls.

## Effects of nanomaterials on plants and soil microorganisms

Plant's morphology and physiology can be impacted by NPs exposure as a function of the NP type, size, concentration, and exposure time.<sup>49,84,85</sup> Aiming at making the comparison on equal terms, Fig. 5(a) presents only data extracted from papers in which nanomaterials concentrations were reported on a weight/weight basis and roots were exposed to the treatments through soil/substrate or sand. It encompasses a total of 1100 treatments reported in 213 original articles. The myriad of plant responses evaluated in these studies was classified and grouped as 'harmful', 'beneficial', and 'beneficial and harmful', the latter comprising treatments that concomitantly presented both positive and negative impacts. Studies that did not observe any changes at all were classified as 'no effect', while those that did not evaluate plant morphological and physiological responses


were represented as 'not applicable'. Some of the plant responses reported by the authors were changes in plant height, root and shoot tissue length, fruit numbers and weight, crop yield, leaf area, protein, lycopene, b-carotene, carotenoids, chlorophyll *a* and *b* contents, and photosynthetic attributes, whose values higher than the control, statistically or suggested by the authors, were considered as beneficial effects. Consequently, their lower values were considered harmful ones. For other features, though, lower values indicate beneficial effects, and higher, in turn, harmful ones (*e.g.*, fruiting and flowering time). Particularly, the increase of antioxidant enzyme activities, such as POD, APX, CAT, SOD was considered beneficial by some studies and harmful by others. Since these results might reflect either enhanced or overwhelmed plant responses to biotic and abiotic stresses, we followed the interpretation given by the authors.

Among the treatments reported in Fig. 5(a), 34.7% resulted in harmful effects, 30.5% pointed out beneficial outcomes, 18.5% presented both harmful and beneficial consequences, 16.3% did not report changes compared to untreated plants, and 15% did not evaluate the effects of NPs on plants.

Notable NP effects on several aspects of agricultural-related research have been reported in recent years. Regarding biofortification, wheat seeds primed with ZnO NPs at 20-1000 mg kg<sup>-1</sup> boosted the Zn content of the harvested grains. Compared with control, the concentration of Zn in grains was up to 3.3-fold higher and ZnO NPs were more effective than ZnSO<sub>4</sub>. In addition to that, at moderate doses (20 mg kg<sup>-1</sup>) biomass and grain yield were also higher.<sup>25</sup>

Besides, some NP-related photocatalytic properties are claimed to benefit plant light-dependent reactions. In this scenario, root ZnO and foliar Fe-based (Fe and Fe<sub>3</sub>O<sub>4</sub>) NPs exposure have also been reported to significantly increase the net photosynthetic rate of maize and tomato, leading to improve plant growth in both species.<sup>86,87</sup> Zinc oxide, Si, Mn, and TiO<sub>2</sub> NPs exposure have also induced higher tolerance either to drought or salinity stress in mango, cucumber, and wheat plants, which might be related to endogenous NPs-driven boosting of plant's antioxidant enzymes.<sup>88-90</sup> Wheat plants grown in Cd-contaminated soils negatively affected photosynthesis yield, and also caused oxidative stress damage in leaves. However, soil application of Fe NPs at 25-100 mg kg<sup>-1</sup> not only decreased Cd content in plant tissues but also increased photosynthesis, growth, and yield parameters.<sup>91</sup>

It is worth mentioning that exposing plant roots to NPs might also influence soil microbial communities. Soil microorganisms play pivotal roles in maintaining soil quality since they carry out the decomposition of organic matter, assist plants in water and nutrient uptake,<sup>92</sup> and fix most of the nitrogen used by legumes<sup>93</sup> and a minor part of nitrogen employed by cereals.<sup>94</sup> Hence, microorganisms living in the rhizosphere improve plant development, yield, and disease resistance.<sup>95</sup> In this scenario, any possibility of NP-induced



**Fig. 5** (a) Frequency of reported effects of NPs on plants whose roots were exposed to the treatments in solid media, *i.e.*, soil or sand. 'Beneficial & harmful' effect comprises treatments reporting overlapped positive and negative effects, whereas the 'No effect' refers to those reporting no changes compared to the negative control group, and 'Not applicable' encompasses studies/treatments that did not evaluate plant biological responses as an effect of NPs exposure. The estimated data uncertainty is *ca.* 7%. (b) Frequency of treatments that reported effects of NPs on soil microbiota. The double-checking did not return a measurable error. In both figures *n* refers to the number of

disruption of these ecological services must be a matter of concern.<sup>96</sup>

Despite their importance, Fig. 5(b) shows that the effect of NPs on soil microorganisms was assessed in only 19.2% of the treatments reported in 41 out of the 213 manuscripts assessed. This is a matter of great concern since the soil is expected to be the major sink of the nanomaterials released into the environment.<sup>97</sup> Nevertheless, 28% of the treatments that assessed this parameter declared the soil microbial community did not change, while 35% reported that either richness or abundance of microorganisms decreased, followed by 15% that increased, and finally 23% reported both increase and decrease in soil microbiota. In this latter case, some papers reported modifications in the structure of the community. According to Simonin & Richaume<sup>98</sup> inorganic NPs (metal and metal oxide) are usually more detrimental to soil microorganisms than organic ones, *e.g.*, fullerenes and carbon nanotubes.

Several studies associate the diversity and abundance of soil microorganisms with the applied NP concentrations. Albeit another key issue relies on the fact that the plant's developmental stages seem to interfere with microbial diversity in NP-amended soils. Carbon nanotubes at 0.1 and 100 mg kg<sup>-1</sup> altered soybean rhizosphere prokaryotic community structure more intensely during the reproductive stage than the vegetative one,<sup>99</sup> whereas Ag NPs at 1 mg kg<sup>-1</sup> showed a transient impact on the structure of the bacterial community during the transition from the seedling to the vegetative stage of wheat, but with a recovery to normal levels 49 days post-treatment.<sup>100</sup>

Several NPs are also known to exhibit antimicrobial effects, their effects on plant pathogens have also been investigated. For example, Cu<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>·3H<sub>2</sub>O nanosheets and CuO NPs resulting in suppressed root fungal disease induced by *Fusarium oxysporum* f. sp. *niveum* fungal in watermelon plants.<sup>101</sup> Moreover, soybean plants inoculated with *Fusarium virguliforme* and foliar sprayed with CuO, B, MoO<sub>3</sub>, or ZnO NPs presented a significant reduction of root rot severity involved in the sudden plant death syndrome.<sup>102</sup> Although the current work did not explore nano-encapsulated formulations, a recent 500-paper review by Wang *et al.*<sup>14</sup> estimates that the overall efficacy of nanopesticides is *ca.* 32% higher compared to bulk-scale ones.


Moreover, *Medicago truncatula* plants cultivated in soil containing 1450, 100, or 2400 mg kg<sup>-1</sup> of Zn, Ag, or Ti NPs, respectively had more significant negative impacts on the microbial abundance in soils more significantly than those with bulk/dissolved treatments.<sup>103</sup> Conversely, Ag<sub>2</sub>S NPs were less harmful to tomato mycorrhizal symbiosis and the soil microbial community than Ag<sup>+</sup>.<sup>104</sup>

In some cases, soil microorganisms might mitigate nanoparticle toxicity to plants. Mycorrhizal fungi colonization in tomatoes cultivated in Ag NPs spiked soil alleviated nanomaterial-induced phytotoxicity by decreasing Ag accumulation in plant tissues.<sup>105</sup> Similarly, the root colonization rate significantly decreased when exposed to ZnO NPs at and above 800 mg kg<sup>-1</sup>, but despite that arbuscular mycorrhizal fungi

mitigated phytotoxic effect by decreasing Zn bioavailability and accumulation in maize.<sup>106</sup>

Plant responses depend on a complex parameter network. The box plots presented in Fig. 6 correlate the effects observed in plants with the concentration of NPs applied into the soil as a function of NPs concentration (a), soil pH (b), and time of exposure (c). Although the highest mean value of the concentration of NPs in the soil comes from treatments that pointed out beneficial effects in plants (*ca.* 1070 mg kg<sup>-1</sup>), median concentration values of the harmful (400 mg kg<sup>-1</sup>) and beneficial & harmful (200 mg kg<sup>-1</sup>) effects are higher than the benefits and no effect (both at 100 mg kg<sup>-1</sup>) (Fig. 6(a)).

As mentioned above, Auffan *et al.*<sup>41</sup> highlight that those non-bulk properties arise mainly for NPs displaying a < 30 nm diameter. Hence, to verify whether this threshold would lead to effects, the data was split into NPs ≥ 30 nm and ≤ 29 nm, then



**Fig. 6** Boxplots of soil-amended NPs concentration (a, n = 992 treatments from 192 articles), soil pH (b, n = 714 treatments from 131 articles), and NPs exposure time (c, n = 1071 treatments from 195 articles) as function of the effects reported. Centrelines point out the median, whereas the blue dots sign the mean values. Due to the high number of variables, the uncertainty values were not recorded. Data computed from original research articles published between 2009 and 2022. The complete data and references are presented as a worksheet public available in a data repository.<sup>23</sup>

the frequency of treatments resulting in beneficial, harmful, beneficial & harmful, and no effect was counted. Supplementary Fig. S1 shows that the type of effect cannot be consistently associated with the 30 nm edge. Figure S2 presents a boxplot showing the size, informed by manufacturer, TEM, and DLS, and the effects, it neither suggests any correlation between the 30 nm size boundary and effects.

Besides, the frequency effects on botanical families and species were also evaluated. The most frequently employed families were Asteraceae, Brassicaceae, Cucurbitaceae, Fabaceae, Poaceae, and Solanaceae (Fig. S3), and the species were *Glycine max*, *Triticum aestivum/durum*, *Zea mays*, and *Solanum lycopersicum* (Fig. S4). A higher frequency of the harmful effects was observed in Brassicaceae and Fabaceae families, whereas the beneficial ones were more noticeable in Asteraceae, Cucurbitaceae, Poaceae, and Solanaceae (Fig. S3). Curiously, a similar trend was observed for *Glycine max* and *Triticum aestivum/durum*, whereas beneficial and both beneficial & harmful ones were seen for *Zea mays* and *Solanum lycopersium* (Fig. S4).

Furthermore, to reduce the dimensionality of the continuous variables explored as a function of the effects, *i.e.*, concentration, NP size, soil pH and exposure time, a principal component analysis (PCA) was applied to observe their possible relationships with the effects (Fig. S5). The PCA analyses did not show any clustering of effects and the first two principal components. This information corroborates with Fig. 6 which shows similar amplitudes for the boxplots of concentration.

In other words, harmful effects might be generally associated with higher doses and vice-versa, however, due to lack of experimental standardization, it does not necessarily configure a cause-effect relationship.

In a range between 50 and 250 mg kg<sup>-1</sup>, Fe<sub>3</sub>O<sub>4</sub> and TiO<sub>2</sub> NPs boosted phosphorus uptake, shoot and root length, and weight of lettuce.<sup>107</sup> At 500 mg kg<sup>-1</sup> CeO<sub>2</sub> NPs increased barley height by 34% and shoot biomass by 330% compared with control, but the plants did not form grains.<sup>108</sup>

Besides concentration, soil characteristics play an important role in plant growth. Particularly, soil pH is one of the main properties which influence microorganism activity<sup>109</sup> as well as nutrient solubility and availability.<sup>110</sup> The box plot shown in Fig. 6(b) indicates that the mean and median values of soil pH were almost the same for all the treatments, close to neutral pH values, which suggests that soil pH might not be correlated with the effects caused by the nanomaterials.

Another important parameter that was not assessed in this data review was the effect of soil organic matter. The presence of humic acids and organic matter can alter the bioavailability of the nanomaterials in soils, as well as the presence of salts or changes of pH that result in the loss of nanomaterials stability.<sup>46</sup>

Small differences were found in the time of exposure (Fig. 6(c)). The median time of exposure in which nanomaterials caused harmful effects to plants was longer than the time of exposure reported for beneficial effects by a small margin (48 and 45 days, respectively). Beneficial and harmful effects presented a median time of exposure of 56 days, while the treatments that did not affect the plants lasted a median of 60 days.

Altogether, the data presented in Fig. 6 lead us to conclude that positive and/or negative effects as a response to nanomaterials treatment are not solely a consequence of the applied concentration, soil pH, or time of exposure, it seems to be more correlated with the nanomaterial type and plant species.

### What to expect for the future?

The properties exhibited by the nanomaterials may contribute to increasing both yield and the quality of the agricultural outputs. In principle, engineered NPs can be targeted for soil amendments, seed coating, and foliar application, pointing to better nutrient use efficiency or crop protection compared to conventional sources.

In this scenario, the one-decade metadata revision herein presented is aimed at understanding the current stage of nanomaterials-based research toward plant systems by quantifying which NPs, and their respective particle sizes and concentrations, have been applied to which plant species. Besides, the experimental conditions, *e.g.*, cultivation environment and NPs exposure way and timeframe, as well as the use of control, were also considered.

Overall, the big picture presented by the quantified results does reveal an urgent need for improving the description of the experimental procedures. Regarding the experimental settings, a high number of NPs assays, frequently used in toxicity studies, must be supported by positive controls as it otherwise challenges establishing a clear correlation between results and NPs exposure.

Furthermore, the size of NPs employed through multiple measurement techniques that capture both the primary particle size as well as agglomerate sizes in the dosing solutions must be included in future studies. The NP suspension media and the surface chemistry of the NPs should be characterized, and controls with the associated agents but with the nanomaterials be performed to allow for the determination of the active agents responsible for the effects observed. Last, but not least, addressing long-term experiments with NPs utilising perennial plants is of the utmost importance.

One should keep in mind that replacing conventional fertilizers with nanosized ones requires experiments employing compatible NP concentration ranges and long exposure times, as the impacts on plant productivity from a commercial perspective, *e.g.*, quantity and quality of edible parts, do need to be reported in future studies. Additionally, the costs involved in transforming either soluble salts or bulk particles into NPs might constitute a barrier unless the use-efficiency of NPs pays it off. Within this framework, it is also crucial to consider the

climate-change-driven menaces to crop production, and in relation to this, explore both biotic and abiotic factors, including soil amendments. Thereby, experiments in large mesocosms and field conditions are a key step that will contribute to fostering and mature nanotechnological-based solutions for agriculture.

Furthermore, the effects of NPs on soil microorganisms must also be more frequently addressed, and specific tests assuring that any commercial technology is innocuous to the soil microbiota should be demanded by regulators. Likewise, concentration thresholds for NPs in food and feed should be established aiming at avoiding possible threats to human and animal health.

The growth rate of agricultural outputs during the 18<sup>th</sup> century raised so much concern that it led to the statement of the "Malthusian Catastrophe". However, the observers at that time could not forecast that the development and adoption of disruptive technologies, made possible by science, would promote the many-fold gains in productivity during the 20<sup>th</sup> century. Perhaps, with the contribution of nanotechnology, the next productivity increase leap will come not by replacing current agricultural inputs, but by creating new ones. Sensing plant metabolism<sup>111</sup> and quickly modulating it, hacking plant photosystems,<sup>112</sup> and stimulating plant functions are some examples of such technologies that will soon be available.

The use of nanotechnologies is often blamed by the literature and media, even though plants naturally take nutrients in the form of colloids and probably NPs.<sup>113</sup> Nevertheless, the summarized conclusions of the studies herein assessed revealed that 50% have employed NPs that exceed the critical nano size (100 nm), as well as concentrations often unrelated to the appropriate agricultural intentions. Despite these shortcomings, 39% of the studies show beneficial effects or no effects at all, against 30% that show negative ones. These negative effects are not often compared to real stresses experienced by plants in the field, leading to the impression that we still don't have a good knowledge of the properties of NPs, under optimized conditions of use.

Altogether, these elements suggest that nanotechnology for agriculture deserves to be further explored. By incorporating the efforts of this very first decade of studies exploring the effects of nanosized materials in plant development, it is now possible to establish enhanced experimental protocols, *i.e.*, including suitable NPs characterisation, reasonable NPs concentrations, and longer exposure times, that will enable properly addressing the real contribution of nanomaterials-based technologies for more productive and sustainable agriculture.

## Author Contributions

This study was conceived by H.W.P.C., M.H.F.G., L.F.F., R.L., C.S., and S.G. The data collection was carried out by E.S., G.S.M., N.M.D., C.G.C., S.Z.R., M.H.F.G., and E.A. Data checking was carried out by H.W.P.C, E.S., G.S.M., N.M.D., C.G.C., S.Z.R., E.A., J.L.O., and A.ES.P. Data discussion and interpretation were performed by H.W.P.C, G.S.M., A.P.L., R.L., S.G, L.F.F, and C.S.

The first manuscript draft was written by H.W.P.C, G.S.M, N.M.D., J.L.O., and A.ES.P. All authors edited the manuscript.

## Data availability statement

The complete data used to the production of all figures has been deposited at Figshare and can be obtained from <https://doi.org/10.6084/m9.figshare.14575185.v1>

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

The authors are grateful to the São Paulo Research Foundation (grants 2020/07721-9, 2020/11178-9, 2020/11546-8, 2017/16375-4, 2015-05942-0, 2016/50014-6, and 2017/21004-5), Brazilian National Council for Scientific and Technological (CNPq) (grants 407380/2016-7 and 306185/2020-2), Agrichem do Brasil S.A. (FEALQ#10245-8), and Vittia Fertilizantes e Biológicos LTDA (FEALQ 10359-8). The authors are also grateful to Dr. Tiago R. Tavares for his assistance during multivariate data analysis.

## References

- 1 L. G. Horlings and T. K. Marsden, Towards the real green revolution? Exploring the conceptual dimensions of a new ecological modernisation of agriculture that could 'feed the world', *Global Environ. Chang.*, 2011, **21**, 441-452.
- 2 H. C. J. Godfray, J. R. Beddington, I. R. Crute, L. Haddad, D. Lawrence, J. F. Muir, J. Pretty, S. Robinson, S. M. Thomas and C. Toulmin, Food security: The challenge of feeding 9 billion people, *Science*, 2010, **327**, 812-818.
- 3 FAO, *How to Feed the World in 2050*, FAO, Rome, 2009.
- 4 FAO, *The future of food and agriculture – Alternative pathways to 2050*, FAO, Rome, 2018.
- 5 D. Tilman, K. G. Cassman, P. A. Matson, R. Naylor and S. Polasky, Agricultural sustainability and intensive production practices, *Nature*, 2002, **418**, 671-677.
- 6 D. Tilman, C. Balzer, J. Hill and B. L. Befort, Global food demand and the sustainable intensification of agriculture, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 20260-20264.
- 7 F. E. Dayan, C. L. Cantrell and S. O. Duke, Natural products in crop protection, *Bioorg. Med. Chem.*, 2009, **17**, 4022-4034.
- 8 A. Van den Putte, G. Govers, J. Diels, K. Gillijns and M. Demuzere, Assessing the effect of soil tillage on crop growth: A meta-regression analysis on European crop yields under conservation agriculture, *Eur. J. Agron.*, 2010, **33**, 231-241.
- 9 F. Shah and W. Wu, Soil and crop management strategies to ensure higher crop productivity within sustainable environments, *Sustainability*, 2019, **11**, 1485.
- 10 G. Lespes, S. Faucher and V. I. Slaveykova, Natural nanoparticles, anthropogenic nanoparticles, where is the frontier?, *Front. Environ. Sci.*, 2020, **8**, 71.
- 11 S. Griffin, M. I. Masood, M. J. Nasim, M. Sarfraz, A. P. Ebokaiwe, K. H. Schafer, C. M. Keck and C. Jacob, Natural

- nanoparticles: A particular matter inspired by nature, *Antioxidants*, 2018, **7**, 3.
- 12 I. Khan and K. Saeed, Nanoparticles: Properties, applications and toxicities, *Arabian J. Chem.*, 2019, **12**, 908-931.
  - 13 E. Hinde, K. Thammasiraphop, H. T. T. Duong, J. Yeow, B. Karagoz, C. Boyer, J. J. Gooding and K. Gaus, Pair correlation microscopy reveals the role of nanoparticle shape in intracellular transport and site of drug release, *Nat. Nanotechnol.*, 2017, **12**, 81-89.
  - 14 D. J. Wang, N. B. Saleh, A. Byro, R. Zepp, E. Sahle-Demessie, T. P. Luxton, K. T. Ho, R. M. Burgess, M. Flury, J. C. White and C. M. Su, Nano-enabled pesticides for sustainable agriculture and global food security, *Nat. Nanotechnol.*, 2022, **17**, 347-360.
  - 15 M. R. Khan, F. Ahamad, T. F. Rizvi, M. Ghorbanpour and S. H. Wani, Effect of nanoparticles on plant pathogens, in *Advances in Phytanotechnology*, ed. M. Ghorbanpour and S. H. Wani, Academic Press, San Diego, 2019, Chapter 8, pp. 215-240.
  - 16 T. A. Swift, T. A. A. Oliver, M. C. Galan and H. M. Whitney, Functional nanomaterials to augment photosynthesis: evidence and considerations for their responsible use in agricultural applications, *Interface Focus*, 2018, **9**, 20180048.
  - 17 R. Tighe-Neira, E. Carmora, G. Recio, A. Nunes-Nesi, M. Reyes-Diaz, M. Alberdi, Z. Rengel and C. Inostroza-Blancheteau, Metallic nanoparticles influence the structure and function of the photosynthetic apparatus in plants, *Plant Physiol. Biochem.*, 2018, **130**, 408-417.
  - 18 M. H. F. Gomes, B. A. Machado, E. S. Rodrigues, G. S. Montanha, M. L. Rossi, R. Otto, F. S. Linhares and H. W. P. Carvalho, In vivo evaluation of Zn foliar uptake and transport in soybean using X-ray absorption and fluorescence spectroscopy, *J. Agric. Food Chem.*, 2019, **67**, 12172-12181.
  - 19 P. M. Kopittke, E. Lombi, P. Wang, J. K. Schjoerring and S. Husted, Nanomaterials as fertilizers for improving plant mineral nutrition and environmental outcomes, *Environ. Sci.: Nano*, 2019, **6**, 3513-3524.
  - 20 K. R. Rakhimol, S. Thomas, N. Kalarikkal and K. Jayachandran, Nanotechnology in controlled-release fertilizers, in *Controlled Release Fertilizers for Sustainable Agriculture*, ed. F. B. Lewu, T. Volova, S. Thomas and K. R. Rakhimol, Academic Press, San Diego, 2021, Chapter 10, pp. 169-181.
  - 21 G. B. Ramirez-Rodriguez, C. Miguel-Rojas, G. S. Montanha, F. J. Carmona, G. Dal Sasso, J. C. Sillero, J. S. Pedersen, N. Masciocchi, A. Guagliardi, A. Perez-de-Luque and J. M. Delgado-Lopez, Reducing nitrogen dosage in *Triticum durum* plants with urea-doped nanofertilizers, *Nanomaterials*, 2020, **10**, 1043.
  - 22 E. P. Perez-Alvarez, G. B. Ramirez-Rodriguez, F. J. Carmona, J. M. Martinez-Vidaurre, N. Masciocchi, A. Guagliardi, T. Garde-Cerdan and J. M. Delgado-Lopez, Towards a more sustainable viticulture: foliar application of N-doped calcium phosphate nanoparticles on Tempranillo grapes, *J. Sci. Food Agric.*, 2021, **101**, 1307-1313.
  - 23 G. Montanha, 2021, METADATA: Are nanomaterials leading to more efficient agriculture? Outputs from 2009-2022 research metadata analysis, Figshare, DOI: 10.6084/m9.figshare.14575185.v1
  - 24 G. S. Montanha, E. S. Rodrigues, J. P. R. Marques, E. de Almeida, M. Colzato and H. W. P. de Carvalho, Zinc nano-coated seeds: an alternative to boost soybean seed germination and seedling development, *SN Appl. Sci.*, 2020, **2**, 857.
  - 25 W. Du, J. Y. Yang, Q. Q. Peng, X. P. Liang and H. Mao, Comparison study of zinc nanoparticles and zinc sulphate on wheat growth: From toxicity and zinc biofortification, *Chemosphere*, 2019, **227**, 109-116.
  - 26 R. C. Marucci, L. M. Freitas, J. R. Santos-Rasera, D. S. Alves, G. A. Carvalho and H. W. P. de Carvalho, Are cerium oxide nanoparticles transferred from plants to the aphid *Myzus persicae* (Hemiptera: Aphididae)?, *Fla. Entomologist*, 2019, **102**, 555-561.
  - 27 E. Spielman-Sun, A. Avellan, G. D. Bland, R. V. Tappero, A. S. Acerbo, J. M. Unrine, J. P. Giraldo and G. V. Lowry, Nanoparticle surface charge influences translocation and leaf distribution in vascular plants with contrasting anatomy, *Environ. Sci.: Nano*, 2019, **6**, 2508-2519.
  - 28 L. Korosi, S. Boudérias, K. Csepregi, B. Bognar, P. Teszlak, A. Scarpellini, A. Castelli, E. Hideg and G. Jakab, Nanostructured TiO<sub>2</sub>-induced photocatalytic stress enhances the antioxidant capacity and phenolic content in the leaves of *Vitis vinifera* on a genotype-dependent manner, *J. Photochem. Photobiol., B*, 2019, **190**, 137-145.
  - 29 M. E. Vance, T. Kuiken, E. P. Vejerano, S. P. McGinnis, M. F. Hochella, D. Rejeski and M. S. Hull, Nanotechnology in the real world: Redeveloping the nanomaterial consumer products inventory, *Beilstein J. Nanotechnol.*, 2015, **6**, 1769-1780.
  - 30 S. Maiti, E. El Fahime, M. Benaissa and S. Kaur Brar, Nanotoxicology of natural and engineered nanoparticles for plants, in *Nanomaterials in the Environment*, ed. S. Kaur Brar, T. C. Zhang, M. Verma, R. Y. Surampalli and R. D. Tyagi, American Society of Civil Engineers, Reston, 2015, Chapter 18, pp. 469-485.
  - 31 A. Krol, P. Pomastowski, K. Rafinska, V. Railean-Plugaru and B. Buszewski, Zinc oxide nanoparticles: Synthesis, antiseptic activity and toxicity mechanism, *Adv. Colloid Interface Sci.*, 2017, **249**, 37-52.
  - 32 P. K. Mishra, H. Mishra, A. Ekielski, S. Talegaonkar and B. Vaidya, Zinc oxide nanoparticles: a promising nanomaterial for biomedical applications, *Drug Discovery Today*, 2017, **22**, 1825-1834.
  - 33 S. Vijayakumar, B. Vaseeharan, B. Malaikozhundan and M. Shobiya, Laurus nobilis leaf extract mediated green synthesis of ZnO nanoparticles: Characterization and biomedical applications, *Biomed. Pharmacother.*, 2016, **84**, 1213-1222.
  - 34 M. S. Waghmode, A. B. Gunjal, J. A. Mulla, N. N. Patil and N. N. Nawani, Studies on the titanium dioxide nanoparticles: biosynthesis, applications and remediation, *SN Appl. Sci.*, 2019, **1**, 310.
  - 35 D. Ziental, B. Czarczynska-Goslinska, D. T. Mlynarczyk, A. Glowacka-Sobotta, B. Stanisz, T. Goslinski and L. Sobotta, Titanium dioxide nanoparticles: prospects and applications in medicine, *Nanomaterials*, 2020, **10**, 387.
  - 36 H. B. Shi, R. Magaye, V. Castranova and J. S. Zhao, Titanium dioxide nanoparticles: a review of current toxicological data, *Part. Fibre Toxicol.*, 2013, **10**, 15.
  - 37 B. Calderon-Jimenez, M. E. Johnson, A. R. M. Bustos, K. E. Murphy, M. R. Winchester and J. R. V. Baudrit, Silver Nanoparticles: Technological Advances, Societal Impacts, and Metrological Challenges, *Front. Chem.*, 2017, **5**, 6.
  - 38 A. Fatima, S. Singh and S. M. Prasad, in *Sustainable Agriculture Reviews 41: Nanotechnology for Plant Growth and Development*, ed. S. Hayat, J. Pichtel, M. Faizan and Q. Fariduddin, Springer, Cham, 2020, Chapter 8, pp. 143-161.
  - 39 J. T. Buchman, N. V. Hudson-Smith, K. M. Landy and C. L. Haynes, Understanding nanoparticle toxicity mechanisms to inform redesign strategies to reduce environmental impact, *Acc. Chem. Res.*, 2019, **52**, 1632-1642.
  - 40 FAOSTAT, Food and Agriculture Organization of the United Nations, <http://www.fao.org/faostat>
  - 41 M. Auffan, J. Rose, J. Y. Bottero, G. V. Lowry, J. P. Jolivet and M. R. Wiesner, Towards a definition of inorganic nanoparticles from an environmental, health and safety perspective, *Nat. Nanotechnol.*, 2009, **4**, 634-641.
  - 42 C. A. David, J. Galceran, C. Rey-Castro, J. Puy, E. Companys, J. Salvador, J. Monne, R. Wallace and A. Vakourov, Dissolution

- kinetics and solubility of ZnO nanoparticles followed by AGNES, *J. Phys. Chem. C*, 2012, **116**, 11758-11767.
- 43 T. N. M. da Cruz, S. M. Savassa, G. S. Montanha, J. K. Ishida, E. de Almeida, S. M. Tsai, J. Lavres and H. W. P. de Carvalho, A new glance on root-to-shoot in vivo zinc transport and time-dependent physiological effects of ZnSO<sub>4</sub> and ZnO nanoparticles on plants, *Sci. Rep.*, 2019, **9**, 10416.
- 44 T. N. M. da Cruz, S. M. Savassa, M. H. F. Gomes, E. S. Rodrigues, N. M. Duran, E. de Almeida, A. P. Martinelli and H. W. P. de Carvalho, Shedding light on the mechanisms of absorption and transport of ZnO nanoparticles by plants via in vivo X-ray spectroscopy, *Environ. Sci.: Nano*, 2017, **4**, 2367-2376.
- 45 K. Kinska, J. Jimenez-Lamana, J. Kowalska, B. Krasnodebska-Ostrega and J. Szpunar, Study of the uptake and bioaccumulation of palladium nanoparticles by *Sinapis alba* using single particle ICP-MS, *Sci. Total Environ.*, 2018, **615**, 1078-1085.
- 46 A. Perez-de-Luque, Interaction of nanomaterials with plants: What do we need for real applications in agriculture?, *Front. Environ. Sci.*, 2017, **5**, 12.
- 47 E. Corredor, P. S. Testillano, M. J. Coronado, P. Gonzalez-Melendi, R. Fernandez-Pacheco, C. Marquina, M. R. Ibarra, J. M. de la Fuente, D. Rubiales, A. Perez-De-Luque and M. C. Risueno, Nanoparticle penetration and transport in living pumpkin plants: in situ subcellular identification, *BMC Plant Biol.*, 2009, **9**, 45.
- 48 E. Etxeberria, P. Gonzalez, P. Bhattacharya, P. Sharma and P. C. Ke, Determining the size exclusion for nanoparticles in citrus leaves, *Hortscience*, 2016, **51**, 732-737.
- 49 M. Kah, N. Tufenkji and J. C. White, Nano-enabled strategies to enhance crop nutrition and protection, *Nat. Nanotechnol.*, 2019, **14**, 532-540.
- 50 A. Avellan, F. Schwab, A. Masion, P. Chaurand, D. Borschneck, V. Vidal, J. Rose, C. Santaella and C. Levard, Nanoparticle Uptake in Plants: Gold Nanomaterial Localized in Roots of *Arabidopsis thaliana* by X-ray Computed Nanotomography and Hyperspectral Imaging, *Environ. Sci. Technol.*, 2017, **51**, 8682-8691.
- 51 C. O. Dimkpa, Soil properties influence the response of terrestrial plants to metallic nanoparticles exposure, *Curr. Opin. Environ. Sci.*, 2018, **6**, 1-8.
- 52 J. Sasse, E. Martinoia and T. Northen, Feed your friends: do plant exudates shape the root microbiome?, *Trends Plant Sci.*, 2018, **23**, 25-41.
- 53 G. Marslin, C. J. Sheeba and G. Franklin, Nanoparticles alter secondary metabolism in plants via ROS burst, *Front. Plant Sci.*, 2017, **8**, 832.
- 54 A. Dev, A. K. Srivastava and S. Karmakar, Nanomaterial toxicity for plants, *Environ. Chem. Lett.*, 2018, **16**, 85-100.
- 55 B. Castro, M. Citterico, S. Kimura, D. M. Stevens, M. Wrzaczek and G. Coaker, Stress-induced reactive oxygen species compartmentalization, perception and signalling, *Nat. Plants*, 2021, **7**, 403-412.
- 56 A. Perez-de-Luque, Guest Edited Collection: Nanotechnology in agriculture, *Sci. Rep.*, 2020, **10**, 15738.
- 57 A. Avellan, M. Simonin, S. M. Anderson, N. K. Geitner, N. Bossa, E. Spielman-Sun, E. S. Bernhardt, B. T. Castellon, B. P. Colman, J. L. Cooper, M. Ho, M. F. Hochella, H. Hsu-Kim, S. Inoue, R. S. King, S. Loughton, C. W. Matson, B. G. Perrotta, C. J. Richardson, J. M. Urnine, M. R. Wiesner and G. V. Lowry, Differential reactivity of copper- and gold-based nanomaterials controls their seasonal biogeochemical cycling and fate in a freshwater wetland mesocosm, *Environ. Sci. Technol.*, 2020, **54**, 1533-1544.
- 58 J. P. Stegemeier, A. Avellan and G. V. Lowry, Effect of initial speciation of copper- and silver-based nanoparticles on their long-term fate and phytoavailability in freshwater wetland mesocosms, *Environ. Sci. Technol.*, 2017, **51**, 12114-12122.
- 59 B. P. Espinasse, N. K. Geitner, A. Schierz, M. Therezien, C. J. Richardson, G. V. Lowry, L. Ferguson and M. R. Wiesner, Comparative persistence of engineered nanoparticles in a complex aquatic ecosystem, *Environ. Sci. Technol.*, 2018, **52**, 4072-4078.
- 60 N. K. Geitner, N. Bossa and M. R. Wiesner, Formulation and validation of a functional assay-driven model of nanoparticle aquatic transport, *Environ. Sci. Technol.*, 2019, **53**, 3104-3109.
- 61 B. P. Colman, B. Espinasse, C. J. Richardson, C. W. Matson, G. V. Lowry, D. E. Hunt, M. R. Wiesner and E. S. Bernhardt, Emerging contaminant or an old toxin in disguise? Silver nanoparticle impacts on ecosystems, *Environ. Sci. Technol.*, 2014, **48**, 5229-5236.
- 62 P. Marschner, *Mineral Nutrition of Higher Plants*, Academic Press, San Diego, 3rd. ed., 2012.
- 63 V. Fernandez, T. Stiropoulos and P. H. Brown, *Foliar fertilization: scientific principles and field practices*, International Fertilizer Industry Association, Paris, 2013.
- 64 A. Karny, A. Zinger, A. Kajal, J. Shainsky-Roitman and A. Schroeder, Therapeutic nanoparticles penetrate leaves and deliver nutrients to agricultural crops, *Sci. Rep.*, 2018, **8**, 7589.
- 65 N. A. Kulikova, D. S. Volkov, A. B. Volikov, D. P. Abroskin, A. I. Krepak and I. V. Perminova, Silver nanoparticles stabilized by humic substances adversely affect wheat plants and soil, *J. Nanopart. Res.*, 2020, **22**, 100.
- 66 P. D. Ittroutwar, G. Kasivelu, V. Raguraman, K. Malaichamy and S. K. Sevathapandian, Effects of biogenic zinc oxide nanoparticles on seed germination and seedling vigor of maize (*Zea mays*), *Biocatal. Agric. Biotechnol.*, 2020, **29**, 101778.
- 67 P. D. Ittroutwar, K. Govindaraju, S. Tamilselvan, M. Kannan, K. Raja and K. S. Subramanian, Seaweed-based biogenic zno nanoparticles for improving agro-morphological characteristics of rice (*Oryza sativa* L.), *J. Plant Growth Regul.*, 2020, **39**, 717-728.
- 68 L. Rossi, L. N. Fedenia, H. Sharifan, X. M. Ma and L. Lombardini, Effects of foliar application of zinc sulfate and zinc nanoparticles in coffee (*Coffea arabica* L.) plants, *Plant Physiol. Biochem.*, 2019, **135**, 160-166.
- 69 F. Perez-Labrada, E. R. Lopez-Vargas, H. Ortega-Ortiz, G. Cadenas-Pliego, A. Benavides-Mendoza and A. Juarez-Maldonado, Responses of tomato plants under saline stress to foliar application of copper nanoparticles, *Plants-Basel*, 2019, **8**, 151.
- 70 D. R. Davis, Declining fruit and vegetable nutrient composition: What is the evidence?, *Hortscience*, 2009, **44**, 15-19.
- 71 R. J. Marles, Mineral nutrient composition of vegetables, fruits and grains: The context of reports of apparent historical declines, *J. Food Compos. Anal.*, 2017, **56**, 93-103.
- 72 E. R. Lopez-Vargas, H. Ortega-Ortiz, G. Cadenas-Pliego, K. D. Romenus, M. C. de la Fuente, A. Benavides-Mendoza and A. Juarez-Maldonado, Foliar Application of copper nanoparticles increases the fruit quality and the content of bioactive compounds in tomatoes, *Appl. Sci.*, 2018, **8**, 1020.
- 73 L. Taiz and E. Zeiger, *Plant Physiology*, Sinauer Associates, Sunderland, 2002.
- 74 S. M. Savassa, N. M. Duran, E. S. Rodrigues, E. de Almeida, C. A. M. van Gestel, T. F. V. Bompadre and H. W. P. de Carvalho, Effects of ZnO nanoparticles on *Phaseolus vulgaris* germination and seedling development determined by X-ray spectroscopy, *ACS Appl. Nano Mater.*, 2018, **1**, 6414-6426.
- 75 J. R. Santos-Rasera, A. S. Neto, R. T. R. Monteiro, C. A. M. van Gestel and H. W. P. de Carvalho, Toxicity, bioaccumulation and biotransformation of Cu oxide nanoparticles in *Daphnia magna*, *Environ. Sci.: Nano*, 2019, **6**, 2897-2906.

- 76 N. M. Duran, S. M. Savassa, R. G. de Lima, E. de Almeida, F. S. Linhares, C. A. M. van Gestel and H. W. P. de Carvalho, X-ray spectroscopy uncovering the effects of Cu based nanoparticle concentration and structure on *Phaseolus vulgaris* germination and seedling development, *J. Agric. Food Chem.*, 2017, **65**, 7874-7884.
- 77 F. Schwabe, R. Schulin, L. K. Limbach, W. Stark, D. Burge and B. Nowack, Influence of two types of organic matter on interaction of CeO<sub>2</sub> nanoparticles with plants in hydroponic culture, *Chemosphere*, 2013, **91**, 512-520.
- 78 F. Schwabe, S. Tanner, R. Schulin, A. Rotzetter, W. Stark, A. von Quadt and B. Nowack, Dissolved cerium contributes to uptake of Ce in the presence of differently sized CeO<sub>2</sub>-nanoparticles by three crop plants, *Metallomics*, 2015, **7**, 466-477.
- 79 J. T. Dahle, K. Livi and Y. Arai, Effects of pH and phosphate on CeO<sub>2</sub> nanoparticle dissolution, *Chemosphere*, 2015, **119**, 1365-1371.
- 80 N. K. Fageria, V. C. Baligar and D. Hillel, in *Encyclopedia of Soils in the Environment*, ed. D. Hillel, Elsevier, Oxford, 2005, pp. 63-71.
- 81 C. Reimann, K. Fabian, M. Birke, P. Filzmoser, A. Demetriades, P. Negrel, K. Oorts, J. Matschullat, P. de Caritat and G. P. Team, GEMAS: Establishing geochemical background and threshold for 53 chemical elements in European agricultural soil, *Appl. Geochem.*, 2018, **88**, 302-318.
- 82 M. Ruffo, R. Olson and I. Daverede, Maize yield response to zinc sources and effectiveness of diagnostic indicators, *Commun. Soil Sci. Plant Anal.*, 2016, **47**, 137-141.
- 83 D. Y. Liu, W. Zhang, P. Yan, X. P. Chen, F. S. Zhang and C. Q. Zou, Soil application of zinc fertilizer could achieve high yield and high grain zinc concentration in maize, *Plant Soil*, 2017, **411**, 47-55.
- 84 T. Hofmann, G. V. Lowry, S. Ghoshal, N. Tufenkji, D. Brambilla, J. R. Dutcher, L. M. Gilbertson, J. P. Giraldo, J. M. Kinsella, M. P. Landry, W. Lovell, R. Naccache, M. Paret, J. A. Pedersen, J. M. Unrine, J. C. White and K. J. Wilkinson, Technology readiness and overcoming barriers to sustainably implement nanotechnology-enabled plant agriculture, *Nat. Food*, 2020, **1**, 416-425.
- 85 N. Zuverza-Mena, D. Martinez-Fernandez, W. C. Du, J. A. Hernandez-Viezcas, N. Bonilla-Bird, M. L. Lopez-Moreno, M. Komarek, J. R. Peralta-Videa and J. L. Gardea-Torresdey, Exposure of engineered nanomaterials to plants: Insights into the physiological and biochemical responses-A review, *Plant Physiol. Biochem.*, 2017, **110**, 236-264.
- 86 M. Faizan, A. Faraz and S. Hayat, Effective use of zinc oxide nanoparticles through root dipping on the performance of growth, photosynthesis and antioxidant system in tomato, *J. Plant Biochem. Biotechnol.*, 2020, **29**, 553-567.
- 87 P. Y. Li, A. D. Wang, W. C. Du, L. Mao, Z. B. Wei, S. F. Wang, H. Y. Yuan, R. Ji and L. J. Zhao, Insight into the interaction between Fe-based nanomaterials and maize (*Zea mays*) plants at metabolic level, *Sci. Total Environ.*, 2020, **738**, 139795.
- 88 N. I. Elsheery, M. N. Helaly, H. M. El-Hoseiny and S. M. Alam-Eldein, Zinc oxide and silicone nanoparticles to improve the resistance mechanism and annual productivity of salt-stressed mango trees, *Agronomy-Basel*, 2020, **10**, 558.
- 89 L. Lu, M. Huang, Y. X. Huang, P. F. X. Corvini, R. Ji and L. J. Zhao, Mn(3)O(4)nanozymes boost endogenous antioxidant metabolites in cucumber (*Cucumis sativus*) plant and enhance resistance to salinity stress, *Environ. Sci.: Nano*, 2020, **7**, 1692-1703.
- 90 J. Faraji and A. Sepehri, Exogenous nitric oxide improves the protective effects of TiO<sub>2</sub> nanoparticles on growth, antioxidant system, and photosynthetic performance of wheat seedlings under drought stress, *J. Soil Sci. Plant Nutr.*, 2020, **20**, 703-714.
- 91 M. Adrees, Z. S. Khan, S. Ali, M. Hafeez, S. Khalid, M. Z. U. Rehman, A. Hussain, K. Hussain, S. A. S. Chatha and M. Rizwan, Simultaneous mitigation of cadmium and drought stress in wheat by soil application of iron nanoparticles, *Chemosphere*, 2020, **238**, 124681.
- 92 D. A. Ferreira, T. F. da Silva, V. S. Pylro, J. F. Salles, F. D. Andreote and F. Dini-Andreote, Soil microbial diversity affects the plant-root colonization by arbuscular mycorrhizal fungi, *Microb. Ecol.*, 2021, **82**, 100-103.
- 93 A. R. Schwember, J. Schulze, A. del Pozo and R. A. Cabeza, Regulation of symbiotic nitrogen fixation in legume root nodules, *Plants-Basel*, 2019, **8**, 333.
- 94 M. Rosenblueth, E. Ormeno-Orrillo, A. Lopez-Lopez, M. A. Rogel, B. J. Reyes-Hernandez, J. C. Martinez-Romero, P. M. Reddy and E. Martinez-Romero, Nitrogen fixation in cereals, *Front. Microbiol.*, 2018, **9**, 1794.
- 95 Q. Qu, Z. Y. Zhang, W. Peijnenburg, W. Y. Liu, T. Lu, B. L. Hu, J. M. Chen, J. Chen, Z. F. Lin and H. F. Qian, Rhizosphere microbiome assembly and its impact on plant growth, *J. Agric. Food Chem.*, 2020, **68**, 5024-5038.
- 96 R. Dinesh, M. Anandaraj, V. Srinivasan and S. Hamza, Engineered nanoparticles in the soil and their potential implications to microbial activity, *Geoderma*, 2012, **173**, 19-27.
- 97 S. Mishra, H. B. Singh and X. Yang, Creating a global database "Nanomaterials in the soil environment": future need for the terrestrial ecosystem, *Energy Ecol. Environ.*, 2019, **4**, 271-285.
- 98 M. Simonin and A. Richaume, Impact of engineered nanoparticles on the activity, abundance, and diversity of soil microbial communities: a review, *Environ. Sci. Pollut. Res.*, 2015, **22**, 13710-13723.
- 99 Y. Ge, C. C. Shen, Y. Wang, Y. Q. Sun, J. P. Schimel, J. L. Gardea-Torresdey and P. A. Holden, Carbonaceous nanomaterials have higher effects on soybean rhizosphere prokaryotic communities during the reproductive growth phase than during vegetative growth, *Environ. Sci. Technol.*, 2018, **52**, 6636-6646.
- 100 G. F. Liu, M. Zhang, Y. J. Jin, X. J. Fan, J. H. Xu, Y. C. Zhu, Z. W. Fu, X. L. Pan and H. F. Qian, The effects of low concentrations of silver nanoparticles on wheat growth, seed quality, and soil microbial communities, *Water, Air Soil Pollut.*, 2017, **228**, 348.
- 101 C. X. Ma, J. Borgatta, R. De La Torre-Roche, N. Zuverza-Mena, J. C. White, R. J. Hamers and W. H. Elmer, Time-Dependent transcriptional response of tomato (*Solanum lycopersicum* L.) to Cu nanoparticle exposure upon infection with *Fusarium oxysporum* f. sp. *lycopersici*, *ACS Sustainable Chem. Eng.*, 2019, **7**, 10064-10074.
- 102 C. D. P. Perez, R. D. Roche, N. Zuverza-Mena, C. X. Ma, Y. Shen, J. C. White, E. A. Pozza, A. A. Pozza and W. H. Elmer, Metalloid and metal oxide nanoparticles suppress sudden death syndrome of soybean, *J. Agric. Food Chem.*, 2020, **68**, 77-87.
- 103 J. D. Judy, D. H. McNear, C. Chen, R. W. Lewis, O. V. Tsyusko, P. M. Bertsch, W. Rao, J. Stegemeier, G. V. Lowry, S. P. McGrath, M. Durenkamp and J. M. Unrine, Nanomaterials in biosolids inhibit nodulation, shift microbial community composition, and result in increased metal uptake relative to bulk/dissolved metals, *Environ. Sci. Technol.*, 2015, **49**, 8751-8758.
- 104 J. D. Judy, J. K. Kirby, C. Creamer, M. J. McLaughlin, C. Fiebigler, C. Wright, T. R. Cavagnaro and P. M. Bertsch, Effects of silver sulfide nanomaterials on mycorrhizal colonization of tomato plants and soil microbial

- 1  
2  
3 communities in biosolid-amended soil, *Environ. Pollut.*,  
4 2015, **206**, 256-263.
- 5 105 A. Noori, J. C. White and L. A. Newman, Mycorrhizal fungi  
6 influence on silver uptake and membrane protein gene  
7 expression following silver nanoparticle exposure, *J.*  
8 *Nanopart. Res.*, 2017, **19**, 66.
- 9 106 F. Y. Wang, X. Q. Liu, Z. Y. Shi, R. J. Tong, C. A. Adams and X.  
10 J. Shi, Arbuscular mycorrhizae alleviate negative effects of  
11 zinc oxide nanoparticle and zinc accumulation in maize  
12 plants - A soil microcosm experiment, *Chemosphere*, 2016,  
13 **147**, 88-97.
- 14 107 Z. Zahra, M. Arshad, R. Rafique, A. Mahmood, A. Habib, I. A.  
15 Qazi and S. A. Khan, Metallic nanoparticle (TiO<sub>2</sub> and Fe<sub>3</sub>O<sub>4</sub>)  
16 application modifies rhizosphere phosphorus availability  
17 and uptake by *Lactuca sativa*, *J. Agric. Food Chem.*, 2015, **63**,  
18 6876-6882.
- 19 108 C. M. Rico, A. C. Barrios, W. J. Tan, R. Rubenecia, S. C. Lee, A.  
20 Varela-Ramirez, J. R. Peralta-Videa and J. L. Gardea-  
21 Torresdey, Physiological and biochemical response of soil-  
22 grown barley (*Hordeum vulgare* L.) to cerium oxide  
23 nanoparticles, *Environ. Sci. Pollut. Res.*, 2015, **22**, 10551-  
24 10558.
- 25 109 S. Stark, M. K. Mannisto and A. Eskelinen, Nutrient  
26 availability and pH jointly constrain microbial extracellular  
27 enzyme activities in nutrient-poor tundra soils, *Plant Soil*,  
28 2014, **383**, 373-385.
- 29 110 J. T. Sims, Soil-pH effects on the distribution and plant  
30 availability of manganese, copper, and zinc, *Soil Sci. Soc. Am.*  
31 *J.*, 1986, **50**, 367-373.
- 32 111 T. T. S. Lew, R. Sarojam, I. C. Jang, B. S. Park, N. I. Naqvi, M.  
33 H. Wong, G. P. Singh, R. J. Ram, O. Shoseyov, K. Saito, N. H.  
34 Chua and M. S. Strano, Species-independent analytical tools  
35 for next-generation agriculture, *Nat. Plants*, 2020, **6**, 1408-  
36 1417.
- 37 112 J. P. Giraldo, M. P. Landry, S. M. Faltermeier, T. P.  
38 McNicholas, N. M. Iverson, A. A. Boghossian, N. F. Reuel, A.  
39 J. Hilmer, F. Sen, J. A. Brew and M. S. Strano, Plant  
40 nanobionics approach to augment photosynthesis and  
41 biochemical sensing, *Nat. Mater.*, 2014, **13**, 400-408.
- 42 113 D. Montavo, F. Degryse and M. J. McLaughlin, Natural  
43 colloidal P and its contribution to plant P uptake, *Environ.*  
44 *Sci. Technol.*, 2015, **49**, 3427-3434.
- 45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60