



**HAL**  
open science

## Modelling Absorption Spectra of Furimamide-Nanoluciferase System

Houda Moumene, Etienne Mangaud, Isabelle Navizet

► **To cite this version:**

Houda Moumene, Etienne Mangaud, Isabelle Navizet. Modelling Absorption Spectra of Furimamide-Nanoluciferase System. Journée du laboratoire MSME, Sep 2022, Marne La Vallée, France. hal-03776478

**HAL Id: hal-03776478**

**<https://hal.science/hal-03776478>**

Submitted on 13 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

# Modelling Absorption Spectra of Furimamide - Nanoluciferase System


**Houda MOUMENE, Etienne MANGAUD and Isabelle NAVIZET**  
Laboratoire Modélisation et Simulation Multi-Echelle – UMR 8208 MSME,  
Université Gustave Eiffel, UPEC, CNRS, F-77454 Marne-la-Vallée, France




Bioluminescence is a biological phenomenon of light production by living species with many applications[1], notably for the detection of cancer cells[2]. The present study deals with a ligand-protein bioluminescent system (furimazine – Nanoluciferase) derived from a shrimp called *Oplophorus gracilirostri* with a very high luminescence intensity[3]. A better understanding of this system will provide insights to tailor new devices with emission of a photon with a red color, and thus a light signal more easily detectable inside the human body. Simulations of the emission and absorption spectra are relevant tools to comply this goal[4]. We focus here on the absorption spectra of the light emitter system furimamide – Nanoluciferase. In order to take into account the protein environment and the system flexibility, we carry out classical molecular dynamics (MM) with AMBER software. Furimamide is a flexible molecule: standard force fields such as GAFF[5] (General Amber Force Field) are unable to reproduce correctly potential energy along torsion angles when compared to in vacuo DFT computations. Thus, we perform a multidimensional fit using 1D and 2D potential energy surface scans with several torsion angles of the molecule in order to parameterize a new force field. Then, we compute electronic transitions at a QM/MM level of theory on a set of 100 snapshots extracted from MM trajectory. As a perspective, we plan to use the same methodology to simulate emission spectra.

## Nanoluc/Furimamide system:

This system derives from a bioluminescent complex of a shrimp species.




The oxidation of the furimazine, in the cavity of the protein called nanoluciferase, leads to furimamide in its excited state : by fluorescence, it emits light.




## Methodology / Spectra simulation:

In order to simulate the absorption spectra, we need to carry out the following steps:




## Adjusting the force field (FF): MDGX program




Initial guess of parameters (GAFF2)

Least-square fitting to minimize the energy difference between DFT and FF




## MD simulation: (Amber20)


50ns Dynamics



## QMMM simulation: (Amber20/Gaussian16)




Molecular mechanic subsystem  
**Protein + WATER**  
Quantum Mechanics subsystem  
**Ligand**  
DFT Method : CAM-B3LYP/6-311G(2d,p)



## Absorption spectra simulation:

Calculated oscillator strengths of excited states for 201 snapshots of furimamide




## Conclusion and perspectives:

We have automatized the absorption spectra simulation, we need to compare our results with experimental data. The emission spectra are yet to be simulated in order to modelize the color of emission of this system.

## How to explore the complex configuration space?

Molecular dynamics allow to calculate the trajectories of a system, the evolution of the positions of each atom of the system over time using Newton's equations of motion.


### 1. Modelling of the system



### 2. The force field

- FF14SB (Force Field 14 Stony Brook) for the protein
- TIP3P for water molecules
- GAFF2 for the ligand ?

The potential energies calculated using the GAFF2 force field doesn't match with the ones calculated using DFT scans of 7 dihedral angles of the molecule. The force field needs to be adjusted.



## References:

- [1] A. J. Syed et J. C. Anderson, Chem. Soc. Rev. 50 (2021) 5668-5705.
- [2] M. A. Paley et J. A. Prescher, Medchemcomm 5 (2014) 255-267.
- [3] M. P. Hall et al., ACS Chem. Biol., 7 (2012) 1848-1857.
- [4] M. Sahihi, J. Sanz García, et I. Navizet, J. Phys. Chem. B 124 (2020) 2539-2548.
- [5] J. Wang, R. M. Wolf, J. W. Caldwell, P. A. Kollman, et D. A. Case, Journal of Computational Chemistry 25 (2014) 18.