



**HAL**  
open science

## Effect of mirror therapy in the treatment of phantom limb pain in amputees: A systematic review of randomized placebo controlled trials does not find any evidence of efficacy

Matthieu Guémann, Émilie Olié, Lea Raquin, Philippe Courtet, Nathan Risch

### ► To cite this version:

Matthieu Guémann, Émilie Olié, Lea Raquin, Philippe Courtet, Nathan Risch. Effect of mirror therapy in the treatment of phantom limb pain in amputees: A systematic review of randomized placebo controlled trials does not find any evidence of efficacy. *European Journal of Pain*, 2023, 27 (1), pp.3-13. 10.1002/ejp.2035 . hal-03777016

**HAL Id: hal-03777016**

**<https://hal.science/hal-03777016>**

Submitted on 14 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Risch Nathan (Orcid ID: 0000-0002-9632-6077)

## TITLE

EFFECT OF MIRROR THERAPY IN THE TREATMENT OF PHANTOM LIMB PAIN IN AMPUTEES: A SYSTEMATIC REVIEW OF RANDOMIZED PLACEBO CONTROLLED TRIALS DOES NOT FIND ANY EVIDENCE OF EFFICACY

## RUNNING HEAD

MIRROR THERAPY FOR PHANTOM LIMB PAIN IN AMPUTEES

## KEYWORDS

Mirror therapy; Phantom limb; Amputee; Disability

## AUTHORS

**Matthieu Guémann (M.G)** PT, PhD; Armed forces Biomedical Research Institute, Physiology of Exercise and Activities in Extreme Conditions Unit, Bretigny-sur-Orge, France

**Emilie Olié (E.O)**, MD, PhD; Institute of Functional Genomics, University of Montpellier, CNRS, INSERM, 34094 Montpellier, France; Department of Emergency Psychiatry and Post-Acute Care, CHU Montpellier.

**Lea Raquin (L.R)**, PT; Clinique du Bourget, Ramsay Santé, 93350 Le Bourget, France

**Philippe Courtet (P.C)**, MD, PhD; Institute of Functional Genomics, University of Montpellier, CNRS, INSERM, 34094 Montpellier, France; Department of Emergency Psychiatry and Post-Acute Care, CHU Montpellier,

**Nathan Risch (N.R)**, PT, MSc; Institute of Functional Genomics, University of Montpellier, CNRS, INSERM, 34094 Montpellier, France; Department of Emergency Psychiatry and Post-Acute Care, CHU Montpellier, 34295 Montpellier, France; Clinique de la Lironde, Clinea Psychiatrie, 34980 Saint-Clément-de-Rivière, France

## CORRESPONDING AUTHOR

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/ejp.2035](https://doi.org/10.1002/ejp.2035)

Nathan Risch, Department of Emergency Psychiatry and Post-Acute Care, CHU Montpellier, 34295 Montpellier, France; [risch.nathan@gmail.com](mailto:risch.nathan@gmail.com)

**CATEGORY OF SUBMISSION:** Review

**FUNDING:** No author received any support for the present manuscript.

**CONFLICTS OF INTEREST:** The authors declare no conflict of interests related to this manuscript.

**ABSTRACT (250 words)**

**Background and Objective:** Phantom limb pain (PLP) concerns >50% of amputees and has a negative impact on their rehabilitation, mental health, and quality of life. Mirror therapy (MT) is a promising strategy, but its effectiveness remains controversial. We performed a systematic review to: (i) evaluate the effectiveness of MT versus placebo in reducing PLP, and (ii) determine MT effect on disability and quality of life.

**Databases and Data Treatment:** We selected randomized controlled trials in five databases (Medline, Cochrane Library, CINAHL, PEDro, and Embase) that included patients with unilateral lower or upper limb amputation and PLP and that compared the effects on PLP of MT versus a placebo technique. The primary outcome was PLP intensity changes and the secondary outcomes were PLP duration, frequency, patients' disability and quality of life.

**Results:** Among the five studies included, only one reported a significant difference between the MT group and control group, with a positive MT effect at week 4. Only one study assessed MT effect on disability, and found a significant improvement in the MT group at week 10 and month 6.

**Conclusions:** Our systematic review did not allow concluding that MT reduces PLP and disability in amputees. This lack of strong evidence is probably due to (i) the low methodological quality of the included studies, and (ii) the lack of statistical power. Future trials should include a higher number of patients, increase the number and frequency of MT sessions, have a long-term follow-up, and improve the methodological quality.

## **SIGNIFICANCE**

Recent meta-analyses concluded that mirror therapy is effective for reducing phantom limb pain. Conversely, the present systematic review that included only studies with the best level of evidence did not find any evidence about its effectiveness for this condition. We identified many ways to improve future randomized controlled trials on this topic: increasing the number of participants, reducing the intra-group heterogeneity, using a suitable placebo, and intensifying the mirror therapy sessions and frequency.

## **INTRODUCTION**

Phantom limb pain (PLP), which affects between 49 and 88% of amputees (Nikolajsen and Jensen, 2001), is a neuropathic pain defined as a painful or unpleasant sensation perceived in the territory of the amputated or deafferented limb (Niraj and Niraj, 2014). Patients describe PLP as a strong pain like electric currents, or as a burning, stinging, cramping, or stabbing pain (Cohen and Hsu, 2013). PLP often occurs suddenly and lasts from few seconds to several hours, and is mostly felt in the distal part of the phantom limb (fingers or toes). PLP affects negatively rehabilitation, mental health, and quality of life (van der Schans et al., 2002; Sinha et al., 2011).

The mechanisms underlying PLP have not been clearly identified and are probably multiple (Flor et al., 2006; Makin et al., 2013). According to the maladaptive cortical plasticity theory, PLP emerges following the progressive occupation of the cortical areas that correspond to the missing limb by adjacent areas. This is due to the absence of sensory afferents signals from the missing limb and spinal dysregulation that increases the transmission of the pain signal (Flor, 2002). Spinal dysregulation contributes to the pro-nociceptive effect due to a cascade mechanism that implicates increased activity of nociceptors in the dorsal horn, degeneration of C-fibres in the lamina II, sprouting of A-fibres, downregulation of GABAergic activity and

opioid receptors, expansion of adjacent receptive fields in deafferented zones, and spinal cord hyperexcitability linked to the altered substance P expression by A $\beta$  fibres (Flor, 2002; Woolf et al., 1992; Woolf and Mannion, 1999). According to another theory, PLP extent is related to the preserved representation of the phantom limb in the cortex that can be activated and to a reduction of inter-regional functional connectivity in the primary sensorimotor cortex (Makin et al., 2013).

Several pharmacological, surgical or non-pharmacological approaches have been proposed for PLP management. The effectiveness of pharmacological treatments remains low (Alviar et al., 2016). Surgical strategies are effective, but invasive and with a long healing time and loss of function (Dumanian et al., 2019; Herr et al., 2020; Knotkova et al., 2012). Therefore, they are not used as first-line treatment. Among non-pharmacological interventions, mirror therapy (MT), first proposed by V.S. Ramachandran in 1996 (Ramachandran, 1996), seems to be a promising strategy. MT consists in placing a mirror in the parasagittal plane between the healthy limb and the amputated limb. The image of the healthy limb is reflected in the mirror and the patient perceives the reflection of the healthy limb instead of the amputated limb. By creating a visual representation of the missing limb, MT might restore the cortical (motor and sensory) areas that correspond to the absent limb. By restoring the body image and body schema disturbed by the amputation, MT might reduce PLP (Flor, 2002; Ramachandran and Altschuler, 2009; Foell et al., 2014).

MT effectiveness remains controversial (Barbin et al., 2016; Batsford et al., 2017; Herrador Colmenero et al., 2018; Plumbe et al., 2013; Rothgangel et al., 2011; Thieme et al., 2016, 2018; Xie et al., 2021) because previous reviews and meta-analyses found inconclusive evidence. This could be due to the fact that the studies included in these reviews assessed MT in various pathologies (Rothgangel et al., 2011; Thieme et al., 2016) or in combination with other techniques (Batsford et al., 2017; Herrador Colmenero et al., 2018). Barbin et al (2016) analysed MT use specifically in amputees with PLP (Barbin et al., 2016). They found only three randomized controlled trials and concluded that the level of evidence was not adequate, due to the low methodological quality of the included studies and the small sample sizes. Two

recent meta-analyses on this question concluded that MT is effective for PLP management (Wang et al., 2021; Xie et al., 2021). One found a positive effect of MT at month 1 of treatment ( $I^2 = 0\%$ , standardized mean difference = -0.46, 95% CI: -0.79 to 0.13,  $p = 0.007$ ), but not at month 3 or 6 of follow-up (Xie et al., 2021). The other meta-analysis showed that MT reduces pain with a larger effect size ( $I^2 = 82\%$ ; standardized mean difference = -0.81; 95% CI = -1.36 to -0.25;  $p = 0.005$ ) compared with other methods (Wang et al., 2021). However, the conclusion of these meta-analyses are subject of debate due to questionable methodological and statistical choices: the use of fixed-effect analyses (Borenstein et al., 2010), the inclusion of low powered and poor quality studies, the strong heterogeneity, and the interpretation of non-clinically relevant effects (Moore et al., 2022). Furthermore, no systematic review has evaluated MT impact on the patients' quality of life and disability. Pain reduction is a good indicator of PLP improvement, but changes in disability and well-being also are important (World Health Organization and World Bank, 2011).

Therefore, we performed a systematic review: (i) to evaluate the effectiveness of MT versus placebo in reducing PLP; and (ii) to determine MT effect on disability and quality of life. To overcome the previously stated limitations, we included only randomized controlled trials that compared MT versus a placebo in unilateral amputees.

## LITERATURE SEARCH METHODS

This systematic review of the literature followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Liberati et al., 2009).

### Eligibility Criteria

We defined the eligibility criteria to include studies according to the Population, Intervention, Comparison, Outcomes and Study (PICOS) model:

- Population: adult patients with unilateral, lower or upper limb amputation due to any cause and with PLP.
- Intervention: MT to reduce PLP. It has been hypothesized that MT restores the cortical (motor and sensory) areas corresponding to the missing limb through the co-occurrence of visual,

Accepted Article

proprioceptive information from the intact limb, and motor inputs from both limbs. This input combination is considered to be the active component of MT (Schone et al., 2022). To avoid heterogeneity and to specifically assess the association of these components, we chose MT only as the intervention. We considered that other techniques, such as mental visualization, tactile discrimination and virtual reality, target a similar process (i.e. maladaptive plasticity), but have different active components. Mental visualization does not induce motor commands or sensory inputs. Tactile discrimination enhances sensory stimulation on the stump; however, it does not implicate movements, motor commands and proprioceptive inputs. Virtual reality can act as MT, but virtual reality approaches are heterogeneous (e.g. using the residual or the intact limb, moving one or two arms, interacting with exergames or avatars).

- Comparison: any other technique considered as placebo (covered mirror or mirrorless visualization). The covered mirror is classically considered as the best controls for MT (Chan et al., 2007; Finn et al., 2017; Ramadugu et al., 2017; Schone et al., 2022). To test the effect of MT active components, an adequate control is a technique as similar as possible. When using the covered mirror and the mirrorless approaches as placebo, the only difference, compared with MT, is the suppression of the visual feedback, while the other inputs remain the same.

- Outcomes:

Primary: PLP intensity (measured with a scale).

Secondary:

- PLP duration and frequency
- Disability and quality of life (measured with questionnaires)

- Study types: randomized controlled trials published between 1996 (first description of MT) and 01 July 2021.

Non-inclusion criteria were:

- Articles without available full text
- Articles published in a language other than English or French

- Articles comparing the effects of MT with another technique (TENS, motor imaging, phantom exercise, tactile discrimination).

#### Information sources and search strategy

We used Medline, Cochrane Library, CINAHL, PEDro and Embase as information sources and searched them using the index terms and syntax listed in Table 1.

Table 1: Index terms and syntax for each database.

**PLEASE INCLUDE TABLE 1**

#### Study selection

This step was carried out by two independent reviewers (L.R., N.R.) following the established inclusion and non-inclusion criteria. Duplicates were removed from the list of results. A first selection was made by reading the titles and abstracts to remove articles not meeting the inclusion criteria. The final selection was made by reading the full text of the selected articles. When no agreement could be reached, a third reviewer (M.G.) intervened to reach an agreement.

#### Data extraction

The data extracted were:

- Participants: sample size, age, sex, location and cause of amputation, PLP intensity and time of onset since amputation.
- Intervention: data were extracted based on the French translation of the Template for Intervention Description and Replication (TIDieR) item list.
- Measurements: tools used to measure PLP and the number of measurements performed.
- Intervention effect: we extracted results of intra- and inter-group assessments. We extracted only results that compared the effect of MT versus placebo. We did not consider comparisons with other techniques (mental visualization, virtual reality).



### Methodological quality of the included studies

We assessed the methodological quality of the retained randomized controlled trials using the PEDro scale (Brosseau et al., 2015). The PEDro scale includes 11 criteria in order to help the PEDro database users to rapidly identify which clinical studies (i.e. randomized clinical trials and controlled clinical trials) are likely to be internally valid (criteria 2-9) and to have sufficient statistical information to make their results interpretable (criteria 10-11). Criterion 1, which relates to the external validity, is not used to calculate the PEDro score. Two reviewers (L.R., N.R.) independently scored the articles, and discussed areas of disagreement. When no agreement could be reached, a third reviewer (M.G.) intervened to reach an agreement.

## **RESULTS**

### Study selection

The literature search identified 155 articles (flowchart in **Fig. 1**). We excluded 43 articles after removal of duplicates and 94 after reading the titles and abstracts. The reasons for exclusions are detailed in Figure 1. We did not find one study (text unavailable) (Perry, 2013). Following this selection, we examined the full text of 17 articles and retained 5 articles: Brodie et al. 2007, Chan et al. 2009, Finn et al. 2017, Ramadugu et al. 2017, and Rothgangel et al. 2018 (Brodie et al., 2007; Chan et al., 2007; Finn et al., 2017; Ramadugu et al., 2017; Rothgangel et al., 2018).

### **Place Figure 1 here**

### Study methodological quality

We assessed the methodological quality of the five studies with the PEDro scale (**Table 2**). Their scores ranged from 4 to 8. Three studies had a score >6 and could be considered as having a limited risk of bias (Brodie et al., 2007; Ramadugu et al., 2017; Rothgangel et al., 2018). The other two studies [27,28] had a score <5 and were at higher risk of bias.

Table 2: PEDro score of the selected studies

**PLEASE INCLUDE TABLE 2**

### Study characteristics

Three studies used a parallel group design (Brodie et al., 2007; Finn et al., 2017; Rothgangel et al., 2018) and two a cross-over design (Chan et al., 2007; Ramadugu et al., 2017). In total, 256 patients were included, with sample sizes varying from 15 to 80 patients. Participant characteristics (age, location and cause of amputation, time of PLP onset since amputation, and PLP intensity at inclusion) varied among studies (Table 3). Conversely, the type of exercise (i.e., the study intervention) and the control intervention (opaque sheet or mirrorless) were similar among studies. However, the duration, number and frequency of MT sessions varied (Table 4). The intervention was performed under the supervision of an examiner in four studies (Brodie et al., 2007; Chan et al., 2007; Finn et al., 2017; Ramadugu et al., 2017). PLP intensity was assessed with different tools: visual analogue scale (VAS) (Brodie et al., 2007; Finn et al., 2017; Ramadugu et al., 2017), numeric rating scale (Rothgangel et al., 2018), and McGill Pain Questionnaire, original (Brodie et al., 2007) or shorter version (Ramadugu et al., 2017). Only one study assessed MT impact on the patients' disability and quality of life (Rothgangel et al., 2018). Disability was assessed with the Patient Specific Functional Scale (PSFS) and the Pain Disability Index (PDI). Quality of life was assessed with the EuroQol 5-Dimensional Questionnaire (EQ-5D-5L) and two numeric rating scales that measure the impact of pain on sleep and mood.

Table 3: Characteristics of the patients included in the studies

**PLEASE INCLUDE TABLE 3**

### Synthesis of the results

The results obtained for each group and the conclusion on the primary outcome in each study are presented in Table 4.

*Primary outcome: phantom limb pain intensity*

Among the four studies with an intergroup comparison (Brodie et al., 2007; Chan et al., 2007; Ramadugu et al., 2017; Rothgangel et al., 2018), only one found a significant difference between MT group and control (placebo) group with a positive effect of MT at week 4 of follow-up (Chan et al., 2007). Furthermore, in the study by Rothgangel et al., a subgroup analysis suggested clinically significant effects of MT on the mean PLP intensity in women and in patients with a motor component (cramps or unnatural phantom limb position) of PLP (Rothgangel et al., 2018).

For intra-group comparisons, three studies showed a significant decrease in pain in the MT group after the intervention (Chan et al., 2007; Finn et al., 2017; Ramadugu et al., 2017).

*Secondary outcomes:*

*Phantom limb pain duration and frequency*

The included studies did not find any inter-group difference in PLP duration and frequency at 4 and 6 weeks of MT (Ramadugu et al., 2017; Rothgangel et al., 2018), but PLP duration was significantly reduced at 6 months in the MT group compared with placebo (Rothgangel et al., 2018). For intra-group comparisons, three studies found that PLP duration and frequency were decreased in both (MT and control) groups (Finn et al., 2017; Ramadugu et al., 2017; Rothgangel et al., 2018).

*Effect on disability and quality of life:*

Only the study by Rothgangel et al. assessed MT effect on the patients' disability level and quality of life. According to some PSFS items (daily living activities), disability was significantly reduced in the MT group at week 10 and 6 months (Rothgangel et al., 2018). Conversely, quality of life (EQ-5D-5L questionnaire) was comparable between groups.

Table 4: Summary of the characteristics, intervention, and outcomes of the selected studies.

**PLEASE INCLUDE TABLE 4**

## **DISCUSSION**

Among the five studies included in this systematic review, three found that MT decreased PLP intensity, but mainly in the intra-group comparison. Only Chan et al. found a specific positive effect of MT on PLP at week 4 compared with the control group (Chan et al., 2007), but this study has a low methodological quality score. One study reported an improvement of disability by MT through the reduction of PLP impact on daily living activities (Rothgangel et al., 2018). Thus, our systematic review did not allow concluding that MT reduces PLP intensity, frequency and duration, disability and quality of life.

The lack of strong evidence of MT efficiency in PLP is probably due to (i) the low methodological quality (Table 2), and (ii) the lack of statistical power of the included studies. Several studies included fewer than 10 patients per group (Chan et al., 2007; Finn et al., 2017) although Rothgangel et al. calculated that at least 30 patients per group were needed to detect a 2-point difference, with high effect size (Cohen's  $d$  close to 0.8). As they did not detect any effect, the effect size for MT might be smaller (Cohen's  $d$  close to 0.5). This would imply that more than 50 patients per group should be recruited to detect an effect with 80% of power. Furthermore, intra-group heterogeneity was high and several factors should have been considered, such as sex, type of prosthesis, ability to move the phantom limb, presence of a PLP motor component (cramps or unnatural position of the phantom limb) (Rothgangel et al., 2018). Future studies should take into account these factors when deciding the inclusion and exclusion criteria.

Our conclusion is similar to that of previous reviews (Aternali and Katz, 2019; Barbin et al., 2016), but differs from that of the two most recent meta-analyses that reported a significant positive effect of MT on PLP (Wang et al., 2021; Xie et al., 2021). However, the methodological quality of these meta-analyses raises questions. In the meta-analysis by Xie (Xie et al., 2021), the risk of bias of the included studies (GRADE method) was not evaluated. Moreover, a fixed effect model was used instead of a random effect model that considers trials as different

parameters. The observed statistical effect is questionable because the study that contributed to 54% of the conclusion had a low methodological quality (PEDro score = 2). A sensitivity analysis should have been carried out to assess the effect of removing low-quality studies (Borenstein et al., 2010). In the meta-analysis by Wang et al. (Wang et al., 2021) the conclusion displayed a large heterogeneity ( $I^2 > 80\%$ ), suggesting the need of a subgroup analysis. When this subgroup analysis was carried out, MT effect, compared with a covered mirror, disappeared. However, the authors concluded that MT is an effective technique to reduce PLP in amputees. On the basis of these different points, we disagree with the interpretation and conclusions of these two meta-analyses.

Although evidence is lacking concerning MT effectiveness for PLP, the beneficial effects of MT in other conditions suggest that it may be useful also for PLP. The latest Cochrane review shows a significant effect of MT on motor function (effect size: 0.47 [0.27, 0.67]) following stroke (Thieme et al., 2018), and a meta-analysis found an effect of MT on pain associated with post-stroke complex regional pain syndrome (Duong et al., 2018). In these reviews, the MT intervention modalities were more intense (30-minute sessions, 5 times per week) than those used in the five studies included in this review (Table 4) (Thieme et al., 2018). Therefore, increasing the MT session frequency and duration (per week and/or for a longer period) might influence brain plasticity and consequently PLP (Andoh et al., 2017; Flor, 2008; Flor et al., 2006; Kuner and Flor, 2016). It has also been shown that MT effects last for several months after treatment, particularly on motor functions, daily living activities, and pain (Sütbeyaz et al., 2007). In agreement, Rothgangel et al. found that MT still modulated pain episode frequency at 6 months of follow-up (Rothgangel et al., 2018). This could be due to its stabilizing effect on brain plasticity.

#### Limitations & strengths

This review presents some limitations. First, we limited the search to five databases, and the selected articles had to be in French or English. Second, we did not register our study in the PROSPERO database, although we followed a pre-established protocol (wrote in French)

before the start of our study that was approved by the ethics committee. Third, during the literature search, we identified many study protocols to address the research question, but the results of these trials have not been published yet, or they might have been negative, and the authors did not try to publish them (publication bias).

The strengths of this review are: (i) we followed the PRISMA reporting recommendations, (ii) we only included studies with a placebo control group, (iii) we assessed the methodology quality of the studies with the PEDro scale, and (iv) we interpreted their results in function of the PEDro scores. We also assessed secondary outcomes: pain frequency and duration, disability, and quality of life.

## **CONCLUSION**

The results obtained in this systematic review do not allow us to conclude that MT reduces PLP intensity, frequency, duration, or disability. Future clinical studies should enrol a larger number of patients to increase the statistical power, use a covered mirror or mirrorless exercises as placebo, intensify MT sessions and frequency (Duong et al., 2018; Thieme et al., 2018), include a long-term follow-up (Rothgangel et al., 2018), reduce intra-group heterogeneity (sex, prosthesis type, ability to move the phantom limb, presence of a PLP motor component), improve their methodological quality (PEDro scale), and evaluate also other criteria (e.g. duration and frequency of painful episodes, disability and quality of life). Other study designs could be considered, for instance single-case experimental design (SCED) studies that are adapted to rare diseases with a variety of specific and clinical situations (Ganz and Ayres, 2018; Krasny-Pacini and Evans, 2018; Smith, 2012; Tate et al., 2008).

**Acknowledgments:** We thank Elisabetta ANDERMARCHER and Kathlyne DUPUIS MAURIN for their careful proofreading.

**Author contributions:** All authors contributed to the study conception and design. NR and LR extracted data from the trials and evaluated the inclusion and exclusion criteria. NR, LR, and

MG assessed the methodological quality of the trials. All authors contributed to the interpretation of the data and drafted the manuscript.

## REFERENCES

- Alviar, M.J.M., Hale, T., Dungca, M. (2016). Pharmacologic interventions for treating phantom limb pain. *Cochrane Database Syst Rev* 10, CD006380.
- Andoh, J., Diers, M., Milde, C., Frobel, C., Kleinböhl, D., Flor, H. (2017). Neural correlates of evoked phantom limb sensations. *Biol Psychol* 126, 89–97.
- Aternali, A., Katz, J. (2019). Recent advances in understanding and managing phantom limb pain [version 1; peer review: 2 approved]. *F1000Research* 8.
- Barbin, J., Seetha, V., Casillas, J.M., Paysant, J., Pérennou, D. (2016). The effects of mirror therapy on pain and motor control of phantom limb in amputees: A systematic review. *Ann Phys Rehabil Med* 59, 270–275.
- Batsford, S., Ryan, C.G., Martin, D.J. (2017). Non-pharmacological conservative therapy for phantom limb pain: A systematic review of randomized controlled trials. *Physiother Theory Pract* 33, 173–183.
- Borenstein, M., Hedges, L.V., Higgins, J.P.T., Rothstein, H.R. (2010). A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods* 1, 97–111.
- Brodie, E.E., Whyte, A., Niven, C.A. (2007). Analgesia through the looking-glass? A randomized controlled trial investigating the effect of viewing a 'virtual' limb upon phantom limb pain, sensation and movement. *Eur J Pain* 11, 428–436.
- Brosseau, L., Laroche, C., Sutton, A., Guitard, P., King, J., Poitras, S., Casimiro, L., Tremblay, M., Cardinal, D., Cavallo, S., Laferrière, L., Grisé, I., Marshall, L., Smith, J.R., Lagacé, J., Pharand, D., Galipeau, R., Toupin-April, K., Loew, L., Demers, C., Sauvé-Schenk, K., Paquet, N., Savard, J., Tourigny, J., Vaillancourt, V. (2015). Une version franco-canadienne de la Physiotherapy Evidence Database (PEDro) Scale : L'Échelle PEDro. *Physiother Can* 67, 232–239.

Chan, B.L., Da, B., Jb, W., Jj, C., Jt, H. (2007). Mirror Therapy for Phantom Limb Pain. *N Engl J Med* 2.

Cohen, S., Hsu (2013). Postamputation pain: epidemiology, mechanisms, and treatment. *J Pain Res* 121.

Dumanian, G.A., Potter, B.K., Mioton, L.M., Ko, J.H., Cheesborough, J.E., Souza, J.M., Ertl, W.J., Tintle, S.M., Nanos, G.P., Valerio, I.L., Kuiken, T.A., Apkarian, A.V., Porter, K., Jordan, S.W. (2019). Targeted Muscle Reinnervation Treats Neuroma and Phantom Pain in Major Limb Amputees: A Randomized Clinical Trial. *Ann Surg* 270, 238–246.

Duong, S., Bravo, D., Todd, K.J., Finlayson, R.J., Tran, D.Q. (2018). Treatment of complex regional pain syndrome: an updated systematic review and narrative synthesis. *Can J Anesth* 65, 658–684.

Finn, S.B., Perry, B.N., Clasing, J.E., Walters, L.S., Jarzombek, S.L., Curran, S., Rouhanian, M., Keszler, M.S., Hussey-Andersen, L.K., Weeks, S.R., Pasquina, P.F., Tsao, J.W. (2017). A Randomized, Controlled Trial of Mirror Therapy for Upper Extremity Phantom Limb Pain in Male Amputees. *Front Neurol* 8, 267.

Flor, H. (2002). Phantom-limb pain: characteristics, causes, and treatment. *Lancet Neurol* 1, 182–189.

Flor, H. (2008). Maladaptive plasticity, memory for pain and phantom limb pain: Review and suggestions for new therapies. *Expert Rev Neurother* 8, 809–818.

Flor, H., Nikolajsen, L., Jensen, T.S. (2006). Phantom limb pain: A case of maladaptive CNS plasticity? *Nat Rev Neurosci* 7, 873–881.

Foell, J., Bekrater-Bodmann, R., Diers, M., Flor, H. (2014). Mirror therapy for phantom limb pain: Brain changes and the role of body representation: Mirror therapy for phantom limb pain. *Eur J Pain* 18, 729–739.

Ganz, J.B., Ayres, K.M. (2018). Methodological standards in single-case experimental design: Raising the bar. *Res Dev Disabil*.



Herr, H.M., Clites, T.R., Srinivasan, S., Talbot, S.G., Dumanian, G.A., Cederna, P.S., Carty, M.J. (2020). Reinventing Extremity Amputation in the Era of Functional Limb Restoration. *Ann Surg* Publish Ahead of Print.

Herrador Colmenero, L., Perez Marmol, J.M., Martí-García, C., Querol Zaldivar, M. de los Á., Tapia Haro, R.M., Castro Sánchez, A.M., Aguilar-Ferrándiz, M.E. (2018). Effectiveness of mirror therapy, motor imagery, and virtual feedback on phantom limb pain following amputation: A systematic review. *Prosthet Orthot Int* 42, 288–298.

Knotkova, H., Cruciani, R.A., Tronnier, V.M., Rasche, D. (2012). Current and future options for the management of phantom-limb pain. *J Pain Res* 5, 39–49.

Krasny-Pacini, A., Evans, J. (2018). Single-case experimental designs to assess intervention effectiveness in rehabilitation: A practical guide. *Ann Phys Rehabil Med* 61, 164–179.

Kuner, R., Flor, H. (2016). Structural plasticity and reorganisation in chronic pain. *Nat Rev Neurosci* 18, 20–30.

Liberati, A., Altman, D.G., Tetzlaff, J., Mulrow, C., Gøtzsche, P.C., Ioannidis, J.P.A., Clarke, M., Devereaux, P.J., Kleijnen, J., Moher, D. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol* 62, e1–e34.

Makin, T.R., Scholz, J., Filippini, N., Henderson Slater, D., Tracey, I., Johansen-Berg, H. (2013). Phantom pain is associated with preserved structure and function in the former hand area. *Nat Commun* 4, 1570.

Moore, R.A., Fisher, E., Eccleston, C. (2022). Systematic reviews do not (yet) represent the ‘gold standard’ of evidence: A position paper. *Eur J Pain* 26, 557–566.

Nikolajsen, L., Jensen, T.S. (2001). Phantom limb pain. *Br J Anaesth* 87, 107–116.

Niraj, S., Niraj, G. (2014). Phantom Limb Pain and Its Psychologic Management: A Critical Review. *Pain Manag Nurs* 15, 349–364.

Perry, B.N. (2013). Twenty-third Meeting of the European Neurological Society, 8–11 June, 2013 Barcelona, Spain. *J Neurol* 260, 1–280.

Plumbe, L., Peters, S., Bennett, S., Vicenzino, B., Coppeters, M.W. (2013). Mirror therapy, graded motor imagery and virtual illusion for the management of chronic pain. In *Cochrane Database of Systematic Reviews*, The Cochrane Collaboration, ed. (Chichester, UK: John Wiley & Sons, Ltd), p. CD010329.

Ramachandran (1996). Synaesthesia in phantom limbs induced with mirrors. 10.

Ramachandran, V.S., Altschuler, E.L. (2009). The use of visual feedback, in particular mirror visual feedback, in restoring brain function. *Brain* 132, 1693–1710.

Ramadugu, S., Nagabushnam, S.C., Katuwal, N., Chatterjee, K. (2017). Intervention for phantom limb pain: A randomized single crossover study of mirror therapy. *Indian J Psychiatry* 59, 457–464.

Rothgangel, A., Braun, S., Winkens, B., Beurskens, A., Smeets, R. (2018). Traditional and augmented reality mirror therapy for patients with chronic phantom limb pain (PACT study): results of a three-group, multicentre single-blind randomized controlled trial. *Clin Rehabil* 32, 1591–1608.

Rothgangel, A.S., Braun, S.M., Beurskens, A.J., Seitz, R.J., Wade, D.T. (2011). The clinical aspects of mirror therapy in rehabilitation: a systematic review of the literature. *Int J Rehabil Res* 34, 1–13.

van der Schans, C.P., Geertzen, J.H.B., Schoppen, T., Dijkstra, P.U. (2002). Phantom Pain and Health-Related Quality of Life in Lower Limb Amputees. *J Pain Symptom Manage* 24, 429–436.

Schone, H.R., Baker, C.I., Katz, J., Nikolajsen, L., Limakatso, K., Flor, H., Makin, T.R. (2022). Making sense of phantom limb pain. *J Neurol Neurosurg Psychiatry* 93, 833–843.

Sinha, R., van den Heuvel, W.J., Arokiasamy, P. (2011). Factors affecting quality of life in lower limb amputees. *Prosthet Orthot Int* 35, 90–96.

Smith, J.D. (2012). Single-case experimental designs: A systematic review of published research and current standards. *Psychol Methods* 17, 510–550.

Sütbeyaz, S., Yavuzer, G., Sezer, N., Koseoglu, B.F. (2007). Mirror Therapy Enhances Lower-Extremity Motor Recovery and Motor Functioning After Stroke: A Randomized Controlled Trial. *Arch Phys Med Rehabil* 88, 555–559.

Tate, R.L., McDonald, S., Perdices, M., Togher, L., Schultz, R., Savage, S. (2008). Rating the methodological quality of single-subject designs and n-of-1 trials: Introducing the Single-Case Experimental Design (SCED) Scale. *Neuropsychol Rehabil* 18, 385–401.

Thieme, H., Morkisch, N., Mehrholz, J., Pohl, M., Behrens, J., Borgetto, B., Dohle, C. (2018). Mirror therapy for improving motor function after stroke. *Cochrane Database Syst Rev* 2018.

Thieme, H., Morkisch, N., Rietz, C., Dohle, C., Borgetto, B. (2016). The Efficacy of Movement Representation Techniques for Treatment of Limb Pain—A Systematic Review and Meta-Analysis. *J Pain* 17, 167–180.

Wang, F., Zhang, R., Zhang, J., Li, D., Wang, Y., Yang, Y.-H., Wei, Q. (2021). Effects of mirror therapy on phantom limb sensation and phantom limb pain in amputees: A systematic review and meta-analysis of randomized controlled trials. *Clin Rehabil* 35, 1710–1721.

Woolf, C.J., Mannion, R.J. (1999). Neuropathic pain: aetiology, symptoms, mechanisms, and management. *The Lancet* 353, 1959–1964.

Woolf, C.J., Shortland, P., Coggeshall, R.E. (1992). Peripheral nerve injury triggers central sprouting of myelinated afferents. *Nature* 355, 75–78.

World Health Organization, World Bank (2011). *World report on disability 2011*.

Xie, H.-M., Zhang, K.-X., Wang, S., Wang, N., Wang, N., Li, X., Huang, L.-P. (2021). Effectiveness of Mirror Therapy for Phantom Limb Pain: A Systematic Review and Meta-analysis. *Arch Phys Med Rehabil* S0003999321013794.

Table 1: Index terms and syntax for each database.

Database	Index terms and syntax
<b>PubMed</b>	("Amputation"[Mesh] OR "Amputees"[Mesh]) AND ("Phantom Limb/therapy"[Mesh]) AND (mirror* OR mirror therap* OR mirror box).
<b>Cochrane Library</b>	amput* AND (phantom limb OR phantom pain) AND (mirror therapy* OR mirror* OR mirror box)
<b>CINHAL</b>	((MH "Phantom Limb") OR (MH "Phantom Pain")) AND ((MH "Amputation") OR (MH "Amputees")) AND ((MH "Mirror Therapy") OR mirror* OR mirror therapy* OR mirror box)
<b>PEDro</b>	amput* AND "phantom limb pain" AND mirror*  <i>The "clinical trial" filter was added to limit the search to clinical trials</i>
<b>Embase</b>	('amputation'/exp OR 'amputee'/exp) AND 'phantom pain'/exp AND ('mirror therapy'/de OR 'mirror'/de)  <i>The filter "Embase" was used to eliminate articles from PubMed and keep only articles that are indexed only in Embase</i>

Table 2: PEDro score of the selected studies

First author	Items											Total score /10
	1	2	3	4	5	6	7	8	9	10	11	
Brodie et al.[28]	1	1	0	1	0	0	0	1	1	1	1	6
Chan et al.[29]	0	1	0	1	0	0	0	1	0	1	0	4
Finn et al.[30]	1	1	0	1	0	0	0	1	1	0	0	4
Ramadugu et al.[31]	1	1	1	1	0	0	1	1	0	1	1	7
Rothgangel et al.[32]	1	1	1	1	0	0	1	1	1	1	1	8

Criterion 1: eligibility criteria were specified; criterion 2: subjects were randomly allocated to groups; criterion 3: allocation was concealed; criterion 4: at baseline, the most important prognostic indicators were similar between groups; criterion 5: blinding of all subjects; criterion 6: blinding of all therapists who administered the therapy; criterion 7: blinding of all assessors who measured at least one key outcome; criterion 8: measures of at least one key outcome were obtained from more than 85% of the subjects initially allocated to groups; criterion 9: all subjects for whom outcome measures were available received the treatment or control condition, as allocated; criterion 10: the results of between-group statistical comparisons are reported for at least one key outcome; criterion 11: the study provides both point measures and variability measures for at least one key outcome.

Table 3: Characteristics of the patients included in the selected studies

Author, years	Pathology	Age (years), sex	PLP intensity and/or number of PLP episodes at recruitment	Interval between amputation and inclusion in the study
Brodie et al. 2007	Unilateral lower limb amputees Reason for amputation: congenital (1%), cancer (5%), accident (33%), medical (61%)	N = 80; ♂ (79%) Mean age: 55 (20-83)	/	<2 weeks: 65% <1 year: 20% >1 year: 6%
Chan et al. 2007	Unilateral lower limb amputees Reason for amputation: not specified	N = 22	/	/
Finn et al. 2017	Unilateral upper extremity amputees Reason for amputation: accident (100%)	N=15; ♂ (100%) Mean age: 28.7 (19 - 68)	Episodes per week: 3	Mean: 4.51 months (0.55 – 24)
Ramadugu et al. 2018	Unilateral lower limb (84%) and unilateral upper extremity amputees (16%) Reason for amputation: not specified  Non-inclusion criteria: traumatic brain injury or major psychiatric illness	N= 64; ♂ (100%) Mean age: NA (17 - 62)	Intensity (VAS): 30 (10.7)	/
Rothgangel et al. 2018	Unilateral lower limb amputees Reason for amputation: accident (33%), diabetes (11%), vascular problems (29%), tumor (13%), other (e.g., infection) (13%)	N = 75; ♂ (69%) Mean age: 61.1 (NP)	Episodes per week: at least one  Intensity (VAS): 5.7	Mean: 3 years

NA, not available; VAS, visual analogue scale.

Author, year, study type	Intervention group	Control intervention	Measurements: - Primary outcome - Secondary outcome(s)	Procedure	Supervised intervention (Yes/No)	Number of sessions and session duration	Fidelity to the programmed intervention
Brodie et al. 2007  RCT - parallel group	Mirror therapy N=41	Covered mirror N=39	- 100-mm VAS  - MPQ	Ten movements with both limbs, each repeated ten times, with a break between each movement type.	Yes	Only 1 session	/
Chan et al. 2007  RCT - crossover for the control group	Mirror therapy N=6 (number of participants who completed the study)	Covered mirror N=6 (number of participants who completed the study)	- 100-mm VAS	Movements with both limbs.	Yes	15 minutes per day, for 8 weeks (for the control group, covered mirror the first 4 weeks then change to mirror therapy).	At 4 weeks, all patients completed the study.
Finn et al. 2017  RCT – parallel group	Mirror therapy N=9	Covered mirror N=3	- 100-mm VAS  - Number of PLP episodes per day  - PLP duration	Both limbs perform slow movements with progressive increase in amplitude.	Yes	15 minutes per day, 5 days per week for 4 weeks.	Due to lack of efficacy of the control treatment, all participants in the control group received the MT intervention after 11 sessions (planned change at 20 sessions initially).
Ramadugu et al. 2018  RCT - crossover for the	Mirror therapy N=32	Covered mirror N=28	- 100-mm VAS  - SF-MPQ  - PLP frequency (Likert scale)	Movements with both limbs	Yes	15 minutes per day, for 4 weeks	All patients in the mirror group completed the study and 86% of participants in the control group completed the study.

control group			- PLP duration (Likert scale)				
Rothgangel et al. 2018  RCT - parallel group	Mirror therapy N=25	Mirrorless N=24	<ul style="list-style-type: none"> <li>- NRS – 10 points</li> <li>- PLP frequency (Likert scale)</li> <li>- PLP duration (Likert scale)</li> <li>- NPSI</li> <li>- PSFS</li> <li>- PDI</li> <li>- 11-point NRS on mood and sleep</li> <li>- EQ-5D–5L</li> <li>- 100-mm VAS overall health</li> <li>- Pain self-efficacy questionnaire</li> <li>- Global perceived effect scale</li> </ul>	<p>Intervention group: exercise program adapted to the patient's preferences with both limbs.</p> <ul style="list-style-type: none"> <li>- Observation of different positions</li> <li>- Motor exercises</li> <li>- Exercises using sensory stimuli (e.g. vibration, heat)</li> <li>- Functional motor exercises with objects.</li> </ul>	Yes, during the first 4 weeks. Then unsupervised for the next 6 weeks (home).	10 sessions of at least 30 minutes/each for the first 4 sessions As many sessions as the patient wishes during the following 6 weeks.	84% of participants in the intervention group completed the study and 80% of participants in the control group completed the study.



RCT randomized controlled trial; VAS visual analogue scale; MPQ McGill pain questionnaire; SF-MPQ Short McGill pain questionnaire; NRS numerical rating scale; NPSI Neuropathic Pain symptom inventory; PSFS patient-specific functional scale; PDI pain disability index.

Table 4: Summary of the characteristics, intervention, and outcomes of the selected studies.




Figure 1: Study selection flowchart according to the PRISMA guidelines.  
RCT, randomized controlled trial.