



Microalgae screening for heterotrophic and mixotrophic growth on butyrate

Julien Lacroux, P. Jouannais, A. Atteia, A. Bonnafous, Eric Trably,
Jean-Philippe Steyer, Robert van Lis

► To cite this version:

Julien Lacroux, P. Jouannais, A. Atteia, A. Bonnafous, Eric Trably, et al.. Microalgae screening for heterotrophic and mixotrophic growth on butyrate. *Algal Research - Biomass, Biofuels and Bioproducts*, 2022, 67, pp.102843. 10.1016/j.algal.2022.102843 . hal-03778414

HAL Id: hal-03778414

<https://hal.science/hal-03778414v1>

Submitted on 27 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Microalgae screening for heterotrophic and mixotrophic growth on butyrate

J. Lacroux¹, P. Jouannais¹, A. Atteia², A. Bonnafe¹, E. Trably¹, J-P. Steyer¹, R. van Lis^{1*}

¹INRAE, Univ Montpellier, LBE, 102 avenue des Etangs, F-11100, Narbonne France

²MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Sète, France.

*Corresponding author: robert.van-lis@inrae.fr

Keywords: microalgae, heterotrophy, mixotrophy, volatile fatty acids, lipids, carbohydrates

Abstract

Volatile fatty acids (VFAs) produced by fermentative bacteria can be used instead of high value carbohydrates like glucose to reduce the costs associated to mixotrophic microalgal cultivation. This process is however limited by a poor assimilation of butyrate and a low tolerance of microalgae to high VFA concentrations. In this study, 10 microalgae strains, isolated from local natural environments or selected from culture collections were screened for their ability to grow on either acetate or butyrate. The non-photosynthetic Chlorophyte *Polytomella* sp. was found to be the fastest growing strain on both VFAs, exhibiting a growth rate of 4.0 d⁻¹ on acetate and 2.5 d⁻¹ on butyrate. The tolerance of the different strains to concentrated VFA was further assessed using increasing acid concentrations (1.5-60 g.L⁻¹ for acetate and 1-40 g.L⁻¹ for butyrate). *Polytomella* sp. appeared as the most resistant, being able to grow up to 38 g.L⁻¹ on acetic acid and to 18 g.L⁻¹ on butyric acid with no inhibitory effect. Finally, biomass yield and productivity, as well as lipid and carbohydrate yields were assessed

on a synthetic VFAs mixture (0.8 g.L⁻¹ acetate and 1.2 g.L⁻¹ butyrate) for the most promising strains. *Polytomella* sp. and *Euglena gracilis* exhibited the highest carbohydrate yield (respectively 0.65 and 0.58 g.g⁻¹) while lipid yield was maximum for *Chlorella sorokiniana* (0.42 g.g⁻¹). This work provides new insights in the trophic metabolism of microalgae of highly divergent lineages using VFAs as substrates.

1. Introduction

Dark fermentation (DF) is an anaerobic biological process allowing concomitant production of energy and treatment of various organic waste sources such as food or crop wastes. During DF, anaerobic bacteria convert complex organic matter into hydrogen and a mixture of end-product metabolites. These metabolites can vary according to fermentation conditions and substrate nature, but are on average mainly composed of volatile fatty acids (VFAs) particularly acetate and butyrate [1]. These VFAs can serve as low-cost organic substrates for the growth of various micro-organisms either non photosynthetic such as yeast and thraustochytrids or photosynthetic such as microalgae [2,3].

Butyrate was identified as a key factor limiting the potential of such a coupling, particularly in the case of microalgae [4,5] due to inhibitory effects [6] or inability to grow on more than 1.25 g.L⁻¹ of butyrate without light [4]. Several attempts have been made to increase the consumption of butyrate by acting on operational parameters such as light and temperature [7], higher acetate content in the medium [6] or pH adjustment [8]. However, even if butyrate inhibition can be alleviated using these strategies, butyrate uptake rates remain about 10-fold lower as compared to acetate [9]. So far, most research published on the coupling of DF and microalgae cultivation has focused on lipid production using strains such as *Chlorella* or *Scenedesmus* species obtained from collections [5,6,9–11]. However, strains

isolated from organic rich environments may be better adapted to the consumption of butyrate. Indeed, collection strains are usually maintained on designed media, leading to an adaptation to laboratory conditions. As a result, these strains may exhibit lower tolerance or lower growth on alternative substrates compared to environmental isolate species [12].

If photosynthetic Chlorophytes seem limited in their ability to consume DF metabolites, purely heterotrophic species could be more adapted to grow on organic acids since they rely entirely on organic carbon for growth. As an illustration, Chalima et al., [13] tested the growth of the heterotrophic marine dinoflagellate *Cryptothecodinium cohnii* on various VFAs (C2-C4). The strain could grow on acetic, propionic and butyric acids at 30 g.L⁻¹, 10 g.L⁻¹ and 15 g.L⁻¹, respectively. Moreover, since most of the aforementioned studies mostly focused on lipid production, other microalgal species which accumulate polysaccharides have received little attention. Some freshwater strains, such as *Euglena gracilis*, *Ochromonas danica* or *Polytomella* sp. possess indeed several traits that make them potential candidates to be cultivated on DF effluents. For example, *E. gracilis* is a photosynthetic protist which accumulates up to 80% of its weight as a valuable polysaccharide resembling starch (paramylon) and that can grow in various nutritional modes such as photo-autotrophy, heterotrophy or mixotrophy [14]. *O. danica*, a chrysophyte (golden-brown microalgae), can grow under mixotrophic condition on organic acids and can also prey on bacteria [15]. Since it is undesirable to sterilize DF effluents due to operational costs, these effluents will contain bacteria that will likely outcompete microalgae for butyrate [16]. These bacteria could be eaten by *O. danica* and therefore increase final algal biomass. Finally, *Polytomella* sp. is a colourless Chlorophyte alga closely related to the model organism *Chlamydomonas reinhardtii*, which has lost its photosynthetic properties [17,18]. This alga grows purely heterotrophically and is able to consume various organic acids and alcohols while accumulating large amount of starch [19].

The objective of this study was to identify microalgal strains that could tolerate high concentrations of butyrate and consume it more efficiently than the most commonly studied species (i.e. Chlorophytes such as *Chlorella* sp. or *Scenedesmus* sp.). To this end, several microalgal strains were isolated from different local natural environments using butyrate as selective factor. In parallel, the pure strains *Euglena gracilis*, *Ochromonas danica*, *Polytomella* sp. and *Polytomella magna* were selected as alternative candidates for their mixo- or heterotrophic capacities. The growth of all these strains on acetate and butyrate as well as their tolerance to VFAs were assessed and compared to that of *C. sorokiniana*. Finally, the biomass, lipid and carbohydrate production potentialities of the candidate strains were evaluated on a synthetic DF effluent.

2. Materials and methods

2.1. Cultivation media and conditions

Microalgae were maintained and precultured on synthetic medium HEPES-Acetate-Phosphate (HAP), which is based on Tris-Acetate-Phosphate (TAP) medium [20] where HEPES buffer replaces Tris. When butyrate was used instead of acetate, the medium is referred to as HBP (HEPES-Butyrate-Phosphate). Buffer concentration was set to 20 mM for maintenance and precultures. To evaluate the growth characteristics, acetate or butyrate were added individually as sodium salts at a concentration of 1.25 g.L⁻¹ or 0.9 g.L⁻¹ respectively to set carbon concentration to 0.5 g.C.L⁻¹ for both substrates. Beijerincks solution (<https://utex.org/products/beijerincks-solution>) was added at 25 mL.L⁻¹ leading to concentration of 7.5 mM NH₄⁺, 0.6 mM MgSO₄ and 0.3 mM CaCl₂. Potassium phosphate (K₂HPO₄) was added at 1 mM and Hutner's trace element solution [21] at 1 mL.L⁻¹. Medium pH was adjusted to 7.0 by addition of NaOH prior to sterilization at 121°C for 20 min. After

cooling, 100 $\mu\text{L.L}^{-1}$ of a vitamin mix (50 mM vitamin B1, 1 mM biotin and 1 mM cyanocobalamin), sterilized by filtration through 0.2 μm filter, was added into the medium. To obtain solid medium, agar was added before sterilization (15 g.L^{-1}). All chemicals were purchased from Sigma Aldrich (Merck).

2.2. Microalgal strains

Chlorella sorokiniana SAG211-8k, *Euglena gracilis* SAG 1224-5/15, *Ochromonas danica* SAG933-7, *Polytomella* sp. Pringsheim SAG 198.80 and *Polytomella magna* SAG 63-9 were purchased from the Culture Collection of Algae in Göttingen (Sammlung von Algenkulturen Göttingen [SAG]; <http://www.epsag.uni-goettingen.de>). *Polytomella* strains were maintained in liquid HAP cultures at 25°C without light control, while the photosynthetic strains were maintained on HAP agar plates under constant illumination ($100 \pm 10 \mu\text{mol.m}^{-2}.\text{s}^{-1}$). Five other strains were obtained through isolation as described in section 2.3.

Algae were inoculated in liquid medium prior to growth studies. The algal suspensions were centrifuged (735 g, 10 min), the supernatant discarded and the biomass dispersed in sterile phosphate buffer saline (PBS) medium. In each experiment, the initial microalgal concentration was set to an optical density (OD_{750}) of 0.05. All cultivations were carried out in triplicates, under constant illumination using cool-white fluorescent lamps ($100 \pm 10 \mu\text{mol.m}^{-2}.\text{s}^{-1}$) at 25°C under agitation of 120 rpm (2.5 cm orbit). Both *Polytomella* strains were cultivated without shaking and light control. Strain screening and undissociated acid tolerance determination were carried out in microplates as described in [8,22]. Cultivations to evaluate the biomass production potential on synthetic DF effluent (composition described in section 2.5) were carried out in 500 mL Erlenmeyer flasks, filled with 200 mL medium.

2.3. Sampling and isolation procedure of strains from environmental habitats

Three natural peat bogs were sampled in February 2020 in the Occitanie region, France. Two of them were acidic (pH 6), named hereinafter A1 (43°38'47.33''N/2°34'36.02''E) and A2 (44°14'45.69''N/2°56'2.53''E) and one of them was alkaline (pH 8), named hereinafter B (44°27'8.51''N/2°0'27.73''E). Samples were taken from both the water surface (Code WS) and from the first 30-cm of the peat (Code P). Besides peat bogs, two aquatic ecosystems rich in organic matter were sampled, namely a wastewater treatment plant (code WWTP) and a microalgal raceway (code RW) fed with synthetic wastewater. Samples were stored at 4°C until processing (maximum a day after sampling).

Environmental aqueous samples (15 mL) were directly streaked onto HBP agar plates and incubated at 25°C under constant illumination ($100 \pm 10 \mu\text{mol.m}^{-2}.\text{s}^{-1}$). After about 10-15 days, green colonies developed. They were picked and streaked on fresh HBP plates. In the case of peat samples, 100 g of peat were mixed for 5 min with 500 mL isotonic water, and prefiltered on coffee filters to remove large solids. Microorganisms in the filtrate were collected via a second filtration step on a 0.7 μM GF/F Whatman® filter. The filters were subsequently used to inoculate two flasks filled with HAP medium (40 mL), either with 100 mg.L^{-1} ampicillin, in order to prevent excessive growth of bacteria, or without to allow any antibiotic sensitive microalgae to grow. Flasks were then placed under constant illumination ($100 \mu\text{mol.m}^{-2}.\text{s}^{-1}$) at 25°C and shaking (130 rpm). After 10 days, these enriched cultures were streaked onto HBP agar plates, incubated at 25°C under constant illumination. Colonies were picked based on colour (green) and streaked until purification. All algal strains were maintained on HAP agar plates.

2.4 Molecular identification and phylogenetic analysis of isolates

Microalgal strains were first identified by 18S rRNA sequencing. Total DNA was first extracted from growing cells using the DNeasy PowerWater Kit (Qiagen). The 18S rRNA (400 bp) amplicon was amplified by PCR using the primers 5'-CTTTCCTACACGACGCTCTTCCATCTGCGGTAATTCCAGCTCCAA-3' (541F) and 5'-GCAGTTCAGACGTGTGCTCTTCCGATCTTTGGCAAATGCTTTCGC-3' (952R). PCR products were purified and sequences by Illumina MiSeq at Genotoul Get-Plage (get.genotoul.fr). After cleaning, sequenced were sorted by OTU with a similarity of 97%. This first identification only allowed identification at the genus level.

To identify strains at the species level, further identification was done. A rapid extraction of total cell DNA from the algal colonies growing on solid media was performed according to Newman et al. [23] with slight modifications. Briefly, cells were collected from plates and resuspended in a 1.5 ml microfuge tube containing 0.2 mL TEN buffer (150 mM NaCl, 10 mM Na-EDTA, 10 mM Tris-HCl, pH 8.0). Cell lysis was done in 0.4 mL of SDS-EB buffer (2% SDS, 400 mM NaCl, 50 mM EDTA, 100 mM Tris-HCl pH 8) containing RNase A at a final concentration of 40 µg.µL⁻¹ and incubated at 37°C for 30 min. DNA was extracted with phenol/chloroform/isoamyl alcohol in a ratio 25:24:1 (v/v/v) and then with chloroform/isoamyl alcohol (24:1 v/v). Total DNA was precipitated by addition of isopropyl alcohol, collected by centrifugation and washed with cold 70% ethanol, air dried and resuspended in TE buffer.

The nuclear sequence spanning the genes for 18S ribosomal RNA-ITS1-5.8S rRNA-ITS2-28S rDNA (500 first bases of 28S) region was amplified by PCR using *Pfu* DNA polymerase (Promega) and the pair of primers EAF3 (TCGACAATCTGGTTCACCTAC)/ITS055R (CTCCTTGGTCCGTGTTTCAAGACGGG) [24]. The PCR products were purified on a QIAquick Gel Extraction Kit (Qiagen) and sequenced directly at Eurofins Genomics (<https://eurofinsgenomics.eu>) with universal and

specific primers. Sequences were submitted to Genbank and accession numbers are provided in Table 1.

A phylogenetic tree was constructed with the MEGA-X program [25] using whole 18S rRNA gene sequences. The MUSCLE module was used to align the DNA sequences using standard parameters. Only the consensus sequence was used, removing highly divergent ends. A Maximum Likelihood tree was produced from this alignment using standard parameters.

2.5. Undissociated acid inhibition thresholds

The tolerance threshold to undissociated acid concentration was determined according to [8]. Briefly, several media were prepared as described in section 2.1, with modifications in pH, buffer and substrate concentration. For each medium, the pH was set at 6.5 and buffered with 100 mM MES buffer. Increasing total concentrations of either acetate or butyrate were added to the media to obtain the desired range of acetic or butyric acid respectively, determined by the Henderson-Hasselbalch equation (1).

$$[RH] = \frac{C_t}{1+10^{(pH-pKa)}} \quad (1)$$

With [RH] the concentration of either acetic or butyric acid (g.L⁻¹), C_t the total concentration of acetate or butyrate (g.L⁻¹) and pKa the dissociation constant of each acid (close to 4.8 for both acetate and butyrate).

2.6. Biomass production potential

Biomass production potential of candidate strains was evaluated by cultivating them on synthetic DF medium in Erlenmeyer flasks. DF medium was prepared as described in section 2.1., with modifications in buffer and substrate concentration. VFA concentration was

1 set to 1.0 g_C.L⁻¹ by mixing 0.33 g_C.L⁻¹ (0.8 g.L⁻¹) acetate and 0.67 g_C.L⁻¹ (1.2 g.L⁻¹) butyrate,
2 resulting in an acetate:butyrate carbon mass ratio of 1:2. This ratio is representative of an
3 average DF effluent as reported [1]. Buffer concentration was set to 100 mM and pH was set
4 to 7.0. Molar C:N:P ratio was set to 106:16:1 by adjusting the amount of ammonium and
5 phosphate salts (N-NH₄ = 225 mg.L⁻¹ and P-PO₄ = 75 mg.L⁻¹). Media were sterilized by
6 autoclave (121°C, 20 min).
7
8
9
10
11
12
13
14
15
16
17

18 *2.7. Biomass measurement*

19
20
21 Biomass growth in microplates was determined via the measurement of the optical
22 density at 750 nm (OD₇₅₀) by an Infinite Nanoquant M200 (TecanR[®]) spectrophotometer.
23
24 Biomass production in flasks was determined by dry weight determination (DW) after
25 filtration of 2 – 20 mL of samples (depending on biomass concentration) over a dried
26 Whatman[®] GF/C glass microfiber filter (1.2 µm) and overnight drying at 105°C.
27
28
29
30
31
32
33
34
35
36

37 *2.8. Volatile fatty acid measurements*

38
39
40 Acetate and butyrate concentrations in the growth media were quantified by gas
41 chromatography. Cultures were sampled at indicated growth times, and centrifuged at 15,000
42 rpm for 10 min. Supernatants were collected, filtered through a 0.45 µm filter and stored at -
43 25°C. Before analysis, 500 µL of supernatant were mixed with 500 µL of internal standard
44 solution (ethyl-2-butyric acid, 1 g.L⁻¹). VFAs were analyzed on a Perkin Clarus 580 with
45 capillary column Elite-FFAP crossbond carbowax[®] (15 m) maintained at 200°C and with N₂
46 as the gas vector (flow rate of 6 mL.min⁻¹) equipped with a flame ionization detector (FID)
47 maintained at 280°C (PerkinElmer, USA).
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2.9. Lipids and sugars measurements

Samples from fresh cultures were immediately centrifuged, the supernatant was discarded and the pellet stored at -20°C until use. Prior to analysis, the pellets were thawed, resuspended in distilled water and transferred to a 10 mL glass tubes for either lipid or sugar measurement.

Total lipids were measured by the phosphovanillin method [26]. Phosphovanillin reagent was freshly prepared before analysis by dissolving 0.6 g vanillin in 10 ml ethanol, 90 ml deionized water and 400 ml of H₃PO₄ (85%). The resulting reagent was stored in the dark until use. First, 2 mL of H₂SO₄ (98%) were added in the tubes containing microalgae samples. The tubes were heated 10 min at 100°C. After cooling on ice, the reaction was initiated by addition of 5 mL of phospho-vanillin reagent prior incubation for 15 min at 37°C. Tubes were periodically shaken by inversion and were then left in the dark at room temperature for 45 min for color development. After cooling, absorbance of suspensions was measured at 530 nm with an Aqualytic[®] spectrometer and compared to distilled water. Calibration curves were obtained using canola oil.

Total sugars were measured by the anthrone method [27]. Anthrone reagent was prepared by dissolving 200 mg of anthrone in 100 mL of H₂SO₄ (98%). Two mL of anthrone reagent were added in the tubes containing microalgae. Tubes were cooled down on ice and then incubated at 100°C for 10 min. After cooling, absorbance of suspensions was measured at 625 nm with an Aqualytic[®] spectrometer and compared to distilled water. Calibration curves were obtained using glucose solution.

3. Results and discussion

3.1. Isolation and identification of microalgae strains

The first strategy followed to isolate environmental microalgae able to use butyrate as carbon source was to explore natural ecosystems featuring characteristics similar to anaerobic fermenters. Therefore, samples were taken from different wetlands (peatbogs), which are organic rich ecosystems flooded by water either seasonally or permanently. The high water table limits gas exchanges and makes the ecosystem largely anoxic. Organic matter is thus degraded through anaerobic bioprocesses leading to fermentative by-products such as acetate, butyrate or hydrogen [28]. In addition, aqueous samples were taken from artificial ecosystems: WWTP activated sludge and biomass from a microalgae raceway fed with synthetic wastewater. Although mostly aerobic, these open ecosystems contain high amount of organic matter. It was expected that microalgae isolated from these environments would exhibit the ability to compete for organic carbon to some extent with bacteria (and thus high growth rate) and tolerance to organic carbon. From all five environmental samples, green algal colonies developed on butyrate-containing plates. Colonies were first identified by metabarcoding (MiSeq). The OTU could not allow identification to the species level but allowed genus discrimination. It could be concluded that each isolate contained a single genus and that all five algae belong to the Chlorophyte division. Thus, a total of five different algal strains could be obtained as pure culture, after enrichment on a butyrate containing medium (Table 1).

Table 1: Summary of the isolated species and their identification. Isolates were sampled from different peatbogs water surface or peats, a local wastewater treatment plant (WWTP) sludge

and a local microalgal raceway. Identification was done by sequencing the nuclear ribosomal DNA operon and BLAST searches.

Strain code	Sampling source	Closest strain	Sequence alignment
<i>Chlamydomonas</i> <i>asymmetrica</i> A2WS (2993 bp) (ON077054)	Peatbog water surface	<i>Chlamydomonas</i> <i>asymmetrica</i> CCAP 11/41 (2557 bp) FR865583	2524/2557
<i>Chlorella sorokiniana</i> A1WS (3070 bp) (ON077055)	Peatbog water surface	<i>Chlorella sorokiniana</i> NIES:2173 (2487 bp)	G1826 T (ITS1)
<i>Chlorella sorokiniana</i> BP (3070 bp) (ON077056)	Peatbog peat	<i>Chlorella sorokiniana</i> SAG 211-8k (2461 bp) FMZ05834	Identical to the lab strain
<i>Desmodesmus armatus</i> WWTP (2955 bp) (ON077057)	WWTP sludge	<i>Desmodesmus armatus</i> GM4k (2591 bp) AB17138	A deletion T 1803 ITS1
<i>Desmodesmus armatus</i> RW (2955 bp) (ON077058)	Microalgal raceway		

However, the 400 bp amplicon did not allow precise identification of each species. The rDNA operon has emerged as a tool for species delimitation, and hence, identification [29]. To identify at the molecular level all isolated algal species, the nuclear sequence of the rDNA operon (spanning the 18S rDNA, ITS1, 5.8S, ITS2 and the 28S rDNA regions) was thus amplified and sequenced. From the peat bogs were isolated two algae closely related to *Chlorella sorokiniana* (Trebouxiophyceae) SAG 211-8k strain, used routinely in the laboratory. The rDNA sequence of the isolate *C. sorokiniana* BP was identical to this reference strain while the isolate *C. sorokiniana* A1WS differs in one single base located in

the ITS1 (Table 1). The third algal species isolated from peat bogs was identified as *Chlamydomonas asymmetrica* (Chlorophyceae). From the artificial ecosystems (WWTP sludge and raceway) were isolated two algae identified as *Desmodesmus armatus*. These two species differ from the *D. armatus* strain GM4k SAG in one nucleotide deletion in the ITS1 region. Due to the very high sequence identity of the isolated strains with published strains over the whole length of the rDNA operon including the highly variable ITS regions, species identification was highly confident (BLAST e-value=0).

Overall, very little diversity was obtained throughout the isolation process since all isolated algae belong to only three species of Chlorophytes. Yet, other species such as desmidiaceae and diatoms were detected during microscope observation of the environmental samples and greater diversity in peat bog samples was already reported elsewhere in the literature [30]. It is thus possible that the media HAP or HBP used for the enrichment from the environmental samples are relatively selective for Chlorophytes. For example, using acetate containing media, similar results were obtained by other authors who isolated four *C. sorokiniana* strains from a local wetland (Lake Massaciuccoli located in Tuscany, Italy) [30,31]. Implementing HAP/HBP media with silica might have allowed the emergence of diatoms. Besides, the other major nutritive elements, N and P, were added in the same form that the one found in DFEs, respectively NH_4^+ and PO_4^{3-} , consequently adding an extra selection pressure. *Chlorella* and *Desmodesmus* are reported as “lab weeds” due to their ability to grow over a wide range of cultivation conditions [32]. These strains are thus often isolated from various ecosystems and with various selection conditions [31,33,34]. It would be undeniably very interesting to use a comprehensive isolation procedure in order to obtain a maximum of species capable of growing or tolerating butyrate. Nevertheless, in the context of microalgae cultivation on DFE effluent, some specific constraints such as the one evoked above (chemical form of the nutrients, absence of certain element) should still be considered.

Therefore, other strains obtained via an extensive screening, which would be extremely efficient at consuming butyrate in specific growth condition, may not be adapted to grow on the more adverse and variable conditions of a DFE.




Figure 1: Maximum Likelihood tree on the investigated species. Since isolate sequences are nearly identical to reported sequences and will not show on the tree, only the reported sequences were included in the tree. Accession numbers of the sequences used were: *Polytomella* sp., FR750564.1; *P. magna*, KP299175.1; *C. asymmetrica*, FR865607.1; *C. reinhardtii*, FR865575.1; *C. sorokiniana*, LK021940.1; *A. protothecoides*, LN610701.1; *A.*

1 *obliquus*, AB917100.1; *D. armatus*, AB917138.1; *A. thaliana*, X16077.1; *O. danica*,
2 JQ281514.1; *E. gracilis*, AJ532426.1.
3
4

5 As can be seen on the Maximum Likelihood tree (Fig. 1), all isolated species were
6 closely related to each other and to other model strains such as *Acutodesmus obliquus*,
7 *Auxenochlorella protothecoïdes* or *Chlamydomonas reinhardtii*. These model strains were
8 previously investigated for their ability to grow on various VFAs and shown to poorly
9 assimilate butyrate [6,8]. Therefore, we also included other species, namely *Euglena gracilis*,
10 *Ochromonas danica*, *Polytomella sp.* and *Polytomella magna* in our screening. Both
11 *Polytomella* strains are non-photosynthetic Chlorophytes closely to *C. reinhardtii* which rely
12 on heterotrophic growth [18]. *E. gracilis* and *O. danica* belong to the very distinct
13 Euglenophyte and Chrysophyte divisions, respectively. *E. gracilis* and *O. danica* have been
14 shown earlier to grow on various organic compounds in heterotrophy [35,36]. Microscopy
15 images of all strains are given in the supplemental figure.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 3.2. *Polytomella sp.* stands out as the fastest butyrate consuming strain

37
38

39 The growth of the environmental strains as well as the four selected strains obtained
40 from the SAG collection was compared to that of our reference strain *C. sorokiniana* SAG
41 211-8k in the light. From the growth curves on each separate substrate (not shown), growth
42 and substrate consumption rates on either acetate (Fig. 2) or butyrate (Fig. 3) were
43 determined.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 2: Strain screening on acetate. Growth rates (μ , d^{-1}) (A) were calculated from the exponential phase while acetate removal (%) (B) was calculated at the end point of cultivation. Strains were cultivated in microplates on 0.5 gC.L^{-1} acetate at pH 7 under constant illumination and 100 rpm agitation for a maximum of 3 days, except for *Polytomella* strains. All data are given as mean and standard deviation of three biological replicates.

Each isolated strain except for *Chlamydomonas asymetrica* A2WS could grow on acetate and reach stationary phase in less than 3 days. The fact that this strain could not grow in liquid media could indicate that it was likely isolated due to its robustness and tolerance to adverse conditions (presence of bacteria and butyrate). Its growth on agar plates may have been favoured by the presence of bacteria, which could consume the VFA and increase local CO_2 concentration but was not due to direct VFA consumption. The other strains exhibited growth rates between 0.84 ± 0.04 to $3.60 \pm 0.07 \text{ d}^{-1}$, averaging at 2.0 d^{-1} (Figure 2A). The strain *C. sorokiniana* SAG 211-8k exhibited a growth rate of $3.60 \pm 0.07 \text{ d}^{-1}$ on acetate, a value in accordance with previous work with this strain [22]. The isolated strains *C. sorokiniana* A1WS and BP show the same growth rate as the reference strain. The growth rate on acetate of the *D. armatus* strains was significantly lower (factor 3-4) than that of the *C. sorokiniana* strains. Similarly, the growth rate of *E. gracilis* and *O. danica* was respectively

5.0- and 6.2-fold lower than that of *C. sorokiniana*. In contrast, the heterotrophic chlorophyte *Polytomella* sp. exhibited a growth rate of $4.01 \pm 0.05 \text{ d}^{-1}$ on acetate, significantly higher than any other strain. However, *Polytomella magna* growth rate was around 2-fold lower than that of *Polytomella* sp.

Acetate was always entirely consumed in 3 days (Fig. 2B). Qualitative consumption of acetate by *E. gracilis*, *O. danica* and *Polytomella* species has been reported by several other authors but few quantitative data are available. Some authors reported for *E. gracilis* a growth rate of 1.08 d^{-1} on glucose and 0.6 d^{-1} on ethanol, comparable to the 0.6 d^{-1} here on acetate [37]. Similarly, a growth rate between 0.45 and 0.75 d^{-1} was determined for *O. danica* on glucose and acetate, in line with our results [35].




Figure 3: Strain screening on butyrate. Growth rates (μ , d^{-1}) (A) were calculated from exponential phase while butyrate removal (%) (B) was calculated at the end point of cultivation. Strains were cultivated in microplates on 0.5 gC.L^{-1} butyrate at pH 7 under constant illumination and 100 rpm agitation for a maximum of 7 days, except for *Polytomella* strains. All data are given as mean and standard deviation of three biological replicates.

On the other hand, use of butyrate as substrate always induced a reduction in growth rate compared to acetate (Fig. 3A). As observed for acetate, no growth could be detected on butyrate for *C. asymetria* A2WS. Growth rate of *C. sorokiniana*, either the isolates or the lab strain, was reduced by 10-fold, in line with previous results [22]. Growth rates of *E. gracilis*, *O. danica* and *Polytomella* sp. were however only reduced by 32%, 40% and 37% respectively. It should still be highlighted that *Polytomella* sp. exhibited a much higher growth rate than any other strain on butyrate, topping at 2.5 d⁻¹. Furthermore, these three strains always consumed butyrate entirely within 7 days, with complete butyrate consumption in less than 2.5 days by *Polytomella* sp. In comparison, all other photosynthetic strains consumed around 50% of butyrate in 7 days (Fig. 3B). Interestingly, *Polytomella magna* did not grow at all in presence of butyrate, indicating that butyrate consumption within the *Polytomella* genus is species specific. Of note, *P. magna* remains evolutionary closer to the *Chlamydomonas*-like photosynthetic ancestor of these colorless species [17], and has retained some metabolic capacities such as β -carotene synthesis in contrast to other *Polytomella* species [38]. Where its closest green relative *C. reinhardtii* can use only acetate efficiently [39], *Polytomella* sp. has evolved towards a heterotrophic lifestyle with the capacity to use a variety of organic substrates, notably C2-C5 VFAs, alcohols, pyruvate and citric acid cycle intermediates [40,41]. Although very little is known on the pathways involved in the assimilation of butyrate or other organic substrates in *Polytomella* sp., major differences in metabolic enzymes have been described especially on the level of the plastid [42,43] and mitochondria [44–47] that likely reflect adaptation to heterotrophy and which may guide further studies on this topic.

3.3. *Polytomella* sp. stands out as the most tolerant strain to undissociated acid

The tolerance to undissociated acid concentration (RH) of the strains able to grow in the initial screening was then assessed since this was shown as a determining factor in growth inhibition on VFAs [8]. For this, the same protocol as described in previous study was followed [8]. Briefly, all strains were cultivated on increasing concentrations of acetic (AcH) or butyric (BuH) acid at a fixed pH and growth rates were monitored. This protocol enabled to determine for each strain an RH concentration threshold above which growth was completely inhibited (Fig. 4 and table 2).




Figure 4: Determination of the undissociated acid tolerance of each strain. The figures represent growth rates (μ , d^{-1}) of (A) *D. armatus* RW, (B) *C. sorokiniana* A1WS, (C) *C.*

1 *sorokiniana* BP, (D) *C. sorokiniana* SAG, (D) *E. gracilis*, (E) *O. danica* and (F) *Polytomella*
2 sp. cultivated on increasing undissociated acid concentration (RH, mg.L⁻¹) of acetate (blue) or
3 butyrate (red). Growth rates were calculated from exponential phase and RH concentration
4 using the Henderson-Hasselbalch equation. Strains were cultivated in microplates under
5 constant illumination for a maximum of 7 days. All data are given as mean and standard
6 deviation of three biological replicates.
7
8
9
10
11
12
13
14
15
16
17

18 The isolated strains, *C. sorokiniana* A1WS and BP presented a very similar behaviour
19 to the lab strain *C. sorokiniana* SAG 211-8k (Fig. 4A, B, D). These three strains were
20 completely inhibited above 72 mg.L⁻¹ AcH (12 g.L⁻¹ total acetate at pH 7) and 26 mg.L⁻¹ BuH
21 (4 g.L⁻¹ butyrate at pH 7) (Table 2). The isolate *D. armatus* RW showed a lower tolerance to
22 the acids, being completely inhibited below 26 mg.L⁻¹ AcH (4 g.L⁻¹ acetate at pH 7) (Fig. 4C).
23 Thus, in line with the fact that isolated strains did not exhibit growth rates much higher than
24 those of the lab strain, the isolates did not exhibit any particular resistance to AcH or BuH
25 compared to the lab strain *C. sorokiniana*. The isolation strategy did thus not allow isolating a
26 strain more resistant to the acids. The medium pH during the isolation process was set
27 between 7 and 8, in order to effectively minimize RH concentrations. This choice was based
28 on the hypothesis that the tolerance of a specific strain to RH concentration is not necessarily
29 correlated to fast substrate consumption. Thus, raising the pH ensured growth of any
30 microalgae strain in the starting inoculum, even at a low cell density. Therefore, although this
31 strategy does select for fast growth as competitive parameter, it did not result in the selection
32 of any strain showing higher tolerance to RH.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54

55 **Table 2:** Maximum undissociated acid (RH) concentration for which microalgal growth is not
56 entirely inhibited as determined in Fig. 4.
57
58
59
60
61
62
63
64
65

Strain	RH threshold (mg.L ⁻¹)	
	Ac	Bu
<i>D. armatus</i> RW	0	26
<i>C. sorokiniana</i> A1WS	72	39
<i>C. sorokiniana</i> BP	72	26
<i>C. sorokiniana</i> SAG	72	26
<i>E. gracilis</i>	72	79
<i>O. danica</i>	96	79
<i>Polytomella</i> sp.	750	352

On the other hand, inhibition threshold of *E. gracilis*, *O. danica* and *Polytomella* sp. exhibited distinct tolerance characteristics compared to *C. sorokiniana* (Fig. 4D-G). *E. gracilis* presented a similar tolerance to AcH as *C. sorokiniana* (72 mg.L⁻¹ AcH) while *O. danica* presented a slightly higher tolerance, up to 96 mg.L⁻¹ AcH (Fig. 4D-F). However, both strains tolerated a concentration of BuH 3-fold higher than that of *C. sorokiniana* (79 mg.L⁻¹ BuH) (Fig. 4D-F) (Table 2). An acetate concentration above 1.0 g.L⁻¹ has been reported to trigger the cell death of *O. danica* [35]. Similarly, when formulating a maintenance medium for *O. danica*, acetate was found inhibitory to its growth [48]. The knowledge of the inhibition threshold allows interpreting these differences. In their studies, pH was set to 5.0. At this pH, the AcH is 387 mg.L⁻¹ for 1 g.L⁻¹ acetate, much higher than the determined threshold of 79 mg.L⁻¹ of the present study. In the same way, optimal growth of *E. gracilis* has been reported to occur at acidic pH, between 3 and 5 [37] but acetate addition at this pH was found to completely inhibit cell growth [49]. Considering the inhibition threshold of *E. gracilis* around 75 mg.L⁻¹, the minimum pH tolerated by the cells for 1.0 g.L⁻¹ total acetate would be 6. Therefore, optimum pH for growth does not necessarily coincide with the

1 minimum pH necessary to avoid intolerance to RHs. The knowledge of this threshold value is
2 thus essential to determine both the adequate growth pH and total acid concentration to avoid
3 inhibition. Lastly, *Polytomella* sp. presented exceptional resistance to both acids (Fig. 4G,
4 Table 2). The strain indeed conserved a stable growth rate around 3.8 d^{-1} up until 750 mg.L^{-1}
5 AcH, which corresponds to 38 g.L^{-1} of total acetate at pH 6.5. Similarly, it tolerates BuH
6 concentrations of 350 mg.L^{-1} (18 g.L^{-1} total butyrate), which is three times higher than those
7 tolerated by *E. gracilis* or *O. danica* and about ten times higher than those tolerated by *C.*
8 *sorokiniana*. The tolerance of *Polytomella* sp. to RH concentrations up to 10 times higher than
9 the other strains is a remarkable property of the strain.
10
11
12
13
14
15
16
17
18
19
20
21
22
23

24 To our knowledge, the organisms most tolerant to RH are fermentative bacteria or
25 yeast, which can tolerate AcH concentration as high as 9 g.L^{-1} . These organisms have evolved
26 various tolerance mechanisms to withstand the deleterious effects of the acids, such as a
27 higher expression of ATP- H^{+} pumps [50,51] or by letting their internal pH decrease [52]. The
28 yeast *S. cerevisiae* can also adapt membrane morphology to reduce acid diffusion during
29 extended stress period [53]. *Polytomella* sp. has been firstly isolated from moderately acidic
30 organic rich environments with high fermentative bacterial activity [54]. It is thus likely that
31 the strain developed similar tolerance mechanisms to the fermentative metabolites such as
32 acetate and butyrate. The higher adaptation to VFAs (and probably other organic acids) may
33 be related to its acidophilic nature. At lower pH the organic acids are more present in the RH
34 form and liable to diffuse into the cell [52,55]. This will exert a higher selective pressure to
35 acquire tolerance mechanisms such as found in yeast, creating favourable conditions to
36 subsequently develop heterotrophic capacities. Their newfound heterotrophic capacities, or
37 pre-existing ones as the case may be, will serve them as competitive advantage since they can
38 occupy a larger (acidic) pH range before organic acids become toxic to them. Among other
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

processes, acidophilic species can grow at low pH by being able to maintain their internal pH close to neutrality due to a higher expression of ATP-H⁺ pumps, which in turn has been shown to be related to a higher tolerance to RH [50,56]. However, our results indicate that the property of a strain to tolerate acidic pH is not necessarily linked to its ability to withstand RH. This was also shown for another algal species, *Dunaliella acidophila*, which can grow at pH levels close to 1. Its photosynthetic activity was still reduced by 50% by AcH concentrations as low as 0.033 mg.L⁻¹ [56]. However, *D. acidophila* is purely autotrophic and cannot assimilate acetate. It is possible that lower tolerance to RH is linked to the photosynthetic activity of a strain, which could partially explain the difference between *E. gracilis*, *O. danica* and *Polytomella* sp. Since no photosynthetic species are known that are as tolerant to RH as *Polytomella* sp., and this alga is the only one that is non-photosynthetic, it should be further investigated whether photosynthetic activity is less compatible with high RH tolerance. If that happens to be the case, it may be proposed that the *Chlamydomonas*-like photosynthetic ancestor of *Polytomella* could only acquire more heterotrophic capacities after losing photosynthesis, instead of before.

3.4. Growth of candidate strains on synthetic DF effluent

Given the results in terms of growth rates and tolerance to butyrate, *E. gracilis*, *O. danica* and *Polytomella* sp. were chosen for further studies to evaluate their biomass production potentialities on synthetic DF effluent and to compare them with *C. sorokiniana* chosen as the reference alga. These four strains were cultivated in 500 mL Erlenmeyer flasks on a synthetic mixture of VFAs composed of acetate and butyrate, at a carbon ratio of 1:2 representing an average DF effluent in terms of VFAs. For each strain, biomass growth, VFA consumption and lipids and sugars productions were determined.




Figure 5: Growth of candidate strains on synthetic DFE. The figures represent biomass dry weight (g_x.L⁻¹, grey circles) and substrate concentration (g_{CS}.L⁻¹, acetate in blue circles and butyrate in red circles) evolution as a function of time (days) in (A) *C. sorokiniana* SAG, (B) *E. gracilis*, (C) *O. danica* and (D) *Polytomella* sp. cultures. Strains were cultivated in flasks in synthetic medium (0.3 g_{CS}.L⁻¹ acetate (Ac) and 0.7 g_{CS}.L⁻¹ butyrate (Bu); pH 7) under constant illumination and shaking. *Polytomella* species were cultivated without light control and shaking. All data are given as mean and standard deviation of three biological replicates.

Each strain could grow on the synthetic DF effluent, and growth curves and substrate consumption of each strain are provided in Figure 5. For each strain, two growth phases could be distinguished, attributable to the substrate assimilated. Acetate was always consumed first, followed by butyrate. This phenomenon, known as diauxic growth or carbon catabolite

repression, was already been reported for *C. sorokiniana* and *Auxenochlorella protothecoïdes* growing on acetate and butyrate mixtures [9,22]. The results further highlight this effect and confirm that it occurs for other microalgal strains. However, it should be noted that *O. danica* did consume a fraction of butyrate ($12.5 \pm 1.6\%$) before complete acetate exhaustion. As a comparison, *C. sorokiniana*, *E. gracilis* et *Polytomella* sp. respectively only consumed 2.0 ± 4.6 , 5.7 ± 4.5 and $1.9 \pm 1.6\%$ initial butyrate, without significant difference, before complete acetate exhaustion. The half inhibitory constant of *C. sorokiniana* associated to diauxic growth on the acetate and butyrate mixture has been determined in a previous study [9]. They found a value of $2.10^{-10} \text{ gC.L}^{-1}$, meaning that the consumption of butyrate for this strain is inhibited for extremely low acetate concentration. The fact that *O. danica* started to consume butyrate slightly before complete acetate exhaustion indicates that its half inhibitory constant is higher than that of *C. sorokiniana*. This suggests differences at the metabolic level in the responses to the presence of both substrates between the various strains.

Considering this diauxic effect, each growth phase was considered individually to calculate growth rates and yields (Table 3). During the acetate phase, *C. sorokiniana* presented the highest growth rate ($3.34 \pm 0.17 \text{ d}^{-1}$) followed by *Polytomella* sp. ($2.21 \pm 0.04 \text{ d}^{-1}$). These two strains consumed acetate within only two days. On the other hand, acetate was exhausted in almost four days by *O. danica*, making it the slowest acetate consuming strain. After acetate exhaustion, butyrate started to be consumed. Growth rate on this substrate was always slower than on acetate. *Polytomella* sp. was the fastest butyrate consuming strain, exhibiting a growth rate of $1.20 \pm 0.04 \text{ d}^{-1}$ during this phase. It should be noted here that the growth rates obtained in this experiment for *Polytomella* sp., in presence of both substrates, are around 2-fold lower than those obtained with the separate substrates (section 3.2). One explanation is that butyrate would inhibit growth on acetate. However, butyrate concentration was here only 1.2 g.L^{-1} , corresponding to $7.5 \text{ mg.L}^{-1} \text{ BuH}$ at pH 7.0. BuH concentration was

therefore around 50-times lower than the inhibitory threshold determined in section 3.3. Furthermore, growth rates obtained for other strains were similar between the two experiments. For *Polytomella* sp., the most probable explanation lies in the different cultivation vessel. The screening was indeed carried out in microplates while the biomass production potential was carried out in Erlenmeyer flasks. Although it was confirmed that growth rates between microplates and shake flasks were similar for *C. sorokiniana* [22], no such validation was done for *Polytomella* sp. *Polytomella* cells are generally cultured without any shaking since the cells are relatively fragile (absence of cell wall) and shaking can inhibit growth [57]. During growth, *Polytomella* cells swim along a water column between the bottom of the vessel and the medium surface in order to oxygenate themselves [19]. These movements will depend on the depth of the vessel. Since medium volume is much lower in microplates, it is possible that cells could oxygenate faster in this vessel, thus exhibiting a higher growth rate. All of this considered, *Polytomella* sp. remained the fastest strain on the synthetic effluent since substrate was entirely consumed after two days against a minimum of six days for other strains. Additional gentle shaking or aeration system might further speed up organic consumption and should be considered for further studies with this strain.

Table 3: Growth characteristics of candidate strains on synthetic DFE (0.3 g_C.L⁻¹ acetate (Ac) and 0.7 g_C.L⁻¹ butyrate (Bu); pH 7). Biomass yields ($Y_{X/S}$, g_X.g_CS⁻¹) and growth rates (μ , d⁻¹) were calculated for each substrate growth phase. Biomass productivities (Q_X , g_X.L⁻¹.d⁻¹) were calculated considering the whole cultivation duration. Values are given as mean and standard deviation for three biological replicates.

Strain	Y _{x/s} (g _x ·g _{CS} ⁻¹)		μ (d ⁻¹)		Q _x (g _x ·L ⁻¹ ·d ⁻¹)
	Ac	Bu	Ac	Bu	Global
<i>C. sorokiniana</i> SAG	1.58 ± 0.16	3.03 ± 0.6	3.34 ± 0.17	0.30 ± 0.01	0.64 ± 0.10
<i>E. gracilis</i>	1.71 ± 0.06	1.66 ± 0.08	0.85 ± 0.01	0.38 ± 0.09	0.30 ± 0.15
<i>O. danica</i>	3.40 ± 0.33	3.40 ± 0.54	0.81 ± 0.04	0.28 ± 0.03	0.18 ± 0.01
<i>Polytomella</i> sp.	1.46 ± 0.12	1.24 ± 0.03	2.20 ± 0.18	1.28 ± 0.01	0.62 ± 0.04

At the end of the acetate phase, *O. danica* exhibited the highest biomass yield (3.40 ± 0.33 g_x·g_{CS}⁻¹). By using the intracellular carbon content of *O. danica* (2.02 pmol per cell growing in mixotrophy) [15] and the dry biomass quantity as a function of cell number (between 3.7 and $5.5 \cdot 10^{-11}$ g·cell⁻¹) [35], it was possible to estimate the C content of *O. danica* per biomass dry weight at an average of 57.9%. The carbon biomass yield on substrate of *O. danica* would thus be as high as 1.96 g_{CX}·g_{CS}⁻¹. This would mean that a large amount of CO₂ was photosynthetically fixed by the cells during growth.

For the three other strains, biomass yields were between 1.46 and 1.71 g_x·g_{CS}⁻¹ without significant difference (Table 3). By estimating the biomass carbon content to be of 53%, close to that estimated for *C. reinhardtii* [58], a carbon yield on substrate was calculated to be of 0.77 g_{CX}·g_{CS}⁻¹ for *Polytomella* sp. Unlike the other strains, *Polytomella* sp. is a pure heterotroph strain unable to perform photosynthesis. Therefore, the amount of biomass produced can only originate from organic carbon assimilation. No quantitative data are available for the growth of *Polytomella* sp. and this value can thus be only compared to other strains grown in heterotrophy. This value is similar to that obtained for the close relative *C. reinhardtii* in heterotrophy on acetate by [58]. By using the same carbon content estimation for *C. sorokiniana*, the carbon biomass yield on substrate for this strain was also calculated to

1 be 0.76 $\text{g}_{\text{CX}}.\text{g}_{\text{CS}}^{-1}$ (Table 3). Since the strain is photosynthetic and can thus photosynthetically
2 fix CO_2 , a higher biomass yield was expected. This could mean that either *C. sorokiniana*
3
4 grew only in heterotrophy on acetate, without fixing CO_2 or that the heterotrophic yield of
5
6 *Polytomella* sp. is higher than that of *C. sorokiniana* on acetate. This latter hypothesis would
7
8 be consistent with previous results [9,59], who reported for *C. sorokiniana* a biomass yield of
9
10 0.4 $\text{g}_{\text{CX}}.\text{g}_{\text{CS}}^{-1}$ and 0.5 $\text{g}_{\text{CX}}.\text{g}_{\text{CS}}^{-1}$ respectively on acetate in heterotrophy. Here, the use of
11
12 mixotrophic condition increased the biomass yield of *C. sorokiniana* to 0.76 $\text{g}_{\text{CX}}.\text{g}_{\text{CS}}^{-1}$ by
13
14 allowing the strain to photosynthetically fix atmospheric CO_2 .
15
16
17
18
19

20 In the case of *E. gracilis*, a biomass yield of 1.71 $\text{g}_{\text{X}}.\text{g}_{\text{CS}}^{-1}$ was obtained on acetate. No
21
22 yield on acetate was reported so far due to the inhibitory effects at low pH discussed
23
24 previously. However, some data are available on ethanol and lactic acid. Fujita et al., [36]
25
26 reported for *E. gracilis* in mixotrophy a value of 0.77 $\text{g}_{\text{X}}.\text{g}_{\text{CS}}^{-1}$ on ethanol but a value of 1.33
27
28 $\text{g}_{\text{X}}.\text{g}_{\text{CS}}^{-1}$ on lactic acid, which is in the same order of magnitude of what was obtained here
29
30 [36]. This would mean that VFAs, such as acetate, butyrate or lactate are better substrates for
31
32 *E. gracilis*, as long as inhibitory concentrations are avoided.
33
34
35
36
37

38 At the end of the butyrate phase, *Polytomella* sp. presented a yield significantly lower
39
40 than obtained on acetate ($1.24 \pm 0.03 \text{ g}_{\text{X}}.\text{g}_{\text{CS}}^{-1}$ against $1.46 \pm 0.12 \text{ g}_{\text{X}}.\text{g}_{\text{CS}}^{-1}$ on acetate). This
41
42 indicates that although the strain can quickly consume butyrate, its assimilation requires more
43
44 energy than acetate since carbon is diverted from biomass production. On the other hand, the
45
46 biomass yield of *E. gracilis* and *O. danica* did not change significantly compared to the
47
48 acetate phase (respectively $1.66 \pm 0.08 \text{ g}_{\text{X}}.\text{g}_{\text{CS}}^{-1}$ and $3.39 \pm 0.54 \text{ g}_{\text{X}}.\text{g}_{\text{CS}}^{-1}$) while the yield of *C.*
49
50 *sorokiniana* drastically increased up to $3.0 \pm 0.6 \text{ g}_{\text{X}}.\text{g}_{\text{CS}}^{-1}$. As these strains can perform
51
52 photosynthesis, it is probable that light could be used as an extra energy source to uptake and
53
54 metabolize butyrate. The carbon biomass yield of *C. sorokiniana* on butyrate, 1.6 $\text{g}_{\text{CX}}.\text{g}_{\text{CS}}^{-1}$,
55
56 indicates that a much higher fraction of CO_2 is photosynthetically fixed by the strain during
57
58
59
60
61
62
63
64
65

1 butyrate consumption compared to growth on acetate. This is due to the low growth rate on
2 butyrate, only slightly higher than the autotrophic growth rate at atmospheric CO₂ levels [22].
3
4
5
6
7

8 9 *3.5. Production of lipids and carbohydrates by the candidate strains*

10
11 As lipids and carbohydrates production is the main incentive for microalgal cultivation
12 on waste streams, the concentration of these two products were measured during growth
13 (Figure 6). The corresponding productivities and yields are reported in table 4. Microalgal
14 lipids and sugars were respectively quantified by the phosphovanilin and the anthrone
15 method, successfully used previously for different microalgal strains, including chlorophytes
16 [26,60,61] or traustochytrids [62], with little deviation compared to standard gravimetric
17 protocols. Some factors may interfere with the accuracy of these colorimetric approaches,
18 such as the type and concentration of pigments, the type of lipid or carbohydrate analyzed or
19 the standard used to establish the calibration curves [61]. Although results may be validated
20 using gravimetric methods, these approaches are generally considered appropriate as a first
21 estimate of the relative amount of each storage compound between strains.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 6: Carbohydrates and lipids production by candidate strains on synthetic DFE. The figures represent biomass dry weight ($\text{g} \cdot \text{L}^{-1}$, grey circles) and product content in each strain ($\% \text{g} \cdot \text{g}^{-1}$, carbohydrates in white squares and lipids in black squares) evolution as a function of time (days) in (A) *C. sorokiniana* SAG, (B) *E. gracilis*, (C) *O. danica* and (D) *Polytomella* sp. cultures. Strains were cultivated in flasks in synthetic medium ($0.3 \text{ g}_{\text{CS}} \cdot \text{L}^{-1}$ acetate (Ac) and $0.7 \text{ g}_{\text{CS}} \cdot \text{L}^{-1}$ butyrate (Bu); pH 7) under constant illumination and shaking. All data are given as mean and standard deviation of three biological replicates. Lack of data points in early phase indicates a product concentration below quantification limit due to lack of sufficient biomass.

The highest carbohydrates content was found for *Polytomella* sp. and *E. gracilis* ($52.6 \pm 4.5\%$ and $53.5 \pm 11.2\%$ respectively) at the end of the acetate phase. This result was

expected since both strains are known for accumulating starch for the former [19] and paramylon for the latter [63]. However, both strains differed in their accumulation dynamics. The carbohydrates content of *Polytomella* sp. remained stable as long as organic substrate was present in the medium. Its content did decrease progressively down to $35.7 \pm 3.9\%$ once organic carbon had been completely exhausted. This is because *Polytomella* sp. can only rely on organic carbon for growth: in its absence, the cells consume stored carbon (*i.e.* starch) to form immobile cysts and assure long term survival. Despite its lower biomass yield compared to other strains, the combination of the highest growth rate on both acetate and butyrate as well as the highest carbohydrate yield made *Polytomella* sp. the most productive strain for carbohydrates (Table 4).

Table 4: Lipid and carbohydrates production of candidate strains on synthetic DFE (0.3 g_C.L⁻¹ acetate (Ac) and 0.7 g_C.L⁻¹ butyrate (Bu); pH 7). Product (either carbohydrates or lipids) yields ($Y_{P/S}$, mg.g_{CS}⁻¹) and product productivities (Q_P , mg.L⁻¹.d⁻¹) were calculated considering the whole cultivation duration. Values are given as mean and standard deviation of three biological replicates.

Strain	$Y_{P/S}$ (mg.g _{CS} ⁻¹)		Q_P (mg.L ⁻¹ .d ⁻¹)	
	Carbohydrates	Lipids	Carbohydrates	Lipids
<i>C. sorokiniana</i> SAG	395.7 ± 49.7	420.7 ± 21.2	50.6 ± 5.6	45.9 ± 0.3
<i>E. gracilis</i>	575.5 ± 145.1	181.2 ± 3.9	78.9 ± 2.7	33.8 ± 1.5
<i>O. danica</i>	226.8 ± 38.0	315.1 ± 5.1	28.4 ± 5.7	43.8 ± 2.12
<i>Polytomella</i> sp.	648.7 ± 66.0	109.8 ± 1.7	309.6 ± 24.5	53.2 ± 1.2

On the other hand, the carbohydrates content of *E. gracilis* significantly declined from 47.3 ± 2.4% after acetate exhaustion down to 26.6 ± 1.7% at the end of the butyrate phase. A paramylon productivity of 80.1 mg.L⁻¹.d⁻¹ has been reported on 2 g.L⁻¹ acetate in line with the global productivity that was obtained here [64]. However, they found that paramylon content of *E. gracilis* could reach up to 70.8% on acetate (without any limitation). The same authors also showed that paramylon content was greatly affected by the carbon sources but did not investigate the effect of butyrate. These results thus complement their findings and indicate that butyrate does not enhance paramylon content as much as acetate. A possible explanation could be that full butyrate metabolization demands more energy to the cells compared to acetate. Indeed, butyrate is more reduced than acetate and thus its conversion to acetyl-CoA requires more metabolic steps contrary to acetate which can directly fuel the tricarboxylic acid or the glyoxylate cycle [65]. In *E. gracilis*, this cycle is directly linked to paramylon synthesis [66]. In the case of butyrate, some more carbon might be diverted toward energy production and not biomass production as has been discussed for *Polytomella* sp.

Lipids and carbohydrates content of both *C. sorokiniana* and *O. danica* remained stable during growth (Figure 6). The carbohydrate and lipid content of *O. danica* remained lower than other strains (respectively 13.5 ± 1.0% and 9.6 ± 1.4%) throughout the whole culture. This lipid content might seem low in regards to previously reported values. For example, intracellular concentration could reach up 79% when cultivating *O. danica* on waste grease [67]. This value was however obtained under nitrogen (N) limitation and in late growth phase. Microalgae tend to not accumulate lipids or carbohydrates when mineral nutrients such as N or phosphorus (P) are not limiting [68]. In the present work, focus was on biomass growth and VFA removal potential of the candidate strains. As such, N and P concentrations were adjusted according to the Redfield ratio 106:16:1, which was found in a previous study to promote biomass growth of *C. sorokiniana* [22]. Carbon was thus the limiting nutrient

1 which probably explains the low carbohydrate and lipid contents of *O. danica*. This alga has
2 also been shown to excrete free fatty acids when growing mixotrophically growth during early
3 and mid-growth phase [69]. It is thus possible that a part of organic carbon was diverted
4 toward fatty acids excretion instead of lipid accumulation. Since the SPV method employed to
5 measure lipids cannot identify free fatty acids, this should be investigated in further studies.
6
7
8
9
10

11
12 Lipid productivities were in the same order of magnitude between all strains despite
13 disparities in lipid content and yield (Table 4). For example, *Polytomella* sp. was the most
14 lipid productive strain thanks to its high growth rate, but it however presented the lowest lipid
15 content of all strains ($8.4 \pm 0.5\%$). Due to its high biomass production, *O. danica* was second
16 to *C. sorokiniana* in terms of lipid yield. Nevertheless, *C. sorokiniana* should remain the most
17 interesting strains in terms of lipid production since it had the highest lipid content (around
18 18%) of all strains, which would be favourable for further extraction. The use of an N-
19 limitation strategy should nonetheless change this result by triggering lipid accumulation by
20 *O. danica*. Still, in the case of dark fermentation effluent treatment, the main objective would
21 remain maximum organic carbon removal, which can only be achieved without N-limitation
22 [22]. Such a strategy could thus only be applied in a post-treatment stage, as has been
23 proposed by [70].
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 4. Conclusions

47
48 A main goal of this study was to identify microalgal strains able to grow on butyrate
49 as fast as on acetate (and ideally simultaneously). This ability was not found to be widespread
50 among the investigated green Chlorophytes. In addition, there was no apparent difference
51 between the natural and the lab strains, indicating that butyrate metabolism did not seem to
52 evolve significantly under different environmental conditions. As such, only three strains,
53
54
55
56
57
58
59
60
61
62
63
64
65

1 namely *E. gracilis*, *O. danica* and *Polytomella* sp. were found to share this characteristic.
2 Among them, *Polytomella* sp. stands out due to its growth properties and tolerance to
3 butyrate. As a close relative of *C. reinhardtii*, it has been used as a model strain to understand
4 the evolutionary events that lead the species from an autotrophic/mixotrophic to a
5 heterotrophic mode of nutrition. The unique physiology of *Polytomella* sp. on acetate and
6 butyrate emphasizes the interest of this strain to identify both fast butyrate assimilation and
7 high tolerance mechanisms actually present in *Polytomella* sp. and absent in photosynthetic
8 strains such as *C. reinhardtii* or *C. sorokiniana*.
9

10 The second objective of this work was to evaluate the biomass production potential of
11 these strains on a synthetic effluent. It was found that the heterotrophic alga *Polytomella* sp.
12 and the photosynthetic species *E. gracilis* accumulated high levels of carbohydrates during
13 growth. This trait is advantageous since it would allow the conversion of VFAs into
14 carbohydrates in continuous culture mode, without limitation strategy needed. It was also
15 found that the cellular composition of both *C. sorokiniana* and *O. danica* remained constant
16 during growth and was more oriented toward proteins production. Overall, this work provides
17 new growth kinetic data of various microalgal strains using VFAs as substrates, which could
18 be further used to design various dark fermentation effluent valorisation strategies.
19

20 **Conflict of interest**

21 The authors declare that the research was conducted in the absence of any commercial or
22 financial relationships that could be construed as a potential conflict of interest.
23

24 **CRedit Author contributions**

JL: conceptualization, investigation, data acquisition, data curation, formal analysis and writing of the original draft. **PJ**: conceptualization, investigation, and writing. **AA** and **AB**: investigation, data curation, validation, review and editing of the original draft. **JPS** and **ET**: supervision, funding acquisition, validation, review and editing of the original draft. **RVL**: conceptualization, funding acquisition, supervision, validation, review and editing of the original draft. All authors approved the final version of the manuscript.

Acknowledgments

JL received a PhD fellowship from European Union from the Occitanie region, France, with complementary funding from FEDER. This study was funded by the National Institute of Agriculture, Alimentation and Environment (INRAE), and partly supported by the FERMALip project, funded by the Carnot institute 3BCAR.

References

- [1] R. Moscoviz, E. Trably, N. Bernet, H. Carrère, The environmental biorefinery: State-of-the-art on the production of hydrogen and value-added biomolecules in mixed-culture fermentation, *Green Chem.* 20 (2018) 3159–3179. <https://doi.org/10.1039/c8gc00572a>.
- [2] A. Patel, O. Sarkar, U. Rova, P. Christakopoulos, L. Matsakas, Valorization of volatile fatty acids derived from low-cost organic waste for lipogenesis in oleaginous microorganisms-A review, *Bioresour. Technol.* 321 (2021) 124457. <https://doi.org/10.1016/j.biortech.2020.124457>.
- [3] V. Turon, E. Trably, E. Fouilland, J.-P. Steyer, Potentialities of dark fermentation effluents as substrates for microalgae growth: A review, *Process Biochem.* 51 (2016) 1843–1854. <https://doi.org/10.1016/J.PROCBIO.2016.03.018>.
- [4] C. Baroukh, V. Turon, O. Bernard, Dynamic metabolic modeling of heterotrophic and mixotrophic microalgal growth on fermentative wastes, *PLoS Comput. Biol.* 13 (2017) 1–18.

<https://doi.org/10.1371/journal.pcbi.1005590>.

- [5] C.-H. Liu, C.-Y. Chang, Q. Liao, X. Zhu, C.-F. Liao, J.-S. Chang, Biohydrogen production by a novel integration of dark fermentation and mixotrophic microalgae cultivation, *Int. J. Hydrogen Energy*. 38 (2013) 15807–15814. <https://doi.org/10.1016/j.ijhydene.2013.05.104>.
- [6] Q. Fei, R. Fu, L. Shang, C.J. Brigham, H.N. Chang, Lipid production by microalgae *Chlorella protothecoides* with volatile fatty acids (VFAs) as carbon sources in heterotrophic cultivation and its economic assessment., *Bioprocess Biosyst. Eng.* 38 (2015) 691–700. <https://doi.org/10.1007/s00449-014-1308-0>.
- [7] V. Turon, E. Trably, E. Fouilland, J.-P. Steyer, Growth of *Chlorella sorokiniana* on a mixture of volatile fatty acids: The effects of light and temperature, *Bioresour. Technol.* 198 (2015) 852–860. <https://doi.org/10.1016/j.biortech.2015.10.001>.
- [8] J. Lacroux, E. Trably, N. Bernet, J.P. Steyer, R. van Lis, Mixotrophic growth of microalgae on volatile fatty acids is determined by their undissociated form, *Algal Res.* 47 (2020) 101870. <https://doi.org/10.1016/j.algal.2020.101870>.
- [9] V. Turon, C. Baroukh, E. Trably, E. Latrille, E. Fouilland, J.-P. Steyer, Use of fermentative metabolites for heterotrophic microalgae growth: Yields and kinetics, *Bioresour. Technol.* 175 (2015) 342–349. <https://doi.org/10.1016/j.biortech.2014.10.114>.
- [10] D. Mu, H. Liu, W. Lin, P. Shukla, J. Luo, Simultaneous biohydrogen production from dark fermentation of duckweed and waste utilization for microalgal lipid production, *Bioresour. Technol.* 302 (2020) 122879. <https://doi.org/10.1016/j.biortech.2020.122879>.
- [11] H.Y. Ren, F. Kong, J. Ma, L. Zhao, G.J. Xie, D. Xing, W.Q. Guo, B.F. Liu, N.Q. Ren, Continuous energy recovery and nutrients removal from molasses wastewater by synergistic system of dark fermentation and algal culture under various fermentation types, *Bioresour. Technol.* 252 (2018) 110–117. <https://doi.org/10.1016/j.biortech.2017.12.092>.
- [12] A.W.D. Larkum, I.L. Ross, O. Kruse, B. Hankamer, Selection, breeding and engineering of microalgae for bioenergy and biofuel production, *Trends Biotechnol.* 30 (2012) 198–205. <https://doi.org/10.1016/j.tibtech.2011.11.003>.

- [13] A. Chalima, A. Hatzidaki, A. Karnaouri, E. Topakas, Integration of a dark fermentation effluent in a microalgal-based biorefinery for the production of high-added value omega-3 fatty acids, *Appl. Energy*. 241 (2019) 130–138. <https://doi.org/10.1016/j.apenergy.2019.03.058>.
- [14] S. Kim, R. Wirasnita, D. Lee, J. Yu, T. Lee, Enhancement of Growth and Paramylon Production of *Euglena gracilis* by Upcycling of Spent Tomato Byproduct as an Alternative Medium, *Appl. Sci.* 11 (2021) 8182. <https://doi.org/10.3390/app11178182>.
- [15] S. Wilken, J.M. Schuurmans, H.C.P. Matthijs, Do mixotrophs grow as photoheterotrophs? Photophysiological acclimation of the chrysophyte *Ochromonas danica* after feeding, *New Phytol.* 204 (2014) 882–889. <https://doi.org/10.1111/nph.12975>.
- [16] V. Turon, E. Trably, A. Fayet, E. Fouilland, J.-P. Steyer, Raw dark fermentation effluent to support heterotrophic microalgae growth: microalgae successfully outcompete bacteria for acetate, *Algal Res.* 12 (2015) 119–125. <https://doi.org/10.1016/j.algal.2015.08.011>.
- [17] F. Round, The evolution of pigmented and unpigmented unicells - a reconsideration of the Protista, *BioSystems*. 12 (1980) 61–69.
- [18] R. van Lis, D. González-Halphen, A. Atteia, Divergence of the mitochondrial electron transport chains from the green alga *Chlamydomonas reinhardtii* and its colorless close relative *Polytomella* sp., *Biochim. Biophys. Acta*. 1708 (2005) 23–34. <https://doi.org/10.1016/j.bbabi.2004.12.010>.
- [19] V.F. de la Cruz, S.M. Gittleson, The genus *Polytomella*: A review of classification, morphology, life cycle, metabolism, and motility, *Arch. Fur Protistenkd.* 124 (1981) 1–28. [https://doi.org/10.1016/S0003-9365\(81\)80001-2](https://doi.org/10.1016/S0003-9365(81)80001-2).
- [20] E.H. Harris, *The Chlamydomonas sourcebook*, 1989.
- [21] S.H. Hutner, Inorganic Nutrition, *Annu. Rev. Microbiol.* 26 (1972) 313–346. <https://doi.org/10.1146/annurev.mi.26.100172.001525>.
- [22] J. Lacroux, J. Seira, E. Trably, N. Bernet, J. Steyer, R. van Lis, Mixotrophic Growth of *Chlorella sorokiniana* on Acetate and Butyrate : Interplay Between Substrate , C : N Ratio and pH, *Front. Microbiol.* 12 (2021) 1–16. <https://doi.org/10.3389/fmicb.2021.703614>.
- [23] S.M. Newman, J.E. Boynton, N.W. Gillham, B.L. Randolph-Anderson, A.M. Johnson, E.H. Harris,

- Transformation of chloroplast ribosomal RNA genes in *Chlamydomonas*: molecular and genetic characterization of integration events., *Genetics*. 126 (1990) 875–888.
<https://doi.org/10.1093/genetics/126.4.875>.
- [24] B. Marin, M. Klingberg, M. Melkonian, Phylogenetic Relationships among the Cryptophyta: Analyses of Nuclear-Encoded SSU rRNA Sequences Support the Monophyly of Extant Plastid-Containing Lineages, *Protist*. 149 (1998) 265–276. [https://doi.org/10.1016/S1434-4610\(98\)70033-1](https://doi.org/10.1016/S1434-4610(98)70033-1).
- [25] S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms, *Mol. Biol. Evol.* 35 (2018) 1547–1549.
<https://doi.org/10.1093/molbev/msy096>.
- [26] S.K. Mishra, W.I. Suh, W. Farooq, M. Moon, A. Shrivastav, M.S. Park, J.W. Yang, Rapid quantification of microalgal lipids in aqueous medium by a simple colorimetric method, *Bioresour. Technol.* 155 (2014) 330–333. <https://doi.org/10.1016/j.biortech.2013.12.077>.
- [27] E.W. Yemm, A.J. Willis, The estimation of carbohydrates in plant extracts by anthrone., *Biochem. J.* 57 (1954) 508–14. <http://www.ncbi.nlm.nih.gov/pubmed/13181867>.
- [28] K.N. Duddleston, M.A. Kinney, R.P. Kiene, M.E. Hines, Anaerobic microbial biogeochemistry in a northern bog: Acetate as a dominant metabolic end product, *Global Biogeochem. Cycles*. 16 (2002) 11-1-11–9. <https://doi.org/10.1029/2001gb001402>.
- [29] A.W. Coleman, ITS2 is a double-edged tool for eukaryote evolutionary comparisons, *Trends Genet.* 19 (2003) 370–375. [https://doi.org/10.1016/S0168-9525\(03\)00118-5](https://doi.org/10.1016/S0168-9525(03)00118-5).
- [30] G. Salazar-Torres, V.L. de Moraes Huszar, Microalgae community of the Huaytire wetland, an Andean high-altitude wetland in Peru, *Acta Limnol. Bras.* 24 (2012) 285–292. <https://doi.org/10.1590/S2179-975X2012005000046>.
- [31] C. Chiellini, L. Guglielminetti, S. Sarrocco, A. Ciurli, Isolation of Four Microalgal Strains From the Lake Massaciuccoli: Screening of Common Pollutants Tolerance Pattern and Perspectives for Their Use in Biotechnological Applications, *Front. Plant Sci.* 11 (2020) 1–18.
<https://doi.org/10.3389/fpls.2020.607651>.
- [32] R.A. Andersen, M. Kawachi, Traditional Microalgae Isolation Techniques, in: *Algal Cult. Tech.*,

Elsevier, 2005: pp. 83–100. <https://doi.org/10.1016/B978-012088426-1/50007-X>.

- [33] A. Bhatnagar, S. Chinnasamy, M. Singh, K.C. Das, Renewable biomass production by mixotrophic algae in the presence of various carbon sources and wastewaters, *Appl. Energy*. 88 (2011) 3425–3431. <https://doi.org/10.1016/j.apenergy.2010.12.064>.
- [34] A. Pandey, S. Srivastava, S. Kumar, Isolation, screening and comprehensive characterization of candidate microalgae for biofuel feedstock production and dairy effluent treatment: A sustainable approach, *Bioresour. Technol.* 293 (2019) 121998. <https://doi.org/10.1016/j.biortech.2019.121998>.
- [35] Z. Lin, C. Li, L.K. Ju, Glycerol and Acetate Additions to Maximize Lipid Content in High-Density Cultures of Phagotrophic Alga *Ochromonas danica*, *JAOCs, J. Am. Oil Chem. Soc.* 96 (2019) 231–238. <https://doi.org/10.1002/aocs.12183>.
- [36] T. Fujita, H. Aoyagi, J.C. Ogbonna, H. Tanaka, Effect of mixed organic substrate on α -tocopherol production by *Euglena gracilis* in photoheterotrophic culture, *Appl. Microbiol. Biotechnol.* 79 (2008) 371–378. <https://doi.org/10.1007/s00253-008-1443-0>.
- [37] J.C. Ogbonna, S. Tomiyama, H. Tanaka, Heterotrophic cultivation of *Euglena gracilis* Z for efficient production of α -tocopherol, *J. Appl. Phycol.* 10 (1998) 67–74. <https://doi.org/10.1023/A:1008011201437>.
- [38] S.R. Asmail, D.R. Smith, Retention, erosion, and loss of the carotenoid biosynthetic pathway in the nonphotosynthetic green algal genus *Polytomella*, *New Phytol.* 209 (2016) 899–903. <https://doi.org/10.1111/nph.13682>.
- [39] Y. Hayashi, N. Sato, A. Shinozaki, M. Watanabe, Increase in peroxisome number and the gene expression of putative glyoxysomal enzymes in *Chlamydomonas* cells supplemented with acetate, *J. Plant Res.* 128 (2015) 177–185. <https://doi.org/10.1007/s10265-014-0681-8>.
- [40] D.L. Wise, Carbon Nutrition and Metabolism of *Polytomella caeca*, *J. Protozool.* 6 (1959) 19–23. <https://doi.org/10.1111/j.1550-7408.1959.tb03921.x>.
- [41] D.L. Wise, The Effect of pH on the Availability of Acid Substrates for *Polytomella caeca* with Unexpected Activity by Succinate, *J. Protozool.* 11 (1964) 246–249. <https://doi.org/10.1111/j.1550-7408.1964.tb01751.x>.

- [42] E.O. Fuentes-Ramírez, M. Vázquez-Acevedo, A. Cabrera-Orefice, S. Guerrero-Castillo, D. González-Halphen, The plastid proteome of the nonphotosynthetic chlorophycean alga *Polytomella parva*, *Microbiol. Res.* 243 (2021). <https://doi.org/10.1016/j.micres.2020.126649>.
- [43] D.R. Smith, R.W. Lee, A plastid without a genome: evidence from the nonphotosynthetic green algal genus *Polytomella*., *Plant Physiol.* 164 (2014) 1812–1819. <https://doi.org/10.1104/pp.113.233718>.
- [44] A. Reyes-Prieto, M. El-Hafidi, R. Moreno-Sánchez, D. González-Halphen, Characterization of oxidative phosphorylation in the colorless chlorophyte *Polytomella* sp., *Biochim. Biophys. Acta - Bioenerg.* 1554 (2002) 170–179. [https://doi.org/10.1016/S0005-2728\(02\)00241-4](https://doi.org/10.1016/S0005-2728(02)00241-4).
- [45] R. van Lis, D. González-Halphen, A. Atteia, Divergence of the mitochondrial electron transport chains from the green alga *Chlamydomonas reinhardtii* and its colorless close relative *Polytomella* sp., *Biochim. Biophys. Acta - Bioenerg.* 1708 (2005) 23–34. <https://doi.org/10.1016/j.bbabbio.2004.12.010>.
- [46] R. van Lis, G. Mendoza-Hernández, G. Groth, A. Atteia, New insights into the unique structure of the F0F1-ATP synthase from the chlamydomonad algae *Polytomella* sp. and *Chlamydomonas reinhardtii*, *Plant Physiol.* 144 (2007) 1190–1199. <https://doi.org/10.1104/pp.106.094060>.
- [47] R. van Lis, Y. Couté, S. Brugière, N.J. Tourasse, B. Laurent, W. Nitschke, O. Vallon, A. Atteia, Phylogenetic and functional diversity of aldehyde-alcohol dehydrogenases in microalgae, *Plant Mol. Biol.* 105 (2021) 497–511. <https://doi.org/10.1007/s11103-020-01105-9>.
- [48] B.L.L. Foster, T.H. Chrzanowski, A simple defined medium for growth and maintenance of the mixotrophic protist *Ochromonas danica*, *J. Microbiol. Methods.* 88 (2012) 322–324. <https://doi.org/10.1016/j.mimet.2011.12.016>.
- [49] R.C. Bates, R.E. Hurlbert, The Effect of Acetate on *Euglena gracilis* var. *bacillaris* as a Function of Environmental Conditions, *J. Protozool.* 17 (1970) 134–138.
- [50] Z. Huang, X. Yu, H. Miao, H. Ren, M. Zhao, W. Ruan, Enzymatic dynamics of microbial acid tolerance response (ATR) during the enhanced biohydrogen production process via anaerobic digestion, *Int. J. Hydrogen Energy.* 37 (2012) 10655–10662. <https://doi.org/10.1016/j.ijhydene.2012.04.116>.
- [51] T. Miwa, H. Esaki, J. Umemori, T. Hino, Activity of H²-ATPase in Ruminant Bacteria with Special Reference to Acid Tolerance, 63 (1997) 2155–2158.

- [52] J.B. Russell, Another explanation for the toxicity of fermentation acids at low pH: anion accumulation versus uncoupling, *J. Appl. Bacteriol.* 73 (1992) 363–370. <https://doi.org/10.1111/j.1365-2672.1992.tb04990.x>.
- [53] N.P. Mira, M.C. Teixeira, I. Sá-Correia, Adaptive response and tolerance to weak acids in *Saccharomyces cerevisiae*: A genome-wide view, *Omi. A J. Integr. Biol.* 14 (2010) 525–540. <https://doi.org/10.1089/omi.2010.0072>.
- [54] E.G. Pringsheim, The Genus *Polytomella**, *J. Protozool.* 2 (1955) 137–145. <https://doi.org/10.1111/j.1550-7408.1955.tb02413.x>.
- [55] S. Giannattasio, N. Guaragnella, M. Ždravčić, E. Marra, Molecular mechanisms of *Saccharomyces cerevisiae* stress adaptation and programmed cell death in response to acetic acid, *Front. Microbiol.* 4 (2013). <https://doi.org/10.3389/fmicb.2013.00033>.
- [56] H. Gimmmler, Acidophilic and Acidotolerant Algae, in: *Algal Adapt. to Environ. Stress.*, Springer Berlin Heidelberg, Berlin, Heidelberg, 2001: pp. 259–290. https://doi.org/10.1007/978-3-642-59491-5_9.
- [57] A. Atteia, R. van Lis, G. Mendoza-Hernández, K. Henze, W. Martin, H. Riveros-Rosas, D. González-Halphen, Bifunctional aldehyde/alcohol dehydrogenase (ADHE) in chlorophyte algal mitochondria, *Plant Mol. Biol.* 53 (2003) 175–188. <https://doi.org/10.1023/B:PLAN.0000009274.19340.36>.
- [58] N.R. Boyle, J.A. Morgan, Flux balance analysis of primary metabolism in *Chlamydomonas reinhardtii*, *BMC Syst. Biol.* 3 (2009) 1–14. <https://doi.org/10.1186/1752-0509-3-4>.
- [59] F. Abiusi, R.H. Wijffels, M. Janssen, Doubling of Microalgae Productivity by Oxygen Balanced Mixotrophy, *ACS Sustain. Chem. Eng.* 8 (2020) 6065–6074. <https://doi.org/10.1021/acssuschemeng.0c00990>.
- [60] K.K. Sharma, H. Schuhmann, P.M. Schenk, High lipid induction in microalgae for biodiesel production, *Energies.* 5 (2012) 1532–1553. <https://doi.org/10.3390/en5051532>.
- [61] Y. Chen, S. Vaidyanathan, Simultaneous assay of pigments, carbohydrates, proteins and lipids in microalgae, *Anal. Chim. Acta.* 776 (2013) 31–40. <https://doi.org/10.1016/j.aca.2013.03.005>.
- [62] A.R. Byreddy, A. Gupta, C.J. Barrow, M. Puri, A quick colorimetric method for total lipid quantification in microalgae, *J. Microbiol. Methods.* 125 (2016) 28–32. <https://doi.org/10.1016/j.mimet.2016.04.002>.

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
 - 61
 - 62
 - 63
 - 64
 - 65
- [63] A. Gissibl, A. Sun, A. Care, H. Nevalainen, A. Sunna, Bioproducts From *Euglena gracilis*: Synthesis and Applications, *Front. Bioeng. Biotechnol.* 7 (2019) 1–16. <https://doi.org/10.3389/fbioe.2019.00108>.
- [64] Rubiyatno, T. Matsui, K. Mori, T. Toyama, Paramylon production by *Euglena gracilis* via mixotrophic cultivation using sewage effluent and waste organic compounds, *Bioresour. Technol. Reports.* 15 (2021) 100735. <https://doi.org/10.1016/j.biteb.2021.100735>.
- [65] V. De Preter, I. Arijs, K. Windey, W. Vanhove, S. Vermeire, F. Schuit, P. Rutgeerts, K. Verbeke, Impaired butyrate oxidation in ulcerative colitis is due to decreased butyrate uptake and a defect in the oxidation pathway*, *Inflamm. Bowel Dis.* 18 (2012) 1127–1136. <https://doi.org/10.1002/ibd.21894>.
- [66] M. Nakazawa, C2 metabolism in *Euglena*, in: 2017: pp. 39–45. https://doi.org/10.1007/978-3-319-54910-1_3.
- [67] S. Xiao, L.-K. Ju, Conversion of wastewater-originated waste grease to polyunsaturated fatty acid-rich algae with phagotrophic capability, *Appl. Microbiol. Biotechnol.* 103 (2019) 695–705. <https://doi.org/10.1007/s00253-018-9477-4>.
- [68] B. Sajjadi, W.Y. Chen, A.A.A. Raman, S. Ibrahim, Microalgae lipid and biomass for biofuel production: A comprehensive review on lipid enhancement strategies and their effects on fatty acid composition, *Renew. Sustain. Energy Rev.* 97 (2018) 200–232. <https://doi.org/10.1016/j.rser.2018.07.050>.
- [69] A.E.F. Abomohra, M. El-Sheekh, D. Hanelt, Extracellular secretion of free fatty acids by the chrysophyte *Ochromonas danica* under photoautotrophic and mixotrophic growth, *World J. Microbiol. Biotechnol.* 30 (2014) 3111–3119. <https://doi.org/10.1007/s11274-014-1738-5>.
- [70] P. Chiranjeevi, S. Venkata Mohan, Diverse acidogenic effluents as feedstock for microalgae cultivation: Dual phase metabolic transition on biomass growth and lipid synthesis, *Bioresour. Technol.* 242 (2017) 191–196. <https://doi.org/10.1016/j.biortech.2017.04.059>.