



HAL
open science

An equivalence between gauge-twisted and topologically conditioned scalar Gaussian free fields

Titus Lupu

► **To cite this version:**

Titus Lupu. An equivalence between gauge-twisted and topologically conditioned scalar Gaussian free fields. 2022. hal-03779011v2

HAL Id: hal-03779011

<https://hal.science/hal-03779011v2>

Preprint submitted on 11 Oct 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

AN EQUIVALENCE BETWEEN GAUGE-TWISTED AND TOPOLOGICALLY CONDITIONED SCALAR GAUSSIAN FREE FIELDS

TITUS LUPU

ABSTRACT. We study on the metric graphs two types of scalar Gaussian free fields (GFF), the usual one and the one twisted by a $\{-1, 1\}$ -valued gauge field. We show that the latter can be obtained, up to an additional deterministic transformation, by conditioning the first on a topological event. This event is that all the sign clusters of the field should be trivial for the gauge field, that is to say should not contain cycles with holonomy -1 . We also express the probability of this topological event as a ratio of two determinants of Laplacians to the power $1/2$, the usual Laplacian and the gauge-twisted Laplacian. As an example, this gives on annular planar domains the probability that no sign cluster of the metric graph GFF surrounds the inner hole of the domain.

Based on our result on the metric graph, and on previous works by Werner and Cai-Ding on the clusters of the metric graph GFF in high dimension, we formulate an intensity doubling conjecture. According to it, if the space dimension is high enough, the cycles in the sign clusters of the metric graph GFF converge in the scaling limit to a Brownian loop soup of intensity parameter $\alpha = 1 = 2 \times \frac{1}{2}$, which is the double of the intensity parameter appearing in isomorphism theorems.

1. INTRODUCTION

In this article we consider two types of scalar Gaussian free fields (GFF), the usual one and the one twisted by a $\{-1, 1\}$ -valued gauge field, and observe that the second is essentially obtained from the first by conditioning on a topological (more precisely homotopical) event.

We will work on an abstract finite electrical network $\mathcal{G} = (V, E)$ endowed with conductances $C(x, y) = C(y, x) > 0$ for $\{x, y\} \in E$. The set of vertices V will be divided into two parts, V_{int} and V_{∂} , with V_{int} being considered as the interior vertices, and V_{∂} as the boundary. The discrete GFF ϕ with 0 boundary conditions on V_{∂} is given by the distribution

$$(1.1) \quad \frac{1}{Z} \exp \left(-\frac{1}{2} \sum_{\{x, y\} \in E} C(x, y) (\phi(y) - \phi(x))^2 \right) \prod_{z \in V_{\text{int}}} d\phi(z).$$

Further, in the language of the gauge theory, we consider $\{-1, 1\}$ as our gauge group, and take a gauge field $\sigma \in \{-1, 1\}^E$. The σ -twisted GFF ϕ_{σ} with 0 boundary conditions on V_{∂} has for distribution

$$(1.2) \quad \frac{1}{Z_{\sigma}} \exp \left(-\frac{1}{2} \sum_{\{x, y\} \in E} C(x, y) (\sigma(x, y) \phi(y) - \phi(x))^2 \right) \prod_{z \in V_{\text{int}}} d\phi(z).$$

The gauge field σ corresponds to disorder operators in the language of the Ising model [KC71].

To see the relation between ϕ and ϕ_{σ} one has to look at the level of metric graphs. The metric graph $\tilde{\mathcal{G}}$ associated to \mathcal{G} is obtained by replacing each discrete edge $\{x, y\}$ by a continuous line of length $C(x, y)^{-1}$ joining x and y . The discrete GFF ϕ has a natural extension $\tilde{\phi}$ to the metric graph $\tilde{\mathcal{G}}$, which is a continuous Gaussian random field satisfying a Markov property. This extension has been introduced in [Lup16]. The field $\tilde{\phi}$, unlike ϕ , is known to satisfy a certain number exact identities, including for instance the probabilities of crossings. These relations have been explored in the articles [Lup16, LW18, DPR22, DPR23]; see Section 2.4 for details. The twisted discrete GFF ϕ_{σ} also has a natural extension $\tilde{\phi}_{\sigma}$ to the metric graph $\tilde{\mathcal{G}}$. It is introduced in this paper in Section 3.1. Unlike $\tilde{\phi}$, the field $\tilde{\phi}_{\sigma}$ is not continuous in general, and has one discontinuity point inside each $e \in E$ for which $\sigma(e) = -1$. However, the




FIGURE 1. Conceptual depiction of the metric graph GFF $\tilde{\phi}$ on an annular domain. The black dots represent the boundary V_∂ . The positive, resp. negative values of the fields are in red, resp. blue. Left: no sign cluster of $\tilde{\phi}$ surrounds the inner hole of the annulus. Right: there is a sign cluster (in red) surrounding the inner hole.

absolute value $|\tilde{\phi}_\sigma|$ is a continuous field on the whole $\tilde{\mathcal{G}}$, since the discontinuities of $\tilde{\phi}_\sigma$ consist in switching to the opposite sign by keeping the same absolute value. By taking a double cover of $\tilde{\mathcal{G}}$ induced by the gauge field σ , one can extend $\tilde{\phi}_\sigma$ to a continuous field on the double cover. This is explained in Section 3.3.

Let \mathcal{T}_σ denote the subset of continuous functions on $\tilde{\mathcal{G}}$, made of functions f such that for every connected component U of the non-zero set $\{f \neq 0\}$, U does not contain loops of holonomy -1 for σ . For the notion of holonomy in this setting (product of the values of σ along the edges of the loop), we refer to Section 2.1. But let us give an example. Consider a planar annular domain (one hole), such as depicted on Figures 3 and 1. One can consider a gauge field σ on this annular domain that gives a holonomy -1 to loops that turn an odd number of times around the inner hole, and holonomy 1 to other loops; see Figure 3. Then $f \in \mathcal{T}_\sigma$ if and only if no connected component of $\{f \neq 0\}$ surrounds the inner hole; see Figure 1.

It is easy to see that $|\tilde{\phi}_\sigma| \in \mathcal{T}_\sigma$ a.s., and this is proved in Lemma 3.15 by relying on the extension of $\tilde{\phi}_\sigma$ to the double cover of $\tilde{\mathcal{G}}$. Our main result is the following.

Theorem 1. *Let be a gauge field $\sigma \in \{-1, 1\}^E$. Then*

$$(1.3) \quad \mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) = \frac{Z_\sigma}{Z},$$

where Z and Z_σ are the partition functions appearing in (1.1) and (1.2). Moreover, conditionally on the event $\{\tilde{\phi} \in \mathcal{T}_\sigma\}$, the field $|\tilde{\phi}|$ has the same distribution as $|\tilde{\phi}_\sigma|$.

To obtain the field $\tilde{\phi}_\sigma$, rather than just the absolute value $|\tilde{\phi}_\sigma|$, from the field $\tilde{\phi}$ conditioned on $\tilde{\phi} \in \mathcal{T}_\sigma$, one has to additionally apply a deterministic sign flipping procedure across the discontinuity points. This is explained in Corollary 3.25.

The identity (1.3) is thus a newcomer to the family of exact identities known to be satisfied by $\tilde{\phi}$. We would like to emphasize that Theorem 1 does not require at all the graph \mathcal{G} to be planar. However, for planar graphs the subset \mathcal{T}_σ is simpler to interpret; see Section 5 and in particular 5.4.

We provide three different proofs for Theorem 1. The first proof is presented in Section 3.4. It proceeds entirely through Gaussian computations at the metric graph level, and in particular

does not rely on isomorphism identities between free fields and Markovian paths. The second proof, presented in Section 3.5, involves an isomorphism identity for $\tilde{\phi}_\sigma$ on the metric graph. The third proof relies on the relation between the metric graph GFF and the FK-Ising random cluster model that has been observed in [LW16]. Actually, a result analogous to our Theorem 1 holds for the FK-Ising model (Theorem 4.2). The latter has been communicated to us by Marcin Lis (TU Wien, Vienna) after the prepublication of the first version of this paper. We are grateful to Marcin Lis for this.

We further derive some consequences of Theorem 1 in the continuum limit in dimension 2. First, Theorem 1 gives an expression for the probability that a certain CLE_4 -type loop ensemble on an annular domain contains only contractible loops. The same probability can be obtained by a different method, through the use of Schramm-Sheffield level lines of the 2D continuum GFF. It is however non-obvious at a first glance that the two expressions obtained by two different methods are actually equal. In Section 5.2.1 we check that the two are indeed equal, and this involves a remarkable computation with Jacobi Theta functions. Then, in Section 5.2.2 we derive a version of the Miller-Sheffield coupling, that involves on one hand the annular CLE_4 conditioned on contractibility, and on the other hand the continuum gauge-twisted GFF. We also describe how the renormalized hyperbolic cosine and the renormalized even powers of the 2D continuum GFF transform under topological conditioning.

We also consider the case of higher dimensions, and in particular formulate an intensity doubling conjecture for high dimensions. Here we will state it only informally. We refer to Section 6 for the precise statement and the heuristic behind.

Conjecture (Intensity doubling). *Assume that the space dimension is $d > 8$. Then the cycles in the sign clusters of the metric graph GFF are described in the scaling limit by a Brownian loop soup of intensity parameter $\alpha = 1 = 2 \times \frac{1}{2}$, which is the double of the intensity parameter appearing in isomorphism theorems (Theorems 2.4 and 2.6).*

The intensity doubling might also hold as soon as $d > 6$.

Let us mention some other works where the ratio Z_σ/Z as (1.3) appeared in a probabilistic context. In [vdBCL18] in a planar setting, the quantities of type Z_σ/Z appeared as correlations of the Kramers-Wannier dual of the discrete GFF. Again in a planar context in [BLV16], quantities of form Z^2/Z_σ^2 entered the expression of the probability that a loop-erased random walk goes through a particular edge.

Let us also mention that the effect of the gauge field has been already studied on the fermionic side, the GFF being of course the Euclidean bosonic free field. The Euclidean fermionic free field (without gauge field) is related to spanning trees and wired spanning forests; see [CJS⁺04]. If one adds a gauge field $\sigma \in \{-1, 1\}^E$, then one gets the cycle-rooted spanning forests introduced by Kenyon [Ken11]. These are spanning sub-graphs where a connected component can contain one cycle of holonomy -1 . So, remarkably, while on the fermionic side the effect of a gauge field is to add cycles of holonomy -1 , on the bosonic side the effect is to remove all possible cycles of holonomy -1 .

This article is organized as follows. Section 2 consists of preliminaries where we recall some background that is maybe not common knowledge. In Section 2.1 we recall the notions of gauge field, gauge equivalence and holonomy in our particular setting where the gauge group is $\{-1, 1\}$. Section 2.2 deals with the discrete GFFs, the usual one and the gauge twisted. Section 2.3 deals with random walk representations of these fields. Section 2.4 recalls the method of the metric graph and some results for the metric graph GFF.

In Section 3 we present the results of this article and provide the corresponding proofs. In Section 3.1 we introduce the natural extrapolation $\tilde{\phi}_\sigma$ of the gauge-twisted GFF ϕ_σ to the metric graph. In Section 3.2 we consider the double cover of the discrete graph induced by the gauge field and observe that the usual discrete GFF and the gauge twisted one are projections on two orthogonal subspaces of the discrete GFF on the double cover. In Section 3.3 we do the same at the level of the metric graph, which in particular provides us a continuous extension of $\tilde{\phi}_\sigma$

to the double cover of the metric graph. In Section 3.4 we give a more detailed statement of Theorem 1 and then prove it through direct Gaussian computations. In Section 3.5 we provide an alternative proof of Theorem 1 through the metric graph version of the isomorphism identity between $\tilde{\phi}_\sigma$ and Markovian loop soups due to Kassel and Lévy [KL21]. We also give a stronger version for this isomorphism identity.

In Section 4 we relate the topological conditioning for the metric graph GFF with the topological conditioning in FK-Ising random cluster model. In Section 4.1 we recall the spin Ising model, the FK-Ising model, and the Edwards-Sokal coupling between the two. In Section 4.2 we present the analogue of Theorem 1 in the Ising setting. In Section 4.3 we recall the relation between the Ising setting and the GFF setting. In Section 4.4 we give the third proof of Theorem 1, where it appears as a consequence of the analogous result for FK-Ising.

Section 5 is a discussion around Theorem 1, where we present some interpretations and implications. In Section 5.1 we explain how metric graph annular domains naturally appear. In Section 5.2 we derive some consequences of Theorem 1 on 2D continuum annular domains in the scaling limit. In Section 5.3 we briefly mention what Theorem 1 provides for planar continuum domains with multiple holes, and what it does not. In Section 5.4 we explain how the non-planar setting differs from the planar setting, and what Theorem 1 provides for the non-planar setting.

In Section 6 we present our intensity doubling conjecture for high dimensions and our reasoning behind it.

2. PRELIMINARIES

2.1. On gauge fields, holonomy and gauge equivalence. Let $\mathcal{G} = (V, E)$ be a finite connected undirected graph. We assume that there are no self-loops or multi-edges. We also assume that the set of vertices consists of two disjoint parts, $V = V_{\text{int}} \cup V_\partial$, $V_{\text{int}} \cap V_\partial = \emptyset$, with both V_{int} and V_∂ non-empty. We see V_{int} as the interior vertices and V_∂ as boundary vertices. Each edge $\{x, y\} \in E$ is endowed with a conductance $C(x, y) = C(y, x) > 0$. Thus, \mathcal{G} is an electrical network.

In this paper, a *gauge field* will mean a family $(\sigma(e))_{e \in E} \in \{-1, 1\}^E$. This is the simplest case when the gauge group is $\{-1, 1\}$. We will also use the notation $\sigma(x, y) = \sigma(\{x, y\})$, when $\{x, y\} \in E$. Given an other collection of signs $(\hat{\sigma}(x))_{x \in V} \in \{-1, 1\}^V$, this time above the vertices, it induces a *gauge transformation* $\sigma \mapsto \hat{\sigma} \cdot \sigma$, where $\hat{\sigma} \cdot \sigma \in \{-1, 1\}^E$ is the gauge field defined by

$$(\hat{\sigma} \cdot \sigma)(x, y) = \hat{\sigma}(x)\sigma(x, y)\hat{\sigma}(y).$$

Two gauge fields $\sigma, \sigma' \in \{-1, 1\}^E$ are said to be *gauge-equivalent* if there is $\hat{\sigma} \in \{-1, 1\}^V$ such that $\sigma' = \hat{\sigma} \cdot \sigma$. A gauge field $\sigma \in \{-1, 1\}^E$ is said *trivial* if it is gauge-equivalent to the gauge field with value 1 on every edge.

Given $\wp = (x_1, x_2, \dots, x_n)$ a discrete nearest-neighbor path in \mathcal{G} , the *holonomy* of $\sigma \in \{-1, 1\}^E$ along \wp is the product

$$\text{hol}^\sigma(\wp) = \sigma(x_1, x_2)\sigma(x_2, x_3) \dots \sigma(x_{n-1}, x_n).$$

If the nearest-neighbor path \wp with finitely many jumps is parametrized by continuous time, then $\text{hol}^\sigma(\wp)$ is the holonomy along the discrete skeleton of \wp .

- Lemma 2.1.** (1) *Assume that two gauge fields $\sigma, \sigma' \in \{-1, 1\}^E$ are gauge-equivalent. Then for every loop (i.e. closed path) $\wp = (x_1, x_2, \dots, x_{n-1}, x_n, x_1)$ in \mathcal{G} , $\text{hol}^\sigma(\wp) = \text{hol}^{\sigma'}(\wp)$.*
(2) *Conversely, assume that there is $x_1 \in V$ such that for every loop $\wp = (x_1, x_2, \dots, x_{n-1}, x_n, x_1)$ rooted in x_1 , $\text{hol}^\sigma(\wp) = \text{hol}^{\sigma'}(\wp)$. Then σ and σ' are gauge equivalent.*
(3) *In particular, a gauge field $\sigma \in \{-1, 1\}^E$ is trivial if and only if for every loop \wp , $\text{hol}^\sigma(\wp) = 1$.*

Proof. (1) If $\sigma' = \hat{\sigma} \cdot \sigma$, then for every path $\wp = (x_1, x_2, \dots, x_{n-1}, x_n, x_{n+1})$,

$$\text{hol}^{\sigma'}(\wp) = \hat{\sigma}(x_1)\text{hol}^\sigma(\wp)\hat{\sigma}(x_{n+1}).$$




FIGURE 2. Top: two trivial gauge fields. The edges with sign -1 are in violet. Bottom: a gauge transformation relating the above gauge fields. In red are the $+1$ vertices and in blue the -1 vertices. The black dots represent the boundary V_∂ .

In particular, if $x_{n+1} = x_1$, $\text{hol}^{\sigma'}(\varphi) = \text{hol}^\sigma(\varphi)$.

(2) For each $x \in V$, take a path $\varphi^{x_1, x}$ from x_1 to x in \mathcal{G} and set

$$\hat{\sigma}(x) = \text{hol}^\sigma(\varphi^{x_1, x})\text{hol}^{\sigma'}(\varphi^{x_1, x}).$$

The value of $\hat{\sigma}(x)$ does not depend on the particular choice of the path $\varphi^{x_1, x}$. Indeed, if $\bar{\varphi}^{x_1, x}$ is another path from x_1 to x , then one can concatenate $\bar{\varphi}^{x_1, x}$ with the time reversal $\overleftarrow{\varphi}^{x_1, x}$ of $\varphi^{x_1, x}$, so as to get a loop $\bar{\varphi}^{x_1, x} \wedge \overleftarrow{\varphi}^{x_1, x}$ from x_1 to x_1 , and then

$$\text{hol}^\sigma(\varphi^{x_1, x})\text{hol}^\sigma(\bar{\varphi}^{x_1, x}) = \text{hol}^\sigma(\bar{\varphi}^{x_1, x} \wedge \overleftarrow{\varphi}^{x_1, x}) = \text{hol}^{\sigma'}(\bar{\varphi}^{x_1, x} \wedge \overleftarrow{\varphi}^{x_1, x}) = \text{hol}^{\sigma'}(\varphi^{x_1, x})\text{hol}^{\sigma'}(\bar{\varphi}^{x_1, x}).$$

With $\hat{\sigma}$ defined in this way, we have that $\sigma' = \hat{\sigma} \cdot \sigma$. Note that $\hat{\sigma}$ is not the only element of $\{-1, 1\}^V$ to relate σ and σ' through induced gauge transformation. With \mathcal{G} being connected, there are exactly two such elements of $\{-1, 1\}^V$, $\hat{\sigma}$ and $-\hat{\sigma}$. \square

Figures 2, 3 and 4 provide examples of gauge-equivalent gauge fields together with the corresponding gauge transformation.

2.2. Discrete scalar GFF twisted by a gauge field. Let $\Delta_{\mathcal{G}}$ denote the discrete Laplacian on \mathcal{G} :

$$(\Delta_{\mathcal{G}}f)(x) = \sum_{y \sim x} C(x, y)(f(y) - f(x)), \quad x \in V,$$

where $y \sim x$ means that y is a neighbor of x , i.e. $\{x, y\} \in E$. Let $(G(x, y))_{x, y \in V}$ be the Green's function of $-\Delta_{\mathcal{G}}$ with 0 boundary conditions on V_∂ .

Further, if $\sigma \in \{-1, 1\}^E$ is a gauge field, the discrete Laplacian twisted by σ is

$$(\Delta_{\mathcal{G}, \sigma}f)(x) = \sum_{y \sim x} C(x, y)(\sigma(x, y)f(y) - f(x)), \quad x \in V.$$




FIGURE 3. Top: two non-trivial and gauge-equivalent gauge fields. The edges with sign -1 are in violet. The holonomy of a loop is -1 to the power the number of turns it performs around the inner hole. Bottom: a gauge transformation relating the above gauge fields. In red are the $+1$ vertices and in blue the -1 vertices. The black dots represent the boundary V_∂ .




FIGURE 4. Top: two non-trivial and gauge-equivalent gauge fields. The edges with sign -1 are in violet. In the holonomy of a loop one multiplies by -1 each time one surrounds one of the two holes but not both. Bottom: a gauge transformation relating the above gauge fields. In red are the $+1$ vertices and in blue the -1 vertices. The black dots represent the boundary V_∂ .

The twisted operator $-\Delta_{\mathcal{G},\sigma}$ is still non-negative in the sense that for every $f \in \mathbb{R}^V$,

$$\mathcal{E}_\sigma(f, f) := \sum_{x \in V} (-\Delta_{\mathcal{G},\sigma} f)(x) f(x) = \sum_{\{x,y\} \in E} C(x,y) (\sigma(x,y) f(y) - f(x))^2 \geq 0.$$

Moreover, if f is a function such that $\mathcal{E}_\sigma(f, f) = 0$ and f is 0 on V_∂ , then f is uniformly 0 on V . Indeed, one can see this by induction on the graph distance of a vertex $x \in V$ from V_∂ . Therefore, one can consider the inverse of the restriction of the operator $-\Delta_{\mathcal{G}, \sigma}$ to the functions that are 0 on V_∂ . This is the gauge-twisted Green's function $(G_\sigma(x, y))_{x, y \in V}$ with 0 boundary conditions on V_∂ . It is defined by

$$\forall x \in V, \forall y \in V_\partial, \quad G_\sigma(x, y) = 0,$$

and

$$\forall x_0 \in V_{\text{int}}, \forall x \in V_{\text{int}}, \quad \sum_{y \sim x} C(x, y)(\sigma(x, y)G(x_0, y) - G(x_0, x)) = -\mathbf{1}_{x=x_0}.$$

The function G_σ is symmetric: $G_\sigma(x, y) = G_\sigma(y, x)$. However, unlike for the usual Green's function $G(x, y)$, some of the values $G_\sigma(x, y)$ may be negative. Still, the operator induced by G_σ is non-negative: for every $f \in \mathbb{R}^V$,

$$\sum_{x, y \in V} f(x)G_\sigma(x, y)f(y) \geq 0.$$

Further, if $\sigma, \sigma' \in \{-1, 1\}^E$ are two gauge-equivalent gauge fields, with a gauge transformation between σ and σ' induced by a $\hat{\sigma} \in \{-1, 1\}^V$, then for every $x, y \in V$,

$$G_{\sigma'}(x, y) = \hat{\sigma}(x)G_\sigma(x, y)\hat{\sigma}(y).$$

The discrete scalar *Gaussian free field* (GFF) on \mathcal{G} with boundary conditions 0 on V_∂ is the random Gaussian field $(\phi(x))_{x \in V}$, with $\phi(x) = 0$ for $x \in V_\partial$, and with the probability distribution

$$(2.1) \quad \frac{1}{Z} \exp\left(-\frac{1}{2} \sum_{\{x, y\} \in E} C(x, y)(\varphi(y) - \varphi(x))^2\right) \prod_{z \in V_{\text{int}}} d\varphi(z).$$

The expectation of ϕ is 0. The covariance function of ϕ is the Green's function $(G(x, y))_{x, y \in V}$.

Further, given $\sigma \in \{-1, 1\}^E$ a gauge field, the GFF twisted by σ is the random Gaussian field $(\phi_\sigma(x))_{x \in V}$, with $\phi_\sigma(x) = 0$ for $x \in V_\partial$, and with the probability distribution

$$(2.2) \quad \frac{1}{Z_\sigma} \exp\left(-\frac{1}{2} \sum_{\{x, y\} \in E} C(x, y)(\sigma(x, y)\varphi(y) - \varphi(x))^2\right) \prod_{z \in V_{\text{int}}} d\varphi(z).$$

The expectation of ϕ_σ is 0. Actually, $\phi_\sigma \stackrel{(\text{law})}{=} -\phi_\sigma$. The covariance function of ϕ_σ is the gauge-twisted Green's function $(G_\sigma(x, y))_{x, y \in V}$. If $\sigma, \sigma' \in \{-1, 1\}^E$ are two gauge-equivalent gauge fields, with a gauge transformation between σ and σ' induced by a $\hat{\sigma} \in \{-1, 1\}^V$, then

$$(2.3) \quad (\phi_{\sigma'}(x))_{x \in V} \stackrel{(\text{law})}{=} (\hat{\sigma}(x)\phi_\sigma(x))_{x \in V}.$$

In particular, if the gauge field σ is trivial, then the field ϕ_σ can be obtained via a deterministic transformation from the usual GFF ϕ .

2.3. Measures on paths. Let $(X_t)_{t \geq 0}$ be the nearest-neighbor Markov jump process on \mathcal{G} with the jump rates given by the conductances $C(x, y)$. Let T_{V_∂} denote the first hitting time of V_∂ . We will denote by $p(t, x, y)$ the transition probabilities of the killed process $(X_t)_{0 \leq t \leq T_{V_\partial}}$, so that

$$\sum_{y \in V_{\text{int}}} p(t, x, y) = \mathbb{P}_x(T_{V_\partial} > t).$$

Since the process is symmetric, $p(t, x, y) = p(t, y, x)$. Moreover,

$$\int_0^{+\infty} p(t, x, y) dt = G(x, y).$$

Given $x, y \in V_{\text{int}}$ and $t > 0$, let $\mathbb{P}_t^{x,y}$ denote the bridge probability measure from x to y of duration t , where one conditions on $T_{V_\partial} > t$. The *excursion measure* from x to y is

$$\mu^{x,y}(d\varphi) = \int_0^{+\infty} \mathbb{P}_t^{x,y}(d\varphi) p(t, x, y) dt.$$

So $\mu^{x,y}$ is a measure on nearest-neighbor paths from x to y , parametrized by continuous time, of finite duration. The total mass of $\mu^{x,y}$ is $G(x, y)$. The measure $\mu^{y,x}$ is the image of $\mu^{x,y}$ by time-reversal. The induced measure on discrete skeletons is

$$\mu^{x,y}(x \rightarrow x_1 \rightarrow x_2 \rightarrow \dots \rightarrow x_{n-1} \rightarrow y) = \frac{C(x, x_1)C(x_1, x_2) \dots C(x_{n-2}, x_{n-1})C(x_{n-1}, y)}{W(x)W(x_1)W(x_2) \dots W(x_{n-1})W(y)},$$

where

$$(2.4) \quad W(z) = \sum_{w \sim z} C(z, w).$$

Consider now $\sigma \in \{-1, 1\}^E$ a gauge field. Kassel and Lévy showed in [KL21, Theorem 5.1] the following.

Theorem 2.2 (Kassel-Lévy, [KL21]). *For every $x, y \in V_{\text{int}}$,*

$$G_\sigma(x, y) = \int \text{hol}^\sigma(\varphi) \mu^{x,y}(d\varphi).$$

In other words,

$$\frac{G_\sigma(x, y)}{G(x, y)} = \mathbb{E}[\text{hol}^\sigma(\varphi^{x,y})],$$

where $\varphi^{x,y}$ is a random excursion from x to y distributed according to the probability measure $\mu^{x,y}/G(x, y)$. In particular, for every $x \in V_{\text{int}}$,

$$\int \text{hol}^\sigma(\varphi) \mu^{x,x}(d\varphi) = G_\sigma(x, x) > 0.$$

The measure on continuous time random walk loops introduced by Le Jan [LJ10, LJ11] is

$$(2.5) \quad \mu^{\text{loop}}(d\varphi) = \sum_{x \in V_{\text{int}}} \int_0^{+\infty} \mathbb{P}_t^{x,x}(d\varphi) p(t, x, x) \frac{dt}{t} = \frac{1}{T(\varphi)} \sum_{x \in V_{\text{int}}} \mu^{x,x}(d\varphi),$$

where $T(\varphi)$ denotes the duration of a path φ . There are two types of loops, those that visit at least two different vertices, and those that stay in one vertex and do not perform jumps. For $n \geq 2$ jumps, the measure induced on discrete skeletons is

$$\mu^{\text{loop}}(x_1 \rightarrow x_2 \rightarrow \dots \rightarrow x_{n-1} \rightarrow x_n \rightarrow x_1) = \frac{1}{n} \frac{C(x_1, x_2) \dots C(x_{n-1}, x_n)C(x_n, x_1)}{W(x_1)W(x_2) \dots W(x_{n-1})W(x_n)}.$$

So this is the same measure on discrete-time loops as the one that appeared in [LTF07] and [LL10, Chapter 9]. The total mass of the loops that visit at least two vertices is

$$(2.6) \quad \mu^{\text{loop}}(\{\text{Loops that visit at least 2 vertices}\}) = \log \left(\det((G(x, y))_{x, y \in V_{\text{int}}}) \prod_{x \in V_{\text{int}}} W(x) \right);$$

see [LJ11, Equation (2.5)]. Besides the loops that visit at least two vertices, μ^{loop} also puts weight on loops that stay in one vertex and do not jump. For every $x \in V_{\text{int}}$, the induced measure on the duration of loops that only stay in x is

$$e^{-t/G(x,x)} \frac{dt}{t}.$$

Given $\sigma \in \{-1, 1\}^E$ a gauge field, the (signed) measure on loops twisted by σ is

$$\mu_\sigma^{\text{loop}}(d\varphi) = \text{hol}^\sigma(\varphi) \mu^{\text{loop}}(d\varphi).$$

The measure μ_σ^{loop} is signed and its total variation measure is μ^{loop} . The measure μ_σ^{loop} is the same for the whole gauge-equivalence class of σ . The signed measure μ_σ^{loop} can be decomposed as

$$\mu_\sigma^{\text{loop}} = \mu_{\sigma,+}^{\text{loop}} - \mu_{\sigma,-}^{\text{loop}},$$

where $\mu_{\sigma,+}^{\text{loop}}$ is the restriction of μ_σ^{loop} to loops φ with $\text{hol}^\sigma(\varphi) = 1$, and $\mu_{\sigma,-}^{\text{loop}}$ is the restriction of μ_σ^{loop} to loops φ with $\text{hol}^\sigma(\varphi) = -1$. According to [KL21, Corollary 3.7], a formula similar to (2.6) also holds in the presence of a gauge field:

$$(2.7) \quad \mu_\sigma^{\text{loop}}(\{\text{Loops that visit at least 2 vertices}\}) = \log \left(\det((G_\sigma(x, y))_{x, y \in V_{\text{int}}}) \prod_{x \in V_{\text{int}}} W(x) \right).$$

So in particular, one gets the following.

Corollary 2.3. *The following identity holds:*

$$\frac{\det((G_\sigma(x, y))_{x, y \in V_{\text{int}}})}{\det((G(x, y))_{x, y \in V_{\text{int}}})} = \exp \left(-2\mu^{\text{loop}}(\{\text{Loops } \varphi \text{ with } \text{hol}^\sigma(\varphi) = -1\}) \right).$$

Proof. By combining the formulas (2.6) and (2.7), one obtains

$$\begin{aligned} \log \left(\frac{\det((G_\sigma(x, y))_{x, y \in V_{\text{int}}})}{\det((G(x, y))_{x, y \in V_{\text{int}}})} \right) &= (\mu_\sigma^{\text{loop}} - \mu^{\text{loop}})(\{\text{Loops that visit at least 2 vertices}\}) \\ &= -2\mu_{\sigma,-}^{\text{loop}}(\{\text{Loops that visit at least 2 vertices}\}). \end{aligned}$$

Further, a loop φ with $\text{hol}^\sigma(\varphi) = -1$ has to visit at least two vertices. Thus,

$$\mu_{\sigma,-}^{\text{loop}}(\{\text{Loops that visit at least 2 vertices}\}) = \mu^{\text{loop}}(\{\text{Loops } \varphi \text{ with } \text{hol}^\sigma(\varphi) = -1\}). \quad \square$$

Given a path φ on \mathcal{G} parametrized by continuous time, $\ell^x(\varphi)$ will denote its total time spent at vertex x , $x \in V$. Given $\alpha > 0$, \mathcal{L}^α will denote the Poisson point process of loops with intensity measure $\alpha\mu^{\text{loop}}$. This is the continuous time *random walk loop soup* [LJ10, LJ11]. For $x \in V$, $\ell^x(\mathcal{L}^\alpha)$ will denote its *occupation field* in x :

$$\ell^x(\mathcal{L}^\alpha) = \sum_{\varphi \in \mathcal{L}^\alpha} \ell^x(\varphi).$$

For the particular value $\alpha = 1/2$, this occupation field is Gaussian. More precisely one has the Le Jan's isomorphism.

Theorem 2.4 (Le Jan, [LJ10, LJ11]). *For $\alpha = 1/2$, the following identity in law holds:*

$$(\ell^x(\mathcal{L}^{1/2}))_{x \in V} \stackrel{(law)}{=} \left(\frac{1}{2} \phi(x)^2 \right)_{x \in V},$$

where ϕ is the GFF (2.1).

Now take $\sigma \in \{-1, 1\}^E$ a gauge field. In [KL21, Theorem 6.6], Kassel and Lévy proved the following extension of Le Jan's isomorphism.

Theorem 2.5 (Kassel-Lévy, [KL21]). *Denote by $\mathcal{L}_{\sigma,+}^{1/2}$, resp. $\mathcal{L}_{\sigma,-}^{1/2}$, a Poisson point process with intensity measure $\frac{1}{2}\mu_{\sigma,+}^{\text{loop}}$, resp. $\frac{1}{2}\mu_{\sigma,-}^{\text{loop}}$. Recall that ϕ_σ denotes the gauge-twisted GFF (2.2). Take ϕ_σ and $\mathcal{L}_{\sigma,-}^{1/2}$ to be independent. Then the following identity in law holds*

$$(2.8) \quad (\ell^x(\mathcal{L}_{\sigma,+}^{1/2}))_{x \in V} \stackrel{(law)}{=} \left(\frac{1}{2} \phi_\sigma(x)^2 + \ell^x(\mathcal{L}_{\sigma,-}^{1/2}) \right)_{x \in V}.$$

In particular, the field ϕ_σ^2 is stochastically dominated by the field ϕ^2 .

The identity (2.8) tells that in some sense the field $\frac{1}{2}\phi_\sigma^2$ is distributed as the occupation field of a Poisson point process with intensity measure $\frac{1}{2}\mu_\sigma^{\text{loop}}$. However, the measure $\frac{1}{2}\mu_\sigma^{\text{loop}}$ is signed, unless the gauge field σ is trivial, and thus cannot be an intensity for a Poisson. Actually, the loops φ with $\text{hol}^\sigma(\varphi) = -1$ have to be counted negatively.

2.4. GFF on metric graphs. Here we will briefly recall the notion of the GFF on metric graphs. For details we refer to [Lup16].

The *metric graph* associated to the electrical network \mathcal{G} , which we will denote by $\tilde{\mathcal{G}}$, is obtained by replacing each edge $e = \{x, y\} \in E$ by a continuous line segment $I_e = I_{\{x, y\}}$, with endpoints x and y . Moreover, $\tilde{\mathcal{G}}$ is endowed with a metric, by setting the length of I_e to be $C(x, y)^{-1}$ (i.e. the resistance, the inverse of the conductance), and with the corresponding length measure, which we will denote by \tilde{m} . So $\tilde{\mathcal{G}}$ is a continuous and connected metric space.

The discrete GFF ϕ on \mathcal{G} (2.1) can be interpolated to a continuous Gaussian field $\tilde{\phi}$. The restriction of $\tilde{\phi}$ to the vertices V is the discrete GFF ϕ . Conditionally on ϕ , the values of $\tilde{\phi}$ along an edge-line $I_{\{x, y\}}$ are distributed as a standard one-dimensional Brownian bridge between $\phi(x)$ and $\phi(y)$ of length $C(x, y)^{-1}$, with values on different edge-lines being independent. The field $\tilde{\phi}$ is the *metric graph GFF*. It satisfies a strong Markov property when cutting not only along the vertices, but also inside edge-lines; see [Lup16, Section 3]. The covariance of $\tilde{\phi}$ is given by the extension of the Green's function to $\tilde{\mathcal{G}} \times \tilde{\mathcal{G}}$, which we will still denote $G(x, y)$.

The Markov jump process $(X_t)_{t \geq 0}$ on \mathcal{G} can be extended to a continuous Markov diffusion process on $\tilde{\mathcal{G}}$, which we will denote by $(\tilde{X}_t)_{t \geq 0}$. Inside an edge-line I_e , \tilde{X}_t behaves as a one-dimensional Brownian motion, and once the process reaches a vertex $x \in V$, it performs Brownian excursions inside each adjacent edge-line. See [Lup16, Section 2] for details. In order to fit with the isomorphism identities, we normalize \tilde{X}_t so that inside every edge-line I_e it behaves like a Brownian motion with quadratic variation $2dt$. Just as a one-dimensional Brownian motion, the process $(\tilde{X}_t)_{t \geq 0}$ admits a family of local times $(\ell_t^x(\tilde{X}))_{x \in \tilde{\mathcal{G}}, t \geq 0}$, continuous in (x, t) , characterized by

$$\int_0^t f(\tilde{X}_s) ds = \int_{\tilde{\mathcal{G}}} f(x) \ell_t^x(\tilde{X}) \tilde{m}(dx).$$

Consider the continuous additive functional $A(t)$ and its inverse $A^{-1}(t)$:

$$(2.9) \quad A(t) = \sum_{x \in V} \ell_t^x(\tilde{X}), \quad A^{-1}(t) = \inf\{s \geq 0 | A(s) > t\}.$$

Then $(\tilde{X}_{A^{-1}(t)})_{t \geq 0}$ is distributed as the Markov jump process $(X_t)_{t \geq 0}$.

Let \tilde{T}_{V_∂} denote the first time \tilde{X}_t hits V_∂ . By construction, $A(\tilde{T}_{V_\partial}) = T_{V_\partial}$. Considered the process $(\tilde{X}_t)_{0 \leq t \leq \tilde{T}_{V_\partial}}$ killed upon hitting V_∂ . Let $\tilde{p}(t, x, y)$ be the transition densities of the killed process, so that

$$\int_{\tilde{\mathcal{G}}} \tilde{p}(t, x, y) \tilde{m}(dy) = \mathbb{P}_x(\tilde{T}_{V_\partial} > t).$$

For $x, y \in \tilde{\mathcal{G}} \setminus V_\partial$ and $t > 0$, let $\tilde{\mathbb{P}}_t^{x, y}$ denote the bridge probability measure from x to y in time t , where one conditions on $\tilde{T}_{V_\partial} > t$. The measure on loops on the metric graph is

$$(2.10) \quad \tilde{\mu}^{\text{loop}}(d\wp) = \int_{\tilde{\mathcal{G}}} \int_0^{+\infty} \tilde{\mathbb{P}}_t^{x, x}(d\wp) \tilde{p}(t, x, x) \frac{dt}{t} \tilde{m}(dx).$$

For $\alpha > 0$, $\tilde{\mathcal{L}}^\alpha$ will denote the Poisson point process of loops on $\tilde{\mathcal{G}}$ of intensity $\alpha \tilde{\mu}^{\text{loop}}$. This is the *metric graph loop soup*. The loops in $\tilde{\mathcal{L}}^\alpha$ can be divided into two: those that visit vertices in V and those that only stay in the interior of an edge-line. If one takes the trace on V of the loops in $\tilde{\mathcal{L}}^\alpha$ that intersect V , by applying the time change A^{-1} (2.9), one gets the continuous time random walk loop soup \mathcal{L}^α , actually up to a rerooting of the loops. See [Lup16, Section 2] for details. Given $\wp \in \tilde{\mathcal{L}}^\alpha$ and $x \in \tilde{\mathcal{G}}$, $\ell^x(\wp)$ will denote the cumulative local time of \wp in x . The *occupation field* of $\tilde{\mathcal{L}}^\alpha$ is

$$\ell^x(\tilde{\mathcal{L}}^\alpha) = \sum_{\wp \in \tilde{\mathcal{L}}^\alpha} \ell^x(\wp).$$

For $x \in V$, we have that $\ell^x(\tilde{\mathcal{L}}^\alpha) = \ell^x(\mathcal{L}^\alpha)$.

Next we will consider the clusters of loops in $\tilde{\mathcal{L}}^\alpha$. Two loops $\wp, \wp' \in \tilde{\mathcal{L}}^\alpha$ belong to the same *cluster* if they are connected by a finite chain of loops in $\tilde{\mathcal{L}}^\alpha$. We will also see a cluster \mathcal{C} of $\tilde{\mathcal{L}}^\alpha$ as a subset of $\tilde{\mathcal{G}}$ by taking the union of the ranges of loops forming the cluster. The clusters of $\tilde{\mathcal{L}}^\alpha$ are exactly the connected components of $\{x \in \tilde{\mathcal{G}} | \ell^x(\tilde{\mathcal{L}}^\alpha) > 0\}$. In [Lup16, Theorem 1] is shown that for $\alpha = 1/2$, there is a one to one correspondence between the clusters of $\tilde{\mathcal{L}}^{1/2}$ and the sign components of $\tilde{\phi}$.

Theorem 2.6 (Lupu, [Lup16]). *Take $\tilde{\mathcal{L}}^{1/2}$ a metric graph loop soup on $\tilde{\mathcal{G}}$ of parameter $\alpha = 1/2$. For each cluster \mathcal{C} additionally sample a conditionally independent sign $\zeta(\mathcal{C})$ with*

$$\mathbb{P}(\zeta(\mathcal{C}) = 1 | \tilde{\mathcal{L}}^{1/2}) = \mathbb{P}(\zeta(\mathcal{C}) = -1 | \tilde{\mathcal{L}}^{1/2}) = 1/2.$$

For $x \in \tilde{\mathcal{G}}$ such that $\ell^x(\tilde{\mathcal{L}}^{1/2}) > 0$, let $\mathcal{C}(x)$ denote the cluster of $\tilde{\mathcal{L}}^{1/2}$ containing x . Then the field

$$(\zeta(\mathcal{C}(x)) \sqrt{2\ell^x(\tilde{\mathcal{L}}^{1/2})})_{x \in \tilde{\mathcal{G}}}$$

is distributed as a metric graph GFF $\tilde{\phi}$.

The result above comes from an application of the Le Jan's isomorphism (Theorem 2.4) at the metric graph level and the use of the intermediate value property of the continuous field $\tilde{\phi}$.

Beside the exact correspondence between the sign components of $\tilde{\phi}$ and the clusters of $\tilde{\mathcal{L}}^{1/2}$, the metric graph GFF $\tilde{\phi}$ satisfies many other exact solvability properties that one does not get for the discrete GFF ϕ . Next we give a brief list of these exact identities, and further in Section 3 we will prove a new one, related to a special kind of topological conditioning; see Theorem 1.

- As observed in [Lup16, Proposition 5.2], given $x, y \in \tilde{\mathcal{G}}$, the probability that x and y belong to the same sign component of $\tilde{\phi}$, or alternatively to the same cluster of $\tilde{\mathcal{L}}^{1/2}$, equals

$$\mathbb{E}[\text{sign}(\tilde{\phi}(x)) \text{sign}(\tilde{\phi}(y))] = \frac{2}{\pi} \arcsin \left(\frac{G(x, y)}{\sqrt{G(x, x)G(y, y)}} \right).$$

- If V_∂ is divided into 3 parts $V_{\partial,0}$, $V_{\partial,1}$ and $V_{\partial,2}$, with $V_{\partial,0}$ allowed to be empty, and if the boundary condition of a metric graph GFF $\tilde{\phi}$ is strictly positive on $V_{\partial,1}$ and $V_{\partial,2}$, and zero on $V_{\partial,0}$ (rather than zero on the whole V_∂), then there is an explicit formula for the existence of a positive crossing from $V_{\partial,1}$ to $V_{\partial,2}$:

$$\mathbb{P}\left(V_{\partial,1} \xrightarrow{\tilde{\phi} > 0} V_{\partial,2}\right) = 1 - \exp\left(-2 \sum_{x \in V_{\partial,1}} \sum_{y \in V_{\partial,2}} \tilde{\phi}(x) H(x, y) \tilde{\phi}(y)\right),$$

where $H(x, y)$ is the boundary Poisson kernel on $V_\partial \times V_\partial$. This is [LW18, Formula (18)]. If the boundary condition mixes values of different sign, no analogous explicit formula is known.

- In the articles [DPR22, DPR23] the authors provide the exact law for the effective conductance (called capacity there) between the boundary V_∂ and the connected component containing x_0 of the level set

$$\{x \in \tilde{\mathcal{G}} | \tilde{\phi}(x) \geq h\}$$

($x_0 \in \tilde{\mathcal{G}}$ and $h \geq 0$ fixed). This also can be derived from [LW18, Proposition 5].

- The Lévy transformation for the one-dimensional Brownian motion can be extended to the general metric graph GFFs, as explained in [LW18].

3. GAUGE-TWISTED GFF ON METRIC GRAPH, DOUBLE COVER AND EQUIVALENCE TO TOPOLOGICAL CONDITIONING

3.1. Gauge-twisted GFF and subdivision of edges. Consider the electrical network $\mathcal{G} = (V, E)$ as in Section 2.1. For $N \in \mathbb{N} \setminus \{0\}$, we will denote by $\mathcal{G}^{(N)} = (V^{(N)}, E^{(N)})$ the electrical

network obtained from $\mathcal{G} = (V, E)$ by subdividing each edge $e \in E$ into N . In this way, $\mathcal{G}^{(1)} = \mathcal{G}$. In general,

$$|E^{(N)}| = N|E|, \quad |V^{(N)}| = |V| + (N-1)|E|.$$

Moreover, if N is a divisor of N' , then $V^{(N)}$ is naturally a subset of $V^{(N')}$. In particular, we always have $V \subset V^{(N)}$. Given $e \in E$, we will denote $E^{(N)}(e)$ the subset of $E^{(N)}$ made of edges obtained by subdivision of e . We also denote by $V^{(N)}(e)$ the endpoints of edges in $E^{(N)}(e)$. So $|E^{(N)}(e)| = N$, $|V^{(N)}(e)| = N+1$ and $|V^{(N)}(e) \setminus V| = N-1$. We endow $\mathcal{G}^{(N)}$ with the following conductances: for every $e \in E$ and $e' \in E^{(N)}(e)$, $C^{(N)}(e') = NC(e)$. Let be the energy

$$\mathcal{E}^{(N)}(f, f) = \sum_{\{x, y\} \in E^{(N)}} C^{(N)}(x, y)(f(y) - f(x))^2.$$

We also see the electrical network $\mathcal{G}^{(N)}$, more precisely $V^{(N)}$, as a subset of the metric graph $\tilde{\mathcal{G}}$. In this way, $V^{(N)}(e)$ is a set of $N+1$ points on I_e with equal spacing $(NC(e))^{-1}$, which includes the two endpoints of e .

If $(\phi^{(N)}(x))_{x \in V^{(N)}}$ is a GFF on $\mathcal{G}^{(N)}$, then its restriction to $V = V^{(1)}$ is a GFF on \mathcal{G} . Indeed,

$$\mathcal{E}^{(N)}(f, f) = \mathcal{E}^{(1)}(f|_V, f|_V) + N(N-1) \sum_{e \in E} C(e) \sum_{i=1}^N (f(x_i) - f(x_{i-1}) - (f(x_N) - f(x_0))/N)^2,$$

where x_0, x_1, \dots, x_N is the ordered set of points in $V^{(N)}(e)$. Moreover, if $(\tilde{\phi}(x))_{x \in \tilde{\mathcal{G}}}$ is a metric graph GFF, then its restriction $\tilde{\phi}|_{V^{(N)}}$ is distributed as a GFF $\phi^{(N)}$ on $\mathcal{G}^{(N)}$. So, in a sense, $\tilde{\phi}$ is the limit of $\phi^{(N)}$ as $N \rightarrow +\infty$.

Now, given a gauge field $\sigma \in \{-1, 1\}^E$, we similarly want a natural extension of a σ -twisted GFF ϕ_σ to the subdivided graphs $\mathcal{G}^{(N)}$, and ultimately to the metric graph $\tilde{\mathcal{G}}$. So, given $\sigma \in \{-1, 1\}^E$, we will denote by $\sigma^{(N)}$ the following element of $\{-1, 1\}^{E^{(N)}}$. If $e \in E$ and $\sigma(e) = 1$, then for every $e' \in E^{(N)}(e)$, $\sigma^{(N)}(e') = 1$. However, if $e \in E$ and $\sigma(e) = -1$, then among the edges in $E^{(N)}(e)$ there is exactly one with sign -1 under $\sigma^{(N)}$, all the $N-1$ other having sign $+1$ (so there are actually choices to make for $\sigma^{(N)}$, N choices for each $e \in E$ with $\sigma(e) = -1$).

Proposition 3.1. *Let $N \in \mathbb{N} \setminus \{0\}$, $\sigma \in \{-1, 1\}^E$ and $\sigma^{(N)} \in \{-1, 1\}^{E^{(N)}}$ as above. Let $(\phi_{\sigma^{(N)}}^{(N)}(x))_{x \in V^{(N)}}$ be the $\sigma^{(N)}$ -twisted GFF on $\mathcal{G}^{(N)}$ with 0 boundary conditions on $V_\partial \subset V \subset V^{(N)}$. Then its restriction to V is distributed as $(\phi_\sigma(x))_{x \in V}$, the σ -twisted GFF on \mathcal{G} with 0 boundary conditions.*

Proof. Consider the energy

$$\mathcal{E}_\sigma^{(N)}(f, f) = \sum_{\{x, y\} \in E^{(N)}} C^{(N)}(x, y)(\sigma^{(N)}(x, y)f(y) - f(x))^2.$$

Let $e \in E$ and let x_0, x_1, \dots, x_N be the ordered set of points in $V^{(N)}(e)$. For $i \in \{1, \dots, N\}$, let be

$$\tilde{\sigma}_i = \sigma^{(N)}(x_0, x_1) \dots \sigma^{(N)}(x_{i-1}, x_i).$$

Then

$$(3.1) \quad \begin{aligned} \sum_{i=1}^N C^{(N)}(x_{i-1}, x_i)(\sigma^{(N)}(x_{i-1}, x_i)f(x_i) - f(x_{i-1}))^2 &= C(x_0, x_N)(\sigma(x_0, x_N)f(x_N) - f(x_0))^2 \\ &+ N(N-1)C(x_0, x_N) \sum_{i=1}^N (\tilde{\sigma}_i f(x_i) - \tilde{\sigma}_{i-1} f(x_{i-1}) - (\sigma(x_0, x_N)f(x_N) - f(x_0))/N)^2. \end{aligned}$$

This induces a decomposition of the energy $\mathcal{E}_\sigma^{(N)}$. This implies that $\phi_{\sigma^{(N)}}^{(N)}$ restricted to V is distributed as ϕ_σ . Moreover, the fields

$$(\tilde{\sigma}_i \phi_{\sigma^{(N)}}^{(N)}(x_i) + i(\sigma(x_0, x_N) \phi_{\sigma^{(N)}}^{(N)}(x_N) - \phi_{\sigma^{(N)}}^{(N)}(x_0))/N)_{1 \leq i \leq N-1, e \in E},$$

are independent from $\phi_{\sigma^{(N)}}^{(N)}$ restricted to V , with also independence across $e \in E$. \square

In the sequel, $\sigma \in \{-1, 1\}^E$ and, for the sake of simplicity, N will be odd and $\sigma^{(N)}$ is chosen such that for $e \in E$ with $\sigma(e) = -1$, the middle edge in $E^{(N)}(e)$ has the sign -1 under $\sigma^{(N)}$. We will define a random Gaussian field $\tilde{\phi}_\sigma$ on the metric graph $\tilde{\mathcal{G}}$ as follows. We consider the following independent objects.

- The twisted discrete GFF $(\phi_\sigma(x))_{x \in V}$.
- For every edge $e \in E$, and independent Brownian bridge $(W_e(u))_{0 \leq u \leq C(e)^{-1}}$ of length $C(e)^{-1}$, from 0 to 0.

We will also chose for each edge $e \in E$ and arbitrary orientation and denote its endpoints e_- and e_+ . For $u \in [0, C(e)^{-1}]$, $x_{e,u}$ will denote the point of I_e at distance u from e_+ . We define $(\tilde{\phi}_\sigma(x))_{x \in \tilde{\mathcal{G}}}$ as follows.

- For $x \in V$, $\tilde{\phi}_\sigma(x) = \phi_\sigma(x)$.
- If $e \in E$ and $\sigma(e) = 1$, then for every $u \in (0, C(e)^{-1})$,

$$(3.2) \quad \tilde{\phi}_\sigma(x_{e,u}) = W_e(u) + C(e)(u\phi_\sigma(e_-) + (C(e)^{-1} - u)\phi_\sigma(e_)).$$

- If $e \in E$ and $\sigma(e) = -1$, then for $u \in (0, C(e)^{-1}/2)$,

$$(3.3) \quad \tilde{\phi}_\sigma(x_{e,u}) = W_e(u) + C(e)(-u\phi_\sigma(e_-) + (C(e)^{-1} - u)\phi_\sigma(e_)),$$

and for $u \in (C(e)^{-1}/2, C(e)^{-1})$,

$$(3.4) \quad \tilde{\phi}_\sigma(x_{e,u}) = -W_e(u) + C(e)(u\phi_\sigma(e_-) - (C(e)^{-1} - u)\phi_\sigma(e_)).$$

The value for $u = C(e)^{-1}/2$ is not specified and can be chosen arbitrary.

Defined in this way, the field $\tilde{\phi}_\sigma$ is discontinuous in the middle of each I_e for $e \in \{e \in E | \sigma(e) = -1\}$, with

$$\lim_{u \rightarrow (C(e)^{-1}/2)_-} \tilde{\phi}_\sigma(x_{e,u}) = - \lim_{u \rightarrow (C(e)^{-1}/2)_+} \tilde{\phi}_\sigma(x_{e,u}).$$

However, the absolute value $|\tilde{\phi}_\sigma|$ extends to a continuous field on $\tilde{\mathcal{G}}$. Note that the law of $\tilde{\phi}_\sigma$ does not depend on the arbitrary choice of orientations for the edges, since $\phi_\sigma \stackrel{(\text{law})}{=} -\phi_\sigma$ and $W_e \stackrel{(\text{law})}{=} -W_e$.

Proposition 3.2. *For every N odd, the restriction of $\tilde{\phi}_\sigma$ to $V^{(N)}$ is distributed as $\phi_{\sigma^{(N)}}^{(N)}$.*

Proof. This is an immediate consequence of the decomposition (3.1). \square

So the metric graph field $\tilde{\phi}_\sigma$ appears as the limit of $\phi_{\sigma^{(N)}}^{(N)}$ as $N \rightarrow +\infty$. However, it is not continuous (except for σ equal to 1 everywhere). To recover continuous fields we will work on a double cover of $\tilde{\mathcal{G}}$; see Section 3.3.

3.2. Double covers induced by gauge fields. Let $\sigma \in \{-1, 1\}^E$. We will introduce a double cover $\mathcal{G}_\sigma^{\text{db}} = (V^{\text{db}}, E_\sigma^{\text{db}})$ of the graph \mathcal{G} , which is as follows. V^{db} consist of two copies of V , $V^{\text{db}} = V_1 \cup V_2$, and the projection $\pi_\sigma : V^{\text{db}} \rightarrow V$ induces a bijection between V_1 and V and between V_2 and V . The set of edges E_σ^{db} is as follows. Let $\{x, y\} \in E$ and let $x_1, y_1 \in V_1$ and $x_2, y_2 \in V_2$ such that $\pi_\sigma(x_1) = \pi_\sigma(x_2) = x$ and $\pi_\sigma(y_1) = \pi_\sigma(y_2) = y$.

- If $e \in E$ and $\sigma(e) = 1$, then $\{x_1, y_1\}, \{x_2, y_2\} \in E_\sigma^{\text{db}}$.
- If $e \in E$ and $\sigma(e) = -1$, then $\{x_1, y_2\}, \{x_2, y_1\} \in E_\sigma^{\text{db}}$.




FIGURE 5. Left: a graph with 4 vertices and 5 edges. Right: its double cover. The edges in violet correspond to $\sigma(e) = -1$.

See Figure 5 for an example. We also have a projection $\pi_\sigma : E_\sigma^{\text{db}} \rightarrow E$ such that

$$\pi_\sigma(\{\hat{x}, \hat{y}\}) = \{\pi_\sigma(\hat{x}), \pi_\sigma(\hat{y})\}.$$

In this way π_σ is a graph covering map from $\mathcal{G}_\sigma^{\text{db}}$ to \mathcal{G} . Given $x_1 \in V_1$ and $x_2 \in V_2$ such that $\pi_\sigma(x_1) = \pi_\sigma(x_2)$, we will define $\psi(x_1) = x_2$ and $\psi(x_2) = x_1$. In this way, ψ is a bijection from V^{db} to V^{db} . It is also a graph automorphism of $\mathcal{G}_\sigma^{\text{db}}$, that is to say if $\{\hat{x}, \hat{y}\} \in E_\sigma^{\text{db}}$ then $\{\psi(\hat{x}), \psi(\hat{y})\} \in E_\sigma^{\text{db}}$. The map ψ is also a covering automorphism, because $\pi_\sigma \circ \psi = \pi_\sigma$.

Lemma 3.3. *Let $\varphi = (x_1, x_2, \dots, x_{n-1}, x_n, x_1)$ be a nearest neighbor loop on \mathcal{G} and let $\hat{\varphi} = (\hat{x}_1, \hat{x}_2, \dots, \hat{x}_{n-1}, \hat{x}_n, \hat{x}_{n+1})$ be a lift of φ on $\mathcal{G}_\sigma^{\text{db}}$ with respect to the covering map π_σ : $\pi_\sigma(\hat{x}_i) = x_i$ for $i \in \{1, 2, \dots, n\}$, $\pi_\sigma(\hat{x}_{n+1}) = x_1$, and $\{\hat{x}_i, \hat{x}_{i+1}\} \in E_\sigma^{\text{db}}$ for $i \in \{1, 2, \dots, n\}$. Then $\text{hol}^\sigma(\hat{\varphi}) = 1$ if and only if $\hat{x}_{n+1} = \hat{x}_1$. Otherwise $\text{hol}^\sigma(\hat{\varphi}) = -1$ and $\hat{x}_{n+1} = \psi(\hat{x}_1)$.*

Proof. By construction, whenever $\sigma(x_i, x_{i+1}) = 1$, then \hat{x}_i and \hat{x}_{i+1} belong to the same copy V_1 or V_2 . Whenever $\sigma(x_i, x_{i+1}) = -1$ then either $\hat{x}_i \in V_1$ and $\hat{x}_{i+1} \in V_2$, or vice-versa. So $\hat{x}_{n+1} = \hat{x}_1$ if and only even there is an even number of transitions from V_1 to V_2 or from V_2 to V_1 , that is to say that the number of $i \in \{1, 2, \dots, n\}$ such that $\sigma(x_i, x_{i+1}) = -1$ is even, which exactly means that $\text{hol}^\sigma(\hat{\varphi}) = 1$. \square

Lemma 3.4. *Let $\sigma, \sigma' \in \{-1, 1\}^E$.*

- (1) *Assume that σ and σ' are gauge equivalent, and let $\hat{\sigma} \in \{-1, 1\}^V$ such that $\sigma' = \hat{\sigma} \cdot \sigma$. Define $\psi_{\hat{\sigma}} : V^{\text{db}} \rightarrow V^{\text{db}}$ as follows. Let $\hat{x} \in V^{\text{db}}$. If $\hat{\sigma}(\pi_\sigma(\hat{x})) = 1$, then $\psi_{\hat{\sigma}}(\hat{x}) = \hat{x}$. If $\hat{\sigma}(\pi_\sigma(\hat{x})) = -1$, then $\psi_{\hat{\sigma}}(\hat{x}) = \psi(\hat{x})$. Then $\psi_{\hat{\sigma}}$ induces a graph isomorphism between $\mathcal{G}_\sigma^{\text{db}}$ and $\mathcal{G}_{\sigma'}^{\text{db}}$, which is moreover a covering isomorphism, i.e. $\pi_{\sigma'} = \pi_\sigma \circ \psi_{\hat{\sigma}}$.*
- (2) *Conversely, if there exists a covering isomorphism between $\mathcal{G}_\sigma^{\text{db}}$ et $\mathcal{G}_{\sigma'}^{\text{db}}$, then σ and σ' are gauge equivalent.*

Proof. (1) The map $\psi_{\hat{\sigma}}$ is clearly a bijection from V^{db} to itself, and it is clear that $\pi_{\sigma'} = \pi_\sigma \circ \psi_{\hat{\sigma}}$. One needs only to check that whenever $\{\hat{x}, \hat{y}\} \in E_\sigma^{\text{db}}$, then also $\{\psi_{\hat{\sigma}}(\hat{x}), \psi_{\hat{\sigma}}(\hat{y})\} \in E_{\sigma'}^{\text{db}}$. But this is clear from the definition of $\psi_{\hat{\sigma}}$ and the fact that

$$\sigma'(\pi_{\sigma'}(\psi_{\hat{\sigma}}(\hat{x})), \pi_{\sigma'}(\psi_{\hat{\sigma}}(\hat{y}))) = \sigma'(\pi_\sigma(\hat{x}), \pi_\sigma(\hat{y})) = \hat{\sigma}(\pi_\sigma(\hat{x}))\sigma(\pi_\sigma(\hat{x}), \pi_\sigma(\hat{y}))\hat{\sigma}(\pi_\sigma(\hat{y})).$$

(2) Let $\hat{\psi}$ be a covering a covering isomorphism between $\mathcal{G}_\sigma^{\text{db}}$ et $\mathcal{G}_{\sigma'}^{\text{db}}$. Since $\pi_{\sigma'} = \pi_\sigma \circ \hat{\psi}$, for every $\hat{x} \in V^{\text{db}}$,

- either $(\hat{\psi}(\hat{x}), \hat{\psi}(\psi(\hat{x}))) = (\hat{\psi}(\hat{x}), \hat{\psi}(\psi(\hat{x})))$,
- or $(\hat{\psi}(\hat{x}), \hat{\psi}(\psi(\hat{x}))) = (\hat{\psi}(\psi(\hat{x})), \hat{\psi}(\hat{x}))$.

Note that here \hat{x} and $\psi(\hat{x})$ play symmetric roles. In the first case we set $\hat{\sigma}(\pi_\sigma(\hat{x})) = 1$. In the second case we set $\hat{\sigma}(\pi_\sigma(\hat{x})) = -1$. Then it is easy to check that $\sigma' = \hat{\sigma} \cdot \sigma$. \square

Lemma 3.5. *Let $\sigma \in \{-1, 1\}^E$ be a gauge field and let $\mathcal{G}_\sigma^{\text{db}}$ be the corresponding double cover. Then the graph $\mathcal{G}_\sigma^{\text{db}}$ is connected if and only if σ is non-trivial. Otherwise $\mathcal{G}_\sigma^{\text{db}}$ consists of two disconnected copies of \mathcal{G} .*

Proof. If $\sigma(e) = 1$ for every $e \in E$, then by construction, V_1 and V_2 are not connected in $\mathcal{G}_\sigma^{\text{db}}$ and one has just two copies \mathcal{G} with vertex sets V_1 and V_2 respectively. In the more general case of σ trivial, one can apply Lemma 3.4 to get that $\mathcal{G}_\sigma^{\text{db}}$ is isomorphic as a covering to the previous case.

Assume now that σ is non-trivial. Let $x \in V$, and let $x_1 \in V_1, x_2 \in V_2$ such that $\pi_\sigma(x_1) = \pi_\sigma(x_2) = x$. According to Lemma 2.1, there is a loop \wp rooted in x such that $\text{hol}^\sigma(\wp) = -1$. Let $\hat{\wp}$ be a lift in $\mathcal{G}_\sigma^{\text{db}}$ of the loop \wp . By Lemma 3.3, $\hat{\wp}$ is a path joining x_1 and x_2 . Thus x_1 and x_2 belong to the same connected component. Further, by construction, if $\{x, y\} \in E$, then the two sets $\pi_\sigma^{-1}(\{x\})$ and $\pi_\sigma^{-1}(\{y\})$ are connected to each other. Since \mathcal{G} is connected, then so is $\mathcal{G}_\sigma^{\text{db}}$. \square

Further, we will endow the double covers $\mathcal{G}_\sigma^{\text{db}}$ with conductances as follows. Given $\{\hat{x}, \hat{y}\} \in E_\sigma^{\text{db}}$,

$$(3.5) \quad C(\hat{x}, \hat{y}) = C(\pi_\sigma(\hat{x}), \pi_\sigma(\hat{y})).$$

Let be $V_\partial^{\text{db}} = \pi_\sigma^{-1}(V_\partial)$ and $V_{\text{int}}^{\text{db}} = \pi_\sigma^{-1}(V_{\text{int}})$. We consider the nearest-neighbor Markov jump process on $\mathcal{G}_\sigma^{\text{db}}$ with the transition rates given by the conductances, and killed upon hitting V_∂^{db} . Let $p_\sigma(t, \hat{x}, \hat{y})$ be the transition probabilities of this killed process. The following is immediate.

Lemma 3.6. (1) *For every $\hat{x}, \hat{y} \in V_{\text{int}}^{\text{db}}$,*

$$p_\sigma(t, \hat{x}, \hat{y}) + p_\sigma(t, \hat{x}, \psi(\hat{y})) = p(t, \pi_\sigma(\hat{x}), \pi_\sigma(\hat{y})).$$

(2) *For every $\hat{x}, \hat{y} \in V_{\text{int}}^{\text{db}}$, such that \hat{x} and \hat{y} are both in V_1 or both in V_2 ,*

$$\mathbb{P}_t^{\pi_\sigma(\hat{x}), \pi_\sigma(\hat{y})}(\text{hol}^\sigma(\wp) = -1) = \frac{p_\sigma(t, \hat{x}, \psi(\hat{y}))}{p(t, \pi_\sigma(\hat{x}), \pi_\sigma(\hat{y}))},$$

where $\mathbb{P}_t^{\pi_\sigma(\hat{x}), \pi_\sigma(\hat{y})}$ is the bridge probability measure for the Markov jump process on \mathcal{G} conditioned on staying in V_{int} .

A *fundamental domain* of π_σ is a subset $\mathcal{D} \subset V^{\text{db}}$ such that π_σ induces a bijection from \mathcal{D} to V . By construction, V_1 and V_2 are both fundamental domains. The following is an immediate consequence of (2.5) and Lemma 3.6.

Corollary 3.7. *Let be a gauge field $\sigma \in \{-1, 1\}^E$. Then*

$$\mu^{\text{loop}}(\{\text{Loops } \wp \text{ with } \text{hol}^\sigma(\wp) = -1\}) = \sum_{\hat{x} \in \mathcal{D} \cap V_{\text{int}}^{\text{db}}} \int_0^{+\infty} p_\sigma(t, \hat{x}, \psi(\hat{x})) \frac{dt}{t},$$

where \mathcal{D} is any fundamental domain.

Let be the space $\mathcal{S}_0^{\text{db}}$:

$$\mathcal{S}_0^{\text{db}} = \{f \in \mathbb{R}^{V^{\text{db}}} \mid \forall \hat{x} \in V_\partial^{\text{db}}, f(\hat{x}) = 0\}.$$

We endow $\mathcal{S}_0^{\text{db}}$ with the positive definite inner product

$$\mathcal{E}_\sigma^{\text{db}}(f, g) = \sum_{\{\hat{x}, \hat{y}\} \in E_\sigma^{\text{db}}} C(\hat{x}, \hat{y})(f(\hat{y}) - f(\hat{x}))(g(\hat{y}) - g(\hat{x})).$$

Let $\mathcal{S}_{0,+}^{\text{db}}$ and $\mathcal{S}_{0,-}^{\text{db}}$ be the following subspaces of $\mathcal{S}_0^{\text{db}}$:

$$\mathcal{S}_{0,+}^{\text{db}} = \{f \in \mathcal{S}_0^{\text{db}} \mid f \circ \psi = f\}, \quad \mathcal{S}_{0,-}^{\text{db}} = \{f \in \mathcal{S}_0^{\text{db}} \mid f \circ \psi = -f\}.$$

Note that $\dim \mathcal{S}_{0,+}^{\text{db}} = \dim \mathcal{S}_{0,-}^{\text{db}} = |V_{\text{int}}|$, and $\dim \mathcal{S}_0^{\text{db}} = 2|V_{\text{int}}|$.

Lemma 3.8. *The space $\mathcal{S}_0^{\text{db}}$ is a direct orthogonal (for $\mathcal{E}_\sigma^{\text{db}}$) sum of the subspaces $\mathcal{S}_{0,+}^{\text{db}}$ and $\mathcal{S}_{0,-}^{\text{db}}$.*

Proof. It is clear that $\mathcal{S}_{0,+}^{\text{db}} \cap \mathcal{S}_{0,-}^{\text{db}} = \{0\}$, and every $f \in \mathcal{S}_0^{\text{db}}$ can be decomposed as

$$f = \frac{1}{2}(f + f \circ \psi) + \frac{1}{2}(f - f \circ \psi).$$

So $\mathcal{S}_0^{\text{db}}$ is a direct sum of $\mathcal{S}_{0,+}^{\text{db}}$ and $\mathcal{S}_{0,-}^{\text{db}}$. To check that the decomposition is orthogonal, consider $f \in \mathcal{S}_{0,+}^{\text{db}}$ and $g \in \mathcal{S}_{0,-}^{\text{db}}$. Given $\{\hat{x}, \hat{y}\} \in E_\sigma^{\text{db}}$, then $\{\psi(\hat{x}), \psi(\hat{y})\} \in E_\sigma^{\text{db}}$ and

$$(f(\hat{y}) - f(\hat{x}))(g(\hat{y}) - g(\hat{x})) = -(f(\psi(\hat{y})) - f(\psi(\hat{x})))(g(\psi(\hat{y})) - g(\psi(\hat{x}))).$$

So $\mathcal{E}_\sigma^{\text{db}}(f, g) = 0$. □

Let $\Delta_\sigma^{\text{db}}$ denote the discrete Laplacian on $\mathcal{G}_\sigma^{\text{db}}$:

$$(\Delta_\sigma^{\text{db}} f)(\hat{x}) = \sum_{\substack{\hat{y} \in V^{\text{db}} \\ \{\hat{x}, \hat{y}\} \in E_\sigma^{\text{db}}}} C(\hat{x}, \hat{y})(f(\hat{y}) - f(\hat{x})), \quad \hat{x} \in V^{\text{db}}.$$

One can see $\Delta_\sigma^{\text{db}}$ as an operator on $\mathcal{S}_0^{\text{db}}$ by taking $((\Delta_\sigma^{\text{db}} f)(\hat{x}))_{\hat{x} \in V_{\text{int}}^{\text{db}}}$ and extending to V_∂^{db} by 0. Then, clearly, the subspaces $\mathcal{S}_{0,+}^{\text{db}}$ and $\mathcal{S}_{0,-}^{\text{db}}$ are stable by $\Delta_\sigma^{\text{db}}$. Therefore,

$$\det_{\mathcal{S}_0^{\text{db}}}(-\Delta_\sigma^{\text{db}}) = \det_{\mathcal{S}_{0,+}^{\text{db}}}(-\Delta_\sigma^{\text{db}}) \det_{\mathcal{S}_{0,-}^{\text{db}}}(-\Delta_\sigma^{\text{db}}),$$

where the subscripts indicate the space considered.

Let $f : V^{\text{db}} \rightarrow \mathbb{R}$, and let f_1 denote the following function from V to \mathbb{R} . Given $x \in V$ and $x_1 \in V_1$ such that $\pi_\sigma(x_1) = x$,

$$f_1(x) = f(x_1).$$

If $f \in \mathcal{S}_{0,+}^{\text{db}}$, then

$$(\Delta_\sigma^{\text{db}} f)(x_1) = (\Delta_{\mathcal{G}} f_1)(x),$$

and if $f \in \mathcal{S}_{0,-}^{\text{db}}$, then

$$(\Delta_\sigma^{\text{db}} f)(x_1) = (\Delta_{\mathcal{G}, \sigma} f_1)(x).$$

The following is an immediate consequence of the above.

Corollary 3.9. *We have that*

$$\det_{\mathcal{S}_{0,+}^{\text{db}}}(-\Delta_\sigma^{\text{db}}) = \det((G(x, y))_{x, y \in V_{\text{int}}})^{-1}, \quad \det_{\mathcal{S}_{0,-}^{\text{db}}}(-\Delta_\sigma^{\text{db}}) = \det((G_\sigma(x, y))_{x, y \in V_{\text{int}}})^{-1}.$$

In particular,

$$\frac{\det_{\mathcal{S}_{0,+}^{\text{db}}}(-\Delta_\sigma^{\text{db}})}{\det_{\mathcal{S}_{0,-}^{\text{db}}}(-\Delta_\sigma^{\text{db}})} = \exp(-2\mu^{\text{loop}}(\{\text{Loops } \wp \text{ with } \text{hol}^\sigma(\wp) = -1\})).$$

Let $(G_\sigma^{\text{db}}(\hat{x}, \hat{y}))_{\hat{x}, \hat{y} \in V^{\text{db}}}$ denote the Green's function on $\mathcal{G}_\sigma^{\text{db}}$ with 0 boundary conditions on V_∂^{db} . Note that $G_\sigma^{\text{db}}(\psi(\hat{x}), \psi(\hat{y})) = G_\sigma^{\text{db}}(\hat{x}, \hat{y})$

Lemma 3.10. *Let $x, y \in V$ and let $x_1, y_1 \in V_1$ and $y_2 \in V_2$ such that $\pi_\sigma(x_1) = x$ and $\pi_\sigma(y_1) = \pi_\sigma(y_2) = y$. Then*

$$G(x, y) = G_\sigma^{\text{db}}(x_1, y_1) + G_\sigma^{\text{db}}(x_1, y_2), \quad G_\sigma(x, y) = G_\sigma^{\text{db}}(x_1, y_1) - G_\sigma^{\text{db}}(x_1, y_2),$$

or equivalently

$$G_\sigma^{\text{db}}(x_1, y_1) = \frac{1}{2}(G(x, y) + G_\sigma(x, y)), \quad G_\sigma^{\text{db}}(x_1, y_2) = \frac{1}{2}(G(x, y) - G_\sigma(x, y)).$$

Proof. We have that

$$G_\sigma^{\text{db}}(x_1, y_i) = \int_0^{+\infty} p_\sigma(t, x_1, y_i) dt, \quad G(x, y) = \int_0^{+\infty} p(t, x, y) dt,$$

and according to Theorem 2.2,

$$G_\sigma(x, y) = \int_0^{+\infty} p(t, x, y) \mathbb{E}_t^{x, y}(\text{hol}^\sigma(\varphi) = -1) dt.$$

We conclude by Lemma 3.6. \square

Let $(\phi_\sigma^{\text{db}}(\hat{x}))_{\hat{x} \in V^{\text{db}}}$ be the discrete GFF on the double cover $\mathcal{G}_\sigma^{\text{db}}$. We have that

$$\mathbb{E}[\phi_\sigma^{\text{db}}(\hat{x})] = 0, \quad \mathbb{E}[\phi_\sigma^{\text{db}}(\hat{x})\phi_\sigma^{\text{db}}(\hat{y})] = G_\sigma^{\text{db}}(\hat{x}, \hat{y}).$$

Let $\phi_{\sigma,+}^{\text{db}}$ and $\phi_{\sigma,-}^{\text{db}}$ be the fields

$$\phi_{\sigma,+}^{\text{db}} = \frac{1}{\sqrt{2}}(\phi_\sigma^{\text{db}} + \phi_\sigma^{\text{db}} \circ \psi), \quad \phi_{\sigma,-}^{\text{db}} = \frac{1}{\sqrt{2}}(\phi_\sigma^{\text{db}} - \phi_\sigma^{\text{db}} \circ \psi).$$

The field $\phi_{\sigma,+}^{\text{db}}$, respectively $\phi_{\sigma,-}^{\text{db}}$, is a random element of $\mathcal{S}_{0,+}^{\text{db}}$, respectively $\mathcal{S}_{0,-}^{\text{db}}$. Since $\mathcal{S}_{0,+}^{\text{db}}$ and $\mathcal{S}_{0,-}^{\text{db}}$ are orthogonal for $\mathcal{E}_\sigma^{\text{db}}$, the fields $\phi_{\sigma,+}^{\text{db}}$ and $\phi_{\sigma,-}^{\text{db}}$ are independent. Lemma 3.10 immediately implies the following.

Corollary 3.11. *Let $(\phi(x))_{x \in V}$ and $(\phi_\sigma(x))_{x \in V}$ be the following fields:*

$$\phi(x) = \phi_{\sigma,+}^{\text{db}}(x_1), \quad \phi_\sigma(x) = \phi_{\sigma,-}^{\text{db}}(x_1),$$

where $x_1 \in V_1$ and $\pi_\sigma(x_1) = x$. Then ϕ is distributed as the GFF on \mathcal{G} with 0 boundary conditions (2.1), and ϕ_σ is distributed as the σ -twisted GFF on \mathcal{G} with 0 boundary conditions (2.2).

3.3. Double covers of metric graphs and related GFFs. Consider a gauge field $\sigma \in \{-1, 1\}^E$. Let be the associated double cover $\mathcal{G}_\sigma^{\text{db}}$, endowed with the conductances (3.5). We will consider the metric graph associated to $\mathcal{G}_\sigma^{\text{db}}$, denoted by $\tilde{\mathcal{G}}_\sigma^{\text{db}}$. The projection π_σ can be extended into a covering map $\pi_\sigma : \tilde{\mathcal{G}}_\sigma^{\text{db}} \rightarrow \tilde{\mathcal{G}}$ which is locally an isometry. In this way, $\tilde{\mathcal{G}}_\sigma^{\text{db}}$ is a double cover of the metric graph $\tilde{\mathcal{G}}$. Similarly, the map $\psi : V^{\text{db}} \rightarrow V^{\text{db}}$ extends into a covering automorphism $\psi_\sigma : \tilde{\mathcal{G}}_\sigma^{\text{db}} \rightarrow \tilde{\mathcal{G}}_\sigma^{\text{db}}$ ($\pi_\sigma \circ \psi_\sigma = \pi_\sigma$) which interchanges the two sheets of the covering. Lemma 3.5 extends to the metric graph setting: the metric space $\tilde{\mathcal{G}}_\sigma^{\text{db}}$ is connected if and only if the gauge field σ is non-trivial. Otherwise it consists of two disjoint copies of $\tilde{\mathcal{G}}$.

Let $\tilde{\phi}_\sigma^{\text{db}}$ be the metric graph GFF on $\tilde{\mathcal{G}}_\sigma^{\text{db}}$ with 0 boundary conditions on V_∂^{db} . The field $\tilde{\phi}_\sigma^{\text{db}}$ is continuous. Its restriction to V^{db} is the field ϕ_σ^{db} introduced previously. Let $\tilde{\phi}_{\sigma,+}^{\text{db}}$ and $\tilde{\phi}_{\sigma,-}^{\text{db}}$ be the fields

$$\tilde{\phi}_{\sigma,+}^{\text{db}} = \frac{1}{\sqrt{2}}(\tilde{\phi}_\sigma^{\text{db}} + \tilde{\phi}_\sigma^{\text{db}} \circ \psi_\sigma), \quad \tilde{\phi}_{\sigma,-}^{\text{db}} = \frac{1}{\sqrt{2}}(\tilde{\phi}_\sigma^{\text{db}} - \tilde{\phi}_\sigma^{\text{db}} \circ \psi_\sigma).$$

The fields $\tilde{\phi}_{\sigma,+}^{\text{db}}$ and $\tilde{\phi}_{\sigma,-}^{\text{db}}$ are again continuous and their restrictions to V^{db} are $\phi_{\sigma,+}^{\text{db}}$ and $\phi_{\sigma,-}^{\text{db}}$ respectively.

Lemma 3.12. *The fields $\tilde{\phi}_{\sigma,+}^{\text{db}}$ and $\tilde{\phi}_{\sigma,-}^{\text{db}}$ are independent.*

Proof. We already know that the fields $\phi_{\sigma,+}^{\text{db}}$ and $\phi_{\sigma,-}^{\text{db}}$ are independent. To conclude, we need to look at what happens inside the edge-lines. Given $(W_1(u))_{0 \leq u \leq C(e)-1}$ and $(W_2(u))_{0 \leq u \leq C(e)-1}$ two independent Brownian bridges from 0 to 0 of the same length, then $(W_1 + W_2)/\sqrt{2}$ and $(W_1 - W_2)/\sqrt{2}$ are indeed independent and again distributed as Brownian bridges from 0 to 0. \square

Since $\tilde{\phi}_{\sigma,+}^{\text{db}} \circ \psi_\sigma = \tilde{\phi}_{\sigma,+}^{\text{db}}$, there is a continuous field $\tilde{\phi}$ on $\tilde{\mathcal{G}}$ (i.e. on the base of the covering) such that

$$\tilde{\phi}_{\sigma,+}^{\text{db}} = \tilde{\phi} \circ \pi_\sigma.$$

Lemma 3.13. *The field $\tilde{\phi}$ is distributed as the metric graph GFF on $\tilde{\mathcal{G}}$ with 0 boundary conditions on V_∂ (see Section 2.4), hence the same notation, and in particular its distribution is the same whatever the gauge field σ .*

Proof. By Corollary 3.11 we already know that the restriction of the field $\tilde{\phi}$ to V is distributed as the discrete GFF with 0 boundary conditions on V_∂ . We need to check that the identity in law also extends inside the edge-lines. Given $(W_1(u))_{0 \leq u \leq C(e)-1}$ and $(W_2(u))_{0 \leq u \leq C(e)-1}$ two independent Brownian bridges from 0 to 0 of the same length, then indeed $(W_1 + W_2)/\sqrt{2}$ is distributed as a Brownian bridge from 0 to 0. \square

Next we will introduce a section $\mathbf{s}_\sigma : \tilde{\mathcal{G}} \rightarrow \tilde{\mathcal{G}}^{\text{db}}$, such that $\pi_\sigma \circ \mathbf{s}_\sigma = \text{Id}_{\tilde{\mathcal{G}}}$. In general, \mathbf{s}_σ will not be continuous and will actually have finitely many discontinuity points. The section \mathbf{s}_σ will be defined by the following conditions:

- (1) $\pi_\sigma \circ \mathbf{s}_\sigma = \text{Id}_{\tilde{\mathcal{G}}}$.
- (2) For every $x \in V$, $\mathbf{s}_\sigma(x) \in V_1$.
- (3) For every edge $e \in E$, the map $u \mapsto \mathbf{s}_\sigma(x_{e,u})$ (see Section 3.1 for the notations) is continuous on the intervals $[0, C(e)^{-1}/2)$ and $(C(e)^{-1}/2, C(e)^{-1}]$, i.e. a discontinuity is possible only in the middle of the edge-line at $u = C(e)^{-1}/2$.

The above entirely specifies \mathbf{s}_σ outside the middle points of edge-lines in $\tilde{\mathcal{G}}$. Further, if $\{x, y\} \in E$ and $\sigma(x, y) = 1$, then $\{\mathbf{s}_\sigma(x), \mathbf{s}_\sigma(y)\} \in E_\sigma^{\text{db}}$, and thus \mathbf{s}_σ extends continuously to the whole edge-line $I_{\{x,y\}}$. However, if $\sigma(x, y) = -1$, then $\{\mathbf{s}_\sigma(x), \mathbf{s}_\sigma(y)\} \notin E_\sigma^{\text{db}}$, and thus \mathbf{s}_σ has a discontinuity in the middle of the edge-line $I_{\{x,y\}}$. We will not specify which is the value assumed by \mathbf{s}_σ at such a discontinuity point, i.e. which of the left or right limit, as this will not be of any importance.

Now, consider on $\tilde{\mathcal{G}}$ the field

$$(3.6) \quad \tilde{\phi}_\sigma = \tilde{\phi}_{\sigma,-}^{\text{db}} \circ \mathbf{s}_\sigma.$$

Lemma 3.14. *The field $\tilde{\phi}_\sigma$ has the same distribution as the Gaussian field on $\tilde{\mathcal{G}}$ introduced in Section 3.1, hence the same notation.*

Proof. By Corollary 3.11 we already know that the restriction of the field $\tilde{\phi}_\sigma$ to V is distributed as the discrete σ -twisted GFF ϕ_σ with 0 boundary conditions on V_∂ . We need only to check what happens inside the edge lines. Given $(W_1(u))_{0 \leq u \leq C(e)-1}$ and $(W_2(u))_{0 \leq u \leq C(e)-1}$ two independent Brownian bridges from 0 to 0 of the same length, then $W_e = (W_1 - W_2)/\sqrt{2}$ is again distributed as a Brownian bridge from 0 to 0. If $\sigma(e) = 1$, then \mathbf{s}_σ is continuous on the edge-line I_e and the restriction of $\tilde{\phi}_\sigma$ to I_e is obtained by interpolating between $\tilde{\phi}_\sigma(e_-)$ and $\tilde{\phi}_\sigma(e_+)$ with W_e (plus the deterministic linear part). If $\sigma(e) = -1$, then \mathbf{s}_σ has a discontinuity in the middle of I_e , and the restriction of $\tilde{\phi}_\sigma$ to I_e is obtained by interpolating between $-\tilde{\phi}_\sigma(e_-)$ and $\tilde{\phi}_\sigma(e_+)$ with W_e (plus the deterministic linear part), then by reflecting the bridge on half of I_e . So this is indeed the same construction as the one given by the formulas (3.2), (3.3) and (3.4). \square

The main point of the Sections 3.2 and 3.3 was actually to introduce the field $\tilde{\phi}_{\sigma,-}^{\text{db}}$. The advantage of $\tilde{\phi}_{\sigma,-}^{\text{db}}$ over $\tilde{\phi}_\sigma$ is that the former is always continuous, and in particular satisfies the intermediate value property. This will be exploited in the forthcoming Section 3.4. See Lemma 3.15.

3.4. Topological events for the metric graph GFF. Here we consider a gauge field $\sigma \in \{-1, 1\}^E$. Given U an open non-empty connected subset of the metric graph $\tilde{\mathcal{G}}$, the inverse image $\pi_\sigma^{-1}(U) \subset \tilde{\mathcal{G}}^{\text{db}}$ is either connected or consists of two isometric connected components. It is easy to verify (see e.g. Lemmas 3.3 and 3.5) that $\pi_\sigma^{-1}(U)$ is connected if and only there is a continuous loop \wp contained in U such that $\text{hol}^\sigma(\wp) = -1$. The holonomy of a continuous loop on the metric graph is given by the parity of the number of crossings of edge-lines I_e with $\sigma(e) = -1$.

In the example of Figure 3, $\pi_\sigma^{-1}(U)$ is connected if and only if U surrounds the inner hole of the domain. In the example of Figure 4, $\pi_\sigma^{-1}(U)$ is connected if and only if U separates one hole from the other. By *separate* we mean separate as a subset of the continuum plane, since the planar metric graph on the figure can be embedded into the plane. However, if U surrounds both holes without separating one from the other, then $\pi_\sigma^{-1}(U)$ is not connected.

So whether $\pi_\sigma^{-1}(U)$ is connected or not is a topological, or rather homotopical property of U . It can be reformulated in a more abstract way as follows. Let $\pi_1(\tilde{\mathcal{G}})$ be the fundamental group of the metric graph $\tilde{\mathcal{G}}$ (the notation π_1 should not be confused with the notation π_σ for the covering map; π_1 is just the standard notation for the fundamental group). Let $\pi_1(U)$ be the fundamental group of U . Since U is a subset of $\tilde{\mathcal{G}}$, there is a natural group homomorphism $\theta_U : \pi_1(U) \rightarrow \pi_1(\tilde{\mathcal{G}})$, obtained by considering the loops in U as loops in $\tilde{\mathcal{G}}$ and rooting all the loops at a point $x_0 \in U$. The gauge field σ induces a group homomorphism $h_\sigma : \pi_1(\tilde{\mathcal{G}}) \rightarrow \{-1, 1\}$ obtained by taking the holonomy of loops. The homomorphism h_σ depends only on the gauge equivalence class of σ . The gauge field σ is trivial if and only if $\ker h_\sigma = \pi_1(\tilde{\mathcal{G}})$. Further, $\pi_\sigma^{-1}(U)$ is **not** connected if and only if $\ker h_\sigma \circ \theta_U = \pi_1(U)$. This depends on σ only through its gauge equivalence class, and if σ is trivial, $\pi_\sigma^{-1}(U)$ cannot be connected.

Let $\mathcal{C}(\tilde{\mathcal{G}})$ denote the space of continuous functions $\tilde{\mathcal{G}} \rightarrow \mathbb{R}$. Given a function $f \in \mathcal{C}(\tilde{\mathcal{G}})$, $\{f \neq 0\}$ will be a short notation for the non-zero set of f :

$$\{f \neq 0\} = \{x \in \tilde{\mathcal{G}} \mid f(x) \neq 0\}.$$

Let \mathcal{T}_σ be the following subset of $\mathcal{C}(\tilde{\mathcal{G}})$:

$$\mathcal{T}_\sigma = \{f \in \mathcal{C}(\tilde{\mathcal{G}}) \mid \forall U \text{ connected component of } \{f \neq 0\}, \pi_\sigma^{-1}(U) \text{ is **not** connected}\}.$$

It is easy to see that \mathcal{T}_σ is a closed subset of $\mathcal{C}(\tilde{\mathcal{G}})$ for the uniform norm. In the example of Figure 3, $f \in \mathcal{T}_\sigma$ if and only if no connected component of $\{f \neq 0\}$ surrounds the inner hole of the domain; see Figure 1. In the example of Figure 4, $f \in \mathcal{T}_\sigma$ if and only if no connected component of $\{f \neq 0\}$ separates one hole from the other. Again, \mathcal{T}_σ depends only on the gauge-equivalence class of σ , and if σ is trivial, then $\mathcal{T}_\sigma = \mathcal{C}(\tilde{\mathcal{G}})$.

We recall that, although the field $\tilde{\phi}_\sigma$ is usually not continuous, its absolute value $|\tilde{\phi}_\sigma|$ can always be continuously extended to $\tilde{\mathcal{G}}$. Thus, we see $|\tilde{\phi}_\sigma|$ as a random element of $\mathcal{C}(\tilde{\mathcal{G}})$.

Lemma 3.15. *Almost surely, $|\tilde{\phi}_\sigma| \in \mathcal{T}_\sigma$.*

Proof. We will show that a.s., for every U connected component of $\{|\tilde{\phi}_\sigma| \neq 0\}$, and for every $\hat{x} \in \pi_\sigma^{-1}(U)$, the points \hat{x} and $\psi_\sigma(\hat{x})$ are not connected inside $\pi_\sigma^{-1}(U)$. If this were not the case, there would be a point $\hat{x} \in \tilde{\mathcal{G}}^{\text{db}}$ and a continuous path $(\hat{\varphi}(t))_{0 \leq t \leq 1}$ joining \hat{x} and $\psi_\sigma(\hat{x})$ such that for every $t \in [0, 1]$, $|\tilde{\phi}_\sigma|(\pi_\sigma(\hat{\varphi}(t))) > 0$. Consider on $\tilde{\mathcal{G}}_\sigma^{\text{db}}$ the field $\tilde{\phi}_{\sigma,-}^{\text{db}}$, related to $\tilde{\phi}_\sigma$ through (3.6). Then, for every $t \in [0, 1]$, $\tilde{\phi}_{\sigma,-}^{\text{db}}(\hat{\varphi}(t)) \neq 0$. But the field $\tilde{\phi}_{\sigma,-}^{\text{db}}$ is continuous, and $\tilde{\phi}_{\sigma,-}^{\text{db}}(\hat{\varphi}(0)) = -\tilde{\phi}_{\sigma,-}^{\text{db}}(\hat{\varphi}(1))$. So this is a contradiction. \square

Now consider the usual metric graph GFF $\tilde{\phi}$, without the gauge field σ .

Lemma 3.16. *We have that $\mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) > 0$. Moreover, if σ is non-trivial, $\mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) < 1$.*

Proof. Let us first prove that $\mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) > 0$. Let A be the event that the field $\tilde{\phi}$ has zeroes inside every edge-line I_e for $e \in E$. With the construction with Brownian bridges, it is clear that $\mathbb{P}(A) > 0$. Let us argue that on the event A , we have that $\tilde{\phi} \in \mathcal{T}_\sigma$, whatever the gauge field σ . Indeed, on the event A , no connected component of $\{\tilde{\phi} \neq 0\}$ contains a full edge-line, and thus cannot contain a loop with holonomy -1 .

Assume now that the gauge field σ is non-trivial. Then there is $(\varphi(t))_{0 \leq t \leq 1}$ a continuous (deterministic) loop in $\tilde{\mathcal{G}}$ such that $\text{hol}^\sigma(\varphi) = -1$. With positive probability, $\tilde{\phi}$ has no zeroes on the range of φ . On this event, the range of φ belongs to the same connected component of $\{\tilde{\phi} \neq 0\}$, and because $\text{hol}^\sigma(\varphi) = -1$ we have that $\tilde{\phi} \notin \mathcal{T}_\sigma$. \square




FIGURE 6. A non-contractible chain in an annular domain formed by contractible loops.

We are ready to state the main result of this paper.

Theorem 1. *Let be a gauge field $\sigma \in \{-1, 1\}^E$. Then*

$$(3.7) \quad \mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) = \exp\left(-\mu^{\text{loop}}(\{\text{Loops } \wp \text{ with } \text{hol}^\sigma(\wp) = -1\})\right).$$

So in particular (see Corollaries 2.3, 3.7 and 3.9),

$$\begin{aligned} \mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) &= \frac{\det((G_\sigma(x, y))_{x, y \in V_{\text{int}}})^{1/2}}{\det((G(x, y))_{x, y \in V_{\text{int}}})^{1/2}} \\ &= \frac{\det_{\mathcal{S}_{0,+}^{\text{db}}}(-\Delta_\sigma^{\text{db}})^{1/2}}{\det_{\mathcal{S}_{0,-}^{\text{db}}}(-\Delta_\sigma^{\text{db}})^{1/2}} \\ &= \exp\left(-\sum_{\hat{x} \in \mathcal{D} \cap V_{\text{int}}^{\text{db}}} \int_0^{+\infty} p_\sigma(t, \hat{x}, \psi(\hat{x})) \frac{dt}{t}\right), \end{aligned}$$

where \mathcal{D} is any fundamental domain of the covering map π_σ . Moreover, conditionally on the event $\{\tilde{\phi} \in \mathcal{T}_\sigma\}$, the field $|\tilde{\phi}|$ has the same distribution as $|\tilde{\phi}_\sigma|$.

Remark 3.17. We would like to emphasize that Theorem 1 is not a direct consequence of the isomorphism identity for $\tilde{\phi}$ (Theorem 2.6). Indeed, if $\tilde{\phi} \in \mathcal{T}_\sigma$, then $\tilde{\mathcal{L}}^{1/2}$ cannot contain loops \wp with $\text{hol}^\sigma(\wp) = -1$. However, even if $\tilde{\mathcal{L}}^{1/2}$ does not contain loops \wp with $\text{hol}^\sigma(\wp) = -1$, it can still form clusters \mathcal{C} out of loops \wp with $\text{hol}^\sigma(\wp) = +1$ such that $\pi_\sigma^{-1}(\mathcal{C})$ is connected. See Figure 6. In particular,

$$\mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) \neq \mathbb{P}(\forall \wp \in \tilde{\mathcal{L}}^{1/2}, \text{hol}^\sigma(\wp) = 1).$$

More precisely,

$$\mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) = \mathbb{P}(\forall \wp \in \tilde{\mathcal{L}}^{1/2}, \text{hol}^\sigma(\wp) = 1)^2.$$

However, this difference in exponents can be explained through the isomorphism identity for the gauge-twisted GFF $\tilde{\phi}_\sigma$: see Section 3.5 and the identity (3.16). This factor 2 in the exponent plays a crucial role in our intensity doubling conjecture; see Section 6 and Conjecture 2.

Remark 3.18. Theorem 1 implies in particular that for every $x \in \tilde{\mathcal{G}}$,

$$\mathbb{E}[\tilde{\phi}(x)^2 | \tilde{\phi} \in \mathcal{T}_\sigma] = G_\sigma(x, x).$$

Further, for any $k \in \mathbb{N} \setminus \{0\}$ and $x_1, x_2, \dots, x_k \in \tilde{\mathcal{G}}$, for the conditional moments (products of squares)

$$\mathbb{E}[\tilde{\phi}(x_1)^2 \tilde{\phi}(x_2)^2 \dots \tilde{\phi}(x_k)^2 | \tilde{\phi} \in \mathcal{T}_\sigma]$$

the Wick's formula with kernel G_σ holds.

Remark 3.19. Theorem 1 immediately extends to interacting bosonic fields on the metric graph where the interaction potential depends only on the absolute value of the field. For instance, one can consider the φ^4 fields on $\tilde{\mathcal{G}}$. Let $g > 0$ be a coupling constant and consider the relative partition functions

$$Z_g^{\varphi^4} = \mathbb{E}\left[\exp\left(-g \int_{\tilde{\mathcal{G}}} \tilde{\phi}(x)^4 \tilde{m}(dx)\right)\right], \quad Z_{g,\sigma}^{\varphi^4} = \mathbb{E}\left[\exp\left(-g \int_{\tilde{\mathcal{G}}} \tilde{\phi}_\sigma(x)^4 \tilde{m}(dx)\right)\right].$$

Let $\tilde{\rho}$ be the field on $\tilde{\mathcal{G}}$ with density

$$\frac{1}{Z_g^{\varphi^4}} \exp\left(-g \int_{\tilde{\mathcal{G}}} \tilde{\phi}(x)^4 \tilde{m}(dx)\right)$$

with respect to the GFF $\tilde{\phi}$. This is the φ^4 field on the metric graph. Similarly, let $\tilde{\rho}_\sigma$ be the field on $\tilde{\mathcal{G}}$ with density

$$\frac{1}{Z_{g,\sigma}^{\varphi^4}} \exp\left(-g \int_{\tilde{\mathcal{G}}} \tilde{\phi}_\sigma(x)^4 \tilde{m}(dx)\right)$$

with respect to the GFF $\tilde{\phi}_\sigma$. This is the σ -twisted φ^4 field. Then Theorem 1 implies that

$$(3.8) \quad \mathbb{P}(\tilde{\rho} \in \mathcal{T}_\sigma) = \frac{\det((G_\sigma(x, y))_{x, y \in V_{\text{int}}})^{1/2} Z_{g,\sigma}^{\varphi^4}}{\det((G(x, y))_{x, y \in V_{\text{int}}})^{1/2} Z_g^{\varphi^4}},$$

and that conditionally on $\tilde{\rho} \in \mathcal{T}_\sigma$, the field $|\tilde{\rho}|$ is distributed as $|\tilde{\rho}_\sigma|$. In (3.8) one again gets a ratio of partition functions, this time for the φ^4 field and its σ -twisted version. One can replace the interaction potential φ^4 by any other potential $\mathcal{V}(|\varphi|)$ bounded from below and depending only on the absolute value of the field.

The rest of this section is dedicated to the proof of Theorem 1. Given a gauge field $\sigma \in \{-1, 1\}^E$, we will denote

$$E_{\sigma,-} = \{e \in E | \sigma(e) = -1\}.$$

The proof will be done under the following additional assumption.

Assumption 1. *Every connected component of the subgraph $(V, E \setminus E_{\sigma,-})$ of \mathcal{G} intersects V_∂ .*

The above assumption is needed because in the proof we will consider GFFs on subgraphs of \mathcal{G} (and sub-metric-graphs of $\tilde{\mathcal{G}}$), and thus each connected component of the subgraph needs to be connected to the boundary V_∂ in order to pin the corresponding field. The above assumption is however not restrictive at all. Indeed, if a couple of graph and gauge field (\mathcal{G}, σ) does not satisfy Assumption 1, then one can enlarge \mathcal{G} by adding an extra site \dagger to V_∂ and connect \dagger to every vertex in $x \in V_{\text{int}}$ by an edge $\{x, \dagger\}$, with a conductance $C(x, \dagger) = \delta > 0$; and extend the gauge field σ with $\sigma(x, \dagger) = 1$ for every $x \in V_{\text{int}}$. Then the extended graph and gauge field now satisfy Assumption 1. Further, if one has Theorem 1 for the extended graph and gauge field, one gets Theorem 1 for the initial graph and gauge field by letting $\delta \rightarrow 0$ (recall that δ is the conductance to the extra boundary vertex \dagger). In essence, adding \dagger is equivalent to considering massive GFFs with a small square-mass δ .

Let be

$$\underline{r} = \min_{e \in E} C(e)^{-1}.$$

Given $e \in E$, x_e^{m} will denote the middle point of the edge-line I_e :

$$x_e^{\text{m}} = x_{e, C(e)^{-1/2}}.$$




FIGURE 7. Inside an edge-line.

For $\varepsilon \leq C(e)^{-1}/2$, we will denote by $x_{e,\varepsilon}^-$ and $x_{e,\varepsilon}^+$ the two points of I_e at distance ε from x_e^m :

$$x_{e,\varepsilon}^- = x_{e,C(e)^{-1}/2+\varepsilon}, \quad x_{e,\varepsilon}^+ = x_{e,C(e)^{-1}/2-\varepsilon}.$$

The points $x_{e,\varepsilon}^-$ and $x_{e,\varepsilon}^+$ will play symmetric roles, the distinction is just for the notations. Let $J_{e,\varepsilon}$ denote the open subinterval of I_e delimited by $x_{e,\varepsilon}^-$ and $x_{e,\varepsilon}^+$:

$$J_{e,\varepsilon} = \{x_{e,u} | C(e)^{-1}/2 - \varepsilon < u < C(e)^{-1}/2 + \varepsilon\}.$$

See Figure 7.

For $\varepsilon \in (0, \underline{r}/2)$, let $\tilde{\mathcal{G}}_{\sigma,\varepsilon}$ denote

$$\tilde{\mathcal{G}}_{\sigma,\varepsilon} = \tilde{\mathcal{G}} \setminus \bigcup_{e \in E_{\sigma,-}} J_{e,\varepsilon}.$$

In this way, $\tilde{\mathcal{G}}_{\sigma,\varepsilon}$ is a closed subset of the metric graph $\tilde{\mathcal{G}}$, and is itself a metric graph, but not necessarily connected. Consider the discrete graph $\mathcal{G}_{\sigma,\varepsilon} = (V_{\sigma,\varepsilon}, E_{\sigma,\varepsilon})$ constructed as follows. The set of vertices is given by

$$V_{\sigma,\varepsilon} = V \cup \{x_{e,\varepsilon}^- | e \in E_{\sigma,-}\} \cup \{x_{e,\varepsilon}^+ | e \in E_{\sigma,-}\},$$

and the set of edges given by

$$E_{\sigma,\varepsilon} = (E \setminus E_{\sigma,-}) \cup \{\{x_{e,\varepsilon}^-, e_-\} | e \in E_{\sigma,-}\} \cup \{\{x_{e,\varepsilon}^+, e_+\} | e \in E_{\sigma,-}\}.$$

The graph $\mathcal{G}_{\sigma,\varepsilon}$ is endowed with the following conductances: for $e \in E \setminus E_{\sigma,-}$, we keep the same conductance $C(e)$. Further, for $e \in E_{\sigma,-}$, we set

$$C(x_{e,\varepsilon}^-, e_-) = C(x_{e,\varepsilon}^+, e_+) = (C(e)^{-1}/2 - \varepsilon)^{-1}.$$

Then $\tilde{\mathcal{G}}_{\sigma,\varepsilon}$ is actually the metric graph associated to the electrical network $\mathcal{G}_{\sigma,\varepsilon}$. We will denote by $\tilde{\phi}_{\sigma,\varepsilon}$ the metric graph GFF on $\tilde{\mathcal{G}}_{\sigma,\varepsilon}$ with 0 boundary conditions on V_∂ . Assumption 1 ensures that $\tilde{\phi}_{\sigma,\varepsilon}$ is well defined. The field $\tilde{\phi}_{\sigma,\varepsilon}$ is a usual metric graph GFF, i.e. **not** gauge-twisted. We will also consider the restrictions of the metric graph GFFs $\tilde{\phi}$ and $\tilde{\phi}_\sigma$ to $\tilde{\mathcal{G}}_{\sigma,\varepsilon}$.

For $x, y \in \tilde{\mathcal{G}}$, we will denote

$$G(x, y) = \mathbb{E}[\tilde{\phi}(x), \tilde{\phi}(y)], \quad G_\sigma(x, y) = \mathbb{E}[\tilde{\phi}_\sigma(x), \tilde{\phi}_\sigma(y)].$$

In this way we have a natural extension of Green's functions G and G_σ beyond $V \times V$. note that G defined in this way is continuous on $\tilde{\mathcal{G}} \times \tilde{\mathcal{G}}$, but G_σ has discontinuities at the middle points x_e^m for $e \in E_{\sigma,-}$. We will also denote by $G_{\sigma,\varepsilon}(x, y)$ the covariance kernel of $\tilde{\phi}_{\sigma,\varepsilon}$ on $\tilde{\mathcal{G}}_{\sigma,\varepsilon} \times \tilde{\mathcal{G}}_{\sigma,\varepsilon}$.

Lemma 3.20. *The restrictions $\tilde{\phi}|_{\tilde{\mathcal{G}}_{\sigma,\varepsilon}}$ and $\tilde{\phi}_\sigma|_{\tilde{\mathcal{G}}_{\sigma,\varepsilon}}$ are both absolutely continuous with respect to $\tilde{\phi}_{\sigma,\varepsilon}$. The Radon-Nikodym derivatives are given by*

$$(3.9) \quad \frac{d\mathbb{P}_{\tilde{\phi}|_{\tilde{\mathcal{G}}_{\sigma,\varepsilon}}}(\tilde{\varphi})}{d\mathbb{P}_{\tilde{\phi}_{\sigma,\varepsilon}}}(\tilde{\varphi}) = \frac{(\det(G_{\sigma,\varepsilon}(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}} \exp\left(-\frac{1}{4\varepsilon} \sum_{e \in E_{\sigma,-}} (\tilde{\varphi}(x_{e,\varepsilon}^+) - \tilde{\varphi}(x_{e,\varepsilon}^-))^2\right),$$

$$(3.10) \quad \frac{d\mathbb{P}_{\tilde{\phi}_\sigma|_{\tilde{\mathcal{G}}_{\sigma,\varepsilon}}}(\tilde{\varphi})}{d\mathbb{P}_{\tilde{\phi}_{\sigma,\varepsilon}}}(\tilde{\varphi}) = \frac{(\det(G_{\sigma,\varepsilon}(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G_\sigma(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}} \exp\left(-\frac{1}{4\varepsilon} \sum_{e \in E_{\sigma,-}} (\tilde{\varphi}(x_{e,\varepsilon}^+) + \tilde{\varphi}(x_{e,\varepsilon}^-))^2\right).$$

Proof. Let us introduce the auxiliary electrical network $\mathcal{G}_{\sigma,\varepsilon}^*$ as follows. Its set of vertices is $V_{\sigma,\varepsilon}$, the same as for $\mathcal{G}_{\sigma,\varepsilon}$. Its set of edges is

$$E_{\sigma,\varepsilon}^* = E_{\sigma,\varepsilon} \cup \{\{x_{e,\varepsilon}^-, x_{e,\varepsilon}^+\} | e \in E_{\sigma,-}\}.$$

The conductances on $E_{\sigma,\varepsilon}^*$ are as follows. If $e \in E_{\sigma,\varepsilon}$, then we keep the same conductance $C(e)$. Moreover, we set

$$C(x_{e,\varepsilon}^-, x_{e,\varepsilon}^+) = \frac{1}{2\varepsilon},$$

for $e \in E_{\sigma,-}$. Then the metric graph associated to the electrical network $\mathcal{G}_{\sigma,\varepsilon}^*$ is again $\tilde{\mathcal{G}}$, the same as for \mathcal{G} . This is because for every $e \in E_{\sigma,-}$,

$$\begin{aligned} C(e_-, x_{e,\varepsilon}^-)^{-1} + C(x_{e,\varepsilon}^-, x_{e,\varepsilon}^+)^{-1} + C(x_{e,\varepsilon}^+, e_+)^{-1} \\ = (C(e)^{-1}/2 - \varepsilon) + 2\varepsilon + (C(e)^{-1}/2 - \varepsilon) = C(e)^{-1}, \end{aligned}$$

as for resistors in series. Therefore, the restriction of $\tilde{\phi}$ to $V_{\sigma,\varepsilon}$ is the discrete GFF on the electrical network $\mathcal{G}_{\sigma,\varepsilon}^*$, with 0 boundary conditions on V_∂ . The right-hand side of (3.9) is then the density of $(\tilde{\phi}(x))_{x \in V_{\sigma,\varepsilon}}$ with respect to $(\tilde{\phi}_{\sigma,\varepsilon}(x))_{x \in V_{\sigma,\varepsilon}}$, both being discrete GFFs, on $\mathcal{G}_{\sigma,\varepsilon}^*$ and $\mathcal{G}_{\sigma,\varepsilon}$ respectively. Indeed, compared to $\mathcal{G}_{\sigma,\varepsilon}$, $\mathcal{G}_{\sigma,\varepsilon}^*$ contains only the extra edges $\{x_{e,\varepsilon}^-, x_{e,\varepsilon}^+\}$ for $e \in E_{\sigma,-}$, with conductance $1/(2\varepsilon)$ each, so the corresponding factors appear in the density. The ratio of determinants of Green's functions to the power $1/2$ is the normalization factor. Finally, the expression of the density is the same when we consider the whole metric graph $\tilde{\mathcal{G}}_{\sigma,\varepsilon}$, because both for $\tilde{\phi}$ and $\tilde{\phi}_{\sigma,\varepsilon}$ we interpolate with the same independent Brownian bridges.

Now let us deal with $\tilde{\phi}_\sigma$. We consider on $\mathcal{G}_{\sigma,\varepsilon}^*$ the gauge-field $\sigma^* \in \{-1, 1\}^{E_{\sigma,\varepsilon}^*}$ defined as follows. For $e \in E \setminus E_{\sigma,-}$, $\sigma^*(e) = \sigma(e) = 1$. Further,

$$\sigma^*(e_-, x_{e,\varepsilon}^-) = \sigma^*(e_+, x_{e,\varepsilon}^+) = 1,$$

and

$$\sigma^*(x_{e,\varepsilon}^-, x_{e,\varepsilon}^+) = -1.$$

In this way, for every \wp nearest-neighbor path in $\mathcal{G}_{\sigma,\varepsilon}^*$ joining two points, $x, y \in V$,

$$\text{hol}^{\sigma^*}(\wp) = \text{hol}^\sigma(\wp_{\mathcal{G}}),$$

where $\wp_{\mathcal{G}}$ is the nearest-neighbor path in \mathcal{G} from x to y obtained by taking the trace of \wp on V . The restriction of $\tilde{\phi}_\sigma$ to $V_{\sigma,\varepsilon}$ is the σ^* -twisted GFF on $\mathcal{G}_{\sigma,\varepsilon}^*$ with 0 boundary conditions on V_∂ . This is similar to Proposition 3.2. Then (3.10) is just the density of $(\tilde{\phi}_\sigma(x))_{x \in V_{\sigma,\varepsilon}}$ with respect to $(\tilde{\phi}_{\sigma,\varepsilon}(x))_{x \in V_{\sigma,\varepsilon}}$. Because of σ^* , there is a $+$ instead of a $-$ in (3.10) compared to (3.9). \square

Note that Lemma 3.20 implies in particular that $\tilde{\phi}_\sigma|_{\tilde{\mathcal{G}}_{\sigma,\varepsilon}}$ is absolutely continuous with respect to $\tilde{\phi}_\sigma|_{\tilde{\mathcal{G}}_{\sigma,\varepsilon}}$. However, if one considers the whole metric graph $\tilde{\mathcal{G}}$, then $\tilde{\phi}_\sigma$ is singular with respect to $\tilde{\phi}$, unless $E_{\sigma,-} = \emptyset$. For instance, $\tilde{\phi}_\sigma$ is a.s. discontinuous at the middle points x_e^m for $e \in E_{\sigma,-}$. So that is the reason why we restrict to ε -far away from these middle points.

Let $\mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon})$ denote the space of continuous real-valued functions on $\tilde{\mathcal{G}}_{\sigma,\varepsilon}$. For $f \in \mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon})$, $\{f \neq 0\}$ will denote the non-zero set of f , with $\{f \neq 0\} \subset \tilde{\mathcal{G}}_{\sigma,\varepsilon}$. Let $\mathcal{T}_{\sigma,\varepsilon}$ be the following subset of $\mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon})$:

$$\mathcal{T}_{\sigma,\varepsilon} = \left\{ f \in \mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon}) \mid \forall U \text{ connected component of } \left(\{f \neq 0\} \bigcup_{e \in E_{\sigma,-}} J_{e,\varepsilon} \right), \pi_\sigma^{-1}(U) \text{ is **not** connected} \right\}.$$

In the above definition, we enlarge $\{f \neq 0\}$ with the intervals $J_{e,\varepsilon}$ for $e \in E_{\sigma,-}$, and then consider the connected components.

Given $f \in \mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon})$, we will define an equivalence relation \sim_f on the set

$$(3.11) \quad \{x_{e,\varepsilon}^- | e \in E_{\sigma,-}\} \cup \{x_{e,\varepsilon}^+ | e \in E_{\sigma,-}\} = V_{\sigma,\varepsilon} \setminus V.$$

Two points x, x' in the above set are in the same class if they are in the same connected component of $\{f \neq 0\}$. If $f(x) = 0$, then x is alone in its class. We will denote by V_f the set (3.11) quotiented by the equivalence relation \sim_f , and by $[x]_f$ the equivalence class of a point x . Next we will introduce an undirected multigraph Γ_f induced by the function f . Note that in general Γ_f is not connected and may have self-loops and multiple edges. The set of vertices of Γ_f is the quotient set V_f . Further, for each $e \in E_{\sigma,-}$, add to the multigraph Γ_f an edge with the ends $[x_{e,\varepsilon}^-]_f$ and $[x_{e,\varepsilon}^+]_f$. So the number of edges of Γ_f is $|E_{\sigma,-}|$.

Lemma 3.21. *Let $f \in \mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon})$. Then the multigraph Γ_f is bipartite, i.e. does not contain cycles with odd number of edges, if and only if $f \in \mathcal{T}_{\sigma,\varepsilon}$.*

Proof. Given a continuous loop φ in the subset

$$(3.12) \quad \left(\{f \neq 0\} \bigcup_{e \in E_{\sigma,-}} J_{e,\varepsilon} \right),$$

it induces a nearest-neighbor loop in the multigraph Γ_f , which we will denote by $[\varphi]_f$ by a slight abuse of notation. If φ does not visit any point in $V_{\sigma,\varepsilon} \setminus V$ (3.11), then $[\varphi]_f$ is just the empty loop. Conversely, every nearest-neighbor loop in Γ_f can be lifted to a continuous loop in (3.12). This follows from the way Γ_f has been constructed. The edges visited by $[\varphi]_f$ correspond to the intervals $J_{e,\varepsilon}$ (with $e \in E_{\sigma,-}$) crossed by φ from one end to the opposite. The holonomy $\text{hol}^\sigma(\varphi)$ is given by the parity of the number of edge-lines I_e , for $e \in E_{\sigma,-}$, crossed by φ . This parity is the same as for the crossings of $J_{e,\varepsilon}$, for $e \in E_{\sigma,-}$, although the number of crossings itself might be different. This is because each time φ crosses an edge-line I_e , it will cross the corresponding subinterval $J_{e,\varepsilon}$ an odd number of times. Thus, $\text{hol}^\sigma(\varphi)$ is also given by the parity of the number of edges of multigraph loop $[\varphi]_f$. Thus, Γ_f is bipartite if and only if the subset (3.12) does not contain loops φ with $\text{hol}^\sigma(\varphi) = -1$, which is the same as $f \in \mathcal{T}_{\sigma,\varepsilon}$. \square

Note that although $\mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon})$ is infinite, the number of different partitions of $V_{\sigma,\varepsilon} \setminus V$ (3.11) induced by the functions f can be only finite. Given $f \in \mathcal{T}_{\sigma,\varepsilon}$, one can choose a ‘‘coloring’’ $\mathbf{b}_f : V_{\sigma,\varepsilon} \setminus V \rightarrow \{-1, 1\}$ satisfying the following two properties.

- (1) The value of \mathbf{b}_f is the same across every equivalence class for \sim_f .
- (2) For every $e \in E_{\sigma,-}$, $\mathbf{b}_f(x_{e,\varepsilon}^-) = -\mathbf{b}_f(x_{e,\varepsilon}^+)$.

The existence of such \mathbf{b}_f is ensured by the fact that Γ_f is bipartite. Moreover, one can choose \mathbf{b}_f to depend only on the partition V_f induced by \sim_f .

Next we define a transformation $\mathbf{T}_{\sigma,\varepsilon}$ acting on $\mathcal{T}_{\sigma,\varepsilon}$. Given $f \in \mathcal{T}_{\sigma,\varepsilon}$, the function $\mathbf{T}_{\sigma,\varepsilon}f$ is defined as follows.

- (1) For every $x \in \tilde{\mathcal{G}}_{\sigma,\varepsilon}$ such that $f(x) = 0$, we also have $(\mathbf{T}_{\sigma,\varepsilon}f)(x) = 0$.
- (2) On every connected component U of $\{f \neq 0\}$ such that U does not intersect the set $V_{\sigma,\varepsilon} \setminus V$, we have $(\mathbf{T}_{\sigma,\varepsilon}f)|_U = f|_U$.
- (3) On every connected component U of $\{f \neq 0\}$ intersecting $V_{\sigma,\varepsilon} \setminus V$, we have $(\mathbf{T}_{\sigma,\varepsilon}f)|_U = \mathbf{b}_f(U)f|_U$, where $\mathbf{b}_f(U)$ is the common value of \mathbf{b}_f on U .

So, in essence, $\mathbf{T}_{\sigma,\varepsilon}f$ is obtained by flipping the signs on some connected components of $\{f \neq 0\}$. In this way, $\mathbf{T}_{\sigma,\varepsilon}f$ is continuous too, and $|\mathbf{T}_{\sigma,\varepsilon}f| \equiv f$. Therefore, $\mathbf{T}_{\sigma,\varepsilon}f \in \mathcal{T}_{\sigma,\varepsilon}$ too. Moreover, the equivalence relation $\sim_{\mathbf{T}_{\sigma,\varepsilon}f}$ is the same as \sim_f . Thus, $\mathbf{T}_{\sigma,\varepsilon}^2 f = \mathbf{T}_{\sigma,\varepsilon}(\mathbf{T}_{\sigma,\varepsilon}f) = f$.

Further, we will extend $\mathbf{T}_{\sigma,\varepsilon}$ to the whole $\mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon})$. For $f \in \mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon}) \setminus \mathcal{T}_{\sigma,\varepsilon}$, we set $\mathbf{T}_{\sigma,\varepsilon}f = f$.

Let us denote by $\tilde{\xi}_\varepsilon$ and $\tilde{\eta}_\varepsilon$ the fields

$$\tilde{\xi}_\varepsilon = \mathbf{T}_{\sigma,\varepsilon}(\tilde{\phi}_{|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}), \quad \tilde{\eta}_\varepsilon = \mathbf{T}_{\sigma,\varepsilon}(\tilde{\phi}_{\sigma,\varepsilon}).$$

Lemma 3.22. *The field $\tilde{\xi}_\varepsilon$ is absolutely continuous with respect to $\tilde{\eta}_\varepsilon$, and the corresponding density is*

$$\frac{d\mathbb{P}_{\tilde{\xi}_\varepsilon}}{d\mathbb{P}_{\tilde{\eta}_\varepsilon}}(\tilde{\varphi}) = \frac{(\det(G_{\sigma,\varepsilon}(x, y))_{x, y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G(x, y))_{x, y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}} \exp\left(-\frac{1}{4\varepsilon} \sum_{e \in E_{\sigma,-}} (\tilde{\varphi}(x_{e,\varepsilon}^+) - (-1)^{\mathbf{1}_{\tilde{\varphi} \in \mathcal{T}_{\sigma,\varepsilon}}} \tilde{\varphi}(x_{e,\varepsilon}^-))^2\right).$$

Proof. Since $\tilde{\phi}_{|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}$ is absolutely continuous with respect to $\tilde{\phi}_{\sigma,\varepsilon}$ (Lemma 3.20) and $\mathbf{T}_{\sigma,\varepsilon}$ is a deterministic and measurable transformation, we get that $\tilde{\xi}_\varepsilon$ is absolutely continuous with respect to $\tilde{\eta}_\varepsilon$, with

$$\frac{d\mathbb{P}_{\tilde{\xi}_\varepsilon}}{d\mathbb{P}_{\tilde{\eta}_\varepsilon}}(\tilde{\varphi}) = \frac{d\mathbb{P}_{\tilde{\phi}_{|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}}}{d\mathbb{P}_{\tilde{\phi}_{\sigma,\varepsilon}}}(\mathbf{T}_{\sigma,\varepsilon}\tilde{\varphi}).$$

Further we use the expression (3.9). Take $e \in E_{\sigma,-}$. For $\tilde{\varphi} \notin \mathcal{T}_{\sigma,\varepsilon}$,

$$((\mathbf{T}_{\sigma,\varepsilon}\tilde{\varphi})(x_{e,\varepsilon}^+) - (\mathbf{T}_{\sigma,\varepsilon}\tilde{\varphi})(x_{e,\varepsilon}^-))^2 = (\tilde{\varphi}(x_{e,\varepsilon}^+) - \tilde{\varphi}(x_{e,\varepsilon}^-))^2.$$

For $\tilde{\varphi} \in \mathcal{T}_{\sigma,\varepsilon}$,

$$((\mathbf{T}_{\sigma,\varepsilon}\tilde{\varphi})(x_{e,\varepsilon}^+) - (\mathbf{T}_{\sigma,\varepsilon}\tilde{\varphi})(x_{e,\varepsilon}^-))^2 = (\tilde{\varphi}(x_{e,\varepsilon}^+) + \tilde{\varphi}(x_{e,\varepsilon}^-))^2.$$

This because $\mathbf{b}_f(x_{e,\varepsilon}^-)\mathbf{b}_f(x_{e,\varepsilon}^+) = -1$. \square

Lemma 3.23. *The field $\tilde{\eta}_\varepsilon = \mathbf{T}_{\sigma,\varepsilon}(\tilde{\phi}_{\sigma,\varepsilon})$ has the same distribution as $\tilde{\phi}_{\sigma,\varepsilon}$.*

Proof. First, by construction, $|\mathbf{T}_{\sigma,\varepsilon}(\tilde{\phi}_{\sigma,\varepsilon})| \equiv |\tilde{\phi}_{\sigma,\varepsilon}|$. Further, the signs $\mathbf{b}_{\tilde{\phi}_{\sigma,\varepsilon}}$ (that is to say \mathbf{b}_f with $f = \tilde{\phi}_{\sigma,\varepsilon}$) are by construction measurable with respect to $|\tilde{\phi}_{\sigma,\varepsilon}|$. Then, we use the fact that the signs of $\tilde{\phi}_{\sigma,\varepsilon}$ (do not confuse them with $\mathbf{b}_{\tilde{\phi}_{\sigma,\varepsilon}}$) are distributed, conditionally on $|\tilde{\phi}_{\sigma,\varepsilon}|$, as independent uniform $\{-1, 1\}$ -valued r.v.s, one for each connected component of $\{\tilde{\phi}_{\sigma,\varepsilon} \neq 0\}$; see [Lup16, Lemma 3.2]. This implies that the product of signs $\mathbf{b}_{\tilde{\phi}_{\sigma,\varepsilon}} \text{sign}(\tilde{\phi}_{\sigma,\varepsilon})$ has the same conditional distribution given $|\tilde{\phi}_{\sigma,\varepsilon}|$ as $\text{sign}(\tilde{\phi}_{\sigma,\varepsilon})$. Therefore, $\mathbf{T}_{\sigma,\varepsilon}(\tilde{\phi}_{\sigma,\varepsilon})$ has the same distribution as $\tilde{\phi}_{\sigma,\varepsilon}$. \square

Lemma 3.24. *The function*

$$\varepsilon \mapsto \frac{(\det(G_\sigma(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}$$

is constant on $(0, \underline{r}/2)$. More precisely, for every $\varepsilon \in (0, \underline{r}/2)$,

$$\begin{aligned} \frac{(\det(G_\sigma(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}} &= \frac{(\det(G_\sigma(x, y))_{x,y \in V_{\text{int}}})^{1/2}}{(\det(G(x, y))_{x,y \in V_{\text{int}}})^{1/2}} \\ &= \exp(-\mu^{\text{loop}}(\{\text{Loops } \varphi \text{ with } \text{hol}^\sigma(\varphi) = -1\})). \end{aligned}$$

Proof. Consider the measure $\tilde{\mu}^{\text{loop}}$ (2.10) on Brownian loops on the metric graph $\tilde{\mathcal{G}}$. Similarly to Corollary (2.3), we have that

$$\frac{(\det(G_\sigma(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}} = \exp(-\tilde{\mu}^{\text{loop}}(\{\text{Loops } \varphi \text{ visiting } V_{\sigma,\varepsilon} \setminus V \text{ and with } \text{hol}^\sigma(\varphi) = -1\})).$$

Moreover,

$$\begin{aligned} \frac{\det((G_\sigma(x, y))_{x,y \in V_{\text{int}}})^{1/2}}{\det((G(x, y))_{x,y \in V_{\text{int}}})^{1/2}} &= \exp(-\mu^{\text{loop}}(\{\text{Loops } \varphi \text{ with } \text{hol}^\sigma(\varphi) = -1\})) \\ &= \exp(-\tilde{\mu}^{\text{loop}}(\{\text{Loops } \varphi \text{ visiting } V \text{ and with } \text{hol}^\sigma(\varphi) = -1\})), \end{aligned}$$

where the second equality is due to the fact that the measure on discrete loops μ^{loop} can be obtained by taking the trace on V of metric graph loops under $\tilde{\mu}^{\text{loop}}$. We conclude by using the fact that any continuous loop φ on $\tilde{\mathcal{G}}$ with $\text{hol}^\sigma(\varphi) = -1$ has to visit both V and $V_{\sigma,\varepsilon} \setminus V$. That is to say a loop not visiting either of two subsets has holonomy 1, because then it cannot cross any of the edge-lines I_e for $e \in E_{\sigma,-}$. Therefore,

$$\begin{aligned} \tilde{\mu}^{\text{loop}}(\{\text{Loops } \varphi \text{ visiting } V_{\sigma,\varepsilon} \setminus V \text{ and with } \text{hol}^\sigma(\varphi) = -1\}) &= \tilde{\mu}^{\text{loop}}(\{\text{Loops } \varphi \text{ with } \text{hol}^\sigma(\varphi) = -1\}) \\ &= \tilde{\mu}^{\text{loop}}(\{\text{Loops } \varphi \text{ visiting } V \text{ and with } \text{hol}^\sigma(\varphi) = -1\}) = \mu^{\text{loop}}(\{\text{Loops } \varphi \text{ with } \text{hol}^\sigma(\varphi) = -1\}). \quad \square \end{aligned}$$

Proof of Theorem 1. By Lemma 3.20, the field $\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}$ is absolutely continuous with respect to the field $\tilde{\phi}_{\sigma,\varepsilon}$. By Lemma 3.23, the field $\tilde{\eta}_\varepsilon$ has the same distribution as $\tilde{\phi}_{\sigma,\varepsilon}$. By Lemma 3.22, the fields $\tilde{\eta}_\varepsilon$ and $\tilde{\xi}_\varepsilon$ are mutually absolutely continuous, the corresponding Radon–Nikodym derivative being positive on $\mathcal{C}(\tilde{\mathcal{G}}_{\sigma,\varepsilon})$. Therefore, the field $\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}$ is absolutely continuous with respect to the field $\tilde{\xi}_\varepsilon$ and the Radon–Nikodym derivative is given by

$$(3.13) \quad \frac{d\mathbb{P}_{\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}}}{d\mathbb{P}_{\tilde{\xi}_\varepsilon}}(\tilde{\varphi}) = \frac{(\det(G(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G_\sigma(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}} \left(\mathbf{1}_{\tilde{\varphi} \in \mathcal{T}_{\sigma,\varepsilon}} + \mathbf{1}_{\tilde{\varphi} \notin \mathcal{T}_{\sigma,\varepsilon}} \exp \left(-\frac{1}{\varepsilon} \sum_{e \in E_{\sigma,-}} \tilde{\varphi}(x_{e,\varepsilon}^+) \tilde{\varphi}(x_{e,\varepsilon}^-) \right) \right).$$

Note that $\tilde{\xi}_\varepsilon \in \mathcal{T}_{\sigma,\varepsilon}$ if and only if $\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}} \in \mathcal{T}_{\sigma,\varepsilon}$. So in particular,

$$\begin{aligned} \mathbb{P}(\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}} \in \mathcal{T}_{\sigma,\varepsilon}) + \mathbb{E} \left[\mathbf{1}_{\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}} \notin \mathcal{T}_{\sigma,\varepsilon}} \exp \left(-\frac{1}{\varepsilon} \sum_{e \in E_{\sigma,-}} \tilde{\phi}(x_{e,\varepsilon}^+) \tilde{\phi}(x_{e,\varepsilon}^-) \right) \right] \\ = \frac{(\det(G_\sigma(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}} = \exp \left(-\mu^{\text{loop}}(\{\text{Loops } \wp \text{ with } \text{hol}^\sigma(\wp) = -1\}) \right), \end{aligned}$$

where the second equality is due to Lemma 3.24. Further,

$$\lim_{\varepsilon \rightarrow 0} \mathbb{P}(\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}} \in \mathcal{T}_{\sigma,\varepsilon}) = \mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma)$$

and

$$\mathbb{E} \left[\mathbf{1}_{\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}} \notin \mathcal{T}_{\sigma,\varepsilon}} \exp \left(-\frac{1}{\varepsilon} \sum_{e \in E_{\sigma,-}} \tilde{\phi}(x_{e,\varepsilon}^+) \tilde{\phi}(x_{e,\varepsilon}^-) \right) \right] = \frac{(\det(G_\sigma(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}}{(\det(G(x, y))_{x,y \in V_{\sigma,\varepsilon} \setminus V})^{1/2}} \mathbb{P}(|\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}| \notin \mathcal{T}_{\sigma,\varepsilon}),$$

with

$$\lim_{\varepsilon \rightarrow 0} \mathbb{P}(|\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}| \notin \mathcal{T}_{\sigma,\varepsilon}) = \mathbb{P}(|\tilde{\phi}_\sigma| \notin \mathcal{T}_\sigma) = 0;$$

see Lemma 3.15. So we get (3.7).

Since $|\tilde{\xi}_\varepsilon| = |\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}|$, the field $|\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}|$ is absolutely continuous with respect to $|\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}|$, and

$$\lim_{\varepsilon \rightarrow 0} \frac{d\mathbb{P}_{|\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}|}}{d\mathbb{P}_{|\tilde{\phi}_{\sigma|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}|}}(\tilde{\varphi}) = \frac{1}{\mathbb{P}(\tilde{\varphi} \in \mathcal{T}_\sigma)} \mathbf{1}_{\tilde{\varphi} \in \mathcal{T}_\sigma},$$

with convergence in L^1 . So $\mathbb{P}(\tilde{\varphi} \in \mathcal{T}_\sigma)^{-1} \mathbf{1}_{\tilde{\varphi} \in \mathcal{T}_\sigma}$ is the density of $|\tilde{\phi}_\sigma|$ with respect to $|\tilde{\phi}|$. \square

Next we explain how to get the field $\tilde{\phi}_\sigma$ given the field $\tilde{\phi}$ conditioned on $\tilde{\phi} \in \mathcal{T}_\sigma$, that is to say how to get the signs, not just the absolute value. In essence, one has to flip the signs on some of the connected components of $\{\tilde{\phi} \neq 0\} \setminus \{x_e^m | e \in E_{\sigma,-}\}$.

To avoid trivialities, we assume that the gauge field σ is not uniformly 1. Let us denote

$$\mathfrak{M}_\sigma = \{x_e^m | e \in E_{\sigma,-}\}.$$

We endow the subset $\tilde{\mathcal{G}} \setminus \mathfrak{M}_\sigma$ with the correspondent length metric d' , which is not the same as the distance inherited from $\tilde{\mathcal{G}}$ since now we are not allowed to cross the points in \mathfrak{M}_σ . The metric space $(\tilde{\mathcal{G}} \setminus \mathfrak{M}_\sigma, d')$ is not complete, but we can consider its completion for d' , which is not $\tilde{\mathcal{G}}$. This completion is

$$(\tilde{\mathcal{G}} \setminus \mathfrak{M}_\sigma) \cup \mathfrak{M}_\sigma^\pm,$$

with

$$\mathfrak{M}_\sigma^\pm = \{x_e^{m,+} | e \in E_{\sigma,-}\} \cup \{x_e^{m,-} | e \in E_{\sigma,-}\},$$




FIGURE 8. Left: the field $\tilde{\phi}$ on the event $\tilde{\phi} \in \mathcal{T}_\sigma$, which in this example means that there are no sign clusters surrounding the inner hole of the domain. Right: the associated $\tilde{\xi}$ field. The black dots represent the boundary V_∂ . The violet dots represent \mathfrak{M}_σ . The positive, resp. negative values of the fields are in red, resp. blue.

where $x_e^{m,-}$ and $x_e^{m,+}$ are to be understood as left and right infinitesimal neighborhoods of x_e^m :

$$x_e^{m,-} = \lim_{\varepsilon \rightarrow 0} x_{e,\varepsilon}^-, \quad x_e^{m,+} = \lim_{\varepsilon \rightarrow 0} x_{e,\varepsilon}^+.$$

The fields $\tilde{\phi}$ and $\tilde{\phi}_\sigma$ can be both seen as continuous fields on $(\tilde{\mathcal{G}} \setminus \mathfrak{M}_\sigma) \cup \mathfrak{M}_\sigma^\pm$, with

$$\tilde{\phi}(x_e^{m,-}) = \tilde{\phi}(x_e^{m,+}), \quad \tilde{\phi}_\sigma(x_e^{m,-}) = -\tilde{\phi}_\sigma(x_e^{m,+}), \quad e \in E_{\sigma,-}.$$

A sample of $\tilde{\phi}$ induces a partition $\mathbf{V}_{\tilde{\phi}}$ of \mathfrak{M}_σ^\pm , where two points $x, x' \in \mathfrak{M}_\sigma^\pm$ are in the same class if they are in the same connected component of $\{\tilde{\phi} \neq 0\}$ seen as a subset of $(\tilde{\mathcal{G}} \setminus \mathfrak{M}_\sigma) \cup \mathfrak{M}_\sigma^\pm$ and not as a subset of $\tilde{\mathcal{G}}$. We will denote by $[x]_{\tilde{\phi}}$ the equivalence class of x for $x \in \mathfrak{M}_\sigma^\pm$. The condition $\tilde{\phi} \in \mathcal{T}_\sigma$ is equivalent to $\mathbf{V}_{\tilde{\phi}}$ being bicolored in the following sense: there is a map $\mathbf{b} : \mathbf{V}_{\tilde{\phi}} \rightarrow \{-1, 1\}$, such that for every $e \in E_{\sigma,-}$, $\mathbf{b}([x_e^{m,-}]_{\tilde{\phi}}) = -\mathbf{b}([x_e^{m,+}]_{\tilde{\phi}})$. This is similar to Lemma 3.21. The number of different bicoloredings is 2^k where k is the number of connected components of $\{\tilde{\phi} \neq 0\}$ in $\tilde{\mathcal{G}}$ (not in $(\tilde{\mathcal{G}} \setminus \mathfrak{M}_\sigma) \cup \mathfrak{M}_\sigma^\pm$!) that intersect \mathfrak{M}_σ . Indeed, there are two different colorings per such connected component, one being the opposite of the other.

Let $\text{Bic}(\mathfrak{M}_\sigma)$ denote the set of bicolored partitions of \mathfrak{M}_σ . To each such partition $\mathbf{p} \in \text{Bic}(\mathfrak{M}_\sigma)$ we will associate (in a deterministic way) a bicoloreding $\mathbf{b}_\mathbf{p}$. We define the random field $\tilde{\xi}$ on $\tilde{\mathcal{G}}$ as follows. On the event $\tilde{\phi} \notin \mathcal{T}_\sigma$, we set $\tilde{\xi} = \tilde{\phi}$. On the event $\tilde{\phi} \in \mathcal{T}_\sigma$, $\tilde{\xi}$ is defined by the following.

- (1) For every $x \in \tilde{\mathcal{G}}$, $|\tilde{\xi}(x)| = |\tilde{\phi}(x)|$.
- (2) On every connected component U of $\{\tilde{\phi} \neq 0\}$, such that $U \cap \mathfrak{M}_\sigma = \emptyset$, we have $\tilde{\xi}|_U = \tilde{\phi}|_U$.
- (3) On every connected component U of $\{\tilde{\phi} \neq 0\} \setminus \mathfrak{M}_\sigma$ such that $\bar{U} \cap \mathfrak{M}_\sigma \neq \emptyset$, $\tilde{\xi}|_U = \mathbf{b}_{\mathbf{V}_{\tilde{\phi}}}(U) \tilde{\phi}|_U$, where $\mathbf{b}_{\mathbf{V}_{\tilde{\phi}}}$ is $\mathbf{b}_\mathbf{p}$ with \mathbf{p} being the random partition $\mathbf{V}_{\tilde{\phi}}$, and $\mathbf{b}_{\mathbf{V}_{\tilde{\phi}}}(U)$ is the common value of $\mathbf{b}_{\mathbf{V}_{\tilde{\phi}}}([x]_{\tilde{\phi}})$ for $x \in \bar{U} \cap \mathfrak{M}_\sigma$.

So $\tilde{\xi}$ is obtain from $\tilde{\phi}$ through a deterministic transformation. It corresponds to flipping the signs of some of the connected components of $\{\tilde{\phi} \neq 0\} \setminus \mathfrak{M}_\sigma$, on the event $\tilde{\phi} \in \mathcal{T}_\sigma$, so as to achieve a bicoloreding. See Figure 8.

Corollary 3.25. *The conditional distribution of $\tilde{\xi}$ on the event $\tilde{\phi} \in \mathcal{T}_\sigma$, is that of $\tilde{\phi}_\sigma$.*

Proof. The couple $(\tilde{\phi}, \tilde{\xi})$ can be obtained as limit in law as $\varepsilon \rightarrow 0$ of $(\tilde{\phi}_{|\tilde{\mathcal{G}}_{\sigma,\varepsilon}}, \tilde{\xi}_\varepsilon)$. Then the result is obtained by passing the density (3.13) to the limit. \square

3.5. Around the isomorphism for $\tilde{\phi}_\sigma$. Recall the measure on metric graph loops $\tilde{\mu}^{\text{loop}}$ (2.10). Let $\sigma \in \{-1, 1\}^E$. Let the signed measure on metric graph loops be

$$\tilde{\mu}_\sigma^{\text{loop}}(d\wp) = \text{hol}^\sigma(\wp) \tilde{\mu}^{\text{loop}}(d\wp),$$

which can be decomposed according to the sign into

$$\tilde{\mu}_\sigma^{\text{loop}} = \tilde{\mu}_{\sigma,+}^{\text{loop}} - \tilde{\mu}_{\sigma,-}^{\text{loop}}.$$

Let $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$, respectively $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$, be the Poisson point processes of metric graph loops on $\tilde{\mathcal{G}}$ with intensity measure $\frac{1}{2}\tilde{\mu}_{\sigma,+}^{\text{loop}}$, respectively $\frac{1}{2}\tilde{\mu}_{\sigma,-}^{\text{loop}}$. The version on Le Jan's isomorphism due to Kassel and Lévy (2.8) extends in a straightforward way to the metric graphs:

$$(3.14) \quad (\ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2}))_{x \in \tilde{\mathcal{G}}} \stackrel{(\text{law})}{=} \left(\frac{1}{2}\tilde{\phi}_\sigma(x)^2 + \ell^x(\tilde{\mathcal{L}}_{\sigma,-}^{1/2}) \right)_{x \in \tilde{\mathcal{G}}},$$

where $\tilde{\phi}_\sigma$ and $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$ are taken independent, and ℓ^x above denotes the Brownian local times. One can use for instance an approximation from discrete through a subdivision of edges as in Section 3.1. We also know that

$$(3.15) \quad (\ell^x(\tilde{\mathcal{L}}^{1/2}))_{x \in \tilde{\mathcal{G}}} = (\ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2}) + \ell^x(\tilde{\mathcal{L}}_{\sigma,-}^{1/2}))_{x \in \tilde{\mathcal{G}}} \stackrel{(\text{law})}{=} \left(\frac{1}{2}\tilde{\phi}(x)^2 \right)_{x \in \tilde{\mathcal{G}}},$$

where $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ and $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$ are taken independent. By combining (3.14) and (3.15) we get the following identity in law:

$$(3.16) \quad \left(\frac{1}{2}\tilde{\phi}(x)^2 \right)_{x \in \tilde{\mathcal{G}}} \stackrel{(\text{law})}{=} \left(\frac{1}{2}\tilde{\phi}_\sigma(x)^2 + \ell^x(\tilde{\mathcal{L}}_{\sigma,-}^1) \right)_{x \in \tilde{\mathcal{G}}},$$

where $\tilde{\mathcal{L}}_{\sigma,-}^1$ is a Poisson point process with intensity measure $\tilde{\mu}_{\sigma,-}^{\text{loop}}$, that is to say with intensity parameter $\alpha = 1 = 2 \times \frac{1}{2}$, and $\tilde{\mathcal{L}}_{\sigma,-}^1$ being independent from $\tilde{\phi}_\sigma$.

The identity (3.16) provides another proof for Theorem 1. Indeed, $\tilde{\phi} \in \mathcal{T}_\sigma$ if and only if $\tilde{\phi}^2 \in \mathcal{T}_\sigma$, which, by (3.16), is equivalent to $(\frac{1}{2}\tilde{\phi}_\sigma(x)^2 + \ell^x(\tilde{\mathcal{L}}_{\sigma,-}^1))_{x \in \tilde{\mathcal{G}}}$ belonging to \mathcal{T}_σ . A necessary condition for the latter is $\tilde{\mathcal{L}}_{\sigma,-}^1 = \emptyset$, since $\tilde{\mathcal{L}}_{\sigma,-}^1$ consists precisely of loops with holonomy -1 . This condition is also sufficient, since $\tilde{\phi}_\sigma^2 \in \mathcal{T}_\sigma$ a.s. (Lemma 3.15). Thus, we get that

$$\mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) = \mathbb{P}(\tilde{\mathcal{L}}_{\sigma,-}^1 = \emptyset),$$

which is precisely the probability appearing in Theorem 1. We also get that conditionally on the event $\{\tilde{\phi} \in \mathcal{T}_\sigma\}$, $\tilde{\phi}^2$ is distributed as $\tilde{\phi}_\sigma^2$.

Let us now return to (3.14). Consider \mathcal{C} a connected component of

$$\left\{ x \in \tilde{\mathcal{G}} \mid \frac{1}{2}\tilde{\phi}_\sigma(x)^2 + \ell^x(\tilde{\mathcal{L}}_{\sigma,-}^{1/2}) \neq 0 \right\}.$$

Then $\pi_\sigma^{-1}(\mathcal{C})$ is either connected or not. If $\pi_\sigma^{-1}(\mathcal{C})$ is not connected, then \mathcal{C} cannot contain any loop with holonomy -1 , and in particular, \mathcal{C} cannot contain any loop in $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$. So in this case, \mathcal{C} is actually a connected component of $\{|\tilde{\phi}_\sigma| \neq 0\}$. We summarize this remark in the corollary below.

Corollary 3.26. *One can couple on the same probability space the metric graph loop soups $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ and $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$, and the field $\tilde{\phi}_\sigma$, such that all of the following conditions hold.*

- (1) *The field $\tilde{\phi}_\sigma$ and the loop soup $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$ are independent.*
- (2) *For every $x \in \tilde{\mathcal{G}}$,*

$$\ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2}) = \frac{1}{2}\tilde{\phi}_\sigma(x)^2 + \ell^x(\tilde{\mathcal{L}}_{\sigma,-}^{1/2}).$$

- (3) For every cluster \mathcal{C} of $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ such that $\pi_\sigma^{-1}(\mathcal{C})$ is not connected, \mathcal{C} is also a connected component of $\{|\tilde{\phi}_\sigma| \neq 0\}$ and $\ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2})$ coincides with $\frac{1}{2}\tilde{\phi}_\sigma(x)^2$ on \mathcal{C} .

In particular, one can couple on the same probability space the metric graph loop soups $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ and the field $\tilde{\phi}_\sigma$, such that the following conditions hold.

- (1) For every $x \in \tilde{\mathcal{G}}$,

$$\frac{1}{2}\tilde{\phi}_\sigma(x)^2 \leq \ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2}).$$

- (2) For every cluster \mathcal{C} of $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ such that $\pi_\sigma^{-1}(\mathcal{C})$ is not connected, \mathcal{C} is also a connected component of $\{|\tilde{\phi}_\sigma| \neq 0\}$ and $\ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2})$ coincides with $\frac{1}{2}\tilde{\phi}_\sigma(x)^2$ on \mathcal{C} . In other words, $\ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2})$ and $\frac{1}{2}\tilde{\phi}_\sigma(x)^2$ can differ only on clusters \mathcal{C} of $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ such that $\pi_\sigma^{-1}(\mathcal{C})$ is connected.

In the example of Figure 3, $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ consists of loops that turn around the inner hole an even number of times, including those that do not surround it, and $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$ consists of loops that turn around the inner hole an odd number of times. Further, $\ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2})$ and $\frac{1}{2}\tilde{\phi}_\sigma(x)^2$ coincide on clusters of $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ that do not surround the inner hole. At the risk of being redundant, let us emphasize that the dichotomy for loops in $\tilde{\mathcal{L}}^{1/2}$ and the dichotomy for clusters of $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ are different. For loops in $\tilde{\mathcal{L}}^{1/2}$ one distinguishes between the loops that turn an even number of times around the hole, and the loops that turn an odd number of time around the hole. In this way, the loops that turn twice around the hole are in the same class as the loops that do not surround the hole at all. For the clusters of loops however, the dichotomy is just surrounding or not surrounding the inner hole.

In view of the above corollary, perhaps it is worth pointing out the difference between the clusters of $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ on the metric graph $\tilde{\mathcal{G}}$ and the clusters of $\mathcal{L}_{\sigma,+}^{1/2}$ on the discrete graph \mathcal{G} .

Proposition 3.27. *The discrete loop soup $\mathcal{L}_{\sigma,+}^{1/2}$ is obtained, up to a rerooting of the loops, from the metric graph loop soup $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ by taking the trace of loops on V with the time change A^{-1} (2.9). By doing this, one only takes into account the loops in $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ that visits at least one vertex. In particular, for every $x \in V$, $\ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2}) = \ell^x(\mathcal{L}_{\sigma,+}^{1/2})$. Further, the crossings of edges-lines I_e by $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ correspond to the jumps through discrete edges e by $\mathcal{L}_{\sigma,+}^{1/2}$.*

If an edge $e \in E$ is visited by $\mathcal{L}_{\sigma,+}^{1/2}$, then $\forall x \in I_e, \ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2}) > 0$ a.s. Moreover, for every $e \in E$,

$$(3.17) \quad \mathbb{P}\left(\forall x \in I_e, \ell^x(\tilde{\mathcal{L}}_{\sigma,+}^{1/2}) > 0 \mid \mathcal{L}_{\sigma,+}^{1/2}, e \text{ not visited by } \mathcal{L}_{\sigma,+}^{1/2}\right) \\ = 1 - \exp\left(-2C(e)(\ell^{e_-}(\tilde{\mathcal{L}}_{\sigma,+}^{1/2})\ell^{e_+}(\tilde{\mathcal{L}}_{\sigma,+}^{1/2}))^{1/2}\right),$$

with conditional independence (given $\mathcal{L}_{\sigma,+}^{1/2}$) across the edges $e \in E$.

Proof. These are the properties already satisfied by the loop soups $\tilde{\mathcal{L}}^{1/2}$ and $\mathcal{L}^{1/2}$, proven in [Lup16]. The fact that $\mathcal{L}^{1/2}$ is the trace of $\tilde{\mathcal{L}}^{1/2}$ on the vertices V immediately implies that $\mathcal{L}_{\sigma,+}^{1/2}$ is the trace of $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ on V , as well as that $\mathcal{L}_{\sigma,-}^{1/2}$ is the trace of $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$ on V .

As for the formula (3.17), given $e \in E$ not visited by $\mathcal{L}_{\sigma,+}^{1/2}$, the connection between the two endpoints of e can be created by a superposition of three objects: the Brownian excursions from e_+ to e_+ inside I_e , the Brownian excursions from e_- to e_- inside I_e , and the Brownian loops of $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$ that stay inside I_e . The formula (3.17) is then the same as for $\tilde{\mathcal{L}}^{1/2}$ and $\mathcal{L}^{1/2}$ appearing in [Lup16] since the same three types of Brownian paths appear in both settings. This is in particular due to the fact that all the Brownian loops in $\tilde{\mathcal{L}}^{1/2}$ that stay inside I_e have

a holonomy 1, and thus also appear in $\tilde{\mathcal{L}}_{\sigma,+}^{1/2}$. Note that for $\tilde{\mathcal{L}}_{\sigma,-}^{1/2}$ and $\mathcal{L}_{\sigma,-}^{1/2}$, a similar formula is no longer true, precisely because the Brownian loops staying inside I_e no longer participate to connecting the two ends of e , as they all have a wrong holonomy. \square

4. RELATION TO DISORDERED ISING MODEL

The goal of this Section is to explain how Theorem 1 can be alternatively derived from a similar result for the FK-Ising model.

4.1. Spin Ising, FK-Ising and Edwards-Sokal coupling. So as to avoid confusion with our notation σ for the $\{-1, 1\}$ -gauge fields, we will denote the Ising spins by ζ . Let $\mathcal{G} = (V, E)$ be a finite connected graph as in Section 2.1. The edges $\{x, y\} \in E$ are endowed with weights $\beta(x, y) = \beta(y, x) \geq 0$. The *spin Ising* field is a random collection of signs $(\hat{\zeta}(x))_{x \in V} \in \{-1, 1\}^V$, with the distribution given by

$$\mathbb{P}((\hat{\zeta}(x))_{x \in V} = (\zeta(x))_{x \in V}) = \frac{1}{Z_{\beta}^{\text{Ising}}} \exp \left(\sum_{\{x, y\} \in E} \beta(x, y) \zeta(x) \zeta(y) \right).$$

Further, the *FK-Ising random cluster model* [Gri06] is a random configuration of edges $(\hat{\omega}(e))_{e \in E} \in \{0, 1\}^E$, where $\hat{\omega}(e) = 1$ corresponds to an open edge and $\hat{\omega}(e) = 0$ corresponds to a closed edge. The probability distribution of the FK-Ising is given by

$$(4.1) \quad \mathbb{P}((\hat{\omega}(e))_{e \in E} = (\omega(e))_{e \in E}) = \frac{1}{Z_{\beta}^{\text{FK-Ising}}} 2^{\# \text{ clusters of } \omega} \prod_{e \in E} (1 - e^{-2\beta(e)})^{\omega(e)} (e^{-2\beta(e)})^{1-\omega(e)},$$

where $\#$ clusters of ω is the number of clusters induced by the open edges of ω .

The spin Ising and the FK-Ising models are related through the Edwards-Sokal coupling [ES88].

Theorem 4.1 (Edwards-Sokal, [ES88]). *The following holds.*

- (1) *The two partition function are related through*

$$Z_{\beta}^{\text{Ising}} = Z_{\beta}^{\text{FK-Ising}} \prod_{e \in E} e^{\beta(e)}.$$

- (2) *The spin Ising configuration $(\hat{\zeta}(x))_{x \in V}$ and the FK-Ising configuration $(\hat{\omega}(e))_{e \in E}$ can be coupled as follows. One first samples the FK-Ising configuration $(\hat{\omega}(e))_{e \in E}$, and then one samples an independent uniform sign for each cluster induced by $\hat{\omega}$.*
- (3) *The coupling above can be alternatively described as follows. One first samples the spin Ising configuration $(\hat{\zeta}(x))_{x \in V}$. Then, for each edge $\{x, y\} \in E$, one sets $\hat{\omega}(x, y) = 0$ if $\hat{\zeta}(x)\hat{\zeta}(y) = -1$, and sets $\hat{\omega}(x, y) = 1$ with conditional probability $1 - e^{-2\beta(x, y)}$ if $\hat{\zeta}(x)\hat{\zeta}(y) = 1$.*

4.2. Disordered Ising and topological probabilities for FK-Ising. Let $\sigma \in \{-1, 1\}^E$ be a gauge field as in Section 2.1. The *disordered spin Ising* field is a random configuration of signs $(\hat{\zeta}_{\sigma}(x))_{x \in V} \in \{-1, 1\}^V$ with the following probability distribution:

$$\mathbb{P}((\hat{\zeta}_{\sigma}(x))_{x \in V} = (\zeta(x))_{x \in V}) = \frac{1}{Z_{\beta, \sigma}^{\text{Ising}}} \exp \left(\sum_{\{x, y\} \in E} \beta(x, y) \zeta(x) \sigma(x, y) \zeta(y) \right).$$

The terminology originates from [KC71]. The field $\hat{\zeta}_{\sigma}$ is covariant under gauge transformations on σ just as in the GFF case (2.3).

Theorem 1 has an analogue in the Ising setting. Recall the notations of Section 3.2: $\pi_{\sigma} : \mathcal{G}_{\sigma}^{\text{db}} \rightarrow \mathcal{G}$ is the double cover induced by σ . Let $\text{Triv}_{\sigma} \subset \{0, 1\}^E$ denote the subset of edge configurations ω such that every \mathcal{C} cluster induced by the open edges of ω , $\pi_{\sigma}^{-1}(\mathcal{C})$ is **not** connected.

Theorem 4.2. *Let $(\hat{\omega}(e))_{e \in E} \in \{0, 1\}^E$ be an FK-Ising configuration distributed according to (4.1). The probability $\mathbb{P}(\hat{\omega} \in \text{Triv}_\sigma)$ can be expressed as a ratio of Ising partition functions:*

$$\mathbb{P}(\hat{\omega} \in \text{Triv}_\sigma) = \frac{Z_{\beta, \sigma}^{\text{Ising}}}{Z_\beta^{\text{Ising}}}.$$

Moreover, the configuration $\hat{\omega}$ conditioned on $\{\hat{\omega} \in \text{Triv}_\sigma\}$ can be sampled as follows.

- (1) First sample a disordered spin Ising configuration $\hat{\zeta}_\sigma$.
- (2) For each edge $e = \{x, y\} \in E$ such that $\zeta(x)\sigma(x,y)\zeta(y) = -1$, set the edge e to closed (i.e. 0).
- (3) For each edge $e = \{x, y\} \in E$ such that $\zeta(x)\sigma(x,y)\zeta(y) = 1$, set the edge e to open (i.e. 1) with conditional probability $1 - e^{-2\beta(x,y)}$.

Theorem 4.2 is a gauge-twisted version of the Edwards-Sokal coupling (Theorem 4.1), and just as the latter, can be derived through an elementary computation. We would like to emphasize that just like Theorem 1, Theorem 4.2 does not rely on planarity at all. This result has been communicated to us by Marcin Lis (TU Wien, Vienna) after the prepublication of the first version of this paper. It also seems to be common knowledge among the Ising community, but we did not find a good reference for it in the literature.

Alternatively, Theorem 4.2 can be derived from the GFF case as in Remark 3.19, by considering double-well φ^4 interactions $g \sum_{x \in V} (\tilde{\phi}(x)^2 - 1)^2$ and letting the coupling constant $g \rightarrow +\infty$.

Note that in the GFF setting (Theorem 1) the partition functions and thus the topological probabilities are way more explicit and tractable than in the Ising setting (Theorem 4.2).

4.3. Relation between Ising and the GFF. Consider an electrical network $\mathcal{G} = (V, E)$ as in Section 2.1 endowed with conductances $C(x, y) = C(y, x) > 0$ for $\{x, y\} \in E$. Given a non-negative function $h : V \rightarrow \mathbb{R}_+$, we will denote by β_h the weights

$$(4.2) \quad \beta_h(x, y) = C(x, y)h(x)h(y), \quad \{x, y\} \in E.$$

In the sequel we will consider ϕ a discrete GFF on \mathcal{G} with 0 boundary conditions, and its metric graph extension $\tilde{\phi}$ ($\tilde{\phi}|_V = \phi$). From the density (2.1) it is clear that conditionally on $(|\phi(x)|)_{x \in V}$, the signs $(\text{sign}(\phi(x)))_{x \in V_{\text{int}}}$ are distributed as an Ising spin field with weights

$$\beta_{|\phi|}(x, y) = C(x, y)|\phi(x)||\phi(y)|.$$

Note that $\beta_{|\phi|}(x, y) = 0$ if x or y is in V_∂ . As observed by Lupu and Werner in [LW16], the metric graph extension $\tilde{\phi}$ also naturally enters this picture and can be interpreted in terms of the FK-Ising. Denote by $\omega_{\tilde{\phi}} \in \{0, 1\}^E$ the following edge configuration. We set $\omega_{\tilde{\phi}}(e) = 1$ if $\tilde{\phi}$ has no zeroes on the edge-line I_e , and $\omega_{\tilde{\phi}}(e) = 0$ otherwise. Note that if e is adjacent to the boundary V_∂ , then $\omega_{\tilde{\phi}}(e) = 0$ a.s.

Proposition 4.3 (Lupu-Werner, [LW16]). *Let $h : V \rightarrow \mathbb{R}_+$ be a random non-negative field distributed as the absolute value of the discrete GFF $(|\phi(x)|)_{x \in V}$. Let $(\hat{\zeta}(x))_{x \in V} \in \{-1, 1\}^V$ be a random spin field distributed, conditionally on h , as spin Ising with weights β_h . Let $(\hat{\omega}(e))_{e \in E} \in \{0, 1\}^E$ be a random edge configuration distributed, conditionally on h , as FK-Ising with weights β_h . We further assume that conditionally on h , $\hat{\zeta}$ and $\hat{\omega}$ are coupled as in Edwards-Sokal coupling (Theorem 4.1). Then $((\hat{\zeta}(x)h(x))_{x \in V}, (\hat{\omega}(e))_{e \in E})$ are jointly distributed as $((\phi(x))_{x \in V}, (\omega_{\tilde{\phi}}(e))_{e \in E})$.*

4.4. From FK-Ising topological probabilities to GFF topological probabilities. Here we will sketch an alternative proof of Theorem 1 that relies on Proposition 4.3 and Theorem 4.2.

The probability distribution of the absolute value of the discrete GFF $(|\phi(x)|)_{x \in V}$ can be written as

$$\frac{1}{Z} Z_{\beta_h}^{\text{Ising}} \prod_{x \in V_{\text{int}}} e^{-\frac{1}{2}W(x)h(x)^2} \prod_{x \in V_{\text{int}}} dh(x),$$

where the weights β_h are given by (4.2), with the convention $h|_{V_\partial} \equiv 0$, $Z_{\beta_h}^{\text{Isg}}$ is the spin Ising partition function for weights β_h , Z is the GFF partition function, and $W(x)$ is given by (2.4). Now, let $\tilde{\phi}$ be a gauge field $\sigma \in \{-1, 1\}$. Then $\tilde{\phi} \in \mathcal{T}_\sigma$ if and only if $\omega_{\tilde{\phi}} \in \text{Triv}_\sigma$. Therefore,

$$\mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma \mid (|\phi(x)|)_{x \in V}) = \mathbb{P}(\omega_{\tilde{\phi}} \in \text{Triv}_\sigma \mid (|\phi(x)|)_{x \in V}) = \frac{Z_{\beta_{|\phi|}, \sigma}^{\text{Isg}}}{Z_{\beta_{|\phi|}}^{\text{Isg}}}.$$

Further,

$$\begin{aligned} \mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma) &= \mathbb{E}[\mathbb{P}(\tilde{\phi} \in \mathcal{T}_\sigma \mid (|\phi(x)|)_{x \in V})] = \mathbb{E}\left[\frac{Z_{\beta_{|\phi|}, \sigma}^{\text{Isg}}}{Z_{\beta_{|\phi|}}^{\text{Isg}}}\right] \\ &= \frac{1}{Z} \int \frac{Z_{\beta_{h, \sigma}}^{\text{Isg}}}{Z_{\beta_h}^{\text{Isg}}} Z_{\beta_h}^{\text{Isg}} \prod_{x \in V_{\text{int}}} e^{-\frac{1}{2}W(x)h(x)^2} \prod_{x \in V_{\text{int}}} dh(x) \\ &= \frac{1}{Z} \int Z_{\beta_{h, \sigma}}^{\text{Isg}} \prod_{x \in V_{\text{int}}} e^{-\frac{1}{2}W(x)h(x)^2} \prod_{x \in V_{\text{int}}} dh(x) = \frac{Z_\sigma}{Z}. \end{aligned}$$

The conditional distribution of $\tilde{\phi}$ on the event $\{\tilde{\phi} \in \mathcal{T}_\sigma\}$ can be obtained by similar arguments.

5. INTERPRETATIONS AND IMPLICATIONS OF THE RESULT

5.1. GFF on annular domains and exploration from inside. In the introduction (Figure 1) we considered the example of planar annular domains and the event that $\tilde{\phi}$ has a sign cluster that surrounds the inner hole. Here we will recall how annular domains naturally arise in a “simply connected” context.

For simplicity, let us consider a two-dimensional discrete box $\mathcal{G} = (V, E)$ with

$$V = \{-n, -n+1, \dots, -1, 0, 1, \dots, n-1, n\}^2,$$

$V_{\text{int}} = \{-n+1, \dots, -1, 0, 1, \dots, n-1\}^2$, $V_\partial = V \setminus V_{\text{int}}$ and the edges formed by $z, w \in V$ such that $|z - w| = 1$ (square lattice). Let $\tilde{\mathcal{G}}$ be the metric graph associated to \mathcal{G} . Let $\tilde{\phi}$ the metric graph GFF (non-twisted) on $\tilde{\mathcal{G}}$ with 0 boundary conditions on V_∂ .

Let \tilde{K}_0 be a deterministic non-empty compact subset of $\tilde{\mathcal{G}}$ such that \tilde{K}_0 is connected and $d(\tilde{K}_0, V_\partial) > 1$. Given a sample of $\tilde{\phi}$, we define the following random subset \tilde{K}_1 of $\tilde{\mathcal{G}}$, depending on \tilde{K}_0 and $\tilde{\phi}$:

$$\tilde{K}_1 = \overline{\tilde{K}_0 \cup \{z \in \tilde{\mathcal{G}} \setminus \tilde{K}_0 \mid \exists \text{ continuous path } \wp \text{ in } \mathcal{G}, z \xleftrightarrow{\wp} \tilde{K}_0 \text{ and } |\tilde{\phi}|_\wp > 0\}}.$$

In other words, \tilde{K}_1 is made of \tilde{K}_0 and all the points that can be connected to \tilde{K}_0 by a path \wp on which $\tilde{\phi}$ does not go through 0, except possibly at the extremity. The random compact subset \tilde{K}_1 is a so-called *stopping set* for the GFF $\tilde{\phi}$: given any deterministic open subset U of $\tilde{\mathcal{G}}$, the event $\{\tilde{K}_1 \subset U\}$ is measurable with respect to the restriction $\tilde{\phi}|_U$. The subset \tilde{K}_1 can be obtained by first discovering $\tilde{\phi}$ on \tilde{K}_0 and then exploring from there in all the directions and stopping an exploration branch whenever the value of $\tilde{\phi}$ on this branch reaches 0. By construction, \tilde{K}_1 is connected and $\tilde{\phi}$ is 0 on $\partial\tilde{K}_1$. It is easy to see that with positive probability, $d(\tilde{K}_1, V_\partial) > 1$. We further introduce another random set \tilde{K}_2 , obtained by filling the inner holes of \tilde{K}_1 :

$$\tilde{K}_2 = \tilde{K}_1 \cup \bigcup_{\substack{O \text{ connected} \\ \text{component} \\ \text{of } \tilde{\mathcal{G}} \setminus \tilde{K}_1, O \cap V_\partial = \emptyset}} O.$$

Then \tilde{K}_2 is again a stopping set for $\tilde{\phi}$. By construction, $d(\tilde{K}_2, V_\partial) = d(\tilde{K}_1, V_\partial)$, and again, $\tilde{\phi}$ is 0 on $\partial\tilde{K}_2$.

On the event $\{d(\tilde{K}_2, V_\partial) > 1\}$, which has a positive probability, the sub-metric-graph $\overline{\tilde{\mathcal{G}} \setminus \tilde{K}_2}$ is annular, that is to say it contains one hole, which is $\tilde{K}_2 \setminus \partial\tilde{K}_2$. The values of $\tilde{\phi}$ on $\partial(\tilde{\mathcal{G}} \setminus \tilde{K}_2)$ are 0 by construction. Since \tilde{K}_2 is a stopping set, by the strong Markov property of $\tilde{\phi}$ (see [Lup16]), conditionally on $(\tilde{K}_2, \tilde{\phi}|_{\tilde{K}_2})$, the field $\tilde{\phi}|_{\overline{\tilde{\mathcal{G}} \setminus \tilde{K}_2}}$ is distributed as the GFF on the metric graph $\overline{\tilde{\mathcal{G}} \setminus \tilde{K}_2}$. Therefore, Theorem 1 gives the conditional probability for a sign cluster of $\tilde{\phi}|_{\overline{\tilde{\mathcal{G}} \setminus \tilde{K}_2}}$ to surround \tilde{K}_2 given $(\tilde{K}_2, \tilde{\phi}|_{\tilde{K}_2})$. The conditional probability given just \tilde{K}_2 is the same, since $\tilde{\phi}|_{\overline{\tilde{\mathcal{G}} \setminus \tilde{K}_2}}$ and $\tilde{\phi}|_{\tilde{K}_2}$ are independent conditionally on \tilde{K}_2 .

5.2. GFF on annular domains: continuum limits. Here we will consider what happens in the scaling limit on doubly connected (annular) domains in the scaling limit.

Here we will call an *annular domain* an open bounded subset $A \subset \mathbb{C}$ such that $\mathbb{C} \setminus A$ has two connected components, one being necessarily unbounded and the other one bounded (the inner hole), with the additional condition that the hole is not reduced to one point. We will denote by $\partial_o A$ and $\partial_i A$ the outer and the inner boundaries of A . The conformal equivalence classes of annular domains are parametrized by the *extremal distance*, or *extremal length*, between the outer and the inner boundary $\text{ED}(\partial_o A, \partial_i A)$. The quantity $\text{ED}(\partial_o A, \partial_i A)$ is really nothing else than the electrical resistance between $\partial_o A$ and $\partial_i A$. Given a circular annulus

$$A_{r_1, r_2} = \{z \in \mathbb{C} | r_1 < |z| < r_2\},$$

the corresponding extremal distance is

$$\text{ED}(\partial_o A_{r_1, r_2}, \partial_i A_{r_1, r_2}) = \frac{1}{2\pi} \log(r_2/r_1).$$

So every annular domain A is conformally equivalent to a circular annulus $A_{r,1}$, where

$$r = \exp(-2\pi \text{ED}(\partial_o A, \partial_i A)).$$

For details, we refer to [Ahl10, Chapter 4].

5.2.1. Probability of non-contractible sign clusters in the scaling limit. Let A be an annular domain as above, and Φ a continuum GFF on A with 0 boundary conditions, both on $\partial_o A$ and $\partial_i A$. In [ALS22], Aru-Lupu-Sepúlveda considered the non-contractible interior *level lines* of Φ with step 2λ , where 2λ is the Schramm-Sheffield *height gap* [SS09, SS13] of the 2D continuum GFF. More precisely, one has a random sequence of simple (Jordan) loops $\mathbf{C}_1, \dots, \mathbf{C}_N$, with N also random, and a sequence of labels $v_1, \dots, v_N \in 2\lambda\mathbb{Z}$, where:

- by convention, $\mathbf{C}_0 = \partial_o A$, $\mathbf{C}_{N+1} = \partial_i A$, $v_0 = v_{N+1} = 0$;
- $N \geq 0$ and N finite and even;
- for all $i \in \{1, \dots, N\}$, \mathbf{C}_i is a simple loop in A that separates $\partial_o A$ and $\partial_i A$;
- for all $i \in \{0, 1, \dots, N\}$, \mathbf{C}_i surrounds \mathbf{C}_{i+1} ;
- for all $i \in \{0, 1, \dots, N\}$, $v_{i+1} - v_i \in \{-2\lambda, 2\lambda\}$;
- the family of random variables $(N, \mathbf{C}_1, \dots, \mathbf{C}_N, v_1, \dots, v_N)$ is measurable with respect to Φ ;
- for all $i \in \{1, \dots, N\}$, v_i is the value of Φ on the outer side of \mathbf{C}_i and v_{i+1} is the value of Φ on the inner side of \mathbf{C}_i .

Each of the loops \mathbf{C}_i locally look like an SLE_4 curve. See [ALS22, Section 4.1] for details.

Now, with positive probability, $N = 0$. On the event $\{N = 0\}$, all the non-contractible level lines $\mathbf{C}_1, \dots, \mathbf{C}_N$ and the labels v_1, \dots, v_N do not exist. The event $\{N = 0\}$ is the continuum analogue of the event of no sign cluster surrounding the inner hole, that is to say of the left side of Figure 1. The probability $\mathbb{P}(N = 0)$ can be expressed explicitly by applying SLE and local set tools, as detailed for instance in [ALS22, Proposition 2.18]. This is also the same probability for a 2D Brownian loop soup $\mathcal{L}_A^{1/2}$ in A (intensity parameter $\alpha = 1/2$) not having a non-contractible cluster. The probability $\mathbb{P}(N = 0)$ can be expressed through one-dimensional

Brownian bridges. Let $(\widehat{W}_t)_{0 \leq t \leq L}$ be a standard Brownian bridge on \mathbb{R} from 0 to 0, of time-length $L = \text{ED}(\partial_o A, \partial_i A)$. Then the following holds.

Proposition 5.1 (Aru-Lupu-Sepúlveda, [ALS22]). *One has the equality*

$$\mathbb{P}(N = 0) = \mathbb{P}((\widehat{W}_t)_{0 \leq t \leq L} \text{ stays in } (-\sqrt{\pi/2}, \sqrt{\pi/2})).$$

Note that $\sqrt{\pi/2}$ is the value of the height gap 2λ in an appropriate normalization of the GFF Φ . The probability $\mathbb{P}(N = 0)$ depends on A only through the extremal distance $L = \text{ED}(\partial_o A, \partial_i A)$, that it to say only through the conformal equivalence class of A . This is consistent with the conformal invariance of Φ .

Now consider $\tilde{A}^{(n)}$ metric graph approximations of the annular domain A in the square lattice $\frac{1}{n}\mathbb{Z}^2$, and let $\tilde{\phi}_n$ be the GFF on $\tilde{A}^{(n)}$ with 0 boundary conditions. We are interested to verify that

$$\lim_{n \rightarrow 0} \mathbb{P}(\tilde{\phi}_n \text{ has no sign clusters surrounding the inner hole of } \tilde{A}^{(n)}) = \mathbb{P}(N = 0).$$

Of course, this can be deduced from the abstract arguments on the convergence of sign clusters. But this is not our goal here. Our goal is to compare the exact formulas given on one hand by Theorem 1, and on the other hand by Proposition 5.1 and check that they indeed match. Consider this as a sanity check.

For this we will need to introduce the Brownian loop measure on A ; see [LW04] and [Law05, Section 5.6]. It is an infinite measure given by

$$(5.1) \quad \mu_A^{\text{loop}}(d\wp) = \int_A \int_0^{+\infty} \mathbb{P}_{A,t}^{z,z}(d\wp) p_A(t, z, z) \frac{dt}{t} d^2z,$$

where $p_A(t, z, w)$ denotes the heat kernel on A with 0 boundary conditions on ∂A , and $\mathbb{P}_{A,t}^{z,z}$ are the 2D Brownian bridge probability measures where the bridge is conditioned on staying in A . Theorem 1 implies that

$$\begin{aligned} & \lim_{n \rightarrow 0} \mathbb{P}(\tilde{\phi}_n \text{ has no sign clusters surrounding the inner hole of } \tilde{A}^{(n)}) \\ &= \exp\left(-\mu_A^{\text{loop}}(\{\text{Loops that wind an odd number of times around the hole of } A\})\right). \end{aligned}$$

For the convergence of the loop measure from discrete to continuum, we refer to [LTF07]. So our goal is to verify the following.

Proposition 5.2. *The following identity holds:*

$$\begin{aligned} & \exp\left(-\mu_A^{\text{loop}}(\{\text{Loops that wind an odd number of times around the hole of } A\})\right) \\ &= \mathbb{P}((\widehat{W}_t)_{0 \leq t \leq L} \text{ stays in } (-\sqrt{\pi/2}, \sqrt{\pi/2})). \end{aligned}$$

Note that in the identity above, the left-hand side involves 2D Brownian bridges, and the right-hand side 1D Brownian bridges. We do not have a direct probabilistic interpretation for this identity, other than relying on the content of this paper and all the knowledge on the level lines of the 2D continuum GFF. So our verification will proceed through explicit computations. These computations are however rather sophisticated and ultimately rely on the Jacobi triple product identity (5.7).

Denote by $p_{\mathbb{R}}$ the heat kernel on \mathbb{R} , and by $p_{(-\sqrt{\pi/2}, \sqrt{\pi/2})}$ the heat kernel on $(-\sqrt{\pi/2}, \sqrt{\pi/2})$ with 0 boundary conditions at $\pm\sqrt{\pi/2}$. Then

$$\mathbb{P}((\widehat{W}_t)_{0 \leq t \leq L} \text{ stays in } (-\sqrt{\pi/2}, \sqrt{\pi/2})) = \frac{p_{(-\sqrt{\pi/2}, \sqrt{\pi/2})}(L, 0, 0)}{p_{\mathbb{R}}(L, 0, 0)} = \sqrt{2\pi L} p_{(-\sqrt{\pi/2}, \sqrt{\pi/2})}(L, 0, 0).$$

The quantity $p_{(-\sqrt{\pi/2}, \sqrt{\pi/2})}(L, 0, 0)$ can be decomposed into series in two different ways. By using the reflection principle, one gets

$$(5.2) \quad p_{(-\sqrt{\pi/2}, \sqrt{\pi/2})}(L, 0, 0) = \frac{1}{\sqrt{2\pi L}} \sum_{k \in \mathbb{Z}} \left(e^{-\frac{4k^2\pi}{L}} - e^{-\frac{(2k+1)^2\pi}{L}} \right).$$

This is for instance Formula 3.0.2 in [BS15, Section 1.3]. By rather using the Fourier decomposition of the heat kernel, we get

$$(5.3) \quad p_{(-\sqrt{\pi/2}, \sqrt{\pi/2})}(L, 0, 0) = \sqrt{\frac{2}{\pi}} \sum_{j \geq 0} e^{-\frac{(2j+1)^2\pi L}{4}}.$$

The two series (5.2) and (5.3) are related by the Poisson summation formula. The quantity $p_{(-\sqrt{\pi/2}, \sqrt{\pi/2})}(L, 0, 0)$ can also be written in terms of Jacobi Theta functions, or rather Theta Nullwert functions; see [AS84, Section 16.27]. With the standard notations, let be

$$\theta_2(q) = \sum_{j \in \mathbb{Z}} q^{(j+1/2)^2}, \quad \theta_4(q) = \sum_{k \in \mathbb{Z}} (-1)^k q^{k^2}.$$

We have that

$$(5.4) \quad \begin{aligned} \mathbb{P}((\widehat{W}_t)_{0 \leq t \leq L} \text{ stays in } (-\sqrt{\pi/2}, \sqrt{\pi/2})) &= \sqrt{2\pi L} p_{(-\sqrt{\pi/2}, \sqrt{\pi/2})}(L, 0, 0) \\ &= \theta_4(q = e^{-\frac{\pi}{L}}) = \sqrt{L} \theta_2(q = e^{-\pi L}). \end{aligned}$$

Now let us perform the computations on the 2D loop side. Let be

$$r = \exp(-2\pi \text{ED}(\partial_o A, \partial_i A)) = e^{-2\pi L}.$$

Let \widehat{A} be the circular annulus $\widehat{A} = A_{r,1}$, so that A is conformally equivalent to \widehat{A} . Let $\mu_{\widehat{A}}^{\text{loop}}$ be the Brownian loop measure on \widehat{A} . The conformal invariance of the Brownian loop measure (see [Law05, Proposition 5.27]) ensures that the following identity holds:

$$\begin{aligned} \mu_A^{\text{loop}}(\{\text{Loops that wind an odd number of times around the hole of } A\}) \\ = \mu_{\widehat{A}}^{\text{loop}}(\{\text{Loops that wind an odd number of times around the hole of } \widehat{A}\}). \end{aligned}$$

Rather than performing computations on the annulus \widehat{A} , we will lift everything up to its universal cover via the log map. However, for doing this, it is much more convenient to endow \widehat{A} with the cylindrical metric $|z|^{-2}d^2z$ rather than the Euclidean metric d^2z . Indeed, consider the strip

$$(5.5) \quad S = \{u + iv \mid -2\pi L < u < 0, v \in \mathbb{R}\}.$$

The exp map induces a covering of \widehat{A} by S , and it sends the Euclidean metric on a fundamental domain in S to the cylindrical metric $|z|^{-2}d^2z$ on \widehat{A} . So consider the time-changed Brownian motion with infinitesimal generator $\frac{1}{2}|z|^2\Delta$, killed upon hitting $\partial\widehat{A}$. This process is symmetric with respect to the cylindrical metric $|z|^{-2}d^2z$. Let $\overset{\circ}{P}_{\widehat{A}}(t, z, w)$ be its transition densities (with condition 0 on $\partial\widehat{A}$) with respect to $|z|^{-2}d^2z$, and $\overset{\circ}{P}_{\widehat{A},t}^{\circ z, z}$ the corresponding bridge probability measures conditioned on staying in \widehat{A} . The Brownian loop measure on \widehat{A} for the cylindrical metric $|z|^{-2}d^2z$ is given by

$$\mu_{\widehat{A}}^{\text{loop}}(d\varphi) = \int_{\widehat{A}} \int_0^{+\infty} \overset{\circ}{P}_{\widehat{A},t}^{\circ z, z}(d\varphi) \overset{\circ}{P}_{\widehat{A}}(t, z, z) \frac{dt}{t} \frac{d^2z}{|z|^2}.$$

Since the cylindrical metric $|z|^{-2}d^2z$ is conformally equivalent to the Euclidean metric d^2z , the conformal invariance of the Brownian loop measure ensures that

$$\begin{aligned} \mu_{\widehat{A}}^{\text{loop}}(\{\text{Loops that wind an odd number of times around the hole of } \widehat{A}\}) \\ = \mu_{\widehat{A}}^{\circ, \text{loop}}(\{\text{Loops that wind an odd number of times around the hole of } \widehat{A}\}). \end{aligned}$$

Now, lifting from \widehat{A} up to S through the log map, one gets that for every $z \in \widehat{A}$ and $t > 0$,

$$\begin{aligned} P_{\widehat{A}, t}^{\circ, z, z}(\text{Bridge winds an odd number of times around the hole of } \widehat{A}) & \overset{\circ}{p}_{\widehat{A}}(t, z, z) \\ & = \sum_{k \in \mathbb{Z}} p_S(t, u + iv, u + i(v + (2k + 1)2\pi)), \end{aligned}$$

where p_S is the heat kernel on the strip S with 0 boundary conditions on ∂S , and $e^{u+iv} = z$, with $v \in [0, 2\pi)$. So, at the end of the day,

$$(5.6) \quad \begin{aligned} \mu_A^{\text{loop}}(\{\text{Loops that wind an odd number of times around the hole of } A\}) \\ = \int_{-2\pi L}^0 \int_0^{2\pi} \int_0^{+\infty} \sum_{k \in \mathbb{Z}} p_S(t, u + iv, u + i(v + (2k + 1)2\pi)) \frac{dt}{t} dv du. \end{aligned}$$

Further, one can factorize the heat kernel on S by separating real and imaginary parts:

$$p_S(t, u + iv, u' + iv') = p_{(-2\pi L, 0)}(t, u, u') p_{\mathbb{R}}(t, v, v') = p_{(-2\pi L, 0)}(t, u, u') \frac{1}{\sqrt{2\pi t}} e^{-\frac{(v'-v)^2}{2t}},$$

where $p_{(-2\pi L, 0)}(t, u, u')$ denotes the heat kernel on $(-2\pi L, 0)$ with boundary condition 0 in $-2\pi L$ and 0. Thus, (5.6) equals

$$2\pi \int_0^{+\infty} \int_{-2\pi L}^0 p_{(-2\pi L, 0)}(t, u, u) du \sum_{k \in \mathbb{Z}} e^{-\frac{2(2k+1)^2\pi^2}{t}} \frac{dt}{\sqrt{2\pi t^{3/2}}}.$$

Further, similarly to (5.3), one can write

$$\begin{aligned} p_{(-2\pi L, 0)}(t, u, u) & = \frac{1}{\pi L} \sum_{j \geq 0} \cos\left(\frac{2j+1}{2L}(u + \pi L)\right)^2 e^{-\frac{(2j+1)^2 t}{8L^2}} \\ & \quad + \frac{1}{\pi L} \sum_{j \geq 1} \sin\left(\frac{j}{L}(u + \pi L)\right)^2 e^{-\frac{j^2 t}{2L^2}}. \end{aligned}$$

Thus,

$$\int_{-2\pi L}^0 p_{(-2\pi L, 0)}(t, u, u) du = \sum_{j \geq 0} e^{-\frac{(2j+1)^2 t}{8L^2}} + \sum_{j \geq 1} e^{-\frac{j^2 t}{2L^2}} = \sum_{j \geq 1} e^{-\frac{j^2 t}{8L^2}}.$$

Thus, (5.6) equals

$$\sqrt{2\pi} \sum_{j \geq 1} \sum_{k \in \mathbb{Z}} \int_0^{+\infty} e^{-\frac{j^2 t}{8L^2} - \frac{2(2k+1)^2\pi^2}{t}} \frac{dt}{t^{3/2}}.$$

Lemma 5.3. For every $a, b > 0$,

$$\int_0^{+\infty} e^{-at-bt^{-1}} \frac{dt}{t^{3/2}} = \sqrt{\frac{\pi}{b}} e^{-2\sqrt{ab}}.$$

Proof. This identity seems to be classic. For instance,

$$\mathbf{1}_{t \geq 0} \sqrt{\frac{\lambda}{2\pi t^3}} \exp\left(-\frac{\lambda(t-\mu)^2}{2\mu^2 t}\right)$$

is the density of the Inverse Gaussian distribution $\text{IG}(\mu, \lambda)$ [FC78], and in particular, its integral on $(0, +\infty)$ equals 1. \square

By applying Lemma 5.3, we get that (5.6) equals

$$\begin{aligned} \sum_{j \geq 1} \sum_{k \in \mathbb{Z}} \frac{1}{|2k+1|} e^{-\frac{j|2k+1|\pi}{L}} &= 2 \sum_{j \geq 1} \sum_{k \geq 0} \frac{1}{2k+1} e^{-\frac{j(2k+1)\pi}{L}} \\ &= - \sum_{j \geq 1} (2 \log(1 - e^{-\frac{j\pi}{L}}) - \log(1 - e^{-\frac{2j\pi}{L}})). \end{aligned}$$

By recombining the terms in the sum, this is turn equals

$$- \sum_{l \geq 1} (\log(1 - e^{-\frac{2l\pi}{L}}) + 2 \log(1 - e^{-\frac{(2l-1)\pi}{L}})).$$

So, we get

$$\begin{aligned} \exp(-\mu_A^{\text{loop}}(\{\text{Loops that wind an odd number of times around the hole of } A\})) \\ = \prod_{l \geq 1} (1 - e^{-\frac{2l\pi}{L}}) \prod_{l \geq 1} (1 - e^{-\frac{(2l-1)\pi}{L}})^2. \end{aligned}$$

By comparing with (5.4), to conclude to Proposition 5.2, we need the following identity. For every $q \in [0, 1)$,

$$(5.7) \quad \prod_{l \geq 1} (1 - q^{2l}) \prod_{l \geq 1} (1 - q^{2l-1})^2 = \sum_{k \in \mathbb{Z}} (-1)^k q^{k^2}.$$

The identity (5.7) is a special case of the Jacobi triple product identity; see [And65].

5.2.2. Decomposition through contractible CLE_4 on annular domains. Here we will see that the Miller-Sheffield coupling has an analogue on annular domains. First, we present the construction of the gauge twisted GFF on the annulus.

We start by considering a circular annulus $\widehat{A} = A_{r,1}$, with as previously $L = \text{ED}(\partial_o \widehat{A}, \partial_i \widehat{A}) = \frac{1}{2\pi} \log r^{-1}$. Given an angle $\theta \in [0, 2\pi)$, we will denote by γ_θ the ray $(ue^{i\theta})_{r \leq u \leq 1}$. Let σ_θ be the $\{-1, 1\}$ -valued gauge field on \widehat{A} corresponding to the *defect line* γ_θ : Given a continuous path $(\varphi(t))_{0 \leq t \leq T}$ in \widehat{A} with $\varphi(0), \varphi(T) \notin \gamma_\theta$, the holonomy $\text{hol}^{\sigma_\theta}(\varphi)$ is defined as follows. Through the log map, one can lift φ to a continuous path $\widehat{\varphi}$ on the universal cover S (5.5) of \widehat{A} , with endpoints $u(0)e^{iv(0)}$ and $u(T)e^{iv(T)}$. If there are an even number of points of $2\pi\mathbb{Z} + \theta$ between $v(0)$ and $v(T)$, then $\text{hol}^{\sigma_\theta}(\varphi) = 1$. If this number is odd, then $\text{hol}^{\sigma_\theta}(\varphi) = -1$. Roughly speaking, $\text{hol}^{\sigma_\theta}(\varphi)$ is given by the parity of the number of crossings of the defect line γ_θ by the path φ , with the caveat that this number may well be infinite, as for instance in the case of φ being a Brownian path; still the parity is well defined even in the case of infinite crossings. Now, if the path φ is a closed loop ($\varphi(0) = \varphi(T)$), then one can remove the restrictive condition $\varphi(0), \varphi(T) \notin \gamma_\theta$. In this case, $\text{hol}^{\sigma_\theta}(\varphi) = (-1)^{(v(T)-v(0))/2\pi}$.

One immediately remarks that for a closed loop φ , the holonomy $\text{hol}^{\sigma_\theta}(\varphi)$ is the same whatever the value of θ . And indeed, all the gauge fields σ_θ belong to the same gauge equivalence class. Given $0 \leq \theta_1 < \theta_2 < 2\pi$, define $\widehat{\sigma}_{\theta_1, \theta_2}$ on $\widehat{A} \setminus (\gamma_{\theta_1} \cup \gamma_{\theta_2})$ as follows: $\widehat{\sigma}_{\theta_1, \theta_2}$ equals 1 on $\{ue^{iv} | r < u < 1, 0 \leq v < \theta_1 \text{ or } \theta_2 < v < 2\pi\}$, and equals -1 on $\{ue^{iv} | r < u < 1, \theta_1 < v < \theta_2\}$. Then for every continuous path $(\varphi(t))_{0 \leq t \leq T}$ in \widehat{A} (not necessarily a closed loop), with $\varphi(0), \varphi(T) \notin \gamma_{\theta_1} \cup \gamma_{\theta_2}$,

$$\text{hol}^{\sigma_{\theta_2}}(\varphi) = \widehat{\sigma}_{\theta_1, \theta_2}(\varphi(0)) \text{hol}^{\sigma_{\theta_1}}(\varphi) \widehat{\sigma}_{\theta_1, \theta_2}(\varphi(T)).$$

Further, the defect line does not have at all to be a straight ray γ_θ . One can take any continuous simple curve $(\gamma(s))_{0 \leq s \leq 1}$ such that $|\gamma(0)| = r$, $|\gamma(1)| = 1$, and $|\gamma(s)| \in (r, 1)$ for every $s \in (0, 1)$. On the universal cover S , the connected components of $(\exp)^{-1}(\widehat{A} \setminus \gamma)$ are naturally ordered: one can define a function $\text{ord} : (\exp)^{-1}(\widehat{A} \setminus \gamma) \mapsto \mathbb{Z}$ which is constant on each connected component of $(\exp)^{-1}(\widehat{A} \setminus \gamma)$ and such that for every $w \in (\exp)^{-1}(\widehat{A} \setminus \gamma)$, $\text{ord}(w + 2\pi i) = \text{ord}(w) + 1$. Then the gauge field σ_γ associated to the defect line φ is defined as

follows. Let \wp be a continuous path $(\wp(t))_{0 \leq t \leq T}$ in \widehat{A} with $\wp(0), \wp(T) \notin \gamma$. Through the log map, we lift \wp to a continuous path $\hat{\wp}$ on the universal cover S of \widehat{A} . Then

$$\text{hol}^{\sigma_\gamma}(\wp) = (-1)^{\text{ord}(\hat{\wp}(t)) - \text{ord}(\hat{\wp}(0))}.$$

The gauge field σ_γ is in the same gauge equivalence class as all the σ_θ .

Let \widehat{A}^{db} be the circular annulus $\widehat{A}^{\text{db}} = A_{\sqrt{r}, 1}$. Then the square map $\pi(w) = w^2$ induces a double cover of \widehat{A} by \widehat{A}^{db} . For this covering map π , the covering automorphism that interchanges the two sheets of the covering is simply given by $\psi(w) = -w$. The maps π and ψ are both holomorphic, a fact that we will need later. Let $\widehat{\Phi}^{\text{db}}$ be the continuum GFF on \widehat{A}^{db} with 0 boundary conditions, normalized so that its covariance function is the Green's function of $-\Delta$ with 0 Dirichlet boundary conditions, denoted $G_{\widehat{A}^{\text{db}}}(z, w)$. As in Sections 3.2 and 3.3, let be

$$\widehat{\Phi}_+^{\text{db}} = \frac{1}{\sqrt{2}}(\widehat{\Phi}^{\text{db}} + \widehat{\Phi}^{\text{db}} \circ \psi), \quad \widehat{\Phi}_-^{\text{db}} = \frac{1}{\sqrt{2}}(\widehat{\Phi}^{\text{db}} - \widehat{\Phi}^{\text{db}} \circ \psi).$$

Note that the fields $\widehat{\Phi}_+^{\text{db}}$ and $\widehat{\Phi}_-^{\text{db}}$ are independent.

From now on we will assume that the defect line γ has zero Lebesgue (area) measure. We want to construct $\widehat{\Phi}_{\sigma_\gamma}$, the continuum GFF on \widehat{A} twisted by the gauge field σ_γ , but we do not want to specify the values of $\widehat{\Phi}_{\sigma_\gamma}$ on the defect line γ itself. So the zero Lebesgue measure ensures that the specification on γ does not matter. Note that this condition is not automatically satisfied by continuous simple curves. As a counterexample, there are the so called Osgood curves; see [Sag94, Chapter 8]. Let $s_\gamma : \widehat{A} \setminus \gamma \rightarrow \widehat{A}^{\text{db}}$ be a section of the covering map π (for all $z \in \widehat{A} \setminus \gamma$, $\pi(s_\gamma(z)) = z$) that is continuous on $\widehat{A} \setminus \gamma$. The section s_γ is a determination of the square root on $\widehat{A} \setminus \gamma$, and in particular it is holomorphic.

Since s_γ and ψ are holomorphic, the field $\widehat{\Phi}_+^{\text{db}} \circ s_\gamma$ is distributed like the usual continuum GFF on \widehat{A} with 0 boundary conditions, $\widehat{\Phi}$. Now define $\widehat{\Phi}_{\sigma_\gamma} = \widehat{\Phi}_-^{\text{db}} \circ s_\gamma$. Then $\widehat{\Phi}_{\sigma_\gamma}$ is a Gaussian field on \widehat{A} with covariance function given by

$$G_{\widehat{A}, \sigma_\gamma}(z, w) = G_{\widehat{A}^{\text{db}}}(s_\gamma(z), s_\gamma(w)) - G_{\widehat{A}^{\text{db}}}(s_\gamma(z), \psi(s_\gamma(w))).$$

The field $\widehat{\Phi}_{\sigma_\gamma}$ is the σ_γ -gauge-twisted GFF on \widehat{A} with 0 boundary conditions. Note the discontinuity of $G_{\widehat{A}, \sigma_\gamma}(z, w)$ when either of the variables crosses the defect line γ .

Regarding the regularity of the field $\widehat{\Phi}_{\sigma_\gamma}$, one can work in the Sobolev space $\mathcal{H}^{-1}(\widehat{A})$; see [She07, Section 2.3] and [Dub09, Section 4.2]. The $\mathcal{H}^{-1}(\widehat{A})$ norm is given by

$$\|f\|_{\mathcal{H}^{-1}(\widehat{A})}^2 = \iint_{\widehat{A} \times \widehat{A}} f(z) G_{\widehat{A}}(z, w) f(w) d^2z d^2w,$$

where $G_{\widehat{A}}$ is the Green's function of $-\Delta$ on \widehat{A} with 0 boundary conditions. Then $\widehat{\Phi}_{\sigma_\gamma}$ can be seen as a random element of $\mathcal{H}^{-1}(\widehat{A})$, with

$$\mathbb{E}[\|\widehat{\Phi}_{\sigma_\gamma}\|_{\mathcal{H}^{-1}(\widehat{A})}^2] \leq \mathbb{E}[\|\widehat{\Phi}\|_{\mathcal{H}^{-1}(\widehat{A})}^2] < +\infty.$$

This is due to the fact that

$$\iint_{\widehat{A} \times \widehat{A}} G_{\widehat{A}}(z, w) |G_{\widehat{A}, \sigma_\gamma}(z, w)| d^2z d^2w \leq \iint_{\widehat{A} \times \widehat{A}} G_{\widehat{A}}(z, w)^2 d^2z d^2w < +\infty.$$

Now consider the case of a general annular domain A that is conformally equivalent to \widehat{A} . Let $\varrho : A \rightarrow \widehat{A}$ be a conformal mapping. Let $(\gamma(s))_{0 \leq s \leq 1}$ be a continuous simple curve such that $\gamma(0) \in \partial_i A$, $\gamma(1) \in \partial_o A$, and $\gamma(s) \in A$ for every $s \in (0, 1)$. We also assume that γ has 0 Lebesgue measure. The gauge-twisted GFF Φ_{σ_γ} on A associated to the defect line γ , with 0 boundary conditions, can be constructed as $\Phi_{\sigma_\gamma} = \widehat{\Phi}_{\sigma_{\varrho(\gamma)}} \circ \varrho$. For the continuous extension of ϱ along γ up to the extremities $\gamma(0)$ and $\gamma(1)$, we refer to [Pom92, Proposition 2.14].

Next we would like to point out that Corollary 3.25 has a continuum version that involves the contractible CLE_4 conformal loop ensemble on annular domains. It can be interpreted as a gauge-twisted version of the Miller-Sheffield coupling. For the original Miller-Sheffield on simply connected domains we refer to [WW17, ASW19]. The contractible version of CLE_4 on annular domains has been first introduced in [SWW17].

So let A be an annular domain as previously. The contractible CLE_4 on A is a random countable collection \mathfrak{C} of Jordan loops in A where each loop is simple, contractible (does not surround the inner hole of A), and any two different loops do not intersect and do not surround one another. Each loop looks locally like and SLE_4 . Given a loop $\mathbf{C} \in \mathfrak{C}$, we will denote by $\text{Int}(\mathbf{C})$ the open domain surrounded by \mathbf{C} . The domain $\text{Int}(\mathbf{C})$ is simply connected by construction.

Now consider the following additional random objects.

- Let $(\Phi_{\mathbf{C}})_{\mathbf{C} \in \mathfrak{C}}$ be a family of fields such that each $\Phi_{\mathbf{C}}$ is distributed, conditionally on \mathfrak{C} , as a GFF on $\text{Int}(\mathbf{C})$ with 0 boundary conditions on \mathbf{C} (and 0 outside $\text{Int}(\mathbf{C})$); and with the fields being conditionally independent given \mathfrak{C} .
- Let $(\zeta_{\mathbf{C}})_{\mathbf{C} \in \mathfrak{C}}$ be random family of sign $(\zeta_{\mathbf{C}} \in \{-1, 1\})$ that are, conditionally on \mathfrak{C} , i.i.d. and uniform $(\mathbb{P}(\zeta_{\mathbf{C}} = 1) = \mathbb{P}(\zeta_{\mathbf{C}} = -1) = 1/2)$, and independent of the fields $(\Phi_{\mathbf{C}})_{\mathbf{C} \in \mathfrak{C}}$.

Recall that 2λ denotes the height gap of the GFF. With our normalization, $2\lambda = \sqrt{\pi/2}$. Let Φ^* be the field on A obtained as

$$\Phi^* = \sum_{\mathbf{C} \in \mathfrak{C}} \mathbf{1}_{\text{Int}(\mathbf{C})} (\Phi_{\mathbf{C}} + \zeta_{\mathbf{C}} 2\lambda).$$

Unlike in the Miller-Sheffield coupling on simply connected domains, on the annular domain A , Φ^* is not distributed as a GFF, but as a conditioned GFF. Given the 0 boundary GFF Φ on A , let $E_{\text{contractible}}$ be the event that Φ has no non-contractible $\pm 2\lambda$ -level lines. The event $E_{\text{contractible}}$ is exactly the event $\{N = 0\}$ from Section 5.2.1. Then the field Φ^* is distributed as the GFF Φ conditioned on the event $E_{\text{contractible}}$; see [ALS22, Section 4.4].

The conditioned GFF Φ^* is *a priori* non-Gaussian. However, we shall see that there is a transformation that makes Φ^* into a Gaussian field. As previously, let γ be a (deterministic) defect line on A with 0 Lebesgue measure. Denote

$$\mathfrak{C}_{\gamma} = \{\mathbf{C} \in \mathfrak{C} \mid \text{Int}(\mathbf{C}) \cap \gamma \neq \emptyset\}.$$

Next we argue that the defect line γ induces a bipartite structure on $\text{Int}(\mathbf{C})$ for each loop $\mathbf{C} \in \mathfrak{C}_{\gamma}$. Let $v : S \rightarrow A$ be a $2\pi i$ -periodic covering map of the annular domain A by the strip S (5.5) (universal cover). For instance, one can take $v = \varrho^{-1} \circ \exp$, where as previously, $\varrho : A \rightarrow \widehat{A}$ is a conformal mapping from the annular domain A to the circular annulus \widehat{A} . Let $\text{ord} : v^{-1}(A \setminus \gamma) \rightarrow \mathbb{Z}$ be an order on the connected components of $v^{-1}(A \setminus \gamma)$ such that $\text{ord}(w + 2\pi i) = \text{ord}(w) + 1$. Given a CLE loop $\mathbf{C} \in \mathfrak{C}$, since \mathbf{C} is contractible, any lift of \mathbf{C} to S is a Jordan loop in S , and the different lifts differ by $2\pi k i$ translations for $k \in \mathbb{Z}$. For each $\mathbf{C} \in \mathfrak{C}$, we choose a particular lift, denoted $\text{Lift}(\mathbf{C})$, given by the condition

$$\inf\{\text{ord}(w) \mid w \in \text{Int}(\text{Lift}(\mathbf{C}))\} = 0,$$

where $\text{Int}(\text{Lift}(\mathbf{C}))$ is the interior surrounded by $\text{Lift}(\mathbf{C})$ in S . This convention is somewhat arbitrary, but we need to choose a particular lift one way or another. For $\mathbf{C} \in \mathfrak{C} \setminus \mathfrak{C}_{\gamma}$, ord is constant equal to 0 on $\text{Int}(\text{Lift}(\mathbf{C}))$. However, for $\mathbf{C} \in \mathfrak{C}_{\gamma}$, ord is not constant on $\text{Int}(\text{Lift}(\mathbf{C})) \setminus v^{-1}(\gamma)$. For $z \in \text{Int}(\mathbf{C})$, we will denote by $v_0^{-1}(z)$ the unique point in $\text{Int}(\text{Lift}(\mathbf{C})) \cap v^{-1}(\{z\})$. Let us define the function $\zeta_{\mathfrak{C}, \gamma} : A \setminus \gamma \rightarrow \{-1, 1\}$ as follows. The function $\zeta_{\mathfrak{C}, \gamma}$ equals 1 on

$$(A \setminus \gamma) \setminus \bigcup_{\mathbf{C} \in \mathfrak{C}_{\gamma}} \text{Int}(\mathbf{C}).$$

For $\mathbf{C} \in \mathfrak{C}_{\gamma}$ and $z \in \text{Int}(\mathbf{C}) \setminus \gamma$, we set

$$\zeta_{\mathfrak{C}, \gamma}(z) = (-1)^{\text{ord}(v_0^{-1}(z))}.$$




FIGURE 9. Conceptual depiction of the signs $\zeta_{\mathfrak{C},\gamma}$. The defect line γ is represented in violet dashed.

Now, this definition of $\zeta_{\mathfrak{C},\gamma}$ may look abstract. What we are simply saying is that the sign $\zeta_{\mathfrak{C},\gamma}(z)$ changes to the opposite as z crosses the defect line γ by moving inside $\text{Int}(\mathfrak{C})$. See Figure 9 for an illustration. The sign function $\zeta_{\mathfrak{C},\gamma}$ only depends on the defect line γ and on the realization of \mathfrak{C} . By taking Corollary 3.25 to the scaling limit, one gets the following.

Corollary 5.4. *The field $\zeta_{\mathfrak{C},\gamma}\Phi^*$ is distributed as the gauge-twisted GFF Φ_{σ_γ} , and in particular, it is Gaussian.*

The convergence from metric graph to the continuum limit in dimension 2 is at this point standard; see [Lup18, ALS20, ALS23].

Corollary 5.4 in particular implies that conditionally on the event $E_{\text{contractible}}$, the even local functions of the GFF Φ are related to even local functions of the gauge-twisted GFF Φ_{σ_γ} . Let us first detail the case of the renormalized hyperbolic cosine.

For $\beta \in (-2\sqrt{2\pi}, 2\sqrt{2\pi})$, the renormalized exponential of Φ (Gaussian multiplicative chaos, GMC) is given by the limit

$$(5.8) \quad : e^{\beta\Phi} := \lim_{\varepsilon \rightarrow 0} e^{\beta\Phi_\varepsilon - \frac{\beta^2}{2} \text{Var}(\Phi_\varepsilon)},$$

where Φ_ε are mollifications of Φ . The renormalized exponential $: e^{\beta\Phi} :$ is actually a random finite Borel measure on A . We refer to [RV14, Ber17, DS11] for details. The unusual range $(-2\sqrt{2\pi}, 2\sqrt{2\pi})$ for β is due to our choice of normalization for Φ , which is different from the one usually used in the GMC literature. Similarly, for $\beta \in (-2\sqrt{2\pi}, 2\sqrt{2\pi})$, one can define the renormalized exponential of Φ_{σ_γ} :

$$(5.9) \quad : e^{\beta\Phi_{\sigma_\gamma}} := \lim_{\varepsilon \rightarrow 0} e^{\beta\Phi_{\sigma_\gamma,\varepsilon} - \frac{\beta^2}{2} \text{Var}(\Phi_{\sigma_\gamma,\varepsilon})},$$

where $\Phi_{\sigma_\gamma,\varepsilon}$ are mollifications of Φ_{σ_γ} . Indeed, Φ_{σ_γ} is just another logarithmically correlated Gaussian field, with

$$G_{A,\sigma_\gamma}(z, w) \stackrel{w \rightarrow z}{\sim} G_A(z, w),$$

and the finite discontinuity of G_{A,σ_γ} on the defect line γ is not an issue. Note however that the renormalization factors in (5.8) and (5.9) differ by a bounded factor. The relevant quantity is given by

$$D_A(z) = \lim_{w \rightarrow z} G_A(z, w) - G_{A,\sigma_\gamma}(z, w), \quad z \in A.$$

In case of a circular annulus \widehat{A} ,

$$D_{\widehat{A}}(\boldsymbol{\pi}(w)) = 2G_{\widehat{A}^{\text{db}}}(w, \psi(w)) = 2G_{\widehat{A}^{\text{db}}}(w, -w) = 2G_{\widehat{A}^{\text{db}}}(|w|, -|w|).$$

Further, given a uniformization $\varrho : A \rightarrow \widehat{A}$,

$$D_A(z) = D_{\widehat{A}}(\varrho(z)).$$

In particular, $D_A(z)$ depends only on the conformal type of the domain with one marked interior point (A, z) , and $D_A(z) > 0$. Also note that $D_A(z)$ does not depend on the particular choice of the defect line γ . Further,

$$D_A(z) = \lim_{\varepsilon \rightarrow 0} \text{Var}(\Phi_\varepsilon(z)) - \text{Var}(\Phi_{\sigma_\gamma, \varepsilon}(z)).$$

Now consider the renormalized hyperbolic cosines $\frac{1}{2}(: e^{\beta\Phi} : + : e^{-\beta\Phi} :)$ and $\frac{1}{2}(: e^{\beta\Phi_{\sigma_\gamma}} : + : e^{-\beta\Phi_{\sigma_\gamma}} :)$. While the fields $\Phi_{\sigma_\gamma} : e^{\beta\Phi_{\sigma_\gamma}} :$ and $: e^{-\beta\Phi_{\sigma_\gamma}} :$ each depend on the particular choice of the defect line γ , the hyperbolic cosine $\frac{1}{2}(: e^{\beta\Phi_{\sigma_\gamma}} : + : e^{-\beta\Phi_{\sigma_\gamma}} :)$ does not. From Corollary 5.4, and also directly from Theorem 1 by taking the scaling limit, we obtain the following

Corollary 5.5. *Conditionally on the event $E_{\text{contractible}}$, the family of random measures $(\frac{1}{2}(: e^{\beta\Phi} : + : e^{-\beta\Phi} :))_{0 \leq \beta < 2\sqrt{2\pi}}$ is distributed as*

$$(5.10) \quad \left(\frac{1}{2} e^{-\frac{\beta^2}{2} D_A} (: e^{\beta\Phi_{\sigma_\gamma}} : + : e^{-\beta\Phi_{\sigma_\gamma}} :) \right)_{0 \leq \beta < 2\sqrt{2\pi}}.$$

We further consider the renormalized (Wick) powers. Let $(H_n)_{n \geq 0}$ be the family of probabilistic Hermite polynomials, that is to say unitary polynomials orthogonal for the Gaussian measure $e^{-\frac{1}{2}x^2} dx$. One can explicitly write

$$H_n(x) = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \frac{n!}{2^k k! (n-2k)!} x^{n-2k}.$$

One obtains the Wick powers as the limits

$$\begin{aligned} : \Phi^n : &= \lim_{\varepsilon \rightarrow 0} \text{Var}(\Phi_\varepsilon)^{n/2} H_n(\text{Var}(\Phi_\varepsilon)^{-1/2} \Phi_\varepsilon), \\ : \Phi_{\sigma_\gamma}^n : &= \lim_{\varepsilon \rightarrow 0} \text{Var}(\Phi_{\sigma_\gamma, \varepsilon})^{n/2} H_n(\text{Var}(\Phi_{\sigma_\gamma, \varepsilon})^{-1/2} \Phi_{\sigma_\gamma, \varepsilon}). \end{aligned}$$

We refer to [Sim74, Jan97, LJ11, LRV17]. The Wick powers are random generalized functions. The degree 0 power is the constant 1, and the degree 1 power is the Gaussian field itself. Note that the even Wick powers are not positive (except for $n = 0$) because of the counterterms in the renormalization. For the gauge-twisted GFF Φ_{σ_γ} , the odd Wick powers $: \Phi_{\sigma_\gamma}^{2k+1} :$ depend on the particular choice of the defect line γ , but the even Wick powers $: \Phi_{\sigma_\gamma}^{2k} :$ do not.

For $|\beta| < \sqrt{2\pi}$ (the L^2 regime), one can expand the renormalized exponential into the Wick powers:

$$: e^{\beta\Phi} : := \sum_{n \geq 0} \frac{\beta^n}{n!} : \Phi^n :, \quad : e^{\beta\Phi_{\sigma_\gamma}} : := \sum_{n \geq 0} \frac{\beta^n}{n!} : \Phi_{\sigma_\gamma}^n :.$$

See [LRV17, Section 2.3.1]. In particular,

$$\frac{1}{2}(: e^{\beta\Phi} : + : e^{-\beta\Phi} :) = \sum_{k \geq 0} \frac{\beta^{2k}}{(2k)!} : \Phi^{2k} :,$$

and similarly for Φ_{σ_γ} . Note however, that in the expansion of (5.10), one needs to take into account the contribution of $e^{-\frac{\beta^2}{2}D_A}$. We have that

$$\frac{1}{2}e^{-\frac{\beta^2}{2}D_A}(: e^{\beta\Phi_{\sigma_\gamma}} : + : e^{-\beta\Phi_{\sigma_\gamma}} :) = \sum_{k \geq 0} \frac{\beta^{2k}}{(2k)!} \sum_{j=0}^k (-1)^{k-j} \frac{(2k)!}{2^{k-j}(k-j)!(2j)!} D_A^{k-j} : \Phi_{\sigma_\gamma}^{2j} : .$$

By identifying the terms in the expansions, we get the following.

Corollary 5.6. *Conditionally on the event $E_{\text{contractible}}$, the family of random fields $(: \Phi^{2k} :)_{k \geq 1}$ is jointly distributed as*

$$\left(\sum_{j=0}^k (-1)^{k-j} \frac{(2k)!}{2^{k-j}(k-j)!(2j)!} D_A^{k-j} : \Phi_{\sigma_\gamma}^{2j} : \right)_{k \geq 1} .$$

Remark 5.7. It would be interesting to have a proof of Corollaries 5.4, 5.5 and 5.6 directly in continuum, that does not rely on approximation by metric graphs.

Remark 5.8. Corollaries 5.5 and 5.6 are also true on more general two-dimensional domains, not necessarily annular, such as the multiply connected domains considered in Section 5.3.

5.3. The planar case with multiple holes. First, consider $D \subset \mathbb{C}$ an open bounded domain with two holes, i.e. $\mathbb{C} \setminus D$ having three connected components, two of them being bounded. We also assume that the holes are not reduced to one point (not punctures). Let $\pi_1(D)$ denote the fundamental group of D . It is isomorphic to \mathbb{F}_2 , the free group with two generators. In particular, there are $4 = 2^2$ group homomorphisms from $\pi_1(D)$ to $\{-1, 1\}$, one being the trivial homomorphism, and the three other being non-trivial. Each of these homomorphisms corresponds to a gauge equivalence class for $\{-1, 1\}$ -gauge fields on D that are locally trivial, but not globally trivial in general. On Figure 10 the three non-trivial equivalence classes are represented through defect lines: the holonomy is multiplied by -1 as one crosses a defect line. Therefore, Theorem 1 provides in the scaling limit the probabilities of three different topological events on the continuum GFF on D with 0 boundary conditions. These topological events deal with the sign excursion clusters of the GFF (in the language of [ALS23]), which are also the clusters of the Brownian loop soup on D with parameter $\alpha = 1/2$. The defect line on top left on Figure 10 detects the sign excursion clusters that separate the first hole of D from the outer boundary of D . The defect line on top right detects the sign excursion clusters that separate the second hole from the outer boundary. The defect line on the bottom detects the sign excursion clusters that separate one hole from the other. Note however that these three probabilities provide only a partial information on the distribution of the homotopical types of the clusters. For instance, Figure 11 depicts a cluster which is non-trivial simultaneously for all the three defect lines. Theorem 1 does not provide the probability of the existence of such a cluster.

More generally, consider $D \subset \mathbb{C}$ an open bounded domain with n holes, the holes not being reduced to points (not punctures). The fundamental group $\pi_1(D)$ is isomorphic to \mathbb{F}_n , the free group with n generators. In particular, there are 2^n group homomorphisms from $\pi_1(D)$ to $\{-1, 1\}$, one of them being the trivial homomorphism. Therefore, by applying Theorem 1 in the scaling limit, one gets access to probabilities of $2^n - 1$ different topological events. By contrast, the number of different homotopical types of a cluster in D grows superexponentially with n . Indeed, given K a connected compact subset of D , the complementary $D \setminus K$ induces a partition on the connected components of the boundary ∂D . Therefore, one looks at the number of partitions of $\{0, 1, \dots, n\}$ which is the Bell number B_{n+1} . We have the asymptotic log $B_{n+1} \sim n \log n$; see [dB70, Section 6.3].

In the works of Dubédat [Dub19] and then Basok-Chelkak [BC21], the authors obtained the probabilities of all possible homotopical events in a nested CLE_4 on a simply connected domain with finitely many punctures, through the use of isomonodromic deformations. To our knowledge, the probabilities of homotopical events in case of macroscopic holes and not punctures were not known. Our method gives some of them. Our method applies in general to compact bordered Riemann surfaces, but we do not detail this here.




FIGURE 10. Depiction of 3 non-equivalent gauge fields on a planar domain with 2 holes. The defect lines are in violet dashed.




FIGURE 11. Depiction in grey of a cluster that is non-trivial for simultaneously all the 3 defect lines on Figure 10.

5.4. **The case of dimensions $d \geq 3$.** As already mentioned several times, Theorem 1 does not require planarity, and for instance holds on lattices in any dimension. However, in a non-planar setting, the interpretation of the event $\{\tilde{\phi} \in \mathcal{T}_\sigma\}$ is more subtle, and this is something to keep in mind. We would like to illustrate this on an example.

Fix $d \geq 3$ and $n \geq 1$. Consider the hypercube $Q = (-4n, 4n)^d$ and the rectangular parallelepiped $Q' = (-n, n)^2 \times (-4n, 4n)^{d-2}$. Let D be the domain $D = Q \setminus \overline{Q'}$. Note that the fundamental group $\pi_1(D)$ is isomorphic to \mathbb{Z} , as a loop in D can surround several times the inner parallelepiped Q' . Let $\mathcal{G} = (V, E)$ be the following graph. We set $V = \overline{D} \cap \mathbb{Z}^d$, with $V_{\text{int}} = D \cap \mathbb{Z}^d$ and $V_\partial = \partial D \cap \mathbb{Z}^d$. The set of edges is given by $E = \{\{x, y\} | x, y \in V, \|y - x\| = 1\}$. The edges are endowed with unit conductance: $C(x, y) = 1$. Now consider the following gauge field σ . We set $\sigma(e) = -1$ for edges $e \in E$ of form

$$\{(k_1, 0, k_3, \dots, k_d), (k_1, 1, k_3, \dots, k_d)\}.$$

For all other edges e , we set $\sigma(e) = 1$. Given a closed loop φ in \mathcal{G} , the holonomy $\text{hol}^\sigma(\varphi)$ equals -1 if φ surrounds the inner parallelotope Q' an odd number of times, and it equals 1 if φ surrounds Q' and even number of times, including if φ does not surround Q' at all. For instance, consider the loop φ corresponding to a square with the following four corners:

$$(2n, -2n, 0, \dots, 0), (2n, 2n, 0, \dots, 0), (-2n, 2n, 0, \dots, 0), (-2n, -2n, 0, \dots, 0).$$

Then $\text{hol}^\sigma(\varphi) = -1$.

Now here is what distinguished the non-planar setting from the planar setting. For instance, on an annular-shaped metric graph as on Figure 1, if there is a connected subset \tilde{K} that surrounds the inner hole, then one can draw a loop φ in \tilde{K} that winds around the inner hole exactly once. But this is a feature of the planarity, that is not true in general. For instance, in our higher dimensional setting, we will give an example of a simple loop φ in \mathcal{G} that winds around Q' twice. Since φ is simple, it cannot have a sub-loop that would wind once around Q' . In particular, we simultaneously have that φ is not contractible in the continuum domain D and $\pi_\sigma^{-1}(\varphi)$ is not connected. Again, such counter-example is impossible in a planar setting: if a loop winds twice around the hole of an annular domain, it cannot be simple.

So here is our example. We will take points x_1, x_2, \dots, x_{10} in V and straight line segments $\wp_1, \wp_2, \dots, \wp_{10}$, where \wp_i is a line from x_i to x_{i+1} for $1 \leq i \leq 9$, and \wp_{10} is a line from x_{10} to x_1 . Our loop φ will be the concatenation

$$\varphi = \wp_1 \wedge \wp_2 \wedge \dots \wedge \wp_{10}.$$

So here are the coordinates of the points x_i :

$$\begin{aligned} x_1 &= (2n, -2n, 0, \dots, 0), & x_2 &= (2n, 2n, 0, \dots, 0), & x_3 &= (-2n, 2n, 0, \dots, 0), \\ x_4 &= (-2n, -3n, 0, \dots, 0), & x_5 &= (3n, -3n, 0, \dots, 0), & & \\ x_6 &= (3n, -3n, n, 0, \dots, 0), & x_7 &= (3n, 3n, n, 0, \dots, 0), & x_8 &= (-3n, 3n, n, 0, \dots, 0), \\ x_9 &= (-3n, -2n, n, 0, \dots, 0), & x_{10} &= (2n, -2n, n, 0, \dots, 0). \end{aligned}$$

The projection of φ on the first two coordinates winds twice around $(-n, n)^2$, but is not a simple loop. We use the third coordinate to avoid self-intersections.

The example above shows that the dichotomy on clusters induced by the gauge field σ is not contractible/non-contractible in D . The contractible clusters are indeed trivial for σ , but σ detects only a certain type of non-contractible clusters, those that allow to wind around Q' an odd number of times.

6. INTENSITY DOUBLING CONJECTURE IN HIGH DIMENSION

Let be a dimension $d \geq 2$. Let be the hypercube $Q = (-1, 1)^d$. Let $\mathcal{G}^{(n)} = (V^{(n)}, E^{(n)})$ be an approximation of Q in the rescaled hypercubic lattice $\frac{1}{n}\mathbb{Z}^d$. More precisely, we set $V^{(n)} = \bar{Q} \cap \frac{1}{n}\mathbb{Z}^d$, $V_{\text{int}}^{(n)} = Q \cap \frac{1}{n}\mathbb{Z}^d$, $V^{(n)} = (\partial Q) \cap \frac{1}{n}\mathbb{Z}^d$, and $E^{(n)} = \{\{x, y\} | x, y \in V^{(n)}, \|y - x\| = \frac{1}{n}\}$. All the conductances are uniform equal to $C(x, y) = \frac{1}{n^{d-2}}$. In this regime, $\tilde{\phi}^{(n)}$, the metric graph GFF on $\tilde{\mathcal{G}}^{(n)}$ with 0 boundary conditions, converges in law towards Φ , the continuum GFF on Q with 0 boundary conditions on ∂Q . Let $\tilde{\mathcal{L}}_n^{1/2}$ denote the metric graph loop soup on $\tilde{\mathcal{G}}^{(n)}$ that is related to $\tilde{\phi}^{(n)}$ through isomorphism (Theorem 2.6).

We will further consider $(d-2)$ -dimensional affine subspaces on \mathbb{R}^d , that is to say like linear, but without the condition of going through 0. Given \mathbf{A} such an affine subspace, and $\varepsilon > 0$, we will denote by $\text{Tube}_\varepsilon(\mathbf{A})$ the hypertube

$$\text{Tube}_\varepsilon(\mathbf{A}) = \{x \in \mathbb{R}^d | d(x, \mathbf{A}) \leq \varepsilon\}.$$

The fundamental group $\pi_1(\mathbb{R}^d \setminus \text{Tube}_\varepsilon(\mathbf{A}))$ is isomorphic to \mathbb{Z} , since a loop can wind around the hypertube $\text{Tube}_\varepsilon(\mathbf{A})$. The fundamental group $\pi_1(Q \setminus \text{Tube}_\varepsilon(\mathbf{A}))$ is also isomorphic to \mathbb{Z} , provided that $Q \cap \mathbf{A} \neq \emptyset$ and ε is small enough. We will denote by $\tilde{\mathcal{G}}_{\mathbf{A}, \varepsilon}^{(n)}$ a sub-metric-graph of $\tilde{\mathcal{G}}^{(n)}$ that reasonably approximates $Q \setminus \text{Tube}_\varepsilon(\mathbf{A})$ as $n \rightarrow +\infty$. We will denote by $\tilde{\phi}_{\mathbf{A}, \varepsilon}^{(n)}$ the metric graph

GFF on $\tilde{\mathcal{G}}_{A,\varepsilon}^{(n)}$ with 0 boundary conditions, that is to say 0 both on ∂Q and $\partial \text{Tube}_\varepsilon(A)$. On $\tilde{\mathcal{G}}_{A,\varepsilon}^{(n)}$, one can choose a $\{-1, 1\}$ gauge field $\sigma_{n,A,\varepsilon}$ that detects the odd number of windings around $\text{Tube}_\varepsilon(A)$ (see also Section 5.4), provided that it is at all possible to wind around $\text{Tube}_\varepsilon(A)$ in Q . By applying Theorem 1 to $\sigma_{n,A,\varepsilon}$, we get the value of

$$(6.1) \quad \mathbb{P}(\exists \mathcal{C} \text{ sign cluster of } \tilde{\phi}_{A,\varepsilon}^{(n)} \text{ with an odd winding around } \text{Tube}_\varepsilon(A)).$$

Further, one can express the limit of (6.1) in terms of Brownian loop soups in Q . For this we first recall the Brownian loop measure on \mathbb{R}^d .

The natural Brownian loop measure on \mathbb{R}^d is given by

$$\mu_{\mathbb{R}^d}^{\text{loop}}(d\wp) = \int_{\mathbb{R}^d} \int_0^{+\infty} \mathbb{P}_{\mathbb{R}^d,t}^{x,x}(d\wp) \frac{dt}{(2\pi)^{d/2} t^{d/2+1}} dx,$$

where $\mathbb{P}_{\mathbb{R}^d,t}^{x,x}$ is the Brownian bridge probability measure in time t from x to x . The measure is obviously invariant by translations and rotations, but it is also scale invariant, provided one scales both space and time according to the Brownian scaling. This scaling invariance imposes the power $t^{-(d/2+1)}$. The total mass of $\mu_{\mathbb{R}^d}^{\text{loop}}$ is infinite. However, it gives a finite mass to families of loops of form $\{\wp \text{ loop} \mid \wp \subset (-R, R)^d, \text{diam}(\wp) \geq \delta\}$. Given D an open subset of \mathbb{R}^d , we will denote by μ_D^{loop} the restriction of the measure $\mu_{\mathbb{R}^d}^{\text{loop}}$ to the loops that stay in D . Note that this is consistent with the previous definition (5.1) in dimension 2. Given $\alpha > 0$, we will denote by \mathcal{L}_D^α the Poisson point process of Brownian loops in D with intensity $\alpha \mu_D^{\text{loop}}$. In particular, for every $\delta > 0$, the Poisson collection $\{\wp \in \tilde{\mathcal{L}}_n^{1/2} \mid \text{diam}(\wp) > \delta\}$ converges in law as $n \rightarrow +\infty$ towards $\{\wp \in \mathcal{L}_Q^{1/2} \mid \text{diam}(\wp) > \delta\}$.

Now, the limit of (6.1) converges as $n \rightarrow +\infty$ towards

$$1 - \exp\left(-\mu_Q^{\text{loop}}\{\wp \text{ loop} \mid \wp \cap \text{Tube}_\varepsilon(A) = \emptyset, \wp \text{ has an odd winding around } \text{Tube}_\varepsilon(A)\}\right).$$

This in turn can be rewritten as

$$(6.2) \quad \mathbb{P}(\exists \wp \in \mathcal{L}_Q^1 \text{ with an odd winding around } \text{Tube}_\varepsilon(A)).$$

Note that the intensity parameter α that appears in (6.2) is not $\alpha = 1/2$ as in the isomorphism theorems (Theorems 2.4 and 2.6), but the double, $\alpha = 1$. This has been already noted in our Remark 3.17, and further developed in the identity (3.16). Now, in general, the sign clusters of a metric graph GFF are not Poisson, as they repel each other. This is for instance understood precisely in dimension 2 with the CLE_4 . However, in high enough dimension d , the sign clusters should be in a sense approximately Poisson, as conjectured by Werner in [Wer21]. Here we will argue that as soon as $d > 8$ (and perhaps as soon as $d > 6$), asymptotically, as $n \rightarrow +\infty$, the clusters formed by the microscopic loops in $\tilde{\mathcal{L}}_n^{1/2}$ give rise to an independent copy of the macroscopic loops of $\tilde{\mathcal{L}}_n^{1/2}$. Previously, the case of dimensions $d > 6$ has been studied both at an heuristic and rigorous level by Werner [Wer21] and Cai and Ding [CD23]. Our conjecture emerged as an attempt to conciliate the above remark with the vision of the high-dimensional picture sketched by Werner and further confirmed by Cai and Ding. It is also natural in view of the identity (3.16), where the intensity $\alpha = 1$ appears explicitly.

To state our conjecture precisely, we will need a notion of size for the loops which is different from the diameter. We will call it the *encompass*. For \wp continuous loop in \mathbb{R}^d , set

$$\text{enc}(\wp) = \sup\{\varepsilon > 0 \mid \exists \mathbf{A} \text{ affine subspace of } \mathbb{R}^d, \dim \mathbf{A} = d - 2, \\ \wp \cap \text{Tube}_\varepsilon(\mathbf{A}) = \emptyset, \wp \text{ non-contractible in } \mathbb{R}^d \setminus \text{Tube}_\varepsilon(\mathbf{A})\}.$$

Note that trivially, $\text{diam}(\wp) \geq 2 \text{enc}(\wp)$. But a converse control does not hold. Even a simple loop with a fixed diameter can have an arbitrarily small encompass. Further, given a compact connected subset $K \subset \mathbb{R}^d$, we will denote

$$\widehat{\text{enc}}(K) = \sup\{\text{enc}(\wp) \mid \wp \text{ continuous loop, } \wp \subset K\}.$$

If K does not contain a loop with positive encompass, then we set $\widehat{\text{enc}}(K) = 0$. Again, $\text{diam}(2) \geq 2\widehat{\text{enc}}(K)$. But in the other direction, a tree, however large, has a zero $\widehat{\text{enc}}$. Note that if $\widehat{\text{enc}}(K) > \varepsilon$, this does not mean that there is an affine subspace A of dimension $d - 2$ such that $K \cap \text{Tube}_\varepsilon(A) = \emptyset$ and K non-contractible in $\mathbb{R}^d \setminus \text{Tube}_\varepsilon(A)$. In particular, for a continuous but non-simple loop φ , the quantities $\text{enc}(\varphi)$ and $\widehat{\text{enc}}(\varphi)$ are not necessarily equal, and $\widehat{\text{enc}}(\varphi)$ may be strictly larger than $\text{enc}(\varphi)$ if φ contains a sub-loop with a larger encompass. As an other example, the $\widehat{\text{enc}}$ of a ball equals its radius.

Conjecture 2 (Intensity doubling). *Assume that $d > 8$. Fix $\varepsilon > 0$. Consider the following two families of random objects. First, the family of loops*

$$(6.3) \quad \{\varphi \in \widetilde{\mathcal{L}}_n^{1/2} \mid \text{enc}(\varphi) > \varepsilon\}.$$

Second, the family of clusters

$$(6.4) \quad \{\mathcal{C} \text{ cluster of } \widetilde{\mathcal{L}}_n^{1/2} \mid \widehat{\text{enc}}(\mathcal{C}) > \varepsilon\}.$$

Note that by definition, the clusters corresponding to the loops in (6.3) belong to the family (6.4). We conjecture that as $n \rightarrow +\infty$, (6.3) and (6.4) jointly converge in law for the Hausdorff distance towards $(\mathcal{L}_{\infty,\varepsilon}, \mathcal{K}_{\infty,\varepsilon})$ with the following properties.

- (1) $\mathcal{L}_{\infty,\varepsilon}$ is a Poisson point process of Brownian loop with intensity measure

$$\frac{1}{2} \times \mathbf{1}_{\text{enc } \varphi > \varepsilon} \mu_Q^{\text{loop}}(d\varphi).$$

This point, concerning the convergence of (6.3) alone, is known.

- (2) $\mathcal{K}_{\infty,\varepsilon}$ is an a.s. finite Poisson point process of compact connected subsets of \overline{Q} .
(3) Each $K \in \mathcal{K}_{\infty,\varepsilon}$ is of dimension 4 (Hausdorff and Minkowski), and in particular is polar for the Brownian motion.
(4) The compacts in $\mathcal{K}_{\infty,\varepsilon}$ are pairwise disjoint.
(5) Each $K \in \mathcal{K}_{\infty,\varepsilon}$ contains a single simple loop which we denote $\text{Loop}(K)$. We will also denote $\text{Loop}(\mathcal{K}_{\infty,\varepsilon}) = \{\text{Loop}(K) \mid K \in \mathcal{K}_{\infty,\varepsilon}\}$.
(6) We have that $\mathcal{L}_{\infty,\varepsilon} \subset \text{Loop}(\mathcal{K}_{\infty,\varepsilon})$.
(7) The family of loops $\text{Loop}(\mathcal{K}_{\infty,\varepsilon})$ is distributed as a Poisson point process with intensity

$$\mathbf{1} \times \mathbf{1}_{\text{enc } \varphi > \varepsilon} \mu_Q^{\text{loop}}(d\varphi).$$

In particular, with positive probability, $\mathcal{L}_{\infty,\varepsilon}$ is a strict subset of $\text{Loop}(\mathcal{K}_{\infty,\varepsilon})$.

- (8) *The two subfamilies $\{K \in \mathcal{K}_{\infty,\varepsilon} \mid \text{Loop}(K) \in \mathcal{L}_{\infty,\varepsilon}\}$ and $\{K \in \mathcal{K}_{\infty,\varepsilon} \mid \text{Loop}(K) \notin \mathcal{L}_{\infty,\varepsilon}\}$ have the same distribution and are independent.*

The dimension 4 is that of the Integrated Super-Brownian Excursions (ISE) [DP98, LG99], which is conjectured to appear in the scaling limit of different critical lattice models in high dimension [Sla99]. Here we took $d > 8 = 4+4$ to ensure that the compacts in $\mathcal{K}_{\infty,\varepsilon}$ stay pairwise disjoint. We believe that the Poisson structure for the clusters in the scaling limit could already emerge in dimensions $d \in \{7, 8\}$. Since the clusters are expected to be asymptotically polar also in these dimensions, and thus invisible for the GFF, there should be no macroscopic repulsion between clusters. However, for $d \in \{7, 8\}$, two disjoint clusters might intersect in the scaling limit, and a cluster might also get additional cycles in the scaling limit that are not there at the metric graph level. Yet, if one focuses solely on the cycles that are already there at the metric graph level, the intensity doubling should also occur in dimensions $d \in \{7, 8\}$.

Next we bring arguments for Conjecture 2. The first question is whether it provides the correct limit (6.2). One should pay attention that in (6.1) appears not the GFF on $\widetilde{\mathcal{G}}^{(n)}$ but the GFF on $\widetilde{\mathcal{G}}_{A,\varepsilon}^{(n)}$. However, if K is a compact in $\mathcal{K}_{\infty,\varepsilon_0}$ such that $\text{Loop}(K) \cap \text{Tube}_\varepsilon(A) = \emptyset$, then the connected component of $\text{Loop}(K)$ in $\overline{K \setminus \text{Tube}_\varepsilon(A)}$ is the scaling limit of a sign cluster of the GFF on $\widetilde{\mathcal{G}}_{A,\varepsilon}^{(n)}$. Indeed, in case $\text{Loop}(K) \in \mathcal{L}_{\infty,\varepsilon_0}$, the metric graph cluster approximating K contains just one macroscopic loop of $\widetilde{\mathcal{L}}_n^{1/2}$, the one approximating $\text{Loop}(K)$, and no macroscopic

loop of $\tilde{\mathcal{L}}_n^{1/2}$ in case $\text{Loop}(K) \notin \mathcal{L}_{\infty, \varepsilon_0}$. The rest of the cluster is made of microscopic loops. By removing the microscopic loops that intersect $\text{Tube}_\varepsilon(\mathbf{A})$, one changes the cluster only on the hypertube $\text{Tube}_\varepsilon(\mathbf{A})$ and its infinitesimal neighborhood. The fact that each cluster in (6.4) asymptotically contains at most one macroscopic loop of $\tilde{\mathcal{L}}_n^{1/2}$ is also consistent with the asymptotic polarity of the clusters.

Let be $d > 6$. For fixed $\delta > 0$, Werner showed that the number of clusters of $\tilde{\mathcal{L}}_n^{1/2}$ of diameter larger than δ is of order at least $n^{d-6+o(1)}$ [Wer21, Proposition 4]. So, in particular, the δ -large clusters proliferate as $n \rightarrow +\infty$. A typical large cluster is believed to look like a tree at a macroscopic scale, with a scaling limit describable in terms of superbrownian excursions. In contrast, according to our Conjecture 2, the number of clusters with large $\widehat{\text{enc}}$ is tight, and therefore such clusters are untypical among clusters of large diameter.

Now let us fix a hypertube $\text{Tube}_\varepsilon(\mathbf{A})$ such that $\pi_1(Q \setminus \text{Tube}_\varepsilon(\mathbf{A})) \cong \mathbb{Z}$. We look at the clusters in $\tilde{\mathcal{G}}_{\mathbf{A}, \varepsilon}^{(n)}$ of diameter larger than 2ε . The number of those diverges as $n \rightarrow +\infty$. As for the number of loops in $\tilde{\mathcal{L}}_n^{1/2}$ with an odd winding around $\text{Tube}_\varepsilon(\mathbf{A})$, it is tight. So the number of clusters in $\tilde{\mathcal{G}}_{\mathbf{A}, \varepsilon}^{(n)}$ of diameter larger than 2ε and that do not contain loops of $\tilde{\mathcal{L}}_n^{1/2}$ with an odd winding around $\text{Tube}_\varepsilon(\mathbf{A})$ also diverges as $n \rightarrow +\infty$. What our Theorem 1 shows is that the clusters of the latter type still can achieve an odd winding around $\text{Tube}_\varepsilon(\mathbf{A})$, and this happens with a probability bounded away from 0 and 1. Since this probability is bounded away from 1, whereas the number of clusters proliferates, we infer that for an individual cluster it is increasingly costly as $n \rightarrow +\infty$ to achieve such an odd winding, and this odd winding is achieved at all only because many clusters try to do so. So heuristically, we interpret things in terms of a Bernoulli scheme as follows.

- The number of clusters that try to achieve an odd winding around $\text{Tube}_\varepsilon(\mathbf{A})$ diverges as $n \rightarrow +\infty$.
- In high enough dimension d , each of these clusters tries to achieve this odd winding almost independently from the other clusters.
- As $n \rightarrow +\infty$, the probability that at least one cluster achieves an odd winding around $\text{Tube}_\varepsilon(\mathbf{A})$ converges to a value in $(0, 1)$.

So we are in the regime of the law of rare events, and the limit distribution of the number of clusters that achieve an odd winding around $\text{Tube}_\varepsilon(\mathbf{A})$ is Poisson. The parameter of the corresponding Poisson distribution is the one given by the Brownian loop soup, in agreement with the limit probability of success.

Now, let us take a second hypertube $\text{Tube}_{\varepsilon'}(\mathbf{A}')$. A typical macroscopic cluster does not contain macroscopic cycles at all. It is already exceptional for such a cluster to have a cycle around $\text{Tube}_\varepsilon(\mathbf{A})$, and it would be exceptional squared for a cluster to have a second cycle around $\text{Tube}_{\varepsilon'}(\mathbf{A}')$, unless it is the same cycle that achieves an odd winding both around $\text{Tube}_\varepsilon(\mathbf{A})$ and $\text{Tube}_{\varepsilon'}(\mathbf{A}')$. So asymptotically, a cluster can contain at most one non-trivial macroscopic cycle.

Now, since the Poisson distribution in the limit is true for an arbitrary choice of the hypertube $\text{Tube}_\varepsilon(\mathbf{A})$, this determines not only the limit law of the number of clusters with prescribed topology, but also the very shape of the non-trivial macroscopic cycles that can be drawn inside the clusters of $\tilde{\mathcal{L}}_n^{1/2}$. In the limit, these macroscopic cycles are distributed as a Brownian loop soup with intensity parameter $\alpha = 1 = 2 \times \frac{1}{2}$, in accordance with (6.2). So this is our heuristic reasoning. Figure 12 depicts our understand of clusters in high dimension.

Remark 6.1. The picture on Figure 12 is somewhat reminiscent of Kenyon's cycle-rooted spanning forests (CRSF) [Ken11]. We expect that in high enough dimension, the cycles in a CRSF also scale to a Poisson point process of Brownian loops, but with a different intensity measure. If the unitary gauge field (values in \mathbb{S}^1) inducing the CRSF in discrete has a continuum fine mesh limit \mathcal{U} , then the intensity measure for the limit Poisson point process should be $(1 - \text{Re hol}^{\mathcal{U}}(\varphi)) \mu_{\mathbb{R}^d}^{\text{loop}}(d\varphi)$.




FIGURE 12. Conceptual depiction of the sign clusters of the metric GFF in high dimension. The trees are not exactly so, but are actually made of microscopic loops at the metric graph level. Most large clusters are macroscopically tree-like, as the trees in green. Their number diverges in the scaling limit. However, there is a tight number per scale of macroscopic cycles, that are represented in orange. The limit distribution of the macroscopic cycles is that of a Brownian loop soup of intensity parameter $\alpha = 1$, that is to say double of the intensity appearing in isomorphism theorems.

ACKNOWLEDGMENTS

The author thanks Marcin Lis (TU Wien, Vienna) for pointing out the topological probabilities for the FK-Ising model (Section 4.2). The author thanks Juhan Aru (EPFL), Adrien Kassel (ENS Lyon) and Thierry Lévy (Sorbonne Université) for inspiring discussions.

REFERENCES

- [Ahl10] Lars Valerian Ahlfors. *Conformal invariants: topics in geometric function theory*, volume 371. American Mathematical Society, 2010.
- [ALS20] Juhan Aru, Titus Lupu, and Avelio Sepúlveda. The first passage sets of the 2D Gaussian free field: convergence and isomorphism. *Communications in Mathematical Physics*, 375:1885–1929, 2020.
- [ALS22] Juhan Aru, Titus Lupu, and Avelio Sepúlveda. Extremal distance and conformal radius of a CLE_4 loop. *The Annals of Probability*, 50(2):509–558, 2022.
- [ALS23] Juhan Aru, Titus Lupu, and Avelio Sepúlveda. Excursion decomposition of the 2D continuum GFF. arXiv:2304.03150, 2023.
- [And65] George E. Andrews. A simple proof of Jacobi’s triple product identity. *Proceedings of the American Mathematical Society*, 16:333–334, 1965.
- [AS84] Milton Abramowitz and Irene A. Stegun, editors. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York; John Wiley & Sons, Inc., New York, 1984. Reprint of the 1972 edition, Selected Government Publications.
- [ASW19] Juhan Aru, Avelio Sepúlveda, and Wendelin Werner. On bounded-type thin local sets of the two-dimensional Gaussian free field. *Journal of the Institute of Mathematics of Jussieu*, 18(3):591–618, 2019.
- [BC21] Mikhail Basok and Dmitry Chelkak. Tau-functions à la Dubédat and probabilities of cylindrical events for double-dimers and $\text{CLE}(4)$. *Journal of the European Mathematical Society*, 23:2787–2832, 2021.
- [Ber17] Nathanaël Berestycki. An elementary approach to Gaussian multiplicative chaos. *Electronic Communications in Probability*, 22(27):1–12, 2017.

- [BLV16] Christian Benes, Gregory F. Lawler, and Fredrik Viklund. Scaling limit of the loop-erased random walk Green’s function. *Probability Theory and Related Fields*, 166:271–319, 2016.
- [BS15] Andrei Borodin and Paavo Salminen. *Handbook of Brownian Motion - Facts and Formulae*. Probability and Its Applications. Birkhäuser, 2nd, corrected edition, 2015.
- [CD23] Zhenhao Cai and Jian Ding. One-arm exponent of critical level-set for metric graph Gaussian free field in high dimensions. arXiv:2307.04434, 2023.
- [CJS⁺04] Sergio Carraciolo, Jesper Lykke Jacobsen, Hubert Saleur, Alan D. Sokal, and Andrea Sportiello. Fermionic field theory for trees and forests. *Physical Review Letters*, 93(8), 2004.
- [dB70] Nicolaas Govert de Bruijn. *Asymptotic methods in analysis*. Dover Publications, 3rd edition, 1970.
- [DP98] Donald Dawson and Edwin Perkins. *Stochastic Partial Differential Equations: Six Perspectives*, volume 64 of *Mathematical Surveys and Monographs*, chapter 2. Measure-Valued Processes and Renormalization of Branching Particle Systems. American Mathematical Society, 1998.
- [DPR22] Alexander Drewitz, Alexis Prévost, and Pierre-François Rodriguez. Cluster capacity functionals and isomorphism theorems for Gaussian free fields. *Probability Theory and Related Fields volume*, 183:255–313, 2022.
- [DPR23] Alexander Drewitz, Alexis Prévost, and Pierre-François Rodriguez. Critical exponents for a percolation model on transient graphs. *Inventiones mathematicae*, 232:229–299, 2023.
- [DS11] Bertrand Duplantier and Scott Sheffield. Liouville quantum gravity and KPZ. *Inventiones Mathematicae*, 185(2):333–393, 2011.
- [Dub09] Julien Dubédat. SLE and the free field: Partition functions and couplings. *Journal of the American Mathematical Society*, 22(4):995–1054, 2009.
- [Dub19] Julien Dubédat. Double dimers, conformal loop ensembles and isomonodromic deformations. *Journal of European Mathematical Society*, 21(1):1–54, 2019.
- [ES88] Robert G. Edwards and Alan D. Sokal. Generalization of the Fortuin-Kasteleyn-Swendsen-Wang representation and Monte Carlo algorithm. *Physical Review D*, 38(6):2009–2012, 1988.
- [FC78] J.L. Folks and R.S. Chhikara. The Inverse Gaussian distribution and its statistical application—a review. *Journal of the Royal Statistical Society: Series B (Methodological)*, 40(3):263–275, 1978.
- [Gri06] Geoffrey R. Grimmett. *The random-cluster model*, volume 333 of *Grundlehren der mathematischen Wissenschaften*. Springer, 2006.
- [Jan97] Svante Janson. *Gaussian Hilbert Spaces*, volume 129 of *Cambridge Tracts in Mathematics*. Cambridge University Press, 1997.
- [KC71] Leo P. Kadanoff and Horacio Ceva. Determination of an operator algebra for the two-dimensional Ising model. *Physical Review B*, 3(3918), 1971.
- [Ken11] Richard Kenyon. Spanning forests and vector bundle Laplacian. *The Annals of Probability*, 39(5):1983–2017, 2011.
- [KL21] Adrien Kassel and Thierry Lévy. Covariant Symanzik identities. *Probability and Mathematical Physics*, 2(3):419–475, 2021.
- [Law05] Gregory F. Lawler. *Conformally invariant processes in the plane*, volume 114 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2005.
- [LG99] Jean-François Le Gal. *Perplexing Problems in Probability*. *Festschrift in Honor of Harry Kesten*, volume 44 of *Progress in Probability*, chapter 16. The Hausdorff Measure of the Range of Super-Brownian Motion. Birkhäuser, 1999.
- [LJ10] Yves Le Jan. Markov loops and renormalization. *The Annals of Probability*, 38(3):1280–1319, 2010.
- [LJ11] Yves Le Jan. *Markov paths, loops and fields*. *École d’Été de Probabilités de Saint-Flour XXXVIII – 2008*, volume 2026 of *Lecture Notes in Mathematics*. Springer, 2011.
- [LL10] Gregory F. Lawler and Vlada Limic. *Random walk: a modern introduction*, volume 123 of *Cambridge studies in advanced mathematics*. Cambridge University Press, 2010.
- [LRV17] Hubert Lacoïn, Rémi Rhodes, and Vincent Vargas. Semiclassical limit of Liouville field theory. *Journal of Functional Analysis*, 273(3):875–916, 2017.
- [LTF07] Gregory F. Lawler and José A. Trujillo-Ferreras. Random walk loop soup. *Transactions of American Mathematical Society*, 359(2):767–787, 2007.
- [Lup16] Titus Lupu. From loop clusters and random interlacements to the free field. *The Annals of Probability*, 44(3):2117–2146, 2016.
- [Lup18] Titus Lupu. Convergence of the two-dimensional random walk loop-soup clusters to CLE. *Journal of the European Mathematical Society*, 21(4):1201–1227, 2018.
- [LW04] Gregory F. Lawler and Wendelin Werner. The Brownian loop soup. *Probability Theory and Related Fields*, 128:565–588, 2004.
- [LW16] Titus Lupu and Wendelin Werner. A note on Ising random currents, Ising-FK, loop-soups and the Gaussian free field. *Electronic Communications in Probability*, 21:1–7, 2016.
- [LW18] Titus Lupu and Wendelin Werner. The random pseudo-metric on a graph defined via the zero-set of the Gaussian free field on its metric graph. *Probability Theory and Related Fields*, 171:775–818, 2018.

- [Pom92] Christian Pommerenke. *Boundary behaviour of conformal maps*, volume 299 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, 1992.
- [RV14] Rémi Rhodes and Vincent Vargas. Gaussian multiplicative chaos and applications: a review. *Probability Surveys*, 11:315–392, 2014.
- [Sag94] Hans Sagan. *Spaces-filling curves*. Universitext. Springer, 1994.
- [She07] Scott Sheffield. Gaussian free fields for mathematicians. *Probability Theory and Related Fields*, 139(3):521–541, 2007.
- [Sim74] Barry Simon. *The $P(\phi)_2$ Euclidean (quantum) field theory*. Princeton Series in Physics. Princeton University Press, 1974.
- [Sla99] Gordon Slade. *Perplexing Problems in Probability. Festschrift in Honor of Harry Kesten*, volume 44 of *Progress in Probability*, chapter 2. Lattice Trees, Percolation and Super-Brownian Motion. Birkhäuser, 1999.
- [SS09] Oded Schramm and Scott Sheffield. Contour lines of the two-dimensional discrete Gaussian free field. *Acta Mathematica*, 202(1):21, 2009.
- [SS13] Oded Schramm and Scott Sheffield. A contour line of the continuum Gaussian free field. *Probability Theory and Related Fields*, 157(1-2):47–80, 2013.
- [SWW17] Scott Sheffield, Samuel S. Watson, and Hao Wu. Simple CLE in doubly connected domains. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques*, 53(2):594–615, 2017.
- [vdBCL18] Tima van de Brug, Federico Camia, and Marcin Lis. Spin systems from loop soups. *Electronic Journal of Probability*, 23:1–17, 2018.
- [Wer21] Wendelin Werner. On clusters of Brownian loops in d dimensions. In Maria Eulália Vares, Roberto Fernández, Luiz Renato Fontes, and Charles M. Newman, editors, *In and Out of Equilibrium 3: Celebrating Vladas Sidoravicius*, volume 77 of *Progress in Probability*, pages 797–817. Birkhäuser, 2021.
- [WW17] Menglu Wang and Hao Wu. Level lines of Gaussian free field I: zero-boundary GFF. *Stochastic Processes and their Applications*, 127(4):1045–1124, 2017.

CNRS AND LPSM, UMR 8001, SORBONNE UNIVERSITÉ, 4 PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE
 Email address: titus.lupu@sorbonne-universite.fr