



HAL
open science

Channel change during catastrophic flood: example of the Alex storm in the Vésubie valley

Frédéric Liébault, Guillaume Piton, Gabriel Melun, Margot Chapuis, Paul Passy

► To cite this version:

Frédéric Liébault, Guillaume Piton, Gabriel Melun, Margot Chapuis, Paul Passy. Channel change during catastrophic flood: example of the Alex storm in the Vésubie valley. 10th IAG International Conference on Geomorphology, Sep 2022, Coimbra, Portugal. hal-03779071

HAL Id: hal-03779071

<https://hal.science/hal-03779071v1>

Submitted on 16 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Channel change during catastrophic flood: example of the Alex storm in the Vésubie valley

Frédéric Liébault¹, Guillaume Piton¹, Gabriel Melun², Margot Chapuis³, and Paul Passy⁴

¹Université Grenoble Alpes, INRAE, ETNA, Grenoble, France (frederic.liebault@inrae.fr)

²Office Français de la Biodiversité, Direction de la Recherche et de l'Appui Scientifique, Vincennes, France (gabriel.melun@ofb.gouv.fr)

³Université Côte d'Azur, CNRS, ESPACE, Nice, France (margot.chapuis@unice.fr)

⁴UMR 8586 PRODIG, Université Paris Cité, Paris, France (paul.passy@u-paris.fr)

Documenting and interpreting channel responses to catastrophic floods help understanding how rapid fluvial metamorphosis can propagate in a catchment under sediment cascading effects. The recent example of the October 2020 Alex storm in SE France (~500 mm of rain in 24 h) provides a unique opportunity to investigate the sudden formation of a braided channel along 35 km of a confined alpine valley (Vésubie) and to link the morphological response with sediment wave initiation and propagation. GIS-based analysis of remote sensing data (high-resolution ortho-imagery and airborne LiDAR data) acquired before and after the flood allowed combining channel changes with sediment erosion and deposition along a 35-km reach of the Vésubie, including the most impacted portions of the valley. Archives of aerial imagery and old pictures were also used to integrate the storm impact in the historical trajectory of the river. Comparison of the Alex storm effect with past major floods that occurred in the valley shows that active channel widening after the storm was one order of magnitude higher than during the 100-yrs flood of November 1997. Dramatic channel aggradation was observed, with net bed-level change exceeding 10 m in some proximal reaches. This case study allows us to discuss the critical role of floodplain and terrace erosion in the formation of the post-flood braided channel, and to compare the geomorphic impact of the storm with similar reported cases in the literature.