



Subcritical water: an original eco-sustainable process for wine lees valorization ?

Benjamin Poulain, Delphine Winstel, Claudia Nioi

► To cite this version:

Benjamin Poulain, Delphine Winstel, Claudia Nioi. Subcritical water: an original eco-sustainable process for wine lees valorization ?. In Vino Analytica Scientia 2022, Jul 2022, Neustadt an der Weinstraße, Germany. 2022. hal-03781910

HAL Id: hal-03781910

<https://hal.science/hal-03781910v1>

Submitted on 20 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Subcritical water : an original eco-sustainable process for wine lees valorization ?

Benjamin POULAIN¹, Delphine WINTEL¹, Claudia NIOI¹

¹Univ. Bordeaux, Bordeaux INP, Bordeaux Science Agro, INRAE, OENO, UMR 1366, ISVV, F-33140 Villenave d'Ornon, France

INTRODUCTION

Wine lees are the second most important wine by-product in terms of quantity after grape stalk and marc [1]. During ageing of white wine on lees, it is well known that lees release compounds that limit oxidation of wine [2]. The antioxidant potential of wine lees is not totally elucidated. Glutathione seems to be one of the most important compounds involved in antioxidant activity [2]. In this way, white wine lees are a promising by-product to obtain biomolecules of interest.

In this study, we propose an eco-sustainable process to extract compounds with antioxidant activity from white wine lees.

METHODS

Raw material :

Sémillon wine lees (2021) from AOC Pessac-Léognan (France).

1. Impact of operating conditions and solvents on extraction yield and antioxidant activity :


- Solvents impact : Water at 20 °C (W), Ethanol 50 % (v/v)(EtOH), Subcritical Water (SW) at 100, 150 and 200 °C at 30 bars (and 60 bars).
- Optimised operating conditions : initial lees concentration 10 g/L and 15 min of extraction.

2. Evaluation of the antioxidant activity (antiradical and ferric reducing power) :

- DPPH and FRAP method (colorimetric).

3. Evaluation of glutathione in extracts :

- DTNB method (colorimetric).




CONCLUSION

Subcritical water is a promising solution for glutathione extraction and to obtain antioxidants extracts from white wine lees.

In conclusion, subcritical water seems to be an original eco-sustainable process for wine lees valorization.

RESULTS

Impact of operating conditions and solvent



Solvent impact :


Subcritical water and water 20 °C allow a better extraction yield compared to conventional solvent.

Subcritical water extraction :


Temperature has a positive impact on extraction yield.

Evaluation of the antioxidant activity

Evaluation of the ferric reducing power of extracts - FRAP




Evaluation of antiradical power of extracts - DPPH



Ethanol 50 % and subcritical water at 200 °C extractions are favourable to obtain extracts with high antiradical and ferric reducing power.


Any differences observed between subcritical water at 30 and 60 bars (data not shown).

Evaluation of glutathione in extracts



Conventional solvent and subcritical water at 200 °C are best solvents for glutathione extraction.

The higher glutathione extraction with SW 200 °C could be one of the reason of its better antioxidant activity.



WHAT IS NEXT ?

- Optimisation of subcritical extraction with DOE.
- Identification of molecules implicated in antioxidant activity of extract and lees.

REFERENCES

- [1] C. Dimou, N. Kopsahelis, A. Papadaki, S. Papanikolaou, I. K. Kooks, I. Mandala, A. A. Koutinas, Food Res. Int. (2015) 81-87.
[2] V. Lavigne et D. Dubourdieu, J. Int. Sci. Vigne Vin. (1996), 30, n°04, 201-206

Acknowledgements : Authors grateful to Region Nouvelle-Aquitaine, Biolaffort (SANSOVIN project) and ANR for financial support.