



HAL
open science

Risk of neonatal hypothyroidism in newborns from mothers exposed to CTPA during pregnancy: Ancillary data from a prospective outcome study

Marc Righini, Helia Robert-Ebadi, Alessio Cremonesi, Antoine Elias, Olivier Sanchez, Emmanuelle Le Moigne, Jeannot Schmidt, Catherine Le Gall, Jacques Cornuz, Drahomir Aujesky, et al.

► To cite this version:

Marc Righini, Helia Robert-Ebadi, Alessio Cremonesi, Antoine Elias, Olivier Sanchez, et al.. Risk of neonatal hypothyroidism in newborns from mothers exposed to CTPA during pregnancy: Ancillary data from a prospective outcome study. *Journal of Thrombosis and Haemostasis*, 2022, 10.1111/jth.15843 . hal-03789161

HAL Id: hal-03789161

<https://hal.science/hal-03789161>

Submitted on 23 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.
















L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

ORIGINAL ARTICLE

Risk of neonatal hypothyroidism in newborns from mothers exposed to CTPA during pregnancy: Ancillary data from a prospective outcome study

Marc Righini¹  | Helia Robert-Ebadi¹  | Alessio Cremonesi²  | Antoine Elias³  |
 Olivier Sanchez^{4,5,6}  | Emmanuelle Le Moigne⁷  | Jeannot Schmidt⁸  |
 Catherine Le Gall⁹ | Jacques Cornuz¹⁰  | Drahomir Aujesky¹¹  | Pierre-Marie Roy¹²  |
 Céline Chaleur^{13,14,15}  | Frédéric Rouyer¹⁶  | Pierre-Alexandre Poletti¹⁷  |
 Caroline Moreau¹⁸  | Grégoire Le Gal^{7,19}  | for the CT-PE-Pregnancy group

¹Division of Angiology and Hemostasis, Geneva University Hospitals and Faculty of Medicine, Geneva, Switzerland

²Division of Clinical Biochemistry and Swiss Newborn Screening, University Children's Hospital Zurich, University of Zurich, Zurich, Switzerland

³Médecine Vasculaire, Centre Hospitalier de Toulon, Toulon, France

⁴Service de Pneumologie et Soins Intensifs, Hôpital Européen Georges Pompidou, AP-HP, Université Paris Descartes, Sorbonne Paris Cité, Paris, France

⁵INSERM UMR S 1140, Paris, France

⁶F-CRIN INNOVTE, Saint-Etienne, France

⁷EA3878 GETBO, Université de Brest, Brest, France

⁸Centre Hospitalier Universitaire de Clermont-Ferrand, Clermont-Ferrand, France

⁹Centre Hospitalier d'Argenteuil, Argenteuil, France

¹⁰Centre Hospitalier Universitaire Vaudois, Lausanne, Switzerland

¹¹Department of General Internal Medicine, Bern University Hospital, University of Bern, Bern, Switzerland

¹²Department of Emergency Medicine, University Hospital of Angers, Angers, France

¹³INSERM U1059, Saint-Etienne, France

¹⁴University of Lyon, Saint-Etienne, France

¹⁵Department of Gynecology and Obstetrics, University Hospital, Saint-Etienne, France

¹⁶Division of Emergency Medicine, Geneva University Hospitals, Geneva, Switzerland

¹⁷Department of Radiology, Geneva University Hospitals, Geneva, Switzerland

¹⁸IRSET UMR 1085, Université de Rennes 1, Rennes, France

¹⁹Department of Hematology, Ottawa Health Research Institute, Ottawa, Ontario, Canada

Correspondence

Marc Righini, Division of Angiology and Hemostasis, Department of Medical Specialties, Geneva University Hospitals and Faculty of Medicine, 4, rue Gabrielle-Perret-Gentil, CH-1211 Geneva 14, Switzerland.

Email: marc.righini@hcuge.ch

Abstract

Background: Neonatal hypothyroidism is often raised as a potential concern for the use of computed tomography pulmonary angiography (CTPA) in pregnant women with suspected pulmonary embolism (PE).

Objectives: To assess the incidence of neonatal hypothyroidism among newborns from mothers exposed to CTPA.

Manuscript handled by: Saskia Middeldorp Final decision: Saskia Middeldorp, 03 August 2022

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Journal of Thrombosis and Haemostasis* published by Wiley Periodicals LLC on behalf of International Society on Thrombosis and Haemostasis.

Funding information

Schweizerischer Nationalfonds zur Förderung der Wissenschaftlichen Forschung, Grant/Award Number: FNS32003B-120760

Patients/Methods: Pregnant women with clinically suspected PE were included in a multicenter, multinational prospective diagnostic management outcome study, based on pretest clinical probability assessment, high-sensitivity D-dimer testing, bilateral lower limb venous compression ultrasonography, and CTPA. Results of Guthrie tests were systematically collected for newborns of all women who required CTPA as part of the diagnostic strategy. A thyroid-stimulating hormone (TSH) level above 15 U/ml was used to define hypothyroidism.

Results: Out of the 166 women included in the Swiss participating centers, 149 underwent a CTPA including 14 with twin pregnancies. Eight women suffered a pregnancy loss and results of the Guthrie test could not be retrieved for four newborns. All TSH levels were reported as being below 15 U/ml. The incidence of neonatal hypothyroidism was 0/151 (0.0%, 95% confidence interval: 0.0%–2.5%).

Conclusions: We did not identify any cases of neonatal hypothyroidism in our cohort of 149 pregnant women investigated for suspected PE using a CTPA. Along with previous literature data, this provides further reassuring data regarding the use of CTPA in this indication.

KEYWORDS

diagnosis, Guthrie test, hypothyroidism, pregnancy, pulmonary embolism

1 | INTRODUCTION

Pulmonary embolism (PE) remains one of the most common causes of maternal death in developed countries.^{1,2} Besides the fact that pregnancy is associated with an increased risk of venous thromboembolism (VTE), one of the potential explanations is that diagnosing PE is particularly challenging during pregnancy. Pregnant women often present symptoms and signs that can suggest PE, such as physiologic shortness of breath or tachycardia.³ There is very limited evidence in the literature to guide clinicians on how to manage pregnant women with suspected PE.^{4,5} Two prospective management outcome studies using computed tomography pulmonary angiography (CTPA) as an imaging test have suggested that this test is safe to rule out PE in this population.^{6,7} There is an ongoing debate on whether lung ventilation/perfusion (V/Q) scan or CTPA should be preferred during pregnancy. In terms of fetal radiation, the dose received is lower with CT during early pregnancy and becomes equivalent by the end of pregnancy.^{8–10} The dose to the mother's breast is significantly higher with CTPA throughout pregnancy.^{11,12}

Potential harmful effects related to intravenous iodinated contrast agents on fetal thyroid function have also been put forward as a reason to prefer the V/Q scan.

While reduced intakes of iodine is the major cause of hypothyroidism, exposure of the fetus or the neonate to excess iodine may also result in hypothyroidism. High iodine levels block thyroid hormone synthesis through inhibition of the organification process. This is known as the Wolff–Chaikoff effect. Also, excess iodine inhibits T4 and T3 secretion. These effects are thought to be present in the thyroid tissues for some weeks. In a normal thyroid

Essentials

- Neonatal hypothyroidism is a potential complication of computed tomography pulmonary angiography (CTPA) use during pregnancy.
- Analysis of this risk was done in a prospective management outcome study including pregnant women with suspected pulmonary embolism (PE).
- Guthrie tests performed in newborns from mothers who had CTPA for suspected PE during pregnancy were collected.
- No case of neonatal hypothyroidism in pregnant women investigated with CTPA for suspected PE was detected.

gland, after some days, the excess iodine induces downregulation of the sodium–iodide symporter and decreased intra-thyroid iodine concentration, leading to an escape from the Wolff–Chaikoff effect. However, the fetal thyroid gland is particularly sensitive to iodine overload because the ability to fully escape from the acute Wolff–Chaikoff effect does not mature until approximately 36 weeks. The risk of fetal hypothyroidism due to long-term maternal ingestion of iodides has been well documented in the past.^{13–16} A theoretical risk of contrast-induced hypothyroidism in neonates exposed to iodinated contrast agents during the antenatal period exists but has never been assessed in a formal prospective study.¹⁷

The CT-PE Pregnancy Study was the first published prospective management outcome study that assessed the safety of a sequential

diagnostic strategy in pregnant women with suspected PE.⁶ This strategy was based on the assessment of clinical probability with the simplified Geneva score, D-dimer measurement, compression ultrasonography of the lower limb veins, and CTPA. In Switzerland, a systematic screening of several neonatal conditions, including congenital hypothyroidism, is performed at birth by the measurement of capillary thyroid-stimulating hormone (TSH) levels. To study the effect of iodinated contrast agents used for CTPA in pregnant women, we prospectively collected TSH measurements in babies born from mothers who underwent CTPA for a suspected PE in the CT-PE Pregnancy Study.

2 | METHODS

2.1 | Study population

We used data from a multicenter, multinational prospective diagnostic management outcome study. We screened outpatient pregnant women presenting at one of the participating centers with a clinical suspicion of PE, as defined as acute onset of new or worsening shortness of breath or chest pain without another obvious cause. Exclusion criteria were: age below 18 years, allergy to iodinated contrast agent, impaired renal function (defined by a creatinine clearance below 30 ml/min as per the Cockcroft–Gault formula), diagnosis made prior to presentation, indication for or already on full dose anticoagulation, and inaccessibility for follow-up. The study was performed in two countries (France and Switzerland) and 11 centers actively included patients. The study was approved by the ethics committee according to legislation at each study site. A written informed consent was obtained from all participating women.

2.2 | Diagnostic work-up

The pretest probability (PTP) of PE was determined using the revised Geneva score. A D-dimer test was performed in all women, using a highly sensitive D-dimer assay (Vidas® D-dimer assay, bioMérieux). PE was deemed excluded in women with a non-high PTP and a negative D-dimer test (i.e., D-dimer < 500 µg/L). Women with a high PTP, and those with a non-high PTP and a positive D-dimer test, underwent bilateral compression ultrasonography (CUS). When a proximal deep vein thrombosis (DVT; popliteal vein and/or above) was found, PE was considered confirmed without further testing. Women with a negative CUS underwent CTPA. The protocol for CTPA consisted of an evaluation of the pulmonary arteries up to and including the subsegmental vessels. Only multidetector CT machines were used. The acquisition parameters for CTPA were: injection of a total volume of 100 ml of non-ionic contrast material (iodine concentration, 300–350 mg/ml) with a power injector at 3–5 ml per second; imaging 9–20 s after initiation of the contrast-material injection; scanning performed at 1.0–1.3 mm per section with a pitch of

1.25–1.75, 120 kV, and 115–260 mA; and reconstruction of images at 0.6–0.8 mm intervals. The complete diagnostic algorithm is depicted in [Figure 1](#).

2.3 | Follow-up

Patients with a negative diagnostic work-up were considered as not having PE and thus left without anticoagulant treatment and underwent clinical follow-up for 3 months. They were instructed to contact the study team in case of any new or worsening symptoms. An independent adjudication committee (three members) reviewed all the suspected VTE events, blinded from the initial diagnostic work-up.

2.4 | Study analysis

Although neonatal screening was performed in both countries, results were not systematically collected in the French participating centers, so this analysis is restricted to babies born from women included in the Swiss participating centers. Results of Guthrie tests were obtained at follow-up or when data were not available at follow-up, through the National Central Swiss Laboratory (Zürich, Switzerland). Results are provided as negative when the TSH measurements were below 15 U/ml. The actual TSH value is given in case of abnormal results. Women using medications that could have an influence on the babies' thyroid function (levothyroxine, amiodarone) were excluded.

The analysis was performed using babies as a unit. We estimated the proportion of newborns with an abnormal TSH result using exact proportions along with the 95% confidence interval (CI).

3 | RESULTS

Between August 2008 and July 2016, 395 pregnant women were included in the study. Of them, 166 were included in the Swiss participating centers. The characteristics of patients are depicted in [Table 1](#).

There were 23 (13.9%) women included during the first trimester of their pregnancy, 69 (41.6%) during the second, and 74 (44.6%) during the third trimester.

CTPA was not performed in 17 women: 14 women with a non-high PTP and negative D-dimer, 1 with a high PTP and proximal DVT on CUS, and 2 due to protocol violations. None of the included mothers were using drugs having a potential impact on the thyroid function of their babies. Therefore, 149 women, including 14 with twin pregnancies, were included in the analysis. The results of the Guthrie test could not be retrieved for four newborns, and there were eight pregnancy losses. Altogether, 151 Guthrie results were available. The study flow chart is depicted in [Figure 1](#).




FIGURE 1 Study flowchart. CI, confidence interval; CTPA, computed tomography pulmonary angiography; CUS, compression ultrasonography; PTP, pretest probability

All Guthrie levels were reported as being below 15U/ml. The proportion of newborns with neonatal hypothyroidism was: 0/151 (0.0%, 95% CI: 0.0%–2.5%).

4 | DISCUSSION

Among women included in the CT-PE Pregnancy Study,⁶ we did not identify any case of neonatal hypothyroidism in newborns of mothers who underwent a CTPA during the pregnancy for a suspected PE.

Congenital hypothyroidism, which can be definitive or transient, has an incidence of 1 in 2000 to 1 in 4000 newborns in the United States, and is the most common and treatable cause of mental retardation.¹⁸ Thyroid function during the intra-uterine period is essential for the neurologic development of the fetus as highlighted by the fact that the fetal gland begins to produce thyroxine under the influence of TSH between 10 and 12 weeks of gestation.¹⁹ Congenital hypothyroidism may be a result of athyreosis, thyroid dysgenesis, or

defects in pathways for biosynthesis of thyroid hormones. Transient congenital hypothyroidism may be due to placental transfer of maternal antibodies, or excessive iodine exposure. As iodine crosses the placenta, a potential adverse effect on the fetal thyroid gland from maternal intravenous administration of iodinated contrast agent resulting in neonatal hypothyroidism has been postulated.²⁰ Indeed, some reports of transient congenital hypothyroidism have been described, for example after amniocentesis with iodinated contrast agents.¹⁵ On the other hand, some small series did not report hypothyroidism in babies born from mothers receiving iodinated contrast medium during pregnancy.¹⁷ Therefore, the issue is still highly debated and has been put forward to avoid CTPA during pregnancy and use a V/Q scan instead.²¹

Depending on the generation of the CTPA and the protocol used, 75–150ml of iodinated contrast medium is administered intravenously. The iodine content of iodinated contrast media varies from 320 to 350mg/ml and is much higher than the recommended daily allowance of 150µg for adults and 220–290µg for pregnant and lactating women.²²

TABLE 1 Characteristics of included patients

Characteristics	Patients (n = 149)
Female sex, n (%)	149, 100
Age in years, median (IQR)	33 (29–36)
Trimester of pregnancy	
First, n (%)	17 (11.4)
Second, n (%)	60 (40.3)
Third, n (%)	72 (48.3)
BMI (kg/m ²)	25.1 (6.0)
Personal history of VTE, n (%)	11 (7.4)
Familial history of VTE, n (%)	15 (10.1)
Active malignancy, n (%)	0 (0.0)
Surgery within 1 month, n (%)	2 (1.0)
Bedridden for >72 h during the last 4 weeks, n (%)	15 (10.1)
Travel >6 h, n (%)	5 (3.4)
Chest pain, n (%)	95 (63.8)
Dyspnea, n (%)	108 (72.5)
Syncope, lipothymia, n (%)	19 (12.8)
Hemoptysis, n (%)	5 (3.4)
Clinical signs or symptoms of DVT, n (%)	14 (9.4)
Heart rate, bpm, mean (SD)	90 (17)
O ₂ saturation, %, mean (SD)	98.4 (1.6)

Abbreviations: BMI, body mass index; DVT, deep vein thrombosis; IQR, interquartile range; SD, standard deviation; VTE, venous thromboembolism.

A V/Q scan has been the cornerstone for PE diagnosis for more than three decades and is often proposed in pregnant women with suspected PE, starting with a perfusion scan and adding a ventilation phase only in case of perfusion abnormality.²³ However, this kind of strategy has never been validated in a properly conducted prospective study. Moreover, the availability of the test is limited to tertiary care centers.

CTPA has become the reference standard test for PE diagnosis outside pregnancy^{24,25} and has recently also been used in pregnant women with suspected PE.^{6,7} Indeed, the sequential use of an assessment of PTP, D-dimer measurement, and CTPA has been tested in two management prospective studies, which confirmed the safety of this diagnostic approach.^{6,7}

Therefore, our prospective data suggesting no detected case of neonatal hypothyroidism in our cohort could further reassure clinicians on the safety of CTPA for PE diagnosis during pregnancy. Of course, chest maternal radiation associated with the use of CTPA is significantly higher than with a V/Q scan, and there is an ongoing debate regarding the long-term risk of such irradiation. However, the risk-benefit balance of using CTPA in pregnant women seems acceptable as misdiagnosing PE during pregnancy may have immediate dramatic consequences for the mother and her baby.

Our study has strengths and limitations. This was a prospective study and we had a priori decided to systematically collect newborn

outcomes. We were able to retrieve results through a centralized laboratory that performs tests for all newborns in Switzerland. This ensures the use of the same pre-analytics variables and the same assay for all newborns. Also, all included pregnant women underwent a standardized diagnostic algorithm, using a similar CTPA protocol and the same contrast agent throughout the conduct of the study.⁶

Despite being the largest prospective study on PE diagnosis in pregnancy collecting newborn outcomes, our sample size remains limited. We were only able to exhaustively collect all Guthrie test results for women included in Switzerland. This limited us to providing very narrow estimates of the risk. However, our results are in line with those of a retrospective study including 344 women who underwent CTPA for suspected PE. In this study, all newborns had a normal T4 at birth, and only one newborn had a transient abnormal TSH, which normalized after few days.¹⁷

In summary, we did not identify any case of neonatal hypothyroidism in our cohort of 149 pregnant women investigated for suspected PE using a CTPA. Along with previous literature data, this provides further reassuring data regarding the use of CTPA in this indication.

AUTHOR CONTRIBUTIONS

Marc Righini, Helia Robert-Ebadi, Grégoire Le Gal: designed research, performed research, collected data, analyzed data, and wrote the paper. Alessio Cremonesi, Antoine Elias, Olivier Sanchez, Emmanuelle Le Moigne, Jeannot Schmidt, Catherine Le Gall, Jacques Cornuz, Drahomir Aujesky, Pierre-Marie Roy, Céline Chauleur, Frédéric Rouyer, Pierre-Alexandre Poletti, Caroline Moreau: performed research, collected data, analyzed data, and wrote the paper.

ACKNOWLEDGMENTS

The study was supported by a grant from the Swiss National Foundation for scientific research (FNS32003B-120760) by a grant from the Groupe d'Etude de la Thrombose de Bretagne Occidentale (GETBO) and by an International Society on Thrombosis and Haemostasis Presidential Grant (2017). GLG holds an Early Researcher Award from the Province of Ontario, a 'CP has heart' cardiovascular clinician scientist award from the Heart and Stroke Foundation of Ontario, and the Chair on Diagnosis of Venous Thromboembolism from the Department of Medicine, Faculty of Medicine, Ottawa. We would like to thank all the residents and physicians from the emergency departments and vascular medicine units of all participating centers. We would also like to thank all study nurses, secretaries, and clinical research technicians for their invaluable help. We extend our thanks to the staff of the Clinical Research Center, Geneva University Hospital, and in particular to Khaled Mostaguir (clinical research associate, no compensation), who developed the electronic eCRF for the study; to the coordinating center in Brest for French study sites: Isabelle Pichon (multicenter coordinator), Elise Poulhazan, Céline Dolou, Nabahats Ibrir, Florence Morvan, Véronique Kouassi, Floriane Masson. And last but not least, we would like to express our gratitude to the patients who made the study possible by accepting the invitation to participate to the trial.

CONFLICTS OF INTEREST

This manuscript represents original work, and it is not under consideration for publication elsewhere. It has never been neither submitted nor published in another scientific journal. All authors meet criteria for authorship and none of the authors have any conflict of interest. All had access to all data in the study, read and approved the final manuscript, and held responsibility for the decision to submit it for publication. The original trial was registered at clinicaltrials.gov (NCT 00740454).

ORCID

Marc Righini  <https://orcid.org/0000-0003-1211-2969>
 Helia Robert-Ebadi  <https://orcid.org/0000-0003-3801-1269>
 Alessio Cremonesi  <https://orcid.org/0000-0002-1524-4169>
 Antoine Elias  <https://orcid.org/0000-0002-1337-1826>
 Olivier Sanchez  <https://orcid.org/0000-0003-1633-8391>
 Emmanuelle Le Moigne  <https://orcid.org/0000-0001-5103-2406>
 Jeannot Schmidt  <https://orcid.org/0000-0003-3424-337X>
 Jacques Cornuz  <https://orcid.org/0000-0003-1276-9717>
 Drahomir Aujesky  <https://orcid.org/0000-0002-3970-2670>
 Pierre-Marie Roy  <https://orcid.org/0000-0003-4811-6793>
 Céline Chaleur  <https://orcid.org/0000-0003-4057-8133>
 Frédéric Rouyer  <https://orcid.org/0000-0002-5727-0859>
 Pierre-Alexandre Poletti  <https://orcid.org/0000-0001-9214-5546>
 Caroline Moreau  <https://orcid.org/0000-0002-5503-4078>
 Grégoire Le Gal  <https://orcid.org/0000-0002-9253-248X>

REFERENCES

- Cantwell R, Clutton-Brock T, Cooper G, et al. Saving Mothers' Lives: reviewing maternal deaths to make motherhood safer: 2006–2008. The Eighth Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom. *BJOG*. 2011;118(Suppl 1):1-203.
- Liu S, Rouleau J, Joseph KS, et al. Epidemiology of pregnancy-associated venous thromboembolism: a population-based study in Canada. *J Obstet Gynaecol Can*. 2009;31:611-620.
- Lee SY, Chien DK, Huang CH, Shih SC, Lee WC, Chang WH. Dyspnea in pregnancy. *Taiwan J Obstet Gynecol*. 2017;56:432-436.
- vanMens TE, Scheres LJ, deJong PG, Leeflang MM, Nijkeuter M, Middeldorp S. Imaging for the exclusion of pulmonary embolism in pregnancy. *Cochrane Database Syst Rev*. 2017;1:CD011053.
- Wan T, Skeith L, Karovitch A, Rodger M, Le Gal G. Guidance for the diagnosis of pulmonary embolism during pregnancy: Consensus and controversies. *Thromb Res*. 2017;157:23-28.
- Righini M, Robert-Ebadi H, Elias A, et al. Diagnosis of pulmonary embolism during pregnancy: a multicenter prospective management outcome study. *Ann Intern Med*. 2018;169:766-773.
- van derPol LM, Tromeur C, Bistervels IM, et al. Pregnancy-adapted YEARS algorithm for diagnosis of suspected pulmonary embolism. *N Engl J Med*. 2019;380:1139-1149.
- Righini M, Le Gal G, Bounameaux H. Venous thromboembolism diagnosis: unresolved issues. *Thromb Haemost*. 2015;113:1184-1192.
- Einstein AJ, Henzlova MJ, Rajagopalan S. Estimating risk of cancer associated with radiation exposure from 64-slice computed tomography coronary angiography. *JAMA*. 2007;298:317-323.
- Chan WS. Diagnosis of venous thromboembolism in pregnancy. *Thromb Res*. 2018;163:221-228.
- McLintock C, Brighton T, Chunilal S, et al. Recommendations for the diagnosis and treatment of deep venous thrombosis and pulmonary embolism in pregnancy and the postpartum period. *Aust N Z J Obstet Gynaecol*. 2012;52:14-22.
- Groves AM, Yates SJ, Win T, et al. CT pulmonary angiography versus ventilation-perfusion scintigraphy in pregnancy: implications from a UK survey of doctors' knowledge of radiation exposure. *Radiology*. 2006;240:765-770.
- Atwell TD, Lteif AN, Brown DL, McCann M, Townsend JE, Leroy AJ. Neonatal thyroid function after administration of IV iodinated contrast agent to 21 pregnant patients. *AJR Am J Roentgenol*. 2008;191:268-271.
- Etling N, Gehin-Fouque F, Vielh JP, Gautray JP. The iodine content of amniotic fluid and placental transfer of iodinated drugs. *Obstet Gynecol*. 1979;53:376-380.
- Rodesch F, Camus M, Ermans AM, Dodion J, Delange F. Adverse effect of amniocentesis on fetal thyroid function. *Am J Obstet Gynecol*. 1976;126:723-726.
- Cosman BC, Schullinger JN, Bell JJ, Regan JA. Hypothyroidism caused by topical povidone-iodine in a newborn with omphalocele. *J Pediatr Surg*. 1988;23:356-358.
- Bourjeily G, Chalhoub M, Phornphutkul C, Alleyne TC, Woodfield CA, Chen KK. Neonatal thyroid function: effect of a single exposure to iodinated contrast medium in utero. *Radiology*. 2010;256:744-750.
- Harris KB, Pass KA. Increase in congenital hypothyroidism in New York State and in the United States. *Mol Genet Metab*. 2007;91:268-277.
- Lee SY, Rhee CM, Leung AM, Braverman LE, Brent GA, Pearce EN. A review: radiographic iodinated contrast media-induced thyroid dysfunction. *J Clin Endocrinol Metab*. 2015;100:376-383.
- Raymond J, LaFranchi SH. Fetal and neonatal thyroid function: review and summary of significant new findings. *Curr Opin Endocrinol Diabetes Obes*. 2010;17:1-7.
- Lim W, Le Gal G, Bates SM, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: diagnosis of venous thromboembolism. *Blood Adv*. 2018;2:3226-3256.
- Rajaram S, Exley CE, Fairlie F, Matthews S. Effect of antenatal iodinated contrast agent on neonatal thyroid function. *Br J Radiol*. 2012;85:e238-e242.
- Chan WS, Ray JG, Murray S, Coady GE, Coates G, Ginsberg JS. Suspected pulmonary embolism in pregnancy: clinical presentation, results of lung scanning, and subsequent maternal and pediatric outcomes. *Arch Intern Med*. 2002;162:1170-1175.
- Righini M, Le Gal G, Aujesky D, et al. Diagnosis of pulmonary embolism by multidetector CT alone or combined with venous ultrasonography of the leg: a randomised non-inferiority trial. *Lancet*. 2008;371:1343-1352.
- Konstantinides SV, Meyer G. The 2019 ESC guidelines on the diagnosis and management of acute pulmonary embolism. *Eur Heart J*. 2019;40:3453-3455.

How to cite this article: Righini M, Robert-Ebadi H, Cremonesi A, et al. Risk of neonatal hypothyroidism in newborns from mothers exposed to CTPA during pregnancy: Ancillary data from a prospective outcome study. *J Thromb Haemost*. 2022;20:2550-2555. doi: [10.1111/jth.15843](https://doi.org/10.1111/jth.15843)