

Modular synthesis of 2-furyl carbinols from 3-benzyldimethylsilylfurfural platforms relying on oxygen-assisted C–Si bond functionalization

Sebastien Curpanen, Per Reichert, Gabriele Lupidi, Giovanni Poli, Julie Oble,

Alejandro Perez-Luna

▶ To cite this version:

Sebastien Curpanen, Per Reichert, Gabriele Lupidi, Giovanni Poli, Julie Oble, et al.. Modular synthesis of 2-furyl carbinols from 3-benzyldimethylsilylfurfural platforms relying on oxygen-assisted C–Si bond functionalization. Beilstein Journal of Organic Chemistry, 2022, 18, pp.1256-1263. 10.3762/bjoc.18.131. hal-03790431

HAL Id: hal-03790431 https://hal.science/hal-03790431

Submitted on 28 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. BEILSTEIN JOURNAL OF ORGANIC CHEMISTRY

Modular synthesis of 2-furyl carbinols from 3-benzyldimethylsilylfurfural platforms relying on oxygen-assisted C–Si bond functionalization

Sebastien Curpanen, Per Reichert, Gabriele Lupidi, Giovanni Poli, Julie Oble^{*} and Alejandro Perez-Luna^{*}

Full Research Paper Open Access Beilstein J. Org. Chem. 2022, 18, 1256-1263. Address Sorbonne Université, CNRS, Institut Parisien de Chimie Moléculaire, https://doi.org/10.3762/bjoc.18.131 IPCM, 4 Place Jussieu, 75005 Paris, France Received: 01 July 2022 Accepted: 31 August 2022 Fmail Julie Oble* - julie.oble@sorbonne-universite.fr; Published: 16 September 2022 Alejandro Perez-Luna* - alejandro.perez_luna@sorbonne-universite.fr Associate Editor: D. Y.-K. Chen * Corresponding author © 2022 Curpanen et al.; licensee Beilstein-Institut. Keywords: License and terms: see end of document. biomass; copper; cyclic siloxanes; 2-furyl carbinols; silicon

Abstract

3-Silylated furfurals, readily prepared in three steps from biomass-derived furfural and 5-methylfurfural, are converted into 3-silylated 2-furyl carbinols upon condensation with organomagnesium or organolithium reagents. The hydroxy unit of the carbinol adducts can be exploited to promote $C3(sp^2)$ –Si bond functionalization through intramolecular activation. Two approaches were contemplated for this purpose. Activation by alkoxides of the C3–SiEt₃ or C3–SiMe₂*t*-Bu bonds was ineffective. Conversely, treatment of the *C*3-benzyldimethylsilyl-appended derivatives with tetrabutylammonium fluoride led to cyclic siloxanes, which revealed to be competent donors for copper-catalyzed cross-coupling reactions, such as arylation reactions catalyzed by $Pd_2(dba)_3/CuI$, as well as allylation and methylation reactions catalyzed by CuI·PPh₃. *C*3-Benzyldimethylsilyl-appended furfurals are thus introduced as versatile platforms, providing a modular access to 3-substituted 2-furyl carbinols from renewable feedstock.

Introduction

Progress towards the use of nonedible renewable feedstock to replace fossil resources as starting material for high-value chemicals is an important endeavor of modern synthesis [1-3]. As part of this effort, exploitation of furfural and corresponding derivatives attracts continuous attention [4-7]. In this area, we have actively investigated catalytic C–H activation reactions [8-14], and as part of this work, we have recently developed an iridium-based protocol for the catalytic C3–H silylation of furfurylimines [15]. This method allows to install a C–Si bond poised for further functionalization on the furfural unit, and thereby leads to synthetic platforms useful to access elaborated furans. This prospect was demonstrated with platforms relying

on the SiMe(OSiMe₃)₂ unit, which were readily converted through Pd- or Cu-catalyzed electrophilic substitution reactions into an array of furfurals decorated at C3 with carbon- or heteroatom-containing substituents (Scheme 1). Conversely, all of our subsequent efforts to achieve cross-coupling reactions from C3-silylated furfurals with triorganosilane units have failed so far.



lated furfural platforms.

2-Furyl carbinols represent a useful class of furanic synthetic intermediates that have given rise to a number of synthetically relevant transformations, such as the Piancatelli and aza-Piancatelli reactions [16,17] or the Achmatowicz rearrangement [18]. Addition reactions of nucleophiles to the C–O double bond of furfurals represent an obvious synthetic approach to 2-furyl carbinols. We reasoned that for carbinols derived from C3-triorganosilyl-substituted furfurals, the OH unit could be exploited to assist C–Si bond activation [19]. Thereby, C3-triorganosilyl-substituted furfurals could be suitable platforms to develop a two-step modular approach to 3-substituted 2-furyl carbinols, entailing nucleophilic addition to the aldehyde function and oxygen-assisted electrophilic substitution of the C–Si bond (Scheme 1).

Results and Discussion Synthesis of 3-silylated 2-furyl carbinols

C3-silylated furfurals **1a–c** and **2c** are accessible from furfural or 5-methylfurfural [20], respectively, according to our previously reported protocol for selective catalytic C3 silylation [15]. The addition of organolithium [21] or Grignard reagents [22,23] to these substrates was uneventful and allowed for the preparation of 2-furylalkyl (see **3a–c**, **4c**), -aryl (see **5c**, **6c**), and -allyl carbinols (see **7c**) having furan rings with various triorganosilyl substituents at C3 in a synthetically useful yield and on an appropriate scale (Scheme 2).

With the 3-silylated 2-furyl carbinol substrates at hand, we then considered C–Si bond activation strategies relying on the assistance of the oxygen atom to promote electrophilic substitution reactions with carbon electrophiles.



C3–Si bond functionalization through intramolecular activation by alkoxides

We first contemplated the possibility to promote C3–Si bond functionalization through intramolecular activation by alkoxides [15]. It was reported that lithium alkoxides **A** undergo 1,4-silyl migration (Brook rearrangement) to generate C2-lithiated furans **C**, which in turn can react in the presence of electrophiles to deliver product **D**, resulting from the overall C2–Si functionalization (Scheme 3, top) [24,25]. The key to the viability of this process is the formation of pentavalent silicon intermediate **B**. This suggested that a related pentavalent intermediate **F** could be similarly accessible from **E**, and thus affording the C3-lithiated furan derivative **G** upon 1,4-silyl migration as well as the electrophilic substitution product **H** in the presence of an appropriate electrophile (Scheme 3, bottom).

However, treatment of aldehyde **1b** with *n*-BuLi, followed by addition of benzaldehyde in THF/DMPU [25], afforded only the addition product **3b** without any detectable formation of prod-

uct **9**, expected from 1,4-silyl migration/electrophilic substitution of **8** (Scheme 4, top).

Conversely, treatment of alcohol **3a** with *t*-BuOCu in the presence of allyl chloride, according to conditions developed by Takeda et al. for the reaction of γ -trimethylsilyl-substituted allylic alcohols [26,27] or *ortho*-silylated aryl carbinols [28], did provide C3-functionalized compound **10**, albeit in very low yield (6%) and along with the O-allylated product **11** (11%), as well as products **12** (4%) and **13** (4%) [29] (Scheme 4, bottom). In spite of extensive experimentation to find better conditions, this result could not be improved.

Nevertheless, important information came from an experiment involving furyl carbinol 4c, having a benzyldimethylsilyl unit (Scheme 5). Here, treatment with *t*-BuOCu-LiI in THF/HMPA in the presence of benzaldehyde led to the formation of adduct 14 (in 75% yield), which arose from the addition of a benzyl carbanion 17 to benzaldehyde. The generation of such a nucleo-



Scheme 3: C-Si bond functionalization of 2,3-disubstituted furyl carbinols by 1,4-silyl migration.



Scheme 4: Attempts of C3-Si bond functionalization promoted by intramolecular activation via alkoxide.

phile strongly suggests the formation of pentavalent silicon intermediate **15** [27], which then produced a (stabilized) benzylic carbanion (by exocyclic cleavage) rather than a 3-furanyl anion (by endocyclic cleavage). As a consequence, the failure to promote C3–Si activation of intermediates **E** (SiR₃ = SiEt₃ or SiMe₂*t*-Bu) according to our initially envisaged scenario depicted in Scheme 3, did not seem to be related to the difficulty to form pentavalent silicon intermediates **F**. Instead, the reason was probably a too low intrinsic stability of the C3-lithiated furans **G**, which thwarted 1,4-silyl migration. Conversely, this experiment provided evidence that cyclic siloxanes (e.g., **16**), which are potential nucleophilic partners for cross-coupling chemistry [30,31], could be accessed from 2-[(3benzyldimethylsilyl)furyl] carbinols.



C3–Si bond functionalization through siloxane formation

In an attempt to isolate cyclic siloxane **16**, we then considered a protocol reported by Anderson and co-workers to prepare cyclic siloxanes by debenzylation/cyclization of benzyldimethysilyl-substituted allylic alcohols [32]. To that end, carbinol **4c** was treated with TBAF·3H₂O (1.0 equiv), which resulted in the formation of **16** in 86% yield (along with toluene, Scheme 6). Alike 5-membered cyclic dimethyl(alkenyl)siloxanes, **16** was found to be highly sensitive towards silica gel column chromatography, and all of our attempts to isolate it failed. For this reason, we went on to consider C3–Si functionalization strategies of alcohols **4c–7c** relying on the formation of cyclic siloxanes and subsequent in situ cross-coupling.

We first briefly investigated Pd-catalyzed arylation reactions (Scheme 7). Fluoride-promoted arylation reactions of benzyldimethyl(alkenyl)silanes have been reported, and it is established that they proceed through the cleavage of the benzyl



moiety from the benzyldimethylsilyl groups, leading to either dimethylsilanols or cyclic siloxanes (as for substrates with pending hydroxy units) [32-35]. To the best of our knowledge, no analogous cross-coupling reactions from aryl- or heteroarylsubstituted benzyldimethylsilanes have been disclosed. We were thus delighted to find that using Pd₂(dba)₃ as precatalyst (2.5 mol %) in combination with CuI (20 mol %), cross-coupling between 4c and iodobenzene was achieved, giving 18 in reasonably good yield (70%). 4-Iodoanisole could also be coupled (giving 19 in 57% yield), but not electron-deficient 1-iodo-4-nitrobenzene. At this point, it should be mentioned that treatment of aldehyde 2c under the same reaction conditions led predominantly to decomposition products and only 5-methylfurfural (resulting from protodesilylation) was recovered in low yield (≈20%). Hence, intramolecular hydroxy activation seems decisive to obtain productive cross-coupling reactions with this system. Also, it is important to underscore that the copper cocatalyst was crucial for the success of this transformation as without, only protodesilylation products were obtained. This behavior is in contrast to that of dimethyl(alkenyl)siloxanes, which do not require copper to undergo cross-coupling [30,32].



This requirement for copper prompted us to test copper-catalyzed $C(sp^2)-C(sp^3)$ cross-coupling reactions, as reported by Takeda et al., to achieve allylation reactions of benzyldimethyl(alkenyl)silanes [36]. Treatment of **4c** with methallyl chloride in the presence of TBAF·(*t*-BuOH)₄ (2.4 equiv), CuI (1.5 equiv), and P(OEt)₃ (1.5 equiv) in DMF at room temperature delivered, after hydrolysis, the cross-coupling compound 21 in 68% yield, along with 9% of protodesilylation product 22 (Table 1, entry 1). As indicated by Takeda et al., the amount and source of TBAF proved crucial: use of either 1.2 equiv of $TBAF \cdot (t-BuOH)_4$ instead of 2.4 equiv (Table 1, entry 2), or use of a (commercially available) 1-molar TBAF solution in THF (Table 1, entry 3), gave an increased amount of protodesilylation product 22 at the expense of 21. By contrast, byproduct 22 was not observed reducing the reaction time from 24 h to 2 h, but the yield of product 21 remained the same (Table 1, entry 4). In spite of the satisfactory yield of 21, these conditions were synthetically impractical as we were unable to conveniently separate the final product from the phosphite ligand. The use of a ligand for copper revealed to be necessary, given that the cross-coupling product 21 was obtained in only 40% yield (along with 23% of 22) in the absence of P(OEt)₃ (Table 1, entry 5). The solution was the replacement of triethylphosphite by triphenylphosphine, which allowed to suppress the formation of the protodesilylated product and to obtain product 21 in 80% yield (Table 1, entry 6). Similar results were obtained using the preformed complex CuI·PPh₃ (1.5 equiv), so that compound 21 could be isolated in 65% yield (Table 1, entry 7). Importantly, the use of copper in a substoichiometric amount (20 mol %) was also suitable, allowing to isolate 21 in an even better 78% yield (Table 1, entry 9). It should also be mentioned that the use of THF instead of DMF as solvent led to worse results (Table 1, entries 7 and 8).

Investigating the scope of this transformation (Scheme 8), we established that the reaction performed well with other 5-methylfuryl carbinols. As such, products 23 and 24, arising from phenyl- and allyl-substituted substrates 6c and 7c, respectively, were isolated in 56% and 65% yield, respectively. Somewhat lower yields were noted with butyl- and phenyl-substituted carbinols 3c and 5c, respectively, bearing C5-unsubstituted furan rings, which gave products 25 and 26 in 42% and 40% yield, respectively. Electrophiles other than methallyl chloride could also be used. C3-Allylation of substrate 4c, leading to 27, was achieved in 52% yield by reaction with allyl chloride and in a better 61% yield using allyl bromide. It should be mentioned that competing protodesilylation could not always be fully suppressed in the above described reactions. Furthermore, purification was troublesome for products 25-27, which could only be isolated in the presence of the protodesilylated side products (see Supporting Information File 1 for details). We also contemplated the use of alkyl iodides as electrophiles. Methylation with methyl iodide was efficient, as shown through the preparation of 28 in 61% yield from 4c. In contrast, higher alkyl iodides, such as ethyl iodide, failed to provide the alkylation product (i.e., 29) and only protodesilylation was observed (even at 50 °C).

Conclusion

In conclusion, we have shown that 3-silylated 2-furyl carbinols are readily accessible in three steps from furfural and 5-methylfurfural, and the hydroxy unit of these adducts can be used to

n-Bu OH H
22
22 ^a (%)
9
32
20
0
23
0
0
20
0

^aYield measured prior to purification by ¹H NMR analysis using Me₂SO₂ as internal standard. ^bTBAF·(*t*-BuOH)₄ (1.2 equiv) was used. ^cTBAF (1 molar solution in THF, 2.4 equiv) was used. ^dIsolated yield.



promote C(sp²)–Si bond functionalization. Although intramolecular activation by alkoxides did not prove useful, C3–Si bond functionalization is achieved from benzyldimethylsilyl units upon siloxane formation in the presence of TBAF. Protocols for fluoride-promoted Pd/Cu-catalyzed arylation, as well as Cu-catalyzed allylation and methylation, have been developed. Overall, we have demonstrated that C3-benzyldimethylsilyl-appended furfurals are useful platforms that offer modular access to 3-substituted 2-furyl carbinols. This strategy represents a new, simple, and selective way to decorate the biomass-derived furan nucleus, allowing to reach synthetically relevant building blocks.

Experimental

Procedure for the addition of *n*-BuLi to C3-silylated furfurals (preparation of compounds **3a–c** and **4c**)

In a flame-dried round-bottom flask under argon was placed the appropriate C3-silylated furfural [15] and dissolved in freshly

distilled THF (0.3 M). The solution was cooled to -78 °C, and then *n*-BuLi (1.2 equiv in hexane) was added dropwise. The reaction mixture was allowed to stir at -78 °C for 30 min and then quenched with aq saturated NH₄Cl/NH₃ 2:1 solution. Et₂O was added, and the aqueous layer was extracted three times. The combined organic layer was washed with brine, dried over MgSO₄, filtered, concentrated under reduced pressure, and purified by silica gel column chromatography.

Procedure for the addition of Grignard reagents to C3-silylated furfurals (preparation of compounds **5c**, **6c**, and **7c**)

In a flame-dried round-bottom flask under argon was placed the appropriate C3-silylated furfural [15] and dissolved in freshly distilled THF (0.2 M solution). The solution was cooled to 0 °C, and then the Grignard reagent (1.3 equiv in Et₂O) was added dropwise (the rate of addition was equal to, or lower than 0.125 mL/min). The mixture was allowed to stir at 0 °C for 1 h and then allowed to reach room temperature. Upon consumption of the starting material, the reaction mixture was quenched with saturated NH₄Cl. CH₂Cl₂ was added, and the aqueous layer was extracted three times using CH₂Cl₂. The combined organic layer was dried over Na₂SO₄, filtered, concentrated under reduced pressure. and purified by silica gel column chromatography.

Procedure for the Pd/Cu-catalyzed arylation of C3-benzyldimethylsilyl-substituted 2-furyl carbinol **4c** (preparation of compounds **18** and **19**)

A flame-dried Schlenk tube was charged with CuI (20 mol %) and $Pd_2(dba)_3$ (2.5 mol %) and heated gently under vacuum using a heat gun. In another Schlenk tube, the appropriate iodoarene (1.5 equiv) and **4c** (1 equiv) were dissolved in freshly distilled THF (0.63 mL). The solution was degassed in three freeze–pump–thaw cycles, and degassed anhydrous TBAF (1 M in THF, 2.2 equiv) was added. The mixture was allowed to stir for 10 min at 0 °C and transferred via cannula to the Schlenk tube containing the catalytic mixture. The resulting mixture was stirred for 2 h at rt, then filtered through a short pad of silica gel, and concentrated. The residue was purified by silica gel column chromatography.

Procedure for the Cu-catalyzed allylation and methylation of C3-benzyldimethylsilylsubstituted 2-furyl carbinols (preparation of compounds **21** and **23–28**)

CuI·PPh₃ [37] (20 or 120 mol %) was introduced to a flamedried microwave vial, which was then placed under an argon atmosphere and sealed. In a Schlenk tube, the appropriate C3-benzyldimethylsilyl-substituted 2-furyl carbinol (0.3 mmol, 1 equiv) was dissolved in CH₂Cl₂ (1 mL), concentrated under reduced pressure, and placed under argon. DMF (3.0 mL) was then added, and the solution was degassed in three freeze–pump–thaw cycles. TBAF·(*t*-BuOH)₄ [38] (402 mg, 0.72 mmol) was then added along with the corresponding electrophile (3 equiv). The mixture was stirred for 2–3 min at rt and then cannulated into the vial containing the copper complex. The mixture was stirred for 2 h at 30 °C and then quenched with aq KOH (6 M, 1 mL). The mixture was extracted with cyclohexane/CH₂Cl₂ 9:1 (4 × 10 mL), and the combined organic layer was washed with water (10 mL), dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography.

Supporting Information

Supporting Information File 1

General information, characterization data, and copies of NMR spectra.

[https://www.beilstein-journals.org/bjoc/content/ supplementary/1860-5397-18-131-S1.pdf]

Acknowledgements

The authors acknowledge MS3U of Sorbonne Université for HRMS analysis.

Funding

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 951996.

ORCID[®] iDs

Sebastien Curpanen - https://orcid.org/0000-0002-7449-8310 Per Reichert - https://orcid.org/0000-0003-2780-9929 Gabriele Lupidi - https://orcid.org/0000-0001-9891-0552 Julie Oble - https://orcid.org/0000-0002-4002-255X Alejandro Perez-Luna - https://orcid.org/0000-0002-5578-9024

References

- Bender, T. A.; Dabrowski, J. A.; Gagné, M. R. Nat. Rev. Chem. 2018, 2, 35–46. doi:10.1038/s41570-018-0005-y
- Bozell, J. J.; Petersen, G. R. Green Chem. 2010, 12, 539–554. doi:10.1039/b922014c
- Mika, L. T.; Cséfalvay, E.; Németh, Á. Chem. Rev. 2018, 118, 505–613. doi:10.1021/acs.chemrev.7b00395
- Chen, S.; Wojcieszak, R.; Dumeignil, F.; Marceau, E.; Royer, S. Chem. Rev. 2018, 118, 11023–11117. doi:10.1021/acs.chemrev.8b00134

- Zhang, X.; Xu, S.; Li, Q.; Zhou, G.; Xia, H. RSC Adv. 2021, 11, 27042–27058. doi:10.1039/d1ra04633k
- Xu, C.; Paone, E.; Rodríguez-Padrón, D.; Luque, R.; Mauriello, F. Chem. Soc. Rev. 2020, 49, 4273–4306. doi:10.1039/d0cs00041h
- Bielski, R.; Grynkiewicz, G. Green Chem. 2021, 23, 7458–7487. doi:10.1039/d1gc02402g
- Pezzetta, C.; Veiros, L. F.; Oble, J.; Poli, G. Chem. Eur. J. 2017, 23, 8385–8389. doi:10.1002/chem.201701850
- Siopa, F.; Ramis Cladera, V.-A.; Afonso, C. A. M.; Oble, J.; Poli, G. Eur. J. Org. Chem. 2018, 6101–6106. doi:10.1002/ejoc.201800767
- Ravasco, J. M. J. M.; Monteiro, C. M.; Siopa, F.; Trindade, A. F.; Oble, J.; Poli, G.; Simeonov, S. P.; Afonso, C. A. M. *ChemSusChem* 2019, *12*, 4629–4635. doi:10.1002/cssc.201902051
- 11. Sala, R.; Roudesly, F.; Veiros, L. F.; Broggini, G.; Oble, J.; Poli, G. Adv. Synth. Catal. 2020, 362, 2486–2493. doi:10.1002/adsc.202000249
- Gupta, K.; Rai, R. K.; Singh, S. K. ChemCatChem 2018, 10, 2326–2349. doi:10.1002/cctc.201701754
- Mariscal, R.; Maireles-Torres, P.; Ojeda, M.; Sádaba, I.; López Granados, M. *Energy Environ. Sci.* 2016, *9*, 1144–1189. doi:10.1039/c5ee02666k
- 14. Karlinskii, B. Y.; Ananikov, V. P. *ChemSusChem* **2021**, *14*, 558–568. doi:10.1002/cssc.202002397
- 15. Curpanen, S.; Poli, G.; Perez-Luna, A.; Oble, J. Asian J. Org. Chem. **2022**, e202200199. doi:10.1002/ajoc.202200199
- 16. Piutti, C.; Quartieri, F. *Molecules* **2013**, *18*, 12290–12312. doi:10.3390/molecules181012290
- Verrier, C.; Moebs-Sanchez, S.; Queneau, Y.; Popowycz, F. Org. Biomol. Chem. 2018, 16, 676–687. doi:10.1039/c7ob02962d
- Ghosh, A. K.; Brindisi, M. RSC Adv. 2016, 6, 111564–111598. doi:10.1039/c6ra22611f
- Curpanen, S.; Poli, G.; Oble, J.; Perez-Luna, A. *Eur. J. Org. Chem.* 2021, 1055–1071. doi:10.1002/ejoc.202001458
- 20. Yang, W.; Sen, A. *ChemSusChem* 2011, *4*, 349–352. doi:10.1002/cssc.201000369
 See for the synthesis of 5-methylfurfural from biomass-derived carbohydrates.
- Kazancioglu, E. A.; Kazancioglu, M. Z.; Fistikci, M.; Secen, H.; Altundas, R. Org. Lett. 2013, 15, 4790–4793. doi:10.1021/ol402163u
- 22. Hashmi, A. S. K.; Häffner, T.; Yang, W.; Pankajakshan, S.; Schäfer, S.; Schultes, L.; Rominger, F.; Frey, W. *Chem. – Eur. J.* **2012**, *18*, 10480–10486. doi:10.1002/chem.201200306
- Lebœuf, D.; Schulz, E.; Gandon, V. Org. Lett. 2014, 16, 6464–6467. doi:10.1021/ol5032987
- 24. Bures, E.; Spinazzé, P. G.; Beese, G.; Hunt, I. R.; Rogers, C.; Keay, B. A. J. Org. Chem. **1997**, *62*, 8741–8749. doi:10.1021/jo971097j
- Devarie-Baez, N. O.; Kim, W.-S.; Smith, A. B., III; Xian, M. Org. Lett.
 2009, 11, 1861–1864. doi:10.1021/ol900434k
- 26. Taguchi, H.; Ghoroku, K.; Tadaki, M.; Tsubouchi, A.; Takeda, T. Org. Lett. 2001, 3, 3811–3814. doi:10.1021/ol016837w
- Taguchi, H.; Ghoroku, K.; Tadaki, M.; Tsubouchi, A.; Takeda, T. J. Org. Chem. 2002, 67, 8450–8456. doi:10.1021/jo025973r
- 28. Taguchi, H.; Takami, K.; Tsubouchi, A.; Takeda, T. *Tetrahedron Lett.* **2004**, *45*, 429–432. doi:10.1016/j.tetlet.2003.10.094
- 29. We rationalized the formation of products 12 and 13 from product 11. The sequence is depicted in Supporting Information File 1 and involves metalation to form an α-furyl anion, which then undergoes a [2,3]-Wittig rearrangement, followed by dehydoxylation through β-hydride transfer.

- Denmark, S. E.; Yang, S.-M. Org. Lett. 2001, 3, 1749–1752. doi:10.1021/ol015950j
- 31. Denmark, S. E.; Yang, S.-M. *Tetrahedron* **2004**, *60*, 9695–9708. doi:10.1016/j.tet.2004.06.149
- 32. Gudmundsson, H. G.; Kuper, C. J.; Cornut, D.; Urbitsch, F.; Elbert, B. L.; Anderson, E. A. J. Org. Chem. 2019, 84, 14868–14882. doi:10.1021/acs.joc.9b01664
- Trost, B. M.; Machacek, M. R.; Ball, Z. T. Org. Lett. 2003, 5, 1895–1898. doi:10.1021/ol034463w
- 34. Denmark, S. E.; Tymonko, S. A. J. Am. Chem. Soc. 2005, 127, 8004–8005. doi:10.1021/ja0518373
- Denmark, S. E.; Liu, J. H.-C. J. Am. Chem. Soc. 2007, 129, 3737–3744. doi:10.1021/ja067854p
- Takeda, T.; Matsumura, R.; Wasa, H.; Tsubouchi, A. Asian J. Org. Chem. 2014, 3, 838–841. doi:10.1002/ajoc.201402060
- Wang, Y.; Xiao, Y.; Yang Tan, T. T.; Ng, S.-C. *Tetrahedron Lett.* 2008, 49, 5190–5191. doi:10.1016/j.tetlet.2008.06.048
- 38. Kim, D. W.; Jeong, H.-J.; Lim, S. T.; Sohn, M.-H. Angew. Chem., Int. Ed. 2008, 47, 8404–8406. doi:10.1002/anie.200803150

License and Terms

This is an open access article licensed under the terms of the Beilstein-Institut Open Access License Agreement (https://www.beilstein-journals.org/bjoc/terms), which is identical to the Creative Commons Attribution 4.0 International License

(<u>https://creativecommons.org/licenses/by/4.0</u>). The reuse of material under this license requires that the author(s), source and license are credited. Third-party material in this article could be subject to other licenses (typically indicated in the credit line), and in this case, users are required to obtain permission from the license holder to reuse the material.

The definitive version of this article is the electronic one which can be found at: https://doi.org/10.3762/bjoc.18.131