



COVID-19 in vaccinated adult patients with hematological malignancies. Preliminary results from EPICOVIDEHA

Livio Pagano, Jon Salmanton-García, Francesco Marchesi, Alberto López-García, Sylvain Lamure, Federico Itri, Maria Gomes-Silva, Giulia Dragonetti, Iker Falces-Romero, Jaap van Doesum, et al.

► To cite this version:

Livio Pagano, Jon Salmanton-García, Francesco Marchesi, Alberto López-García, Sylvain Lamure, et al.. COVID-19 in vaccinated adult patients with hematological malignancies. Preliminary results from EPICOVIDEHA. *Blood*, 2022, 139 (10), pp.1588-1592. <10.1182/blood.2021014124>. <hal-03791235>

HAL Id: hal-03791235

<https://hal.science/hal-03791235v1>

Submitted on 29 Sep 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



American Society of Hematology
2021 L Street NW, Suite 900,
Washington, DC 20036
Phone: 202-776-0544 | Fax 202-776-0545
editorial@hematology.org

COVID-19 in vaccinated adult patients with hematological malignancies. Preliminary results from EPICOVIDEHA

Tracking no: BLD-2021-014124R2

Livio Pagano (Hematology, Università Cattolica del Sacro Cuore, Rome, Italy, Italy) Jon Salmanton-García (University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), Cologne, Germany, Germany) Francesco Marchesi (IRCCS Regina Elena National Cancer Institute, Italy) Alberto Lopez-Garcia (Fundacion Jimenez Diaz University Hospital, Health Research Institute IIS-FJD, Spain) Sylvain Lamure (Montpellier University Hospital, France) Federico Itri (San Luigi Gonzaga Hospital - Orbassano, Orbassano, Italy, Italy) Maria Gomes da Silva (Portuguese Institute of Oncology, Portugal) Giulia Dragonetti (Hematology, Università Cattolica del Sacro Cuore, Rome, Italy, Italy) Iker Falces-Romero (La Paz University Hospital, Spain) Jaap van Doesum (University Medical Center Groningen, Groningen, the Netherlands, Netherlands) Uluhan Sili (School of Medicine, Marmara University, Istanbul, Turkey, Turkey) Jorge Labrador (Hospital Universitario de Burgos, Spain) Maria Calbacho (Hospital Universitario 12 De Octubre, Spain) Yavuz Bilgin (Admiraal de Ruijter Hospital, Netherlands) Barbora Weinbergerova (Masaryk University and University Hospital Brno, Czech Republic) Laura Serrano (Hospital Universitario de Cabueñes, Gijón, Spain, Spain) Josep-Maria Ribera (Institut Català d'Oncologia-Hospital Germans Trias i Pujol. Josep Carreras Research Institute. Badalona. Universitat Autònoma de Barcelona, Spain) Sandra Malak (Institut Curie - Saint-Cloud, France) Jose Loureiro-Amigo (Hospital de Sant Joan Despi Moisès Broggi, Sant Joan Despi, Spain, Spain) Andreas Glenthøj (Rigshospitalet, Denmark) Raul Cordoba (Fundación Jimenez Diaz University Hospital, Spain) Raquel Nunes Rodrigues (Departamento de Hematologia, Instituto Português de Oncologia, Lisboa, Portugal, Portugal) Tomas Jose Gonzalez-Lopez (Hospital Universitario de Burgos, Spain) Linda Katharina Karlsson (Department of Hematology Herlev and Gentofte Hospital, Denmark) Maria-Jose Jimenez (Institut Català d'Oncologia, Hospital Germans Trias i Pujol, Josep Carreras Leukemia Research Instit, Spain) José-Ángel Hernández-Rivas (Hospital Universitario Infanta Leonor, Spain) Ozren Jaksic (Dubrava University Hospital, Croatia) Zdenek Racil (Institute of Hematology and Blood Transfusion Prague, Czech Republic) Alessandro Busca (A.O.U. Città della Salute e della Scienza, Italy) Paolo Corradini (University of Milan & Fondazione IRCCS Istituto Nazionale dei Tumori, Italy) Martin HOENIGL (Division of Infectious Diseases, Medical University of Graz, Austria) Nikolai KLIMKO (30) Department of Clinical Mycology, Allergy and Immunology, North Western State Medical University, Russian Federation) Philipp Koehler (University Hospital Cologne, Germany) Antonio Pagliuca (Kings College London and Kings College Hospital,) Francesco Passamonti (University of Insubria, Italy) Oliver Cornely (University Hospital of Cologne, Germany)

Abstract:

Conflict of interest: No COI declared

COI notes:

Preprint server: No;

Author contributions and disclosures: LP invented and EPICOVIDEHA, conceived the study idea, provided clinical details from local patients interpreted the data, interpreted the data, wrote the initial draft of the manuscript, and revised and approved the final manuscript. JSG enrolled patients and performed formal validation of the clinical details, extracted data from EPICOVIDEHA patients, performed the statistical analysis and interpreted the data, wrote the initial draft of the manuscript, created tables and revised and approved the final manuscript. FM provided clinical details from local patients interpreted the data, interpreted the data, wrote the initial draft of the manuscript, and revised and approved the final manuscript. ALG, SL, FI, MGS, GD, IFR, JVD, US, JL, MC, YMB, BW, LS, JMRSS, SM, JLA, AG, RCM, RNR, TJGL, LKK, MJJL, JAHR, OJ, ZR and the researchers listed in the study group provided clinical details from local patients and revised and approved the final manuscript. AB, PC, MH, NK, PK, AP, FP, and OAC invented and EPICOVIDEHA, conceived the study idea, and revised and approved the final manuscript.

Non-author contributions and disclosures: Yes; The researchers listed below belong to the EPICOVIDEHA study group and provided clinical details from local patients and revised and approved the final

manuscript: - Florian REIZINE, University Hospital Rennes, Centre Hospitalier Régional Universitaire Pontchaillou, France - Malgorzata MIKULSKA, Clinica Malattie Infettive Genova, Genoa, Italy - Hytham K. S. HAMID , AOU Caserta, Caserta, Italy - Nicola S. FRACCHIOLLA, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy - Francesca FARINA, Hematology and Bone Marrow Transplantation, IRCCS San Raffaele Scientific Institute, Milan, Italy - Nicola COPPOLA, University of Campania, Luigi Vanvitelli, Naples, Italy - Caterina BUQUICCHIO, Haematology Dept., Dimiccoli Hospital, Barletta, Italy - Avinash AUJAYEB, Northumbria Healthcare NHS Foundation Trust, Cramlington, United Kingdom - Przemyslaw ZDZIARSKI, Specialist of Infectious Disease and Clinical Immunology Lower Silesian Center Hirsztfeld Institute of Immunology and Experimental , Wrocław, Poland - Maria Chiara TISI, Cell Therapy and Hematology, San Bortolo Hospital, Vicenza, Italy - Martin SCHÖNLEIN, Department of Oncology, Hematology and Bone Marrow Transplantation with Section of Pneumology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany - Gianpaolo NADALI, Policlinico Borgo Roma Verona, Verona, Italy - Martin KOLDITZ, University Hospital Dresden, Dresden, Germany - Michaela HANAKOVA, Institute of Hematology and Blood Transfusion, Prague, Czech Republic - Monica FUNG, University of California San Francisco, Division of Infectious Diseases, San Francisco, USA - Maureen CHBAT, CHU Versailles, Châtenay, France - Caroline BESSON, Centre Hospitalier de Versailles, Versailles, France - Valentina BONUOMO, Policlinico Borgo Roma Verona, Verona, Italy - Ghaith ABU-ZEINAH, Division of Hematology and Oncology, Weill Cornell Medicine, New York, United States

Agreement to Share Publication-Related Data and Data Sharing Statement: na

Clinical trial registration information (if any):

COVID-19 in vaccinated adult patients with hematological malignancies.

Preliminary results from EPICOVIDEHA

Short title: COVID-19 in vaccinated hematological malignancies

Authors:

Livio **PAGANO**,^{1*} Jon **SALMANTON-GARCÍA**,^{2*} Francesco **MARCHESI**,^{3*} Alberto **LÓPEZ-GARCÍA**,⁴ Sylvain **LAMURE**,⁵ Federico **ITRI**,⁶ Maria **GOMES-SILVA**,⁷ Giulia **DRAGONETTI**,⁸ Iker **FALCES-ROMERO**,⁹ Jaap **VAN DOESUM**,¹⁰ Uluhan **SILI**,¹¹ Jorge **LABRADOR**,¹² Maria **CALBACHO**,¹³ Yavuz M. **BILGIN**,¹⁴ Barbora **WEINBERGEROVÁ**,¹⁵ Laura **SERRANO**,¹⁶ José-María **RIBERA-SANTA SUSANA**,¹⁷ Sandra **MALAK**,¹⁸ José **LOUREIRO-AMIGO**,¹⁹ Andreas **GLENTHØJ**,²⁰ Raúl **CÓRDOBA-MASCUÑANO**,²¹ Raquel **NUNES-RODRIGUES**,²² Tomás-José **GONZÁLEZ-LÓPEZ**,²³ Linda Katharina **KARLSSON**,²⁴ María-Josefa **JIMÉNEZ-LORENZO**,²⁵ José-Ángel **HERNÁNDEZ-RIVAS**,²⁶ Ozren **JAKSIC**,²⁷ Zdeněk **RÁČIL**,²⁸ Alessandro **BUSCA**,²⁹ Paolo **CORRADINI**,³⁰ Martin **HOENIGL**,³¹ Nikolai **KLIMKO**,³² Philipp **KÖHLER**,³³ Antonio **PAGLIUCA**,³⁴ Francesco **PASSAMONTI**,³⁵ Oliver A. **CORNELY**³⁶ on behalf of the EPICOVIDEHA working group

* These authors share first authorship

Affiliations:

- 1 Hematology, Fondazione Policlinico Universitario Agostino Gemelli – IRCCS, Rome, Italy
- 2 Hematology, Università Cattolica del Sacro Cuore, Rome, Italy
- 3 University of Cologne, Faculty of Medicine and University Hospital Cologne, Department I of Internal Medicine, Excellence Center for Medical Mycology (ECMM), Cologne, Germany
- 4 University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), Cologne, Germany
- 5 Hematology and Stem Cell Transplant Unit, IRCCS Regina Elena National Cancer Institute, Rome, Italy
- 6 Fundacion Jimenez Diaz University Hospital, Health Research Institute IIS-FJD, Madrid, Spain
- 7 Departement d'Hematologie Clinique, CHU de Montpellier, UMR-CNRS 5535, Universite de Montpellier, Montpellier, France
- 8 San Luigi Gonzaga Hospital – Orbassano, Orbassano, Italy
- 9 Portuguese Institute of Oncology, Lisbon, Portugal
- 10 Hematology, Fondazione Policlinico Universitario Agostino Gemelli – IRCCS, Rome, Italy
- 11 Hematology, Università Cattolica del Sacro Cuore, Rome, Italy

- 38 9 La Paz University Hospital, Madrid, Spain
- 39 10 University Medical Center Groningen, Groningen, Netherlands
- 40 11 Marmara University, Istanbul, Turkey
- 41 12 Department of Hematology, Hospital Universitario de Burgos, Burgos, Spain
- 42 13 Hospital Universitario 12 de Octubre, Madrid, Spain
- 43 14 Admiraal de Ruijter Hospital, Goes, Netherlands
- 44 15 Masaryk University and University Hospital Brno – Department of Internal Medicine –
45 Hematology and Oncology, Brno, Czech Republic
- 46 16 Hospital Universitario de Cabueñes, Gijón, Spain
- 47 17 Hospital Germans Trias i Pujol-Institut Català d'Oncologia Badalona Servicio de Hematología
48 Clínica, Badalona (Barcelona), Spain
- 49 18 Departement d'Oncologie medicale, Hematologie, Institut Curie, Saint Cloud, France
- 50 19 Hospital de Sant Joan Despí Moisès Broggi, Sant Joan Despí, Spain
- 51 20 Rigshospitalet, Copenhagen, Denmark
- 52 21 Fundacion Jimenez Diaz University Hospital, Health Research Institute IIS-FJD, Madrid, Spain
- 53 22 Departamento de Hematologia, Instituto Português de Oncologia, Lisboa, Portugal
- 54 23 Department of Hematology, Hospital Universitario de Burgos, Burgos, Spain
- 55 24 Center for Hemoglobinopathies, Department of Hematology Herlev and Gentofte Hospital,
56 Herlev, Denmark
- 57 25 Hospital Germans Trias i Pujol-Institut Català d'Oncologia Badalona Servicio de Hematología
58 Clínica, Badalona (Barcelona), Spain
- 59 26 Servicio de Hematología y Hemoterapia, Hospital Universitario Infanta Leonor, Madrid, Spain
- 60 27 Department of Hematology, Dubrava University Hospital, Zagreb, Croatia
- 61 28 Institute of Hematology and Blood Transfusion, Prague, Czech Republic
- 62 29 Stem Cell Transplant Center, AOU Citta' della Salute e della Scienza, Turin, Italy
- 63 30 University of Milan and Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy
- 64 31 Division of Infectious Diseases and Global Public Health, Department of Medicine, University of
65 California San Diego, San Diego, CA, United States
- 66 Clinical and Translational Fungal-Working Group, University of California San Diego, La Jolla,
67 CA, United States
- 68 Division of Infectious Diseases, Department of Internal Medicine, Medical University of Graz,
69 Graz, Austria
- 70 32 North-Western State Medical University named after Iliá Ilich Méchnikov, Saint-Petersburg,
71 Russia
- 72 33 University of Cologne, Faculty of Medicine and University Hospital Cologne, Department I of
73 Internal Medicine, Excellence Center for Medical Mycology (ECMM), Cologne, Germany
- 74 University of Cologne, Faculty of Medicine and University Hospital Cologne, Cologne
75 Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD),
76 Cologne, Germany

Department of Hematological Medicine, King's College Hospital NHS Foundation Trust, London, United Kingdom

Department of Medicine and Surgery, University of Insubria and ASST Sette Laghi, Ospedale di Circolo of Varese, Varese, Italy

University of Cologne, Faculty of Medicine and University Hospital Cologne, Department I of Internal Medicine, Excellence Center for Medical Mycology (ECMM), Cologne, Germany

University of Cologne, Faculty of Medicine and University Hospital Cologne, Chair Translational Research, Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), Cologne, Germany

University of Cologne, Faculty of Medicine and University Hospital Cologne, Clinical Trials Centre Cologne (ZKS Köln), Cologne, Germany

University of Cologne, Faculty of Medicine and University Hospital Cologne, Center for Molecular Medicine Cologne (CMMC), Cologne, Germany

German Centre for Infection Research (DZIF), Partner Site Bonn-Cologne, Cologne, Germany

ORCID number of authors:

Livio PAGANO	0000-0001-8287-928X
Jon SALMANTON-GARCÍA	0000-0002-6766-8297
Francesco MARCHESI	0000-0001-6353-2272
Alberto LÓPEZ-GARCÍA	0000-0002-5354-5261
Sylvain LAMURE	0000-0001-5980-305X
Federico ITRI	0000-0002-3532-5281
Maria GOMES-SILVA	0000-0002-6993-2450
Giulia DRAGONETTI	0000-0003-1775-6333
Iker FALCES-ROMERO	0000-0001-5888-7706
Jaap VAN DOESUM	0000-0003-0214-3219
Uluhan SILI	0000-0002-9939-9298
Jorge LABRADOR	0000-0002-3696-0287
Maria CALBACHO	0000-0001-8106-4863
Yavuz M. BILGIN	0000-0003-4854-5424
Barbora WEINBERGEROVÁ	0000-0001-6460-2471
José-María RIBERA-SANTA SUSANA	0000-0003-1042-6024
Sandra MALAK	0000-0001-8823-5055
José LOUREIRO-AMIGO	0000-0002-6451-8971
Andreas GLENTHØJ	0000-0003-2082-0738
Raúl CÓRDOBA-MASCUÑANO	0000-0002-7654-8836
Raquel NUNES-RODRIGUES	0000-0002-8347-4281
Tomás-José GONZÁLEZ-LÓPEZ	0000-0001-9575-1816
Linda Katharina KARLSSON	0000-0003-3317-7550
María-Josefa JIMÉNEZ-LORENZO	0000-0002-5469-5237
Zdeněk RÁČIL	0000-0003-3511-4596
Alessandro BUSCA	0000-0001-5361-5613
Paolo CORRADINI	0000-0002-9186-1353
Martin HOENIGL	0000-0002-1653-2824
Nikolai KLIMKO	0000-0001-6095-7531
Philipp KOEHLER	0000-0002-7386-7495

Antonio PAGLIUCA	0000-0003-2519-0333
Francesco PASSAMONTI	0000-0001-8068-5289
Oliver A. CORNELLY	0000-0001-9599-3137

Collaborators (to be listed in PubMed):

Florian **REIZINE**, Malgorzata **MIKULSKA**, Hytham K. S. **HAMID**, Nicola S. **FRACCHIO**
 Francesca **FARINA**, Nicola **COPPOLA**, Caterina **BUQUICCHIO**, Avinash **AUJAYEB**, Przemyslaw
ZDZIARSKI, Maria Chiara **TISI**, Martin **SCHÖNLEIN**, Gianpaolo **NADALI**, Martin **KOLDITZ**,
 Michaela **HANAKOVA**, Monica **FUNG**, Maureen **CHBAT**, Caroline **BESSON**, Valentina
BONUOMO, Ghaith **ABU-ZEINAH**

Corresponding author data:

Prof. Livio Pagano, MD
 Fondazione Policlinico Universitario A. Gemelli – IRCCS – Università Cattolica del Sacro Cuore
 Largo Francesco Vito 1, 00168 Roma
 Italia
 E-mail: Livio.Pagano@unicatt.it

Text word count: 1232

Number of figures and tables: 2 tables

Supplementary material: 1 figure, 2 Tables

Number of references: 25 references

Scientific Category: Letter to Blood

Coronavirus disease 2019 (COVID-19) is a life-threatening condition of high relevance for comorbid patients, such as those with baseline hematological malignancies (HM).¹⁻³ In April 2020, the European Hematology Association - Infectious Diseases Working Party (EHA-IDWP) opened an open web-based registry to collect all cases of HM adult patients that developed COVID-19 infections (EPICOVIDEHA survey).⁴ This registry aimed to describe the epidemiology, risk factors, and mortality rates of HM patients. Overall, we collected 3801 valid cases and we observed an overall mortality rate of 31%.⁵

Nearly one year after the first described COVID-19 case, in December 2020, the first vaccines against the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) were available^{6,7} and administration to the highest risk populations including HM patients started.^{8,9} From January 1, 2021, we prospectively collected registry data on adult fully or partially vaccinated HM patients that developed COVID-19, to assess the vaccine efficacy and potentially identify categories of patients that may be less protected by vaccines. With this report we share our findings of the first 113 patients included in the registry.

EPICOVIDEHA survey has been approved centrally by the Institutional Review Board and Ethics Committee of Fondazione Policlinico Universitario A. Gemelli – IRCCS – Università Cattolica del Sacro Cuore (Rome, Italy), and by the respective local partners as appropriate. EPICOVIDEHA has been registered at www.clinicaltrials.gov with the identifier NCT04733729.⁴ From January 1, 2021, until December 31, 2021, all participating institutions document episodes of COVID-19 in their patients with baseline HM that received a vaccination against SARS-CoV-2. Data are collected via the EPICOVIDEHA electronic case report form (eCRF), available at www.clinicalsurveys.net. This online survey is provided by EFS Fall 2018 (Questback, Cologne, Germany). Clinical and epidemiological data from patients with the laboratory-based diagnosis of SARS-CoV-2 infection after partial or complete vaccination are collected. Data captured included underlying conditions before SARS-CoV-2, HM status and management before SARS-CoV-2, SARS-CoV-2 vaccination, and infection details and mortality. The diagnosis of COVID-19 accords to the international recommendations of the WHO.¹⁰ The severity of COVID-19 at admission is graded according to the China Centers for Disease Control and Prevention definitions.¹¹ Patients

157 are considered fully vaccinated if the final dose was administered at least 14 days before symptom
158 onset or a positive PCR test for SARS-CoV-2.

159 As of 31 August 2021, 113 COVID-19 episodes among partially or completely vaccinated
160 patients with HM have been registered in EPICOVIDEHA. These patients have been reported from
161 42 out of 163 centres in 14 out of 38 European and non-European countries participating in the
162 survey. The clinical characteristics of these patients are reported in Table 1. The majority of them
163 were males (61.1%) and over 50 years of age (85.8%). More than 80% of patients had underlying
164 lymphoproliferative malignancies (chronic lymphoid leukemia [CLL], non-Hodgkin lymphoma [NHL],
165 acute lymphoblastic leukemia [ALL], Hodgkin's lymphoma [HL], and multiple myeloma [MM]).
166 Seventy-eight (68.1%) patients received active treatment for underlying HM at the time of COVID-
167 19 or within the prior 3 months. Following the recommendations of major international scientific
168 societies,^{8,9} the majority of our patients received an mRNA vaccine (BioNTech/Pfizer N=79
169 (69.9%), Moderna N=20 (17.7%)), whereas the remaining 14 (12.4%) received a vector-based
170 vaccine (AstraZeneca Oxford, N=10) or an inactivated vaccine (Sinovac CoronaVac, N=4); overall,
171 the median time from the last dose of vaccine and COVID-19 diagnosis was of 64 days (IQR: 33.5-
172 108). Eighty-seven patients (77%) were considered fully vaccinated, whereas the remaining 26
173 received only one shot; in all fully vaccinated patients, COVID-19 was diagnosed more than two
174 weeks after the second vaccine dose. Viral genomes of infection were analyzed in only 37 (32.7%)
175 cases and the alpha-variant was the most frequently observed (Supplemental Figure 1). Post-
176 vaccine IgG levels against SARS-CoV-2 spike protein were analyzed in 40 (35.4%) fully vaccinated
177 patients, 2-4 weeks from the last vaccine dose. Among these patients, only 13 (32.5%) presented
178 an antibody response to vaccine (optimal: 8; weak: 5), whereas the remaining 27 (67.5%) were
179 considered no responders (Binding Antibody Units, BAU < 30/ml). Overall, 79 (60.4%) patients had
180 a severe or critical infection. Seventy-five patients (66.4%) were admitted to the hospital, 16
181 (21.3%) of them to an ICU, and 10/16 required mechanical ventilation (Table 2); detailed data
182 about COVID-19 symptoms and severity according HM diagnosis have been described in
183 Supplemental Table 1. After a follow-up of 30 days post COVID-19 diagnosis, the overall mortality
184 rate was 12.4% (N=14). COVID-19 was the main or a secondary cause of death for all but one

patients; interestingly, we did not observe any statistical difference in terms of mortality between partially or fully vaccinated patients (15.4% vs 11.5%; $P=0.734$) and between patients achieving a serological response to vaccine vs non responders (13.3% vs 15.6%; $P=1$). In addition, we did not find any significant differences in terms of age or comorbidities comparing responder vs non responder patients. Moreover, our multivariable analysis showed that the only factor independently related to the risk of death in our cohort of vaccinated patients was the age ($P=0.035$; HR 1.053, 95%CI: 1.004-1.105) (Supplemental Table 2). Ten of 14 (71.4%) patients who died had underlying lymphoproliferative malignancies. With the caution due to the limited number of reported cases, it is worth to underline that none of the patients who died had underlying acute myeloid leukemia, which in our previous analysis in non-vaccinated patients was the category with one of the highest mortality rates.⁵

A generalized anti-SARS-CoV-2 vaccination policy has allowed a marked reduction in the incidence of severe COVID-19 in the general population. However, some reports indicates the occurrence of the infection in a limited number of vaccinated subjects.¹²⁻¹⁴ These are mostly subjects who have not developed protective immunity. Our survey, involving 42 hematology departments around the world, provides some preliminary insights. The majority of patients who do not respond to vaccination are patients with lymphoproliferative diseases, mainly CLL and NHL. This has also been observed for other vaccinations (e.g., influenza).¹⁵⁻¹⁶ Our results suggest that the low serologic response rate to anti-SARS-CoV-2 vaccines in patients with HM may translate to higher rates of infections. This has previously been described following monoclonal antibody treatment.¹⁷⁻²³ Unfortunately, only little data is available on the genomic characterization of the virus, we expect having a greater proportion of genotyping cases while continuing the survey. Given policies that differ between sites, post-vaccination serology results were available in only about 35% of patients and of those about two thirds were serologically non responders. It should be taken into account that the methods differ from a center to anotherone, even though our attempt of reducing inter-laboratory variation by referring to the WHO standardized method (<https://www.who.int/news-room/feature-stories/detail/standardization-of-vaccines-for-coronavirus-disease-covid-19>). Importantly, the overall mortality observed in our patients, although lower than

in the pre-vaccination period (~31%), remained high (12.4%). This percentage, on one hand remains quite worrying for hematologists, but on the other hand should be interpreted as a significant achievement following the spread of vaccination programs around the world. The hospitalization and mortality are still higher than the one observed in the fully vaccinated general population where the hospitalization rates of 2-3% have been reported.^{12,13,24,25} Our study reports preliminary observations and the low number of vaccinated patients is the main weakness, for now limiting the possibility to define the real incidence of breakthrough COVID-19 in HM.

Recruitment to this survey continues and larger numbers of cases will enable us to draw more conclusions in order to develop strategies to prevent severe COVID-19 in this frail population.

Informed consent was collected as applicable.

Acknowledgments: none.

Funding: Università Cattolica del Sacro Cuore contributed to the funding of this research project and its publication.

Author contributions:

LP invented and EPICOVIDEHA, conceived the study idea, provided clinical details from local patients interpreted the data, interpreted the data, wrote the initial draft of the manuscript, and revised and approved the final manuscript.

JSG enrolled patients and performed formal validation of the clinical details, extracted data from EPICOVIDEHA patients, performed the statistical analysis and interpreted the data, wrote the initial draft of the manuscript, created tables and revised and approved the final manuscript.

FM provided clinical details from local patients interpreted the data, interpreted the data, wrote the initial draft of the manuscript, and revised and approved the final manuscript.

ALG, SL, FI, MGS, GD, IFR, JVD, US, JL, MC, YMB, BW, LS, JMRSS, SM, JLA, AG, RCM, RNR, TJGL, LKK, MJJL, JAHR, OJ, ZR and the researchers listed in the study group provided clinical details from local patients and revised and approved the final manuscript.

AB, PC, MH, NK, PK, AP, FP, and OAC invented and EPICOVIDEHA, conceived the study idea, and revised and approved the final manuscript.

Disclosure of conflict of interest:

All the authors have no disclosures to declare for this submitted paper.

References

1. Wood WA, Neuberg DS, Thompson JC, et al. Outcomes of patients with hematologic malignancies and COVID-19: a report from the ASH Research Collaborative Data Hub. *Blood Adv.* 2020;4:5966-5975.
2. Ljungman P, de la Camara R, Mikulska M, et al. COVID-19 and stem cell transplantation; results from an EBMT and GETH multicentre prospective study. *Leukemia.* 2021 doi: 10.1038/s41375-021-01302-5. Online ahead of print.
3. Passamonti F, Cattaneo C, Arcaini L, et al. Clinical characteristics and risk factors associated with COVID-19 severity in patients with haematological malignancies in Italy: a retrospective, multicentre, cohort study. *Lancet Haematol.* 2020;7:e737-e745.
4. Salmanton-García J, Busca A, Cornely OA, et al. EPICOVIDEHA: a ready-to use platform for epidemiological studies in hematological patients with COVID-19. *Hemasphere.* 2021;5:e612.
5. Pagano L, Salmanton-García J, Marchesi F, et al. COVID-19 Infection in Adult Patients with Hematological Malignancies: A European Hematology Association Survey (EPICOVIDEHA). *J Hematol Oncol.* 2021. In press.
6. Anderson EJ, Rouphael NG, Widge AT, et al. Safety and immunogenicity of SARS-CoV-2 mRNA-1273 vaccine in older adults. *N Eng J Med.* 2020;383:2427–2438.
7. Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *N Engl J Med.* 2020;383:2603–2615.
8. <https://ehaweb.org/covid-19/eha-statement-on-covid-19-vaccines/recommendations-forcovid-19-vaccination-in-patients-with-hematologic-cancer/>.
9. Committee NCCNC-VA. Preliminary recommendations of the NCCN-COVID-19 Vaccination Advisory Committee. 2020 https://www.nccn.org/covid-19/pdf/COVID-19_Vaccination_Guidance_V1.0.pdf.
10. COVID-19 clinical management. Living guidance World Health Organization. January 15, 2021. WHO/2019-nCoV/clinical/2021.1.
11. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020;323:1239-42.
12. Bergwerk M, Gonen T, Lustig Y, et al. Covid-19 Breakthrough Infections in Vaccinated Health Care Workers. *N Engl J Med.* 2021 doi: 10.1056/NEJMoa2109072. Online ahead of print.
13. Brown CM, Vostok J, Johnson H, et al. Outbreak of SARS-CoV-2 Infections, Including COVID-19 Vaccine Breakthrough Infections, Associated with Large Public

- Gatherings - Barnstable County, Massachusetts, July 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70:1059-1062.
14. Juthami PV, Gupta A, Borges KA et al. Hospitalisation among vaccine breakthrough COVID-19 infections. *Lancet Infect Dis.* 2021. S1473-3099(21)00558-2 <https://doi.org/10.1016>. Online ahead of print.
 15. Ljungman P, Nahi H, Linde A. Vaccination of patients with haematological malignancies with one or two doses of influenza vaccine: a randomised study. *Br J Haematol.* 2005;130:96-98.
 16. Cordonnier C, Mikulska M, Einarsdottir S, Cesaro S, Ljungman P. 2017 ECIL 7 vaccine guidelines. ECIL vaccine group. *Lancet Infect Dis.* 2019;19:694-695.
 17. Greenberger LM, Saltzman LA, Senefeld JW, et al. Antibody response to SARS-CoV-2 vaccines in patients with hematologic malignancies. *Cancer Cell.* 2021;39:1031-1033.
 18. Herishanu Y, Avivi I, Aharon A, et al. Efficacy of the BNT162b2 mRNA COVID-19 vaccine in patients with chronic lymphocytic leukemia. *Blood.* 2021;137:3165-3173.
 19. Pimpinelli F, Marchesi F, Piaggio G, et al. Fifth-week-immunogenicity and safety of anti-SARS-CoV-2 BNT162b2 vaccine in patients with multiple myeloma and myeloproliferative malignancies on active treatment: preliminary data from a single Institution. *J Hematol Oncol.* 2021;14:81.
 20. Herzog-Tzarfati K, Gutwein O, Apel A, et al. BNT162b2 vaccine is significantly less effective in patients with hematologic malignancies. *Am J Hematol.* 2021 doi: 10.1002/ajh.26284.
 21. Lim SH, Campbell N, Johnson M, et al. Antibody responses after SARS-CoV-2 vaccination in patients with lymphoma. *Lancet Haematol.* 2021;8:e542-544.
 22. Parry H, McIlroy G, Bruton R, et al. Antibody responses after first and second Covid-19 vaccination in patients with chronic lymphocytic leukaemia. *Blood Cancer J.* 2021;30:136.
 23. Terpos E, Gavriatopoulou M, Ntanas-Stathopoulos I, et al. The neutralizing antibody response post COVID-19 vaccination in patients with myeloma is highly dependent on the type of anti-myeloma treatment. *Blood Cancer J.* 2021;11:138.
 24. Tenforde MW, Patel MM, Ginde AA, et al. Effectiveness of SARS-CoV-2 mRNA Vaccines for Preventing Covid-19 Hospitalizations in the United States. *Clin Infect Dis.* 2021. doi: 10.1093/cid/ciab687. Online ahead of print.
 25. Griffin JB, Haddix M, Danza P, et al. SARS-CoV-2 Infections and Hospitalizations Among Persons Aged ≥16 Years, by Vaccination Status - Los Angeles County, California, May 1-July 25, 2021. *MMWR Morb Mortal Wkly Rep.* 2021;70:1170-1176.

Table 1. Clinical characteristics of 113 vaccinated HM patients that developed COVID-19 infection

	N patients	%
Sex		
Female/male	44/69	38.9/61.1
Age (y.o.) (IQR) [range]	66 (58 - 78) [21 - 94]	
<50/>50 y.o.	16/97	14.2/85.8
Comorbidities		
None/ 1-2-3 comorbidities	36/77	31.9/68.1
Smoking history	17	15.0
Malignancy		
Acute lymphoid leukemia	3	2.6
Chronic lymphoid leukemia	28	24.8
Acute myeloid leukemia	5	4.4
Chronic myeloid leukemia	1	0.9
Myelodysplastic syndrome	7	6.2
Hodgkin lymphoma	4	3.5
Non-Hodgkin lymphoma	36	31.9
Myelofibrosis	3	2.7
Polycythemia vera	2	1.8
Systemic mastocytosis	2	1.8
Multiple myeloma	20	17.7
Aplastic anemia	2	1.8
Malignancy status before COVID-19		
Controlled disease (^)	51	45.1
Active disease	60	53.1
Not reported	2	
Last malignancy treatment (in the last 3 months)		
alloHSCT (in the last 6 months)	1	0.9
Chemotherapy	77	68.1
<i>Conventional chemotherapy</i>	13	11.5
<i>Hypomethylating agents</i>	4	3.5
<i>Immunotherapy</i>	9	8.0
<i>Immunotherapy</i>	30	26.5
<i>Targeted therapy</i>	21	18.6
No treatment	35	31.0
Patients with previous COVID-19 infections	2	1.8
y/n	2/111	1.8/98.2
Vaccination		
One dose	25	22.1
Two doses	88	77.8
<i>Patient that received vaccination at least 14 days before COVID-19 infection</i>	87	77.0
Type of vaccine		
mRNA + LNP		
<i>BioNTech/Pfizer</i>	79	69.9
<i>Moderna COVE</i>	20	17.7
Vector-based		
<i>AstraZeneca Oxford</i>	10	8.8
Inactivated		
<i>Sinovac</i>	4	3.5

	N patients	%
Anti-spike protein Ig dosage after vaccination (referring to WHO international standards, BAU/mL)		
No response (< 30)	27	23.9
Weak response (31-250)	5	4.4
Optimal response (> 250)	8	7
Unknown/not measured	73	64.7
COVID-19 infection		
Wild type - <i>WT</i>	11	9.7
English - <i>Alpha</i> (α)	16	14.2
South African - <i>Beta</i> (β)	1	0.9
Indian - <i>Delta</i> (δ)	9	8.0
Not tested	76	67.3
Severity		
Asymptomatic	22	19.5
Mild infection	12	10.6
Severe infection	63	55.8
Critical infection	16	14.2
Symptomatology at onset		
Asymptomatic	23	20.4
Pulmonary symptoms	37	32.7
Extrapulmonary symptoms	14	12.4
Pulmonary and extrapulmonary	39	34.5
Neutrophils count		
$\geq 500/\text{mm}^3$	98	86.7
Lymphocytes count		
$\geq 200/\text{mm}^3$	92	81.4

alloHSCT, allogeneic hematopoietic stem cell transplantation; **BAU**: binding antibody units; **COVE**, Coronavirus Efficacy and Safety Study; **COVID-19**, coronavirus disease 2019; **HM**, patients with hematological malignancy; **IQR**, interquartile range; **LNP**, lipid nanoparticles; **mm³**, cubic millimetre; **mRNA**, messenger ribonucleic acid; **N**, number; **WT**, wild type; **y.o.**, years old

^ Controlled disease: partial remission or better.

Table 2. Outcome of vaccinated patients that developed COVID-19 infection

	N patients	%
Stay during COVID-19		
Hospital	75	66.4
COVID-19 ward	59	83.8
ICU	16	14.2
of which, invasive mechanical ventilation	10	8.8
Home	38	33.6
Overall mortality at 30 days	14	12.4
Attributable to COVID-19	9/14	64.3
+ Hematological malignancy	3/14	21.4
Contributable by COVID-19	4/14	28.6
+ Other reasons*	2/14	14.3
Not related to COVID-19	1/14	7.1
+ Hematological malignancy	1/14	7.1
Mortality according to severity		
Asymptomatic	1/14	7.1
Mild infection	1/14	7.1
Severe infection	7/14	50.0
Critical infection	5/14	35.7
Mortality for stay		
Hospital	13/14	11.5
ICU	5/14	35.7
of which, invasive mechanical ventilation	5/5	100.0
Home	1/14	7.1
Mortality according to type of vaccine		
BioNTech/Pfizer	12/79	15.2
Moderna COVE	1/20	5.0
AstraZeneca Oxford	1/10	10.0
Sinovac	0/4	0.0
Mortality according to SARS-CoV-2 variant		
Wild type - <i>WT</i>	0/14	0.0
English - <i>Alpha</i> (α)	4/14	28.6
South African - <i>Betha</i> (β)	0/14	0.0
Indian - <i>Delta</i> (δ)	0/14	0.0
Not tested	10/14	71.4
Mortality according to vaccine scheme		
One dose	4/25	28.6
Full dose	10/78	71.4
Mortality according to type of hematological malignancy		
Acute lymphoid leukemia	0/3	0.0
Chronic lymphoid leukemia	2/28	7.1
Acute myeloid leukemia	0/5	0.0
Chronic myeloid leukemia	0/1	0.0
Myelodysplastic syndrome	2/7	28.6
Hodgkin lymphoma	1/4	25.0
Non-Hodgkin lymphoma	6/36	16.7
Myelofibrosis	1/3	33.3
Polycythemia vera	0/2	0.0
Systemic mastocytosis	1/2	50.0
Multiple myeloma	1/20	5.0
Aplastic anemia	0/2	0.0
Mortality for patients with active hematological malignancy		
y/n	7/7	50.0/50.0

	N patients	%
Mortality for patients with chemo-immuno or radiotherapy		
in the last 3 months	10/14	71.4
more than 3 months/ w&w	4/14	28.6

* Renal impairment + Bacterial infection; Intestinal subocclusion

alloHSCT, allogeneic hematopoietic stem cell transplantation; **COVE**, Coronavirus Efficacy and Safety Study; **COVID-19**, coronavirus disease 2019; **HM**, patients with hematological malignancy; **ICU**, intensive care unit; **w&w**, watch and wait