



HAL
open science

Impact of non-pharmaceutical interventions, weather, vaccination, and variants on COVID-19 transmission across departments in France: a modelling study

Juliette Paireau, Marie-Laure Charpignon, Sophie Larrieu, Clémentine Calba, Nathanaël Hozé, Pierre-Yves Boëlle, Rodolphe Thiébaud, Mélanie Prague, Simon Cauchemez

► To cite this version:

Juliette Paireau, Marie-Laure Charpignon, Sophie Larrieu, Clémentine Calba, Nathanaël Hozé, et al.. Impact of non-pharmaceutical interventions, weather, vaccination, and variants on COVID-19 transmission across departments in France: a modelling study. 2022. hal-03791761v1

HAL Id: hal-03791761

<https://hal.science/hal-03791761v1>

Preprint submitted on 29 Sep 2022 (v1), last revised 11 May 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

Impact of non-pharmaceutical interventions, weather, vaccination, and variants on COVID-19 transmission across departments in France: a modelling study

Authors

Juliette Paireau^{1,2*}, Marie-Laure Charpignon^{3,4,5}, Sophie Larrieu⁶, Clémentine Calba⁷, Nathanaël Hozé¹, Pierre-Yves Boëlle⁸, Rodolphe Thiebaut⁵, Mélanie Prague⁵, Simon Cauchemez¹

Affiliations

¹Mathematical Modelling of Infectious Diseases Unit, Institut Pasteur, Université Paris Cité, CNRS UMR 2000, Paris, France

²Infectious Diseases Department, Santé publique France, Saint Maurice, France

³Institute for Data, Systems, and Society (IDSS), Cambridge, MA, USA

⁴Computational Health Informatics Program, Boston Children's Hospital, Boston, MA, USA

⁵University of Bordeaux, Inria Bordeaux Sud-Ouest, Inserm, Bordeaux Population Health Research Center, SISTM Team, UMR1219, Bordeaux, France

⁶Regions Department, Regional Office Nouvelle-Aquitaine, Santé publique France, Bordeaux, France

⁷Regions Department, Regional Office Provence-Alps-French Riviera and Corsica, Santé publique France, Marseille, France

⁸INSERM, Sorbonne Université, Institut Pierre Louis d'Epidémiologie et de Santé Publique, Paris, France

*Corresponding author. Email: juliette.paireau@pasteur.fr

Abstract

Background: Multiple factors shape the temporal dynamics of the COVID-19 pandemic. Quantifying their relative contributions is key to guide future control strategies. Our objective was to disentangle the individual effects of non-pharmaceutical interventions (NPIs), weather, vaccination, and variants of concern (VOC) on local SARS-CoV-2 transmission.

Methods: We developed a log-linear model for the weekly reproduction number (R) of hospital admissions in 92 French metropolitan departments. We leveraged (i) the homogeneity in data collection and NPI definitions across departments, (ii) the spatial heterogeneity in the timing of NPIs, and (iii) an extensive observation period (14 months) covering different meteorological conditions, VOC proportions, and vaccine coverage levels.

Results: Three lockdowns reduced R by 72.9% (95%CI: 71.4-74.2), 70.4% (69.2-71.6) and 60.4% (56.1-64.3), respectively. Curfews implemented at 6/7pm and 8/9pm reduced R by 34.5% (28.1-40.4) and 18.4% (11.4-24.8), respectively. School closures reduced R by only 4.6% (1.6-7.4). We estimated that vaccination of the entire population would have reduced R by 74.0% (59.4-83.3), whereas the emergence of VOC (mainly Alpha during the study period) increased transmission by 46.9% (38.2-56.0) compared with the historical variant. Winter weather conditions (lower temperature and absolute humidity) increased R by 41.7% (37.0-46.7) compared to summer weather conditions. Additionally, we explored counterfactual scenarios (absence of VOC or vaccination) to assess their impact on hospital admissions.

Conclusions: Our study demonstrates the strong effectiveness of NPIs and vaccination and quantifies the role of meteorological factors while adjusting for other confounders. It highlights the importance of retrospective evaluation of interventions to inform future decision-making.

Keywords: COVID-19 ; SARS-CoV-2 ; non-pharmaceutical interventions ; vaccination ; variants ; climate ; reproduction number ; multivariable regression model

Background

Since the beginning of the coronavirus disease 2019 (COVID-19) pandemic, several factors have contributed to the transmission dynamics of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in time and space. First, many countries around the world have implemented non-pharmaceutical interventions (NPIs), such as lockdowns, curfews, and school closures (1). Before the introduction of vaccines, NPIs were the primary means to control disease spread. By the end of 2020 onwards, mass vaccination campaigns have helped mitigate the transmission of SARS-CoV-2 (2), while the concurrent emergence of more transmissible and immune escape variants of concern (VOC) has fostered virus spread (3). Finally, the weather may also have modulated disease transmission (4). Quantifying the relative contributions of each of these factors is key to better anticipate epidemic trends and guide future control strategies. However, this is challenging due to potential confounding, interaction effects, and a lack of identifiability of single effects when interventions or other factors are concomitant.

Several studies have investigated the effectiveness of NPIs at reducing SARS-CoV-2 transmission. The vast majority consisted of meta-analyses that combined data from multiple countries (1,5–10). However, conclusions drawn from such international comparisons may be affected by differences in local settings, data quality, NPI definitions, and population adherence to NPIs. In addition, most of these studies have only estimated the effect of NPIs during the first pandemic wave. Only a few have examined how the magnitude of NPI effects may have changed over time and in subsequent COVID-19 waves (11,12). In particular, the effect of NPIs that were only applied later in the pandemic, such as curfews, is still unclear. In addition to NPIs, the influence of meteorological factors on disease spread has been

much debated (13,14). Previous statistical studies investigating the role of weather variables generally relied on single estimates of the reproduction number R , measured at different locations (4,15–17) and early in the pandemic only. Thus, although these studies were timely and informative, they only covered a limited time period, when meteorological factors were likely less important to disease spread than governmental restrictions (18). Importantly, only a limited number of meteorological studies controlled for other factors such as NPIs (16,19,20); yet, not adjusting for sources of confounding may lead to spurious associations between weather and transmission. Now that data from a longer timespan are available, the role of weather conditions can be better elucidated.

To disentangle the effects of NPIs, weather, vaccination, and VOC on local SARS-CoV-2 transmission, we developed a statistical model to explain the time-varying reproduction number R reconstructed from the dynamics of hospital admissions, at the departmental level in metropolitan France, from March, 2020 to May, 2021. First, we leveraged the homogeneity in data collection and NPI definitions across departments. Indeed, in France, the number of patients hospitalized with COVID-19 was monitored through a single surveillance system implemented in all departments. In addition, most of the decisions on NPI implementation were made in a centralized manner. Such a standardized approach enabled harmonization of both data collection processes and NPI definition across departments, which benefited our study. Second, we leveraged spatial heterogeneity in the timing of NPI implementation. For example, lockdowns, curfews, and school closures were not systematically implemented at the same time in all departments, depending on the phase of the pandemic. This pattern in the timing of NPIs allowed us to circumvent the difficulty of assessing the impact of NPIs arising when measures are applied simultaneously across locations. Third, our study spanned a long observation period (14 months) that included varying meteorological conditions, VOC proportions, and vaccine coverage levels. This extensive study period covered three pandemic waves, thereby allowing us to examine the

impact of successive NPIs, vaccine distribution, and the emergence of VOC. Capturing a full seasonal cycle allowed us to quantify the role of meteorological factors.

Material and Methods

COVID-19 data

Hospital data were obtained from the SI-VIC database, which is the national inpatient surveillance system used during the pandemic. This database is maintained by the ANS (Agence du Numérique en Santé) and provides real-time data on COVID-19 patients hospitalized in French public and private hospitals. All cases are either biologically confirmed or present with a PET scan image highly suggestive of SARS-CoV-2 infection. New daily hospital admissions were defined as the incremental number of patients admitted to a general ward or intensive care unit, indexed by date of admission (rather than date of reporting). Data were aggregated by department (administrative unit), based on hospital location. Of note, metropolitan France consists of 96 departments, with a median population size of 600,000 inhabitants.

Covariates

To build covariates relative to NPIs, we collected data on the timeline of curfews, lockdowns, reopening periods following the lockdowns and during which restrictions were progressively lifted, as well as periods of more moderate restrictions (in between lockdowns), using a combination of governmental websites, press articles, and Wikipedia pages. We also included data on pandemic-related school closures (full or partial) and regular school holidays. Official dates were extracted from the Ministry of Education website (www.education.gouv.fr/calendrier-scolaire-100148). We measured an overall effect of school closures (whether pandemic-related or regular), with separate effects for summer and Christmas holidays.

The time-varying proportion of VOC was estimated using SIDEPE (Système d'Information de Dépistage Populationnel - Information system for population-based testing) database, which is the national surveillance system describing RT-PCR and antigen test results arising from all private and public French laboratories. Test results are reported by date of nasopharyngeal swab and include patient information such as residential zip code. Aggregated data are publicly available ([https://www.data.gouv.fr/fr/datasets /donnees-de-laboratoires-pour-le-depistage-indicateurs-sur-les-variants/](https://www.data.gouv.fr/fr/datasets/donnees-de-laboratoires-pour-le-depistage-indicateurs-sur-les-variants/)). The proportion of VOC was assessed among positive RT-PCR or antigen test results, using RT-PCR screening kits. The three main VOC circulating during the study period were Alpha, Beta, and Gamma. Data on variants were available from February 15, 2021 onwards. Before this date, the proportion of VOC was imputed by fitting a separate logistic regression model for each department. We assumed absence of VOC before December 15, 2020 (Additional file 1: Figure S1).

Vaccination data were obtained from the VAC-SI database, the national information system developed by the French Health Insurance to monitor the implementation of vaccination campaigns. Data are publicly available (www.data.gouv.fr/fr/datasets/donnees-relatives-aux-personnes-vaccinees-contre-la-covid-19-1/) and include both daily first-dose and full vaccine coverage time series, stratified by age group, and department, since the start of vaccine distribution in December 2020.

To account for increasing immunity at the population level, we included estimates of the proportion of adults previously infected with SARS-CoV-2, by day and department, using the method of Hozé et al (21). Specifically, estimates of age-stratified infection–hospitalisation ratios (IHR) were derived from joint analysis of hospitalisation and serological data (22). Daily infection counts were then reconstructed from deconvolution of daily hospitalisation counts and the infection-to-hospitalisation delay distribution, divided by the IHR (21).

Meteorological data –including temperature, absolute humidity, and relative humidity– were obtained from Météo France/PREDICT Services for 112 weather stations nationally. We also included the IPTCC index (*Index PREDICT de transmissivité climatique de la COVID-19*) which characterizes weather conditions favoring SARS-CoV-2 transmission (23). Data were averaged by department.

Mobility data were obtained from Google mobility reports (www.google.com/covid19/mobility/). They describe the change in time spent at points of interest compared to a five-week baseline period (Jan 3 – Feb 6, 2020). The six points of interest are: residential (time spent at home), workplaces, grocery and pharmacy, retail and recreation, parks, and transit stations. Data were available by department.

Finally, we included demographic data by department (population count and density), as obtained from the National Institute of Statistics and Economic Studies (<https://www.insee.fr/fr/statistiques/4989753?sommaire=4989761>).

Statistical analyses

We analysed data collected from week 11-2020 (March 9-15, 2020) to week 20-2021 (May 17-23, 2021), in 92 of the 96 departments of metropolitan France (Figure 1A). Four departments (Maine-et-Loire, Manche, Corse-du-Sud, and Haute-Corse) were excluded due to missing covariates. To remove random noise, daily hospital admission time series were smoothed using local polynomial regression (Figure 1B). Using the package EpiEstim of the R software, we computed the reproduction number R on the smoothed series of each department, over seven-day rolling windows. The reproduction number is the average number of secondary cases caused by an infected individual. We used a gamma distribution with a mean of 7 days and a standard deviation of 5.2 days for the generation time (24).

We developed a log-linear mixed-effects model for the reproduction number R_{ij} in department i in week j :

$$\log(R_{ij}) = \alpha + \beta_k \cdot X_{ijk} + \delta_i + \varepsilon_{ij}$$

where α is an intercept, X_{ijk} are k covariates, β_k are the associated regression coefficients (fixed effects), δ_i are department-level random effects following a Gaussian distribution, and ε_{ij} is a Gaussian error term. We performed the analysis on a weekly scale to reduce temporal autocorrelation. Covariates with daily granularity were averaged by week. Meteorological covariates were introduced into the model either linearly or as cubic B-splines. The model was fitted by maximum likelihood using the R package nlme. Confidence intervals (CI) for the parameters were obtained using a normal approximation to the distribution of the maximum likelihood estimators.

To account for the delayed effects of covariates on hospital admission dynamics, we applied an 11-day lag (5 days for the incubation period and 6 days for the delay between symptom onset and hospital admission (24)) for NPIs, proportion of the population infected, weather conditions, and mobility. For first-dose vaccination, we applied a 20-day lag (14 days for the build-up of immunity (25) and 6 days for the delay between symptom onset and hospital admission). For full vaccination, we reduced this delay to 13 days (25). For VOC proportion (based on testing data), we applied a three-day lag (mean delay between test and hospital admission observed in French data). In a sensitivity analysis, we tested for additional lags (+/- two days) and selected the value leading to the lowest Akaike Information Criteria (AIC).

We first built a baseline model that only included lockdowns, reopenings and moderate restrictions. Then, we incorporated additional covariates using a forward selection procedure. At each step, the covariate leading to the lowest AIC was introduced in the model, until no additional covariate improved the AIC.

To further characterize the individual effects of key covariates on transmission and hospital admissions, we determined the expected R for an average department under two distinct counterfactual scenarios: (i) without the effect of vaccination, and (ii) without the effect of VOC. We then projected the expected number of new hospital admissions at the national level under such scenarios, from January 11, 2021 onwards.

Results

COVID-19 dynamics, NPIs and holidays in France

The median reproduction number R was above 2.5 during the first two weeks and oscillated between 0.6 and 1.9 during the rest of the period (Figure 1C). Three national lockdowns were implemented. The first started on March 17, 2020 and lasted approximately two months. The second and third lockdowns were initiated on October 30, 2020 and April 3, 2021, respectively, and lasted one month each (Figure 1C). The third lockdown started one or two weeks earlier in 19 departments than in the rest of the country (Figure 1D). During the first lockdown, local movements were restricted to a maximum of 1 km around the place of residence for no more than one hour, gatherings in public space were forbidden, and non-essential shops, parks, bars, and restaurants were closed (Additional file 1: Table S1). During the second and third lockdowns, similar measures were imposed, but gatherings of up to 6 people were allowed in public space and parks remained open. During the third lockdown, local movements around the place of residence were allowed up to 10 km. Each lockdown was followed by a reopening phase, during which some (but not all) of the restrictions were lifted. For instance, during the reopening period that followed the first lockdown, local movements were allowed and non-essential shops were open, but inter-regional movements were limited to 100 km around the place of residence and bars and restaurants remained closed. Between lockdowns, several restrictions were applied, including public events limited to 5000 persons and partial closing of cultural places

(Additional file 1: Table S1). An overnight curfew starting at 9pm was first implemented in 16 departments (9 metropolises) on October 17, 2020, followed by 38 other departments on October 24, 2020. At the end of the reopening period that followed the second lockdown, on December 15, 2020, a curfew starting at 8pm was implemented in all departments. This curfew was then moved to an earlier start at 6pm. The measure was first applied in 15 departments on January 2, 2021 and in 10 more departments on January 12, 2021, before extension to the whole country on January 16, 2021 (Figure 1E). On March 20, 2021, the national curfew was pushed to 7pm. There were five regular holiday periods: 2 weeks in February/March, 2 weeks in April, 2 months in July-August, 2 weeks in November and 2 weeks in December/January (“Christmas holidays”). The timing of holiday periods may vary by department (Figure 1F). In addition to regular school holidays, schools remained fully or partially closed during the first lockdown through June 22, 2020, and during the third lockdown (Figure 1C). However, they remained open during the second lockdown.

Multivariable model

In addition to lockdowns, reopenings, and moderate restrictions, the final multivariable model included curfews, school closures, first-dose vaccine coverage, proportion of VOC, temperature, and absolute humidity (Additional file 1: Figure S2). The lowest AIC was obtained for the following lags: 11 days for NPIs and meteorological variables, 22 days for vaccine coverage, and 5 days for the proportion of VOC. The correlation between the observed and fitted values of R was relatively high, although the fitted values presented lower variability than the observed values: the proportion of the variance explained by the full model, including fixed and random effects, reached 63.6%, as estimated based on the conditional R^2 (26) (Additional file 1: Figure S3). Except for the small peak observed in September 2020, the average trajectory of R was well captured by the model (Figure 2A). School closures, excluding summer and Christmas holidays, reduced R by 4.6% (95% CI 1.6-7.4) (Figure 2B). Covariates characterizing summer and Christmas holidays were not statistically significant and therefore not included in the final model. The earlier overnight

curfews started, the stronger their effect on transmission: R was reduced by 18.4% (11.4-24.8) for the 8/9pm curfews and by 34.5% (28.1-40.4) for the 6/7pm curfews. The first lockdown reduced R by 72.9% (71.4-74.2). Combined with school closures, it yielded a reduction in R of 74.0% during the corresponding time period (non-linear effect). The second lockdown reduced R by 70.4% (69.2-71.6), with schools remaining open. The third lockdown reduced R by 60.4% (56.1-64.3). Combined with school closures and the nightly curfew starting at 7pm, it yielded a reduction in R of 75.2%. Reductions in R observed during reopening periods following lockdowns were similar to those measured during lockdowns. The reduction in transmission associated with moderate restrictions ranged from 45.9% to 65.0%, depending on the time period. Furthermore, we estimate that 100% first-dose vaccine coverage would have reduced R by 74.0% (59.4-83.3). In practice, this effect induced a 18.0% and 34.8% reduction in transmission in the departments with the lowest and highest first-dose vaccine coverage at the end of the observation period, respectively. In contrast, a 100% proportion of VOC (mainly Alpha) increased transmission by 46.9% (38.2-56.0) compared with the period during which the historical strain was predominant. Finally, among the weather conditions that we considered, temperature was the factor that improved the model the most, followed by absolute humidity. The AIC was lower when these covariates were included as splines rather than linear effects. We found that R was the lowest at 23.6°C and the highest at 4.7°C. Between the minimum and the maximum values, it increased by up to 31.9% (24.5-39.7) (Figure 2C). For absolute humidity, R was the lowest at 12.0 g/m³ and increased by up to 9.3% (3.2-15.6) to reach a maximum at 4.3 g/m³ (Figure 2D). Considering a national average of weather conditions, we predicted that the transmission rate was the highest in December-February and the lowest in July-August, with an overall amplitude of 41.7% (37.0-46.7) (Figure 2E). Estimated department-level random effects were small, ranging between -1.10^{-8} and 1.10^{-8} (Additional file 1: Figure S4). A model without random effects only had a slightly higher AIC (difference of 2 points).

Counterfactual scenarios

In a first counterfactual scenario, we showed that, for the set of NPIs that were implemented at the time, R would have remained above 1 for three additional weeks (until week 18 vs week 15 in reality) in the absence of vaccination (Figure 3A). Such a scenario would have resulted in a peak of 25,000 new weekly hospital admissions in May 2021 (Figure 3B), higher than observed during the first wave in March 2020.

In a second counterfactual scenario, we showed that, in the absence of VOC, the epidemic could have been contained earlier, with R remaining below 1 in February-April 2021 (Figure 3A) and the observed increase in hospital admissions (third wave for France) would not have occurred (Figure 3B).

Discussion

The methodology used in this study allowed us to disentangle the effects of multiple factors on the reproduction number across French departments, over an extensive observation period spanning three distinct pandemic waves. In particular, our multivariable model demonstrated the strong beneficial effect of NPIs and vaccination on COVID-19 transmission. It also highlighted the detrimental role of emerging variants. Importantly, it enabled quantifying the impact of weather conditions on local transmission while adjusting for other covariates.

The final mixed-effects model presented in this study was able to replicate the temporal dynamics of the reproduction number observed in metropolitan France during the first 14 months of the COVID-19 pandemic. The model closely matched the data but for a short period in September 2020, where the temporary increase in the reproduction number was not captured. This sporadic increase coincided with the end of the summer holidays and a period when the avoidance of social gatherings fell to a low level (27). Yet information on

such individual behaviours at the department level was not available for inclusion in the regression model. However, both the return-to-school and the resurgence of social gatherings presumably contributed to shape the change in transmission, in addition to weather conditions and decisions made by health authorities.

The multivariable model yielded a gradient in the effectiveness of lockdown measures. The first lockdown had the largest impact, followed by the second and the third, which is consistent with international data (11). This could be explained by the more restrictive measures implemented during the first lockdown, but also potentially by increasing pandemic fatigue, which may have resulted in mobility rebounds, more frequent social interactions, and decreasing compliance with preventive measures (28). We found that transmission during the reopening phases following the lifting of lockdowns remained similar to that observed during lockdowns. One possible interpretation is that population behavior did not change immediately with policies: the population may have continued to adhere to public health measures, such as physical distancing during the reopening phase, e.g., due to fear of a COVID-19 rebound and preventive habits taken during lockdowns. Early-pandemic association studies spanning over 131 countries also reported that more time was needed to observe the effects of relaxing NPIs than to detect those resulting from the introduction of new restrictions (7). Moreover, during reopening phases, restrictions were only partially lifted and remained quite intense (Additional file 1: Table S1). Even during intermediate periods of moderate restrictions, the reduction in R was substantial, ranging from 45.9% to 65.0%. This likely reflects the additional contributions of other NPIs such as mask-wearing, hygiene measures, contact tracing and case isolation. Our framework did not allow evaluation of individual effects of such NPIs, which were applied throughout the study period. Moreover, due to collinearity, the effects of specific policies described in Additional file 1-Table S1 (e.g. shop closures, restaurant closures etc.) could not be evaluated separately and were therefore collapsed into broader categories (lockdowns, reopenings and moderate restrictions). Interestingly, overnight curfews considerably reduced SARS-CoV-2

transmission, corroborating results obtained in French Guiana (up to 35% reduction in transmission rates) (29) and in Quebec, Canada (similar reductions in human mobility) (30). Moreover, we found that curfews starting earlier in the evening (6/7pm) had a larger impact on transmission than curfews starting later (8/9pm).

Perhaps more surprisingly, school closures were found to have only a limited effect on transmission. Importantly, school closures did not uniformly affect all households and instead led to disparities in childcare across families that may potentially hinder their effect. As highlighted in other studies, policy decisions about school closures or hybrid school schedules often need to be weighted against the risks of disease transmission to elderly populations associated with increased intergenerational contact rates that exacerbate their vulnerability due to weaker immune systems (31). Our result differs from that of Nader *et al* (1), who found that school closures were one of the most important NPIs in the 60 days following their implementation. In the French context, we believe that the enforcement of mask-wearing and barrier gestures at school was also an impactful NPI that might explain such a difference in magnitude: because these restrictions were applied concurrently, the effect of school closures may have been partially occulted. This observation is consistent with simulation-based scenarios tested in Saudi Arabia (32), where mask-wearing and physical distancing applied in schools were able to drastically reduce the effect of in-person education on SARS-CoV-2 transmission.

The role of weather conditions in COVID-19 transmission has been debated in the literature (15). Here, after controlling for other confounding factors such as NPIs, we found a substantial effect of temperature (up to 31.9% variation in R), followed by absolute humidity (up to 9.3% variation), which led to a 41.7% variation in R between summer and winter months in France. Although both temperature and humidity were associated with SARS-CoV-2 transmission, the overall goodness-of-fit was found to be lower when using the compound IPTCC index. In the future, a different parametrization of this index may yield

better performance. The estimate of the joint effect of temperature and absolute humidity was similar in magnitude and range to individual contributions reported by prior observational (17,33,34), in-vitro (35), and physio-mechanical studies (36). Similar to Sera et al (16), we found a non-linear relationship between weather variables and R. However, in our study, the peak of transmission was identified at both lower temperature and lower absolute humidity values. Pursuing such research in larger countries with more spatial heterogeneities in climatic conditions would be particularly valuable. Interestingly, we found that the temporal seasonality of SARS-CoV-2 transmission was similar to the known seasonality of influenza epidemics in temperate climates (37).

We found that variants of concern (mainly Alpha) increased the reproduction number by 46.9% (38.2-56.0). This is consistent with the effect of Alpha variant on transmission reported in the literature, ranging from about 25% to more than 90% (3,38,39). Notably, the strength of our study is that it estimates the effect of VOC while simultaneously adjusting for weather conditions. Considering that VOC appeared in winter 2021 period and that their proportion substantially increased in February-March 2021, when temperatures were still low and favoured SARS-CoV-2 transmission, such an adjustment was deemed necessary. Our counterfactual scenario analysis showed that, in the absence of VOC, vaccination associated with moderate restrictions and curfew would have been sufficient to contain the historical virus. In the other counterfactual scenario, we showed that, without vaccination, the spread of VOC would have resulted in a peak of hospital admissions higher than observed during the first wave. However, it should be noted that these scenarios were not based on a dynamical transmission model. For instance, we did not account for the fact that if herd immunity was reached in the scenario without vaccination, the number of hospital admissions at the peak would have been lower than predicted by our model. In addition, in practice, it is likely additional measures would have been implemented that would have limited the impact on healthcare.

Although the multivariable model successfully captured the overall temporal dynamics of R , unexplained variability across departments remained. Spatial variation in the reproduction number may arise from underlying socio-demographic determinants such as age distribution, degree of urbanicity, or job market structure (40), which differ between departments. Therefore, we accounted for geographical variation through a department-level random effect on the reproduction number. However, the estimated magnitude of this effect was extremely small. Further, we did not find any statistically significant effect when additionally testing for a potential contribution of population density or population count, suggesting limited impact of population structure. Apart from spatial variation in the reproduction number, small deviations from the average effect of a given intervention may exist among geographical units, due to spatial variations in determinants of population adherence to preventative health measures. Yet introducing the possibility of such variation brings challenges in parameter identifiability. As a result, model sparsity was preferred and department-specific effects on explanatory variables were not considered.

Importantly, our results regarding the effects of NPIs, first-dose vaccination, the proportion of VOC, and weather conditions on the reproduction number are based on a retrospective observational study in which interactions, collinearity and mediation effects may occur. Therefore, the effects estimated here only reflect statistical associations and do not necessarily imply causal mechanisms. In addition, given the differential timeline of interventions, interpretation of their absolute effect should only be made within a specific context of implementation and cannot be directly extrapolated to other settings.

Conclusions

In summary, through a multivariable analysis across 92 French departments, this study allowed disentangling the individual contribution of NPIs, weather, first-dose vaccination, and

VOC proportion on local SARS-CoV-2 transmission during three successive pandemic waves. Our findings highlight the importance of retrospective evaluation of past interventions to inform future decision-making for better epidemic control.

Acknowledgements

We thank Alix Roumagnac, Raphaël Bertrand, and Eurico de Carvalho Filho (PREDICT Services) for providing meteorological data.

Funding

We acknowledge financial support from the Investissement d'Avenir program, the Laboratoire d'Excellence Integrative Biology of Emerging Infectious Diseases program (grant ANR-10-LABX-62-IBEID), Santé publique France, the INCEPTION project (PIA/ANR16-CONV-0005), the European Union's Horizon 2020 research and innovation program under grants 101003589 (RECOVER) and 874735 (VEO), AXA, Groupama, the French Agency for Research on AIDS and Emerging Infectious Diseases via the EMERGEN project (ANRS0151), and the National Research Agency (ANR) through the ANR-Flash call for COVID-19 (grant ANR-20-COVI-0018).

References

1. Nader IW, Zeilinger EL, Jomar D, Zauchner C. Onset of effects of non-pharmaceutical interventions on COVID-19 infection rates in 176 countries. BMC Public Health. 2021 Jul 28;21(1):1472.

2. Haas EJ, Angulo FJ, McLaughlin JM, Anis E, Singer SR, Khan F, et al. Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data [Internet]. Vol. 397, *The Lancet*. 2021. p. 1819–29. Available from: [http://dx.doi.org/10.1016/s0140-6736\(21\)00947-8](http://dx.doi.org/10.1016/s0140-6736(21)00947-8)
3. Davies NG, Abbott S, Barnard RC, Jarvis CI, Kucharski AJ, Munday JD, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England [Internet]. Available from: <http://dx.doi.org/10.1101/2020.12.24.20248822>
4. Landier J, Paireau J, Rebaudet S, Legendre E, Lehot L, Fontanet A, et al. Cold and dry winter conditions are associated with greater SARS-CoV-2 transmission at regional level in western countries during the first epidemic wave [Internet]. Vol. 11, *Scientific Reports*. 2021. Available from: <http://dx.doi.org/10.1038/s41598-021-91798-9>
5. Brauner JM, Mindermann S, Sharma M, Johnston D, Salvatier J, Gavenčiak T, et al. Inferring the effectiveness of government interventions against COVID-19. *Science* [Internet]. 2021 Feb 19;371(6531). Available from: <http://dx.doi.org/10.1126/science.abd9338>
6. Flaxman S, Mishra S, Gandy A, Unwin HJT, Mellan TA, Coupland H, et al. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. *Nature*. 2020 Aug;584(7820):257–61.
7. Li Y, Campbell H, Kulkarni D, Harpur A, Nundy M, Wang X, et al. The temporal association of introducing and lifting non-pharmaceutical interventions with the time-varying reproduction number (R) of SARS-CoV-2: a modelling study across 131 countries. *Lancet Infect Dis*. 2021 Feb;21(2):193–202.
8. Liu Y, Morgenstern C, Kelly J, Lowe R, CMMID COVID-19 Working Group, Jit M. The

- impact of non-pharmaceutical interventions on SARS-CoV-2 transmission across 130 countries and territories. *BMC Med.* 2021 Feb 5;19(1):40.
9. Askitas N, Tatsiramos K, Verheyden B. Estimating worldwide effects of non-pharmaceutical interventions on COVID-19 incidence and population mobility patterns using a multiple-event study. *Sci Rep.* 2021 Jan 21;11(1):1972.
 10. Haug N, Geyrhofer L, Londei A, Dervic E, Desvars-Larrive A, Loreto V, et al. Ranking the effectiveness of worldwide COVID-19 government interventions. *Nat Hum Behav.* 2020 Dec;4(12):1303–12.
 11. Ge Y, Zhang WB, Liu H, Ruktanonchai CW, Hu M, Wu X, et al. Impacts of worldwide individual non-pharmaceutical interventions on COVID-19 transmission across waves and space. *Int J Appl Earth Obs Geoinf.* 2022 Feb;106:102649.
 12. Hale T, Angrist N, Hale AJ, Kira B, Majumdar S, Petherick A, et al. Government responses and COVID-19 deaths: Global evidence across multiple pandemic waves. *PLoS One.* 2021 Jul 9;16(7):e0253116.
 13. Mecnas P, Bastos RT da RM, Vallinoto ACR, Normando D. Effects of temperature and humidity on the spread of COVID-19: A systematic review. *PLoS One.* 2020 Sep 18;15(9):e0238339.
 14. Briz-Redón Á, Serrano-Aroca Á. The effect of climate on the spread of the COVID-19 pandemic: A review of findings, and statistical and modelling techniques. *Progress in Physical Geography: Earth and Environment.* 2020 Oct 1;44(5):591–604.
 15. Poirier C, Luo W, Majumder MS, Liu D, Mandl KD, Mooring TA, et al. The role of environmental factors on transmission rates of the COVID-19 outbreak: an initial assessment in two spatial scales. *Sci Rep.* 2020 Oct 12;10(1):17002.
 16. Sera F, Armstrong B, Abbott S, Meakin S, O'Reilly K, von Borries R, et al. A cross-

sectional analysis of meteorological factors and SARS-CoV-2 transmission in 409 cities across 26 countries. *Nat Commun.* 2021 Oct 13;12(1):5968.

17. Smith TP, Flaxman S, Gallinat AS, Kinoshian SP, Stemkovski M, Unwin HJT, et al. Temperature and population density influence SARS-CoV-2 transmission in the absence of nonpharmaceutical interventions. *Proc Natl Acad Sci U S A* [Internet]. 2021 Jun 22;118(25). Available from: <http://dx.doi.org/10.1073/pnas.2019284118>
18. Baker RE, Yang W, Vecchi GA, Metcalf CJE, Grenfell BT. Assessing the influence of climate on wintertime SARS-CoV-2 outbreaks. *Nat Commun.* 2021 Feb 8;12(1):846.
19. Runkle JD, Sugg MM, Leeper RD, Rao Y, Matthews JL, Rennie JJ. Short-term effects of specific humidity and temperature on COVID-19 morbidity in select US cities. *Sci Total Environ.* 2020 Oct 20;740:140093.
20. Rubin D, Huang J, Fisher BT, Gasparrini A, Tam V, Song L, et al. Association of Social Distancing, Population Density, and Temperature With the Instantaneous Reproduction Number of SARS-CoV-2 in Counties Across the United States. *JAMA Netw Open.* 2020 Jul 1;3(7):e2016099.
21. Hozé N, Paireau J, Lapidus N, Tran Kiem C, Salje H, Severi G, et al. Monitoring the proportion of the population infected by SARS-CoV-2 using age-stratified hospitalisation and serological data: a modelling study. *Lancet Public Health.* 2021 Jun;6(6):e408–15.
22. Lapidus N, Paireau J, Levy-Bruhl D, de Lamballerie X, Severi G, Touvier M, et al. Do not neglect SARS-CoV-2 hospitalization and fatality risks in the middle-aged adult population. *Infect Dis Now.* 2021 Jun;51(4):380–2.
23. Roumagnac A, de Carvalho Filho E, Bertrand R, Banchereau AK, Lahache G. Étude de l'influence potentielle de l'humidité et de la température dans la propagation de la pandémie COVID-19. *Méd Catastr - urgences collect.* 2021 Mar;5(1):87–102.

24. Salje H, Tran Kiem C, Lefrancq N, Courtejoie N, Bosetti P, Paireau J, et al. Estimating the burden of SARS-CoV-2 in France. *Science*. 2020 Jul 10;369(6500):208–11.
25. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N Engl J Med*. 2020 Dec 31;383(27):2603–15.
26. Nakagawa S, Johnson PCD, Schielzeth H. The coefficient of determination and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *J R Soc Interface [Internet]*. 2017 Sep;14(134). Available from: <http://dx.doi.org/10.1098/rsif.2017.0213>
27. CoviPrev : une enquête pour suivre l'évolution des comportements et de la santé mentale pendant l'épidémie de COVID-19 [Internet]. [cited 2022 Jun 24]. Available from: <https://www.santepubliquefrance.fr/etudes-et-enquetes/coviprev-une-enquete-pour-suivre-l-evolution-des-comportements-et-de-la-sante-mentale-pendant-l-epidemie-de-covid-19#block-249165>
28. Di Domenico L, Sabbatini CE, Boëlle PY, Poletto C, Crépey P, Paireau J, et al. Adherence and sustainability of interventions informing optimal control against the COVID-19 pandemic. *Commun Med [Internet]*. 2021 Dec;1(1). Available from: <https://www.nature.com/articles/s43856-021-00057-5>
29. Andronico A, Tran Kiem C, Paireau J, Succo T, Bosetti P, Lefrancq N, et al. Evaluating the impact of curfews and other measures on SARS-CoV-2 transmission in French Guiana. *Nat Commun*. 2021 Mar 12;12(1):1634.
30. Ghasemi A, Daneman N, Berry I, Buchan SA, Soucy JP, Sturrock S, et al. Impact of a nighttime curfew on overnight mobility [Internet]. *bioRxiv*. medRxiv; 2021. Available from: <http://medrxiv.org/lookup/doi/10.1101/2021.04.04.21254906>
31. Bayham J, Fenichel EP. Impact of school closures for COVID-19 on the US health-care

- workforce and net mortality: a modelling study. *Lancet Public Health*. 2020 May;5(5):e271–8.
32. Bisanzio D, Reithinger R, Alqunaibet A, Almudarra S, Alsukait RF, Dong D, et al. Estimating the effect of non-pharmaceutical interventions to mitigate COVID-19 spread in Saudi Arabia. *BMC Med*. 2022 Feb 7;20(1):51.
 33. Raines KS, Doniach S, Bhanot G. The transmission of SARS-CoV-2 is likely comodulated by temperature and by relative humidity. *PLoS One*. 2021 Jul 29;16(7):e0255212.
 34. Collin A, Hejblum BP, Vignals C, Lehot L, Thiébaud R, Moireau P, et al. Using population based Kalman estimator to model COVID-19 epidemic in France: Estimating the effects of non-pharmaceutical interventions on the dynamics of epidemic [Internet]. *bioRxiv*. medRxiv; 2021. Available from: <http://medrxiv.org/lookup/doi/10.1101/2021.07.09.21260259>
 35. Morris DH, Yinda KC, Gamble A, Rossine FW, Huang Q, Bushmaker T, et al. Mechanistic theory predicts the effects of temperature and humidity on inactivation of SARS-CoV-2 and other enveloped viruses. *Elife* [Internet]. 2021 Jul 13;10. Available from: <http://dx.doi.org/10.7554/eLife.65902>
 36. Wang CC, Prather KA, Sznitman J, Jimenez JL, Lakdawala SS, Tufekci Z, et al. Airborne transmission of respiratory viruses. *Science* [Internet]. 2021 Aug 27;373(6558). Available from: <http://dx.doi.org/10.1126/science.abd9149>
 37. Tamerius J, Nelson MI, Zhou SZ, Viboud C, Miller MA, Alonso WJ. Global influenza seasonality: reconciling patterns across temperate and tropical regions. *Environ Health Perspect*. 2011 Apr;119(4):439–45.
 38. Campbell F, Archer B, Laurenson-Schafer H, Jinnai Y, Konings F, Batra N, et al. Increased transmissibility and global spread of SARS-CoV-2 variants of concern as at

June 2021. Euro Surveill [Internet]. 2021 Jun;26(24). Available from:
<http://dx.doi.org/10.2807/1560-7917.ES.2021.26.24.2100509>

39. Gaymard A, Bosetti P, Feri A, Destras G, Enouf V, Andronico A, et al. Early assessment of diffusion and possible expansion of SARS-CoV-2 Lineage 20I/501Y.V1 (B.1.1.7, variant of concern 202012/01) in France, January to March 2021. Euro Surveill [Internet]. 2021 Mar;26(9). Available from: <http://dx.doi.org/10.2807/1560-7917.ES.2021.26.9.2100133>
40. Valdano E, Lee J, Bansal S, Rubrichi S, Colizza V. Highlighting socio-economic constraints on mobility reductions during COVID-19 restrictions in France can inform effective and equitable pandemic response. J Travel Med [Internet]. 2021 Jun 1;28(4). Available from: <http://dx.doi.org/10.1093/jtm/taab045>

Figures




Figure 1: Dynamics of the COVID-19 epidemic and timing of non-pharmaceutical interventions across departments in metropolitan France, week 11-2020 to week 20-2021. (A) Map of departments, colored by region. The two departments marked by an asterisk were excluded, as well as Corsica (not shown), due to missing covariates. (B) Time

series of new daily hospital admissions by department (logarithmic scale). (C) Temporal evolution of the reproduction number R by department and by week, overlaid with the timeline of non-pharmaceutical interventions. Boxplots feature the 2.5th, 25th, 50th, 75th, and 97.5th percentiles. Non-pharmaceutical interventions are shown taking the Rhône department as an example. (D) Timing of the third lockdown by department. (E) Timing of the 6pm curfew by department. (F) Start dates of 2021 winter holidays by department.




Figure 2: Goodness-of-fit and estimated effects of covariates included in the multivariable model on the reproduction number R (in percentage of variation). (A) Trajectory of R estimated by the full model for an average department (red) compared to mean R (black). (B) Effects of linear and categorical covariates. Of note, the effect shown for first-dose vaccine coverage and the proportion of VOC corresponds to a covariate value of 100% (i.e., reflecting a fully vaccinated population and maximum prevalence of variants). (C) Non-linear effect of temperature. (D) Non-linear effect of absolute humidity. (E) Estimated seasonality of COVID-19 based on average temperature and absolute humidity observed over 1981-2010 in metropolitan France. For C, D and E, fitted lines and their 95% confidence intervals show the estimated percentage of variation in R with respect to a baseline set to the value at the trough of the corresponding curve for meteorological variables. For C and D, the range of the x-axis is determined by the 2.5–97.5th percentile of the meteorological variable distribution.




Figure 3: Projected trajectory of R and hospital admissions from January 11, 2021, under different scenarios: the full model, a model without vaccination and a model without VOC. (A) Mean trajectory of R. (B) Number of national-level weekly hospital admissions (summed across all departments).

Supplementary material

NPI	Lockdown 1	Reopening 1	Moderate restrictions 1	Moderate restrictions 2	Lockdown 2	Reopening 2	Moderate restrictions 3	Lockdown 3	Reopening 3
Inter-regional travels	forbidden	100 km or within department	authorized	authorized	forbidden	forbidden	authorized	forbidden	authorized
Local movements	1km 1h	authorized	authorized	authorized	1km 1h	20km 3h	authorized	10km	authorized
Public events	forbidden	5000 persons	5000 persons	5000 persons	forbidden	forbidden	forbidden	forbidden	forbidden
Gatherings in public space	forbidden	10 persons	10 persons	10 persons then authorized	6 persons	6 persons	6 persons	6 persons	6 persons
Non-essential shops	closed	open	open	open	closed	open	open	closed	closed
Cultural places	closed	closed	partial	open	closed	closed	closed	closed	closed
Gym	closed	closed	area-specific	open	closed	closed	closed	closed	closed
Parks and beaches	closed	area-specific	open	open	open	open	open	open	open
Restaurants	closed	closed	open	open	closed	closed	closed	closed	closed
Bars	closed	closed	open	open	closed	closed	closed	closed	closed

Table S1: Description of the main non-pharmaceutical interventions (NPIs) applied during each lockdown, reopening, and period of moderate restrictions. The darker the color, the more restrictive the measure.




Figure S1: Imputation of the proportion of VOC using a logistic regression. The points represent available data and the line represents the logistic fit, taking the Rhône department as an example from December 2020 to May 2021.




Figure S2: Time series of continuous covariates included in the final multivariable model. Each line represents a department and is colored by region. Regions: Auvergne-Rhône-Alpes (ARA), Bourgogne-Franche-Comte (BFC), Bretagne (BRE), Centre-Val de Loire (CVL), Grand Est (GES), Hauts-de-France (HDF), Île-de-France (IDF), Nouvelle-Aquitaine (NAQ), Normandie (NOR), Occitanie (OCC), Provence-Alpes-Côte d’Azur (PAC), and Pays de la Loire (PDL).





Figure S3: Calibration performance: correlation of observed R vs fitted R. Each point represents a department-week pair and is colored by region.

A



B




Figure S4: Map (A) and histogram (B) of department-level random effects.