



**HAL**  
open science

# Energetic processing of hydrocarbons molecular species by ionizing radiation towards the formation of cyclic species

S Indrajith, J Kočíšekt, A Domaracka, P Rousseau, B Huber

► **To cite this version:**

S Indrajith, J Kočíšekt, A Domaracka, P Rousseau, B Huber. Energetic processing of hydrocarbons molecular species by ionizing radiation towards the formation of cyclic species. *Journal of Physics: Conference Series*, 2020, 1412 (14), pp.142018. 10.1088/1742-6596/1412/14/142018 . hal-03792743

**HAL Id: hal-03792743**

**<https://hal.science/hal-03792743>**

Submitted on 9 Jan 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PAPER • OPEN ACCESS

## Energetic processing of hydrocarbons molecular species by ionizing radiation towards the formation of cyclic species

To cite this article: S Indrajith *et al* 2020 *J. Phys.: Conf. Ser.* **1412** 142018

View the [article online](#) for updates and enhancements.

You may also like

- [Self-organized nanopatterning of polycarbonate surfaces by argon ion sputtering](#)  
Meetika Goyal, Divya Gupta, Sanjeev Aggarwal et al.
- [Ion guiding in macro-size insulating capillaries: straight, tapered, and curved shapes](#)  
Takao M Kojima
- [Ion irradiation \(low & high energy ion\) induced surface plasmon resonance in Cu\(10%\)C<sub>70</sub> nanocomposite thin films](#)  
Pooja Sharma and Rahul Singhal

**PRIME**  
PACIFIC RIM MEETING  
ON ELECTROCHEMICAL  
AND SOLID STATE SCIENCE

HONOLULU, HI  
Oct 6–11, 2024

Abstract submission deadline:  
**April 12, 2024**

Learn more and submit!

**Joint Meeting of**  
The Electrochemical Society  
•  
The Electrochemical Society of Japan  
•  
Korea Electrochemical Society

## Energetic processing of hydrocarbons molecular species by ionizing radiation towards the formation of cyclic species

S Indrajith<sup>1</sup>\*, J Kočíšek<sup>2</sup>†, A Domaracka<sup>1</sup>, P Rousseau<sup>1</sup> and B A Huber<sup>1</sup>

<sup>1</sup>Normandie Université, ENSICAEN, UNICAEN, CEA, CNRS, CIMAP, Caen, 14000, France

<sup>2</sup>J. Heyrovský Institute of Physical Chemistry v.v.i., the Czech Academy of Sciences, Prague, 18223, Czech Republic

**Synopsis** Here we report on experimental studies of the collisions between butadiene clusters and three different ionizing radiation sources: 70 eV electrons, 270 nm photons and 3 keV Ar<sup>+</sup> ions. New molecular species are produced by growth mechanism inside of clusters.

Cassini-Huygens mission brought interesting results concerning one of Saturn's biggest satellites, Titan. In fact, it has been shown that this moon's typical orange-brownish haze layers could be due to the presence of aerosols. These aerosols are essential for Titan's climate, radiation balance and atmosphere chemistry. The formation of such aerosols is a complex process ranging from the molecular scale to large particles scale. Polycyclic Aromatic Hydrocarbons (PAHs) and their clusters are important components in terrestrial soot formation. Given this, their presence in Titan's atmosphere can be considered as a nucleation agent, which triggers aerosol formation in Saturn's moon's low temperature atmosphere. However, since it is mainly composed of nitrogen, methane and a small amount of more complex hydrocarbons, understanding the way such PAHs are formed is mandatory [1].


Ionizing radiation sources (photons, electrons and ions) can trigger gas-phase polymerization, ion-molecule reactions and nanoparticle grain surfaces catalysis. These processes contribute to the formation of complex organic synthesis pathways in atmosphere and space. Simple ionization may induce chemical reactions and isomerization processes.

To understand them, simple linear hydrocarbons like butadiene (a non-aromatic molecule) have been chosen to study the formation of more complex cyclic species (aromatic). The irradiation of hydrocarbons clusters has been performed using different radiation sources (electrons, UV photons and ions) (Fig.1). This allowed the de-

\*E-mail: [suvasthika.indrajith@ganil.fr](mailto:suvasthika.indrajith@ganil.fr)

†E-mail: [jaroslav.kocisek@jh-inst.cas.cz](mailto:jaroslav.kocisek@jh-inst.cas.cz)

termination of the balance between ionization induced reactivity and collision induced one. The latter is the typical collision occurring in Titan's atmosphere involving slow and heavy ions (few keV O cations). It has been shown that this triggers reactivity, as observed in C<sub>60</sub> clusters [2] or PAH ones [3]. These experiments were performed at ARIBE, the low energy ion beam facility of GANIL in Caen and in J. Heyrovský Institute of Physical Chemistry in Prague.



**Figure 1.** Mass spectra of butadiene clusters in collision with electrons, photons and ions showing growth products between the monomer and the dimer.

### References

- [1] Long Z *et al* 2018 *Nature Astronomy* **2** 973
- [2] Delaunay R *et al* 2018 *Carbon* **129** 766
- [3] Delaunay R *et al* 2015 *J. Phys. Chem. Lett.* **6** 1536

