



HAL
open science

External Validation and Comparison of Early Readmission's Risk Scores in Older Adults: Prospective Observational Study

Schwab Camille, Hindlet Patrick, Duverger Clarisse, Nazoiri Charifa, Brigitte Sabatier, Fernandez Christine, Korb-Savoldelli Virginie

► **To cite this version:**

Schwab Camille, Hindlet Patrick, Duverger Clarisse, Nazoiri Charifa, Brigitte Sabatier, et al.. External Validation and Comparison of Early Readmission's Risk Scores in Older Adults: Prospective Observational Study. *Journal of the American Medical Directors Association*, 2020, 21 (4), pp.556-557.e4. 10.1016/j.jamda.2019.12.016 . hal-03794018

HAL Id: hal-03794018

<https://hal.science/hal-03794018v1>

Submitted on 30 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EXTERNAL VALIDATION AND COMPARISON OF EARLY READMISSION'S RISK SCORES IN THE ELDERLY: PROSPECTIVE OBSERVATIONAL STUDY

READMISSION SCORES: EXTERNAL VALIDATION & COMPARISON

Schwab Camille¹² (PharmD, PhD Student), Hindlet Patrick¹³⁴ (PharmD, PhD), Duverger Clarisse⁴ (Pharmacy student), Nazoiri Charifa² (Pharmacy student), Sabatier Brigitte²⁵ (PharmD, PhD), Fernandez Christine¹³⁴ (PharmD, PhD), Korb-Savoldelli Virginie²³ (PharmD, PhD)

Authors affiliations:

¹GHU APHP.Sorbonne Université, Sorbonne Université, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique, F75012, Paris, France

²GHU APHP.Centre-Université de Paris, Hôpital européen Georges Pompidou, Service Pharmacie, F-75015, Paris, France

³Université Paris-Sud, Faculté de Pharmacie, Département de Pharmacie Clinique, F-92296, Châtenay-Malabry, France

⁴GHU APHP.Sorbonne Université, Hôpital Saint Antoine, Service Pharmacie, F75012, Paris, France

⁵INSERM UMR 1138 Team 22: Information Sciences to Support Personalized Medicine, Cordelier Research Center, F75006, Paris, France

Corresponding author: Dr Camille SCHWAB

camille.schwab@aphp.fr

GHU APHP.Centre-Université de Paris, Hôpital européen Georges Pompidou

Service Pharmacie

20-40 rue Leblanc, Paris, France

Tel : +331 56 09 51 ; Fax : +331 56 09 25 59

Word counts:

Main text: 748

Number of tables: 1

Number of references: 10

Key words:

Elderly

Readmission

Clinical risk scores

Funding sources: This research did not receive any funding from agencies in the public, commercial or not-for-profit sectors.

Brief summary: We conducted a prospective observational study to externally validate and statistically compare eight clinical risk scores in order to identify elderly at high risk of unplanned hospital readmission within 30 days.

Acknowledgments: The authors would like to acknowledge Dr Germain Perrin for his informed advice for this study and Stella Ghouti for the correction of the manuscript.

Author contributions: Concept, Design: CS, VK-S and PH. Data collection: CD, CD and CN. Interpretation of results: CS, VK-S, PH and CF. Drafting of manuscript: CS. Critical revision of manuscript for important intellectual contents: BS, CS, VK-S, PH and CF

Competing interests: None declared

RESEARCH LETTER

Introduction

Unplanned readmissions in older patients are a prominent and current issue for healthcare professionals and regulators. They can indeed be very deleterious for patients, leading to an increased risk of dependence and functional or psychosocial decline¹ and for hospitals as these readmissions are costly². Moreover, the reduction of readmissions was introduced into guidelines by health system organizations as quality indicators^{3,4}

Several clinical care interventions have been shown to reduce the risk of early unplanned readmission⁵ but they may be difficult to implement and costly in terms of time, money and caregiving if delivered to all patients. Thus, it is necessary to predict patient readmission risk to target the patients who will most likely benefit from these interventions. Clinical risk scores can help physicians and pharmacists select these patients.

Eight clinical risk scores identifying inpatients at risk of unplanned readmission have been published to date (ISAR, TRST, Nomogram, LACE, VIP, SST, HOSPITAL, 80+ score⁶⁻⁹). They were externally validated in diverse populations and with different readmission time lapses.

In order to have a reliable comparison of these eight scores, we aimed to externally validate them in a homogeneous older population and with a unique definition of early unplanned readmission (i.e. emergency department (ED) visit or hospitalization 30 days after discharge).

Methods

An observational prospective cohort study was conducted in seven medical wards of two French university hospitals (February-August 2018). This study was approved by an ethic committee (*Comité de Protection des Personnes Ouest 6*, number 1086 RNI).

All patients aged 65 and older, discharged home (or nursing home) after hospitalization and reachable by telephone the 31st day after discharge, were included.

For each included patient, the eight scores were calculated from the electronic medical records (EMR). On the 31st day after discharge, medical records were checked and/or a phone call was made to ascertain unplanned readmission (ED visit or hospitalization).

Scores contain 3 to 8 questions divided into five categories: demographics, functional impairment, hospitalization, medication, comorbidities.

The discrimination of each score was assessed with the area under the receiver operating characteristic curve (AUC ROC) and metrological characteristics were calculated.

Results

Among 503 included patients, 134 (26.64%) had an early unplanned readmission. All the AUC ROCs of the scores were between 0.56 and 0.65. The VIP score had the lowest AUC ROC (0.56, 95%IC: 0.51-0.61) and LACE had the highest (0.65 95%CI: 0.60-0.70). The metrological characteristics of each score are presented in table I. The ISAR score had the highest sensitivity (76.12%) but the lowest specificity (40.38%). The SST, the VIP and the HOSPITAL scores had the highest specificity but the lowest sensitivity. Furthermore, the SST had also the highest positive predictive value and accuracy.

The HOSPITAL score and LACE index could not be calculated before discharge, because one of the items is the length of stay.

The ISAR, TRST and VIP scores might not be automatable since they call for a reading of the EMR, or even a face to face interview.

Discussion

In this observational prospective study, we externally validated eight clinical risk scores for 30-day unplanned readmissions in an older population. These eight scores had all fair discrimination (AUC ROC greater than 0.5, but lower than 0.7), which is consistent with the literature^{6,10}.

In order to implement an intervention, an early and ideally automatable identification of patients is essential. Three scores (80+, Nomogram and SST) meet these requirements but have very scattered metrological characteristics. The SST has the highest specificity and, therefore, the best ability to correctly rank patients who will not be readmitted as they are at low risk for readmission.

As it seems more convenient to choose a specific test and intervene with patients classified as low risk of readmission but who will be readmitted (i.e. not missing false negative) rather than a sensitive test, the SST, which has the highest specificity, might be preferable. Moreover, it has good negative predictive values (76.52%) and low positive predictive values (50%).

The limitations of our study were that we included only certain medical wards and not surgical specialties, we included only persons discharged home (no rehabilitation) and 5.45% of the participants were lost to follow up.

Conclusion and implications

In this external validation study, we showed that the SST score is a useful tool to predict the risk of early unplanned readmission of the elderly patients. The perspectives of this study are to integrate the score into medical software in order to assess its ability to be automatically calculated, thus facilitating its implementation.

References

1. Arendts G, Fitzhardinge S, Pronk K, Donaldson M, Hutton M, Nagree Y. The impact of early emergency department allied health intervention on admission rates in older people: a non-randomized clinical study. *BMC Geriatr.* 20 mars 2012;12:8.
2. Zook CJ, Savickis SF, Moore FD. Repeated hospitalization for the same disease: a multiplier of national health costs. *Milbank Mem Fund Q Health Soc.* 1980;58(3):454-71.
3. Haute Autorité de Santé. Comment réduire le risque de réhospitalisations évitables des personnes âgées ? 2013.
4. Fischer C, Lingsma HF, Marang-van de Mheen PJ, Kringos DS, Klazinga NS, Steyerberg EW. Is the Readmission Rate a Valid Quality Indicator? A Review of the Evidence. *PLoS ONE* [Internet]. 7 nov 2014 [cité 26 févr 2018];9(11). Disponible sur: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4224424/>
5. Hansen LO, Young RS, Hinami K, Leung A, Williams MV. Interventions to reduce 30-day rehospitalization: a systematic review. *Ann Intern Med.* 18 oct 2011;155(8):520-8.

6. Schwab C, Hindlet P, Sabatier B, Fernandez C, Korb-Savoldelli V. Risk scores identifying elderly inpatients at risk of 30-day unplanned readmission and accident and emergency department visit: a systematic review. *BMJ Open*. 29 juill 2019;9(7):e028302.
7. Donzé JD, Williams MV, Robinson EJ, Zimlichman E, Aujesky D, Vasilevskis EE, et al. International Validity of the HOSPITAL Score to Predict 30-Day Potentially Avoidable Hospital Readmissions. *JAMA Intern Med*. avr 2016;176(4):496-502.
8. Alassaad A, Melhus H, Hammarlund-Udenaes M, Bertilsson M, Gillespie U, Sundström J. A tool for prediction of risk of rehospitalisation and mortality in the hospitalised elderly: secondary analysis of clinical trial data. *BMJ Open*. 18 févr 2015;5(2):e007259.
9. Ben-Chetrit E, Chen-Shuali C, Zimran E, Munter G, Nesher G. A simplified scoring tool for prediction of readmission in elderly patients hospitalized in internal medicine departments. *Isr Med Assoc J IMAJ*. déc 2012;14(12):752-6.
10. Braes T, Moons P, Lipkens P, Sterckx W, Sabbe M, Flamaing J, et al. Screening for risk of unplanned readmission in older patients admitted to hospital: predictive accuracy of three instruments. *Aging Clin Exp Res*. août 2010;22(4):345-51.

Table : Metrological characteristics of the scores

Score	Cut-off (≥)	Specificity (%)	Sensitivity (%)	npv* (%)	ppv* (%)	Accuracy (%)
ISAR	3	40.38	76.12	82.32	31.68	49.90
80+ Score	4	43.63	71.64	80.90	31.58	51.09
LACE	11	50.68	69.40	82.02	33.82	55.67
Nomogram	100	50.95	64.18	79.66	32.21	54.47
TRST	3	72.09	38.81	76.44	33.55	63.22
VIP	2	87.26	22.39	75.68	38.96	69.98
HOSPITAL	7	90.79	11.19	73.79	30.61	69.58
SST	3	91.87	22.39	76.52	50	73.36

*npv = negative predictive value, ppv = positive predictive value