



On the Saito basis and the Tjurina number for plane branches

Yohann Genzmer, M.E. Hernandez

► To cite this version:

Yohann Genzmer, M.E. Hernandez. On the Saito basis and the Tjurina number for plane branches. Transactions of the American Mathematical Society, 2020, 373 (5), pp.3693-3707. 10.1090/tran/8019 . hal-03794511

HAL Id: hal-03794511

<https://hal.science/hal-03794511>

Submitted on 3 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE SAITO BASIS AND THE TJURINA NUMBER FOR PLANE BRANCHES

Y. GENZMER AND M. E. HERNANDES

ABSTRACT. We introduce the concept of a good Saito basis for a plane curve and we explore it to obtain a formula for the minimal Tjurina number in a topological class. In particular, we give a lower bound for the Tjurina number in terms of the Milnor number that allow us to present a positive answer for a question of Dimca and Greuel.

Mathematics Subject Classification: Primary 14H50; Secondary 14B05, 32S05.

keywords: Plane curves, Tjurina number, Saito basis.

1. INTRODUCTION.

Let $S : \{f = 0\}$ be a germ of an irreducible analytic plane curve. An important analytic invariant of S is the Tjurina number $\tau(S) = \dim_{\mathbb{C}} \frac{\mathbb{C}\{x,y\}}{(f)+J(f)}$ where $J(f)$ denotes the Jacobian ideal of f .

In general, the computation of $\tau(S)$ is not easy. For instance, we can obtain it considering a Gröbner basis for the ideal $(f) + J(f)$, or alternatively, it is possible to compute τ by the dimension of $\frac{J(f):(f)}{J(f)}$ (see Theorem 1 in [7]) that is related with the $\mathbb{C}\{x,y\}$ -module $\Omega^1(S)$ of all germs of 1-holomorphic forms $\omega \in \mathbb{C}\{x,y\}dx + \mathbb{C}\{x,y\}dy$ such that f divides $\omega \wedge df$. More precisely, according to K. Saito [9], $\Omega^1(S)$ is freely generated by two elements $\{\omega_1, \omega_2\}$. It will be shown that $\tau(S)$ can be expressed from, among other invariants, the codimension of the ideal (g_1, g_2) where $\omega_i \wedge df = g_i f dx \wedge dy$.

If L denotes a topological class of a plane curve - for instance, given by the characteristic exponents - then the Milnor number $\mu = \dim_{\mathbb{C}} \frac{\mathbb{C}\{x,y\}}{J(f)}$ is constant for any $S : \{f = 0\} \in L$ and $\tau_{\min} \leq \tau(S) \leq \mu$. Generically, an element $S \in L$ is such that $\tau(S) = \tau_{\min}$, so τ_{\min} can be expressed using the topological data that characterizes L . Delorme in [3], presented a formula to compute the generic dimension $d(\beta_0, \beta_1)$ of the moduli space for an irreducible plane curve with characteristic exponents (β_0, β_1) . As $d(2, \beta_1) = 0$ and $d(\beta_0, \beta_1) = \frac{(\beta_0-3)(\beta_1-3)}{2} + \left\lceil \frac{\beta_0}{\beta_1} \right\rceil - 1 - \mu + \tau_{\min}$ (see [6]) we can compute the minimal Tjurina number for this topological class. On the other hand, Peraire in [8] developed an algorithm to compute τ_{\min} by means of a flag of $J(f)$.

This work has been partially supported by the Réseau de Coopération France-Brésil. M. E. Hernandez was also partially supported by CNPq.

In this paper we present a way to express the difference $\mu - \tau$ for a singular irreducible plane curve S when $\Omega^1(S)$ admits a basis $\{\omega_1, \omega_2\}$ of special kind, that we call a *good* Saito basis (see Definition 2).

More specifically, we present a formula (see Theorem 11) to compute the difference between $\mu(S) - \tau(S)$ and $\mu(\tilde{S}) - \tau(\tilde{S})$ where \tilde{S} denotes the strict transform of S . If S is generic in L , then, according to [5], S admits a good basis and this fact allows us to obtain a formula to compute τ_{\min} in L by the sole topological data: the sequence of multiplicities in the canonical resolution or the characteristic exponents for instance. In particular, for irreducible plane curves, we are able to present a lower bound for the minimum Tjurina number in L in terms of the Milnor number that allow us to give an affirmative answer to a question of Dimca and Greuel [4] about the inequality $4\tau > 3\mu$ and obtained simultaneously by Alberich-Carramiñana *et al.* in [1] published in ArXiv a few days before the first version of this paper.

The paper is organized as follows. In the section 2 we present some general properties of a Saito basis. The concept of a good Saito basis is introduced in the section 3 and its properties as well. The section 4 is devoted to the formula for the minimal Tjurina number, a lower bound for the Tjurina number using the Milnor number and consequently an answer to the Dimca-Greuel question.

2. THE SAITO BASIS.

Let $S : \{f = 0\}$ be a germ of an analytic plane curve and consider the $\mathbb{C}\{x, y\}$ -module $\Omega^1(S)$ of all germs of 1-holomorphic forms $\omega \in \mathbb{C}\{x, y\}dx + \mathbb{C}\{x, y\}dy$ such that f divides $\omega \wedge df$. It is equivalent to require that the foliation induced by ω lets invariant S . Saito in [9] shows that $\Omega^1(S)$ is a free module of rank 2 and a basis of $\Omega^1(S)$ is called a Saito basis.

It is not trivial to obtain a Saito basis, but there is a simple criterion to verify if $\{\omega_1, \omega_2\}$ is a basis for $\Omega^1(S)$ (see Theorem, page 270 in [9]).

Theorem (Saito criterion). *The set $\{\omega_1, \omega_2\}$ is a Saito basis for $S : \{f = 0\}$ if and only if $\omega_1 \wedge \omega_2 = u f dx \wedge dy$, where u is a unit in $\mathbb{C}\{x, y\}$.*

This criterion can be interpreted as follows : $\{\omega_1, \omega_2\}$ is a basis for $\Omega^1(S)$ if the tangency locus between the two forms reduces to S .

Below, we present some examples of Saito basis for $S : \{f = 0\}$. All of them will illustrate, in the sequel, various sensitivities of the Saito basis with respect to small perturbations of the curve S . In the whole article, we will keep the same numbering of the examples for the convenience of the reader.

Example (1). The simplest case is when $f = y^p - x^q$, that is $S_1 : \{f = 0\}$ is quasi-homogeneous. In fact, if $\omega_1 = qydx - pxdy$ and $\omega_2 = df$, then

$$\omega_1 \wedge \omega_2 = pqf dx \wedge dy$$

and $\{\omega_1, \omega_2\}$ is a basis for $\Omega^1(S_1)$.

Example (2). If $f = y^5 - x^6 + x^4y^3$ then $S_2 : \{f = 0\}$ is topologically quasi-homogeneous, that is, S_2 presents characteristic exponents $(5, 6)$, but not analytically equivalent to $y^5 - x^6 = 0$. One can show that the set $\{\omega_1, \omega_2\}$ where

$$\begin{aligned}\omega_1 &= \left(-6xy + \frac{16}{15}x^3y^2 - \frac{8}{5}xy^5\right)dx + \left(5x^2 + \frac{4}{3}y^3 + \frac{4}{5}x^2y^4\right)dy \\ \omega_2 &= \left(-6y^2 + \frac{8}{5}x^4 - \frac{12}{5}x^2y^3\right)dx + \left(5xy + \frac{6}{5}x^3y^2\right)dy\end{aligned}$$

satisfy $\omega_1 \wedge \omega_2 = 8f dx \wedge dy$, so $\{\omega_1, \omega_2\}$ is a Saito basis for $\Omega^1(S_2)$.

Example (3). The curve $S_3 : \{f = 0\}$ with $f = y^5 - x^{11} + x^6y^3$ is topologically equivalent to the any curve with characteristic exponents $(5, 11)$ and its strict transform is S_2 . The set $\{\omega_1, \omega_2\}$ where

$$\begin{aligned}\omega_1 &= (605y^2 + 198xy^3 - 88x^6)dx - (275xy + 66x^2y^2)dy \\ \omega_2 &= (605x^4y + 150x^5y^2)dx - (40y^3 + 275x^5 + 90x^6y)dy\end{aligned}$$

satisfy $\omega_1 \wedge \omega_2 = (-24200 - 7920xy)f dx \wedge dy$, so $\{\omega_1, \omega_2\}$ is a Saito basis for $\Omega^1(S_3)$.

Example (4). The class of curve with characteristic exponents the form $(n, n+1)$ has been extensively studied by Zariski [11]. The curve S_4 given by

$$f = y^7 - x^8 - 7x^6y^2 - \frac{147}{8}x^4y^4$$

that, belongs to the latter class, will be shown of a peculiar interest. The forms

$$\begin{aligned}\omega_1 &= \left(8x^2y - \frac{147}{8}x^4 - \frac{3087}{4}x^2y^2 - \frac{21609}{16}y^4\right)dx + \\ &\quad + \left(-7x^3 + \frac{7}{4}xy^2 + \frac{64827}{64}xy^3 + \frac{5145}{8}x^3y\right)dy \\ \omega_2 &= \left(8xy^2 + \frac{1029}{8}x^3y\right)dx + \left(-7x^2y + \frac{7}{4}y^3 - \frac{1029}{8}x^4\right)dy.\end{aligned}$$

produce a Saito basis for $\Omega^1(S_4)$ because $\omega_1 \wedge \omega_2 = -\frac{151263}{64}f dx \wedge dy$.

Given a 1-form $\omega = Adx + Bdy$ we denote by $\nu(\omega) = \min\{\nu(A), \nu(B)\}$ its algebraic multiplicity, where $\nu(H)$ indicates the multiplicity of $H \in \mathbb{C}\{x, y\}$ at $(0, 0) \in \mathbb{C}^2$. Among all the possible basis $\{\omega_1, \omega_2\}$ for $\Omega^1(S)$ we choose some that maximizes the sum $\nu(\omega_1) + \nu(\omega_2)$ that, following the Saito criterion, cannot be bigger than $\nu = \nu(f) = \nu(S)$. For such basis we denote

$$\nu_1 := \nu(\omega_1) \quad \nu_2 := \nu(\omega_2).$$

The following result is immediate and identifies a new analytical invariant of S .

Proposition 1. *The couple (ν_1, ν_2) , up to order, is an analytical invariant of S .*

Remark that the pair (ν_1, ν_2) is not a topological invariant. For instance, following the examples above, for S_1 with $p = 5$ and $q = 6$ we have $(\nu_1, \nu_2) = (1, 4)$. But the curve S_3 which is topological equivalent to S_1 has corresponding pair of multiplicities $(2, 2)$.

From now on, we consider $S : \{f = 0\}$ singular and irreducible (a plane branch) with a Saito basis $\{\omega_1, \omega_2\}$ such that $\omega_i = A_i dx + B_i dy$. In particular, we have

$$(2.1) \quad A_1 B_2 - A_2 B_1 = u f \quad \text{and} \quad A_i \frac{\partial f}{\partial y} - B_i \frac{\partial f}{\partial x} = g_i f$$

where $u(0,0) \neq 0$ and $g_i \in \mathbb{C}\{x, y\}$ is called the cofactor of ω_i .

Applying a generic linear change of coordinates if necessary, we can suppose that for $i = 1, 2$, one has $\nu(A_i) = \nu(B_i) = \nu_i$ and in this coordinates (x, y) the tangent cone of f , i.e. its ν -jet, is $(y + \epsilon x)^\nu$. From now on, we denote $f^\nu = (y + \epsilon x)^\nu$.

Example (1). Consider the irreducible curve S_1 . Suppose by symmetry that $p < q$, we have $\nu(A_1) = \nu(B_1) = \nu_1 = 1$ but $q - 1 = \nu(A_2) > p - 1 = \nu(B_2) = \nu_2$. Consider the change of coordinates $T(x, y) = (x, y - \epsilon x)$ with $\epsilon \neq 0$ we obtain $f_1 = T^*(f) = (y - \epsilon x)^p - x^q$ and the Saito basis $\eta_1 = T^*(\omega_1)$ and df_1

$$\begin{aligned} \eta_1 &= (q(y - \epsilon x) + \epsilon p x) dx - p x dy \\ df_1 &= (-\epsilon p(y - \epsilon x)^{p-1} - q x^{q-1}) dx + p(y - \epsilon x)^{p-1} dy \end{aligned}$$

satisfying the above condition. In addition, $\eta_1 \wedge df_1 = pq f_1 dx \wedge dy$, that is, $g_1 = pq$ and $g_1 = 0$.

Example (2). For the curve S_2 , we have

$$\omega_1 \wedge df = (-30x - 8xy^4) f dx \wedge dy \quad \text{and} \quad \omega_2 \wedge df = (-30y - 12x^2 y^2) f dx \wedge dy,$$

that is, $g_1 = -30x - 8xy^4$ and $g_2 = -30y - 12x^2 y^2$.

Example (3). Considering the curve S_3 , we have $\nu(A_1) = \nu(B_1) = 2$ but $5 = \nu(A_2) > \nu(B_2) = 3$. By the change of coordinates $T(x, y) = (x, x + y)$ we obtain $f_1 = T^*(f) = (y + x)^5 - x^{11} + x^6(y + x)^3$ and $\eta_i = T^*(\omega_i) = (A_i + B_i) dx + B_i dy$ with $\nu(A_1 + B_1) = \nu(B_1) = 2$ and $\nu(A_2 + B_2) = \nu(B_2) = 3$. In addition,

$$\begin{aligned} \eta_1 \wedge df &= (3025(x + y) + 990x(y + x)^2) f_1 dx \wedge dy \\ \eta_2 \wedge df &= (3025x^4 + 990x^5(y + x)) f_1 dx \wedge dy, \end{aligned}$$

consequently, $g_1 = 3025(x + y) + 990x(y + x)^2$ and $g_2 = 3025x^4 + 990x^5(y + x)$.

Example (4). Finally, for S_4 we find

$$\begin{aligned} \omega_1 \wedge df &= \left(56x^2 - \frac{151263}{16} y^3 - \frac{21609}{4} x^2 y \right) f dx \wedge dy \\ \omega_2 \wedge df &= (56xy + 1029x^3) f dx \wedge dy. \end{aligned}$$

Notice that any generator ω_i in a Saito basis $\{\omega_1, \omega_2\}$ has an isolated singularity, that is, $\gcd(A_i, B_i) = 1$. In addition, by (2.1), we have that $\nu(g_i) \geq \nu_i - 1$.

3. GOOD SAITO BASIS AND THE TJURINA NUMBER FOR S .

As we mentioned before, given a Saito basis $\{\omega_1, \omega_2\}$ for $\Omega^1(S)$ we get $\nu_1 + \nu_2 \leq \nu$. In [5], the first author shows the following theorem:

Theorem (Generic Basis Theorem). *In a fixed topological class L , generically any curve S admits a Saito basis satisfying*

$$\nu_1 = \nu_2 = \frac{\nu}{2} \quad \text{if } \nu = \nu(S) \text{ is even}$$

$$\nu_1 = \nu_2 - 1 = \frac{\nu-1}{2} \quad \text{if } \nu = \nu(S) \text{ is odd.}$$

Notice that, generically $\nu_1 + \nu_2$ is maximum. Of course, Example 1 shows that we can obtain $\nu_1 + \nu_2 = \nu$ in other cases. This motivates the following definition.

Definition 2. We say that S (or $\Omega^1(S)$) admits a *good basis* if $\nu_1 + \nu_2 = \nu$.

This section is devoted to present some properties of a good basis. One of them is related with the index $i(S)$ we introduce in the sequel.

Let E be the standard blowing-up of the origin in \mathbb{C}^2 with coordinates (x, y) and suppose that, in the chart (x_1, y_1) such that $E(x_1, y_1) = (x_1, x_1 y_1)$, the strict transform of S goes through $(0, y_1)$.

As before, we consider $f^\nu = (y + \epsilon x)^\nu$.

Definition 3. For any $\omega = A dx + B dy \in \Omega^1(S)$, we denote by $i(\omega) \in \mathbb{N} \cup \{\infty\}$ the valuation given by

$$i(\omega) = \nu_{y_1 = -\epsilon} \left(A^{(\nu(\omega))}(1, y_1) + y_1 B^{(\nu(\omega))}(1, y_1) \right)$$

where $\nu_{y_1 = -\epsilon}(G)$ denotes the multiplicity of $G \in \mathbb{C}\{y_1\}$ at $-\epsilon \in \mathbb{C}$.

Moreover, we denote by $i(S) \in \mathbb{N}$ the integer

$$i(S) = \min_{\omega \in \Omega^1(S)} i(\omega).$$

The value $i(\omega)$ is nothing but the index $\text{Ind}(\mathcal{F}, C, 0)$ introduced in [2] for a germ of foliation \mathcal{F} having C as a smooth invariant curve.

Notice that for a given ω , the index $i(\omega)$ is infinite if and only if ω is dicritical, that is, $A^{(\nu(\omega))}(1, y_1) + y_1 B^{(\nu(\omega))}(1, y_1) = 0$.

However, for any curve $i(S)$ is finite. Indeed, if f is a reduced equation for S then df belongs to $\Omega^1(S)$ and it is not dicritical, thus $i(S) \leq i(df) < \infty$. In particular, if $\omega \in \Omega^1(S)$ is non dicritical, then $i(\omega) \leq \nu(\omega) + 1$.

Example (1). For S_1 , with $p < q$, the considered Saito basis is a good basis. Moreover, $i(\omega_1) = 1$ and $i(\omega_2) = p$.

Example (2). Having a good basis is a property sensitive to perturbation. Indeed, for instance, the basis $\{\omega_1, \omega_2\}$ of S_2 computed in the example is not good. Besides that, we have $i(\omega_1) = 1$ and $i(\omega_2) = 2$.

Actually S_2 does not admit any good basis. In fact, if S_2 admits a good basis $\{\varpi_1, \varpi_2\}$, then we can suppose that $2 = \nu(\varpi_1), 3 = \nu(\varpi_2)$, $\omega_i^{(2)} = c_i \varpi_1^{(2)}$ where $c_i \in \mathbb{C} \setminus \{0\}$ and $\omega_i^{(2)} = A_i^2 dx + B_i^2 dy$. In this way, $\omega_1^{(2)} = \frac{c_2}{c_1} \omega_2^{(2)}$, an absurd.

Example (3). Good basis is not preserved by blowing-up. In fact, S_3 has a good basis, but its strict transform is analytically equivalent to S_2 that does not admit good basis. For S_3 we have $i(\omega_1) = 2$ and $i(\omega_2) = 4$.

Example (4). Finally, S_4 does not have a good basis. We find $i(\omega_1) = 1$ and $i(\omega_2) = 2$.

The next result shows that if S admits a good basis, the index $\mathbf{i}(S)$ is achieved for one of its elements.

Proposition 4. *If S admits a good basis $\{\omega_1, \omega_2\}$ then*

$$\mathbf{i}(S) = \min \{i(\omega_1), i(\omega_2)\}.$$

Proof. By Saito criterion, one has $\omega_1 \wedge \omega_2 = uf$ with $u(0,0) \neq 0$. Since $\nu_1 + \nu_2 = \nu$, one has $\omega_1^{(\nu_1)} \wedge \omega_2^{(\nu_2)} \neq 0$, where $\omega_i^{(\nu_i)} = A_i^{\nu_i} dx + B_i^{\nu_i} dy$. In particular, both forms ω_1 and ω_2 cannot be dicritical and therefore $\min \{i(\omega_1), i(\omega_2)\} < \infty$. Now, consider any form $\omega = P_1\omega_1 + P_2\omega_2 \in \Omega^1(S)$ with $P_i \in \mathbb{C}\{x, y\}$ and $m_i = \nu(P_i)$. Since $P_1^{(m_1)}\omega_1^{(\nu_1)} + P_2^{(m_2)}\omega_2^{(\nu_2)}$ cannot vanish identically, it is the homogeneous part of smallest degree of ω . Therefore

$$\begin{aligned} i(\omega) &= \nu_{y_1=-\epsilon} \left(P_1^{(m_1)}(1, y_1) (A_1^{\nu_1}(1, y_1) + y_1 B_1^{\nu_1}(1, y_1)) \right. \\ &\quad \left. + P_2^{(m_2)}(1, y_1) (A_2^{\nu_2}(1, y_1) + y_1 B_2^{\nu_2}(1, y_1)) \right) \\ &\geq \min \{i(\omega_1), i(\omega_2)\}. \end{aligned}$$

□

In the previous section, we remark that for an element ω_i in a Saito basis we get $\nu(g_i) \geq \nu_i - 1$ and $i(\omega_i) \leq \nu(\omega_i) + 1$. For good basis it is possible to obtain the following result.

Lemma 5. *Given a good basis $\{\omega_1, \omega_2\}$ for S , if $\nu(g_i) \geq \nu_i$ then $i(\omega_i) = \nu_i + 1$.*

Proof. By symmetry let us consider $i = 1$ and suppose that $\nu(g_1) \geq \nu_1$. The $(\nu_1 - 1 + \nu)$ -jet of

$$A_1 \frac{\partial f}{\partial y} - B_1 \frac{\partial f}{\partial x} = g_1 f \quad \text{is} \quad A_1^{(\nu_1)} \nu(y + \epsilon x)^{\nu-1} - B_1^{(\nu_1)} \nu \epsilon(y + \epsilon x)^{\nu-1} = 0,$$

thus $A_1^{(\nu_1)} = \epsilon B_1^{(\nu_1)}$. On the other hand the ν -jet of $A_1 B_2 - A_2 B_1 = uf$ where $u(0,0) \neq 0$ reduces to

$$A_1^{(\nu_1)} B_2^{(\nu_2)} - A_2^{(\nu_2)} B_1^{(\nu_1)} = B_1^{(\nu_1)} \left(\epsilon B_2^{(\nu_2)} - A_2^{(\nu_2)} \right) = u(0,0) (y + \epsilon x)^\nu.$$

Thus, there exists some constant $c \neq 0$ such that $B_1^{(\nu_1)} = c(y + \epsilon x)^{\nu_1}$. Therefore, ω_1 can be written

$$\omega_1 = \frac{c}{\nu_1 + 1} d \left((y + \epsilon x)^{\nu_1 + 1} \right) + \text{h.o.t.}$$

thus $i(\omega_1) = \nu_1 + 1$. □

Notice that the above proof ensures that the inequality $\nu(g_i) \geq \nu_i$ cannot hold for both elements in a good basis. Moreover, given a good basis for $\Omega^1(S)$ we can always get a good basis with some nice properties. To do this we present the following lemmas.

Lemma 6. *If $\Omega^1(S)$ admits a good basis $\{\omega_1, \omega_2\}$, then we can suppose that*

$$i(\omega_1) = \mathbf{i}(S) \quad \text{and} \quad \nu(g_1) = \nu_1 - 1.$$

Proof. By symmetry we can suppose that $i(\omega_1) = \mathbf{i}(S)$.

Case 1. If $i(\omega_2) = i(\omega_1)$, then, as mentioned above, for $i = 1$ or 2 , one has $\nu(g_i) = \nu_i - 1$. Switching maybe the two forms, we can suppose that ω_1 satisfies the conclusion of the lemma.

Case 2. Suppose now that $i(\omega_1) < i(\omega_2)$.

Subcase 2.a if $\nu_1 \leq \nu_2$, we consider, the family $\{\omega_1, \overline{\omega_2}\}$, where $\overline{\omega_2} = \omega_2 + cx^{\nu_2-\nu_1}\omega_1$ and $c \in \mathbb{C}$. For a generic value of c , we still have a good basis for S . Moreover, the ν_2 -jet of $\overline{\omega_2}$ is

$$\left(A_2^{(\nu_2)} + cx^{\nu_2-\nu_1}A_1^{(\nu_1)}\right)dx + \left(B_2^{(\nu_2)} + cx^{\nu_2-\nu_1}B_1^{(\nu_1)}\right)dy.$$

Thus, to evaluate its index, one writes

$$\begin{aligned} i(\overline{\omega_2}) &= \nu_{y=-\epsilon} \left(A_2^{(\nu_2)}(1, y) + cA_1^{(\nu_1)}(1, y) + y \left(B_2^{(\nu_2)}(1, y) + cB_1^{(\nu_1)}(1, y) \right) \right) \\ &= \nu_{y=-\epsilon} \left(A_2^{(\nu_2)}(1, y) + yB_2^{(\nu_2)}(1, y) + c \left(A_1^{(\nu_1)}(1, y) + yB_1^{(\nu_1)}(1, y) \right) \right) = i(\omega_1). \end{aligned}$$

Thus we are led to the previous case (1).

Subcase 2.b. Finally, if $\nu_1 > \nu_2$, suppose that $\nu(g_1) \geq \nu_1$, then by Lemma 5 we have $i(\omega_1) = \nu_1 + 1$. Consequently $i(\omega_1) > \nu_2 + 1$ and then $i(\omega_2) > \nu_2 + 1$. If ω_2 is not dicritical, the inequality above leads to a contradiction, thus ω_2 is dicritical. Therefore, it can be seen that $\nu(g_2) = \nu_2 - 1$. Let us consider now $\overline{\omega_1} = \omega_1 + x^{\nu_1-\nu_2}\omega_2$. Then, the family $\{\overline{\omega_1}, \omega_2\}$ is still a good basis and one has

$$\begin{aligned} \overline{\omega_1} \wedge df &= \overline{g_1} f dx \wedge dy \quad \text{with } \nu(\overline{g_1}) = \nu_1 - 1 \\ i(\overline{\omega_1}) &= i(\omega_1) = \mathbf{i}(S). \end{aligned}$$

□

In addition, from a basis for $\Omega^1(S)$ we can get a basis satisfying the following lemma.

Lemma 7. *Given a basis $\{\omega_1, \omega_2\}$ for $\Omega^1(S)$ with $i(\omega_1) \leq i(\omega_2)$ we can suppose that*

$$\gcd\left(B_i, \frac{\partial f}{\partial y}\right) = 1, \quad \text{for } i = 1, 2.$$

Proof. Suppose that $H = \gcd\left(B_1, B_2, \frac{\partial f}{\partial y}\right)$. Since by (2.1) $A_1B_2 - A_2B_1 = uf$, H would divide f . As $\frac{\partial f}{\partial y}$ and f are relatively prime, we get

$$(3.1) \quad \gcd\left(B_1, B_2, \frac{\partial f}{\partial y}\right) = 1.$$

Now consider the family $\{\overline{\omega_1} = \omega_1 + P_1\omega_2, \overline{\omega_2} = \omega_2 + P_2\omega_1\}$ where $P_i \in \mathbb{C}\{x, y\}$ with $\nu(P_i) \gg 1$. Note that for P_i of algebraic multiplicity big enough, the forms

$$\begin{aligned} \overline{\omega_1} &= (A_1 + P_1A_2)dx + (B_1 + P_1B_2)dy = \overline{A_1}dx + \overline{B_1}dy \\ \overline{\omega_2} &= (A_2 + P_2A_1)dx + (B_2 + P_2B_1)dy = \overline{A_2}dx + \overline{B_2}dy \end{aligned}$$

satisfy $\nu(\overline{\omega_i}) = \nu(\overline{A_i}) = \nu(\overline{B_i}) = \nu_i$, and $i(\omega_1) = i(\overline{\omega_1}) \leq i(\overline{\omega_2})$.

Moreover, $\{\overline{\omega_1}, \overline{\omega_2}\}$ is a basis for $\Omega^1(S)$. Now the relation (3.1) ensures that for a generic choice of the P'_i s, $i = 1, 2$ - in the sense of Krull -, one has

$$\gcd\left(\overline{B_i}, \frac{\partial f}{\partial y}\right) = 1.$$

□

As a consequence we obtain the following.

Corollary 8. *For any basis $\{\omega_1, \omega_2\}$ for $\Omega^1(S)$ satisfying the previous Lemma we have*

$$\gcd(B_i, g_i) = \gcd\left(\frac{\partial f}{\partial y}, g_i\right) = 1.$$

Proof. As $A_i \frac{\partial f}{\partial y} - B_i \frac{\partial f}{\partial x} = g_i f$, if $1 \neq H = \gcd(B_i, g_i)$ then H must divide $A_i \frac{\partial f}{\partial y}$. By the previous lemma, $\gcd(B_i, \frac{\partial f}{\partial y}) = 1$ so H divides A_i , a contradiction because ω_i has an isolated singularity.

Suppose $H' = \gcd\left(\frac{\partial f}{\partial y}, g_i\right)$, so H' divides $B_i \frac{\partial f}{\partial x}$. As $\gcd\left(\frac{\partial f}{\partial y}, \frac{\partial f}{\partial x}\right) = \gcd(B_i, g_i) = 1$, we must have $H' = 1$. □

In particular, the above lemma allow us to consider a good Saito basis $\{\omega_1, \omega_2\}$ with $i(S) = i(\omega_1)$ and $\gcd\left(B_i, \frac{\partial f}{\partial y}\right) = \gcd(B_i, g_i) = \gcd\left(\frac{\partial f}{\partial y}, g_i\right) = 1$.

Lemma 9. *If $S : \{f = 0\}$ admits a good basis satisfying the previous conditions, then the intersection of the tangent cone of*

- (1) g_1 and g_2 ,
- (2) B_i and g_i , for $i = 1, 2$,
- (3) B_i and $\frac{\partial f}{\partial y}$, for $i = 1, 2$

is empty or equal to $y + \epsilon x = 0$.

Proof. The ν -jet of (2.1) is

$$(3.2) \quad A_1^{(\nu_1)} B_2^{(\nu_2)} - A_2^{(\nu_2)} B_1^{(\nu_1)} = c(y + \epsilon x)^\nu.$$

where $c \neq 0$ and $\epsilon \in \mathbb{C}$. Now, for $i = 1, 2$, both following relations $A_i^{(\nu_i)} - \epsilon B_i^{(\nu_i)} = 0$ cannot be true all together since it would yield a contradiction with the relation (3.2). Suppose the relation above is not true for at least $i = 1$, then the cofactor relations ensures that

$$A_1^{(\nu_1)} - \epsilon B_1^{(\nu_1)} = \frac{1}{\nu} g_1^{(\nu(g_1))} (y + \epsilon x).$$

Combining the above relations yields

$$g_1^{(\nu(g_1))} B_2^{(\nu_2)} - g_2^{(\nu(g_2))} B_1^{(\nu_1)} = c\nu (y + \epsilon x)^{\nu-1}, \quad \text{or} \quad g_1^{(\nu(g_1))} B_2^{(\nu_2)} = c\nu (y + \epsilon x)^{\nu-1}$$

from which is derived (1) and (2). The point (3) follows from the fact that the tangent cone of $\frac{\partial f}{\partial y}$ and f are the same. □

In what follows we denote by $I_P(G, H)$ the intersection multiplicity of $G, H \in \mathbb{C}\{x, y\}$ at the point $P \in \mathbb{C}^2$. If $P = (0, 0)$ then we write $I(G, H) := I_P(G, H)$, that is, $I(G, H) = \dim_{\mathbb{C}} \frac{\mathbb{C}\{x, y\}}{(G, H)}$.

An important topological invariant for $S : \{f = 0\}$ is the Milnor number μ which can be computed by

$$(3.3) \quad \mu := I\left(\frac{\partial f}{\partial y}, \frac{\partial f}{\partial x}\right) = \sum_{i=1}^N \nu_{(i)}(\nu_{(i)} - 1)$$

where $\nu_{(i)}$; $i = 1, \dots, N$ denote the sequence of multiplicities in the canonical resolution of S . In addition, by Zariski (see (2.4) in [11]), we have

$$(3.4) \quad I\left(\frac{\partial f}{\partial y}, f\right) = \mu + \nu - 1.$$

Combining the Lemma 6 and the above result we can obtain an expression for $I(g_1, g_2)$.

Lemma 10. *If g_1 and g_2 are the cofactors for a good basis for $\Omega^1(S)$, then $I(g_1, g_2)$ is finite and*

$$I(g_1, g_2) = I\left(\frac{\partial f}{\partial y}, B_1\right) - I(B_1, g_1) - \nu + 1.$$

Proof. By Lemma 6 we have $\nu(g_1) = \nu_1 - 1 < \nu$. As f is irreducible it follows that $\gcd(f, g_1) = 1$ and $I\left(f \frac{\partial f}{\partial y}, g_1\right) < \infty$. So, from (2.1) that

$$I\left(f \frac{\partial f}{\partial y}, g_1\right) = I\left(A_1 B_2 \frac{\partial f}{\partial y} - A_2 B_1 \frac{\partial f}{\partial y}, g_1\right) = I\left(B_1 B_2 \frac{\partial f}{\partial x} - A_2 B_1 \frac{\partial f}{\partial y}, g_1\right) = I(B_1 g_2 f, g_1).$$

Hence,

$$(3.5) \quad I(g_1, g_2) = I\left(\frac{\partial f}{\partial y}, g_1\right) - I(B_1, g_1).$$

The Corollary 8 insures that $\frac{\partial f}{\partial y}$ and g_1 are coprime. So, by (3.5) and using (3.4) we obtain

$$\begin{aligned} I(g_1, g_2) &= I\left(\frac{\partial f}{\partial y}, g_1\right) + I\left(\frac{\partial f}{\partial y}, f\right) - I\left(\frac{\partial f}{\partial y}, f\right) - I(B_1, g_1) \\ &= I\left(\frac{\partial f}{\partial y}, g_1 f\right) - I\left(\frac{\partial f}{\partial y}, f\right) - I(B_1, g_1) \\ &= I\left(\frac{\partial f}{\partial y}, A_1 \frac{\partial f}{\partial y} - B_1 \frac{\partial f}{\partial x}\right) - (\mu + \nu - 1) - I(B_1, g_1) \\ &= I\left(\frac{\partial f}{\partial y}, B_1\right) - \nu + 1 - I(B_1, g_1). \end{aligned}$$

□

Let us consider the Tjurina number τ of a plane curve $S : \{f = 0\}$, that is,

$$\tau := \dim_{\mathbb{C}} \frac{\mathbb{C}\{x, y\}}{\left(f, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial x}\right)}.$$

Zariski (see Theorem 1 in [10]) considered the torsion submodule $T\Omega_{\mathcal{O}/\mathbb{C}}^1$ of the Kähler differential module $\Omega_{\mathcal{O}/\mathbb{C}}^1$ over $\mathcal{O} = \frac{\mathbb{C}\{x,y\}}{(f)}$ and he showed that $\tau = \dim_{\mathbb{C}} T\Omega_{\mathcal{O}/\mathbb{C}}^1$. On the other hand, Michler (Theorem 1 in [7]) proved that $T\Omega_{\mathcal{O}/\mathbb{C}}^1$ is isomorphic as \mathcal{O} -module, to $\frac{(\frac{\partial f}{\partial y}, \frac{\partial f}{\partial x}) : (f)}{(\frac{\partial f}{\partial y}, \frac{\partial f}{\partial x})}$. As $(\frac{\partial f}{\partial y}, \frac{\partial f}{\partial x}) : (f)$ is precisely the cofactor ideal of S , that is, (g_1, g_2) , one has

$$\tau = \dim_{\mathbb{C}} \frac{(g_1, g_2)}{(\frac{\partial f}{\partial y}, \frac{\partial f}{\partial x})} = \dim_{\mathbb{C}} \frac{\mathbb{C}\{x, y\}}{(\frac{\partial f}{\partial y}, \frac{\partial f}{\partial x})} - \dim_{\mathbb{C}} \frac{\mathbb{C}\{x, y\}}{(g_1, g_2)} = \mu - I(g_1, g_2),$$

that is,

$$\mu - \tau = I(g_1, g_2).$$

Denoting $\tilde{\mu}$ the Milnor number of \tilde{S} , the strict transform of S under a standard blowing up of the origin of \mathbb{C}^2 , we provide in the next theorem a precise relation between $\mu - \tau$ and $\tilde{\mu} - \tilde{\tau}$ by means of the analytic invariants we have introduced previously for curves that admit a good basis.

Theorem 11. *If S admits a good basis, then*

$$\mu - \tau = \tilde{\mu} - \tilde{\tau} + (\nu_1 - 1)(\nu_2 - 1) + i(S) - 1.$$

Proof. By symmetry, one can suppose $i(S) = \min\{i(\omega_1), i(\omega_2)\} = i(\omega_1)$. By Lemma 9 and the Max-Noether formula one has,

$$\mu - \tau = I(g_1, g_2) = I_{(0, -\epsilon)}(\tilde{g}_1, \tilde{g}_2) + \nu(g_1)\nu(g_2),$$

where $\tilde{H} := E^*(H)$ and E denotes the standard blowing-up of the origin in \mathbb{C}^2 . In addition, the previous lemma and Lemma 9, yield

$$\begin{aligned} I(g_1, g_2) &= I\left(\frac{\partial f}{\partial y}, B_1\right) - I(B_1, g_1) - \nu + 1 \\ &= I_{(0, -\epsilon)}\left(\frac{\partial \tilde{f}}{\partial y}, \tilde{B}_1\right) - I_{(0, -\epsilon)}(\tilde{B}_1, \tilde{g}_1) + \nu\left(\frac{\partial f}{\partial y}\right)\nu(B_1) - \nu(B_1)\nu(g_1) - \nu + 1. \end{aligned}$$

If $\tilde{\omega}_i = \frac{E^*\omega_i}{x^{\nu_i}}$, then the Saito criterion yields $x^{\nu_1}\tilde{\omega}_1 \wedge x^{\nu_2}\tilde{\omega}_2 = \tilde{u}x^{\nu}\tilde{f}dx \wedge dy$. Since we have a good basis, that is, $\nu_1 + \nu_2 = \nu$, one has $\tilde{\omega}_1 \wedge \tilde{\omega}_2 = u\tilde{f}dx \wedge dy$. Locally around $(0, -\epsilon)$ for $i = 1, 2$ we have

$$\tilde{\omega}_i = (A_i^{\nu_i}(1, y) + yB_i^{\nu_i}(1, y) + x(\cdots))dx + x(B_i^{\nu_i}(1, y) + (\cdots))dy$$

We notice that the form

$$\bar{\omega}_2 = \frac{1}{x}\left(\tilde{\omega}_2 - \frac{A_2^{\nu_2}(1, y) + yB_2^{\nu_2}(1, y)}{A_1^{\nu_1}(1, y) + yB_1^{\nu_1}(1, y)}\tilde{\omega}_1\right)$$

is holomorphic at $(0, -\epsilon)$ and $\{\tilde{\omega}_1, \bar{\omega}_2\}$ is a Saito basis for $\tilde{S} : \{\tilde{f} = 0\}$. A computation shows that the cofactor associated to $\tilde{\omega}_1$ is written $g'_1 = \tilde{g}_1 + \nu\tilde{B}_1$. Moreover, one has $\tilde{\omega}_1 = (\tilde{A}_1 + y\tilde{B}_1)dx + x\tilde{B}_1dy = A'dx + B'dy$. Now,

$$(A_1^{\nu_1}(1, y) + yB_1^{\nu_1}(1, y) + x(\cdots))\frac{\partial \tilde{f}}{\partial y} - x\tilde{B}_1\frac{\partial \tilde{f}}{\partial x} = g'_1\tilde{f}.$$

If x divides g'_1 then $\tilde{\omega}_1$ would be dicritical and this is not possible. Therefore,

$$I_{(0,-\epsilon)}\left(x, g'_1 \tilde{f}\right) = I_{(0,-\epsilon)}\left(A_1^{\nu_1}(1, y) + yB_1^{\nu_1}(1, y), x\right) + I_{(0,-\epsilon)}\left(x, \frac{\partial \tilde{f}}{\partial y}\right).$$

and, by Corollary 8, $I_{(0,-\epsilon)}(x, g'_1) = i(\omega_1) - 1 = i(S) - 1$.

Notice that \tilde{B}_1 and g'_1 cannot have a common divisor, since it would be a common divisor of \tilde{g}_1 and \tilde{B}_1 that is impossible by Lemma 7. So,

$$I_{(0,-\epsilon)}(\tilde{B}_1, \tilde{g}_1) = I_{(0,-\epsilon)}(x\tilde{B}_1, g'_1) - i(S) + 1 = I_{(0,-\epsilon)}(B'_1, g'_1) - i(S) + 1.$$

Moreover,

$$\begin{aligned} I_{(0,-\epsilon)}\left(\frac{\partial \tilde{f}}{\partial y}, \tilde{B}_1\right) &= I_{(0,-\epsilon)}\left(\frac{\partial \tilde{f}}{\partial y}, B'_1\right) - I_{(0,-\epsilon)}\left(\frac{\partial \tilde{f}}{\partial y}, x\right) \\ &= I_{(0,-\epsilon)}\left(\frac{\partial \tilde{f}}{\partial y}, B'_1\right) - I_{(0,-\epsilon)}(\tilde{f}, x) + 1. \end{aligned}$$

So, as $\frac{\partial \tilde{f}}{\partial y} = \frac{\partial f}{\partial y}$ and combining all the above relation yields

$$\begin{aligned} \mu - \tau &= I_{(0,-\epsilon)}\left(\frac{\partial \tilde{f}}{\partial y}, B'_1\right) - I_{(0,-\epsilon)}(\tilde{f}, x) + 1 - \left(I_{y=-\epsilon}(B'_1, g'_1) - i(S) + 1\right) \\ &\quad + \nu\left(\frac{\partial f}{\partial y}\right)\nu(B_1) - \nu(B_1)\nu(g_1) - \nu + 1 \\ &= I_{(0,-\epsilon)}(g'_1, g'_2) + (\nu - 1)\nu_1 - \nu_1\nu(g_1) - \nu + i(S). \end{aligned}$$

As $I_{(0,-\epsilon)}(g'_1, g'_2) = \tilde{\mu} - \tilde{\tau}$ and $\nu(g_1) = \nu_1 - 1$, we obtain finally

$$\mu - \tau = \tilde{\mu} - \tilde{\tau} + (\nu_1 - 1)(\nu_2 - 1) + i(S) - 1.$$

□

Let us analyze the examples previously considered.

Example (1). For S_1 we have a good basis with $\nu_1 = 1$, $\nu_2 = p - 1$ and $i(S_1) = 1$, then $\mu - \tau = 0$ as classically known.

Example (2). Notice that for S_2 we have $i(S_2) = i(\omega_1) = 1$, $\nu_1 = \nu_2 = 2$ and \tilde{S}_3 is regular, so $\tilde{\mu} - \tilde{\tau} = 0$. In this way,

$$1 = I(g_1, g_2) = \mu - \tau = 0 + (2 - 1)(2 - 1) + 1 - 1.$$

So, the formula in the previous theorem holds although S_2 does not admit any good basis.

Example (3). For S_3 we get $i(S_3) = i(\omega_1) = 2$, $\nu_1 = 2$, $\nu_2 = 3$ and \tilde{S}_3 is analytically equivalent to S_2 , so $\tilde{\mu} - \tilde{\tau} = 1$. In this way,

$$4 = I(g_1, g_2) = \mu - \tau = 1 + (2 - 1)(3 - 1) + 2 - 1.$$

Example (4). As we presented above, S_4 does not have a good basis. We have $i(S_4) = i(\omega_1) = 1$, $\nu_1 = \nu_2 = 3$ and $\tilde{\mu} - \tilde{\tau} = 0$, but in this case,

$$5 = I(g_1, g_2) = \mu - \tau \neq 4 = 0 + (3 - 1)(3 - 1) + 1 - 1.$$

A more detailed analysis shows that Lemma 9 is not valid in this case because the intersection of the tangent cone of g_1 and g_1 is $x = 0$ that is distinct to the tangent cone $y = 0$ of S_4 .

4. THE MINIMAL TJURINA NUMBER AND THE DIMCA-GREUEL QUESTION FOR PLANE BRANCHES.

Given a curve S , we denote by $L = L(S)$ its topological class. Although the Milnor number is constant in L , the same is not true for the Tjurina number $\tau(S)$. On the other hand, as $\tau(S)$ is upper semicontinuous, the minimum value τ_{\min} for curves in L is achieved generically and it should be computed by the sole topological data (see Chapitre III, Appendice of [11] by Teissier).

For a topological class L given by characteristic exponents (β_0, β_1) , Delorme in [3] presented a formula for the dimension of the generic component of the Moduli space that allow us to compute τ_{\min} . For an arbitrary topological class, Péraire (see [8]) presented an algorithm to compute the τ_{\min} using the flag of the Jacobian ideal.

In this section, using the last theorem and results of [5], we give an alternative method to compute τ_{\min} in a fixed topological class L and as a bonus we are able to answer a question of Dimca-Greuel for the irreducible plane curves.

If S admits a good basis we can not insure that the same is valid for \tilde{S} (see Example (3)). However, this property is true generically.

Theorem 12. *Let L the topological class of plane branch given by the characteristic exponents $(\beta_0, \beta_1, \dots, \beta_s)$, τ_{\min} the minimal Tjurina number in L and $\tilde{\tau}_{\min}$ the minimal Tjurina number in \tilde{L} . If S is generic in L , then*

$$(4.1) \quad \mu - \tau_{\min} = \tilde{\mu} - \tilde{\tau}_{\min} + \left(\left\lfloor \frac{\beta_0}{2} \right\rfloor - 1 \right) \left(\beta_0 - \left\lfloor \frac{\beta_0}{2} \right\rfloor - 1 \right) + i(S) - 1.$$

Moreover, if $n = \left\lfloor \frac{\beta_1}{\beta_1 - \beta_0} \right\rfloor$, then $i(S) = \left\lfloor \frac{\beta_0}{2} \right\rfloor + 1 - p_1(S)$, where $p_1(S)$ can be computed in the following way:

$$\begin{aligned} \bullet \text{ if } \beta_0 \text{ is even then } p_1(S) &= \begin{cases} 1 & \text{if } n = 2 \\ 1 & \text{if } \beta_1 \text{ is even} \\ \frac{n-1}{2} & \text{if } \beta_1 \text{ is odd and } n \text{ odd} \\ \frac{n-2}{2} & \text{if } \beta_1 \text{ is odd and } n \text{ is even} \end{cases} \\ \bullet \text{ if } \beta_0 \text{ is odd then } p_1(S) &= \begin{cases} 0 & \text{if } n = 2 \\ 1 & \text{if } \beta_1 \text{ is odd} \\ \frac{n-3}{2} & \text{if } \beta_1 \text{ is even and } n \text{ odd} \\ \frac{n-2}{2} & \text{if } \beta_1 \text{ is even and } n \text{ is even.} \end{cases} \end{aligned}$$

Proof. Suppose that $\beta_0 = \nu(S)$ is even. According to the Generic Basis Theorem, S admits a good basis $\{\omega'_1, \omega'_2\}$ with $\nu(\omega'_1) = \nu(\omega'_2) = \frac{\beta_0}{2}$. For generic $\alpha_1, \alpha_2 \in \mathbb{C}$ $\{\omega_1 = \omega'_1 + \alpha_2 \omega'_2, \omega_2 = \omega'_2 + \alpha_1 \omega'_1\}$ remain a good basis with $\nu_1 = \nu_2 = \frac{\beta_0}{2}$ and

$$i(\omega_1) = i(\omega_2).$$

Now, according to [5] - using the notations of the mentioned paper, it refers to the case $\delta_1 = 0$ and $\delta_2 = 1$ - we obtain $\nu_1 + 1 = \frac{\beta_0}{2} + 1 = \sum_{q \in \mathbb{P}^1} \text{Ind}(\tilde{\mathcal{F}}, C, q) = i(\omega_1) + p_1(S)$, that is,

$$i(S) = i(\omega_1) = \nu_1 + 1 - p_1(S) = \frac{\beta_0}{2} + 1 - p_1(S) = \left\lfloor \frac{\beta_0}{2} \right\rfloor + 1 - p_1(S)$$

where $p_1(S)$ is described in [5].

Now, suppose β_0 is odd and let $\{\omega'_1, \omega'_2\}$ be a Saito basis for $S \cup l$ with l a generic line that without loss of generality can be considered $x = 0$. As $\nu(S \cup l)$ is even, by the same above argument, we can suppose that

$$\nu(\omega'_1) = \nu(\omega'_2) = \frac{\beta_0 + 1}{2} = \left\lfloor \frac{\beta_0}{2} \right\rfloor + 1 \quad \text{and} \quad i(S \cup l) = i(\omega'_1) = i(\omega'_2).$$

Denoting $\omega'_i = (a_i(y) + x(\cdots))dx + x(\cdots)dy$ and considering generic $\alpha_1, \alpha_2 \in \mathbb{C}$ we obtain a good Saito basis $\{\omega_1 = \omega'_1 + \alpha_2 \omega'_2, \omega_2 = \omega'_2 + \alpha_1 \omega'_1\}$ such that $\nu(a_1(y) + \alpha_2 a_2(y)) = \nu(a_2(y) + \alpha_1 a_1(y))$,

$$i(\omega_1) = i(\omega'_1) = i(\omega'_2) = i(\omega_2) \quad \text{and} \quad \nu(\omega_1) = \nu(\omega'_1) = \nu(\omega'_2) = \nu(\omega_2).$$

Now the family

$$\left\{ \omega_1, \frac{1}{x} \left(\omega_2 - \frac{a_2(y) + \alpha_1 a_1(y)}{a_1(y) + \alpha_2 a_2(y)} \omega_1 \right) \right\}$$

is a good Saito basis for S . Finally, since $i\left(\frac{1}{x} \left(\omega_2 - \frac{a_2(y) + \alpha_1 a_1(y)}{a_1(y) + \alpha_2 a_2(y)} \omega_1 \right)\right) \geq i(\omega_1)$, one has $i(S) = i(\omega_1)$. By the description of $p_1(S \cup l)$ given in [5] - using the notations of the article, it refers to the case $\delta_1 = 1$ and $\delta_2 = 1$ - we get

$$i(S) = i(\omega_1) = \frac{\nu(S) + 1}{2} - p_1(S) = \left\lfloor \frac{\beta_0}{2} \right\rfloor + 1 - p_1(S).$$

Thus, the proof of the formula is a consequence of Theorem 11 noticing that by the Generic Basis Theorem we have $\nu_1 = \left\lfloor \frac{\beta_0}{2} \right\rfloor$ and $\nu_2 = \beta_0 - \left\lfloor \frac{\beta_0}{2} \right\rfloor$. \square

Example (5). In [8], Peraire computed the minimum Tjurina number for the topological class whose characteristic exponents are $(9, 12, 17)$. After five blowing-ups, we obtain a curve with multiplicity 2. The corresponding characteristics exponents of the sequence of blown-up curves are $(3, 14)$, $(3, 11)$, $(3, 8)$, $(3, 5)$, $(2, 3)$. Applying inductively the formula (4.1), one accumulates contribution to the difference $\mu - \tau_{\min}$. Actually, it can be seen that the respective contributions are 15, 1, 1, 1, 0, 0. Thus $\tau_{\min} = \mu - 18 = 98 - 18 = 80$ which coincides with the computation of Peraire.

The last theorem allow us obtain a formula to compute the minimum Tjurina number in a topological class using the multiplicity sequence.

Corollary 13. *Let L a topological class of a singular plane branch determined by the multiplicity sequence $\nu_{(1)}, \nu_{(2)}, \dots, \nu_{(N)}, \nu_{(N+1)} = 1, \dots$. The minimal Tjurina number achieved in L is*

$$\tau_{\min} = \sum_{i=1}^N \left(\nu_{(i)}^2 + \left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor \left(\left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor - \nu_{(i)} + 1 \right) - 1 + p_1(S_{(i)}) \right)$$

where $S_{(i)}$ denote the curve with multiplicity $\nu_{(i)}$ in the canonical resolution process for a generic curve in L .

Proof. Applying inductively the formula presented in the last theorem and using that $i(S_{(i)}) = \left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor + 1 - p_1(S_{(i)})$ yields

$$\begin{aligned} \tau_{\min} &= \mu - \sum_{i=1}^N \left(\left(\left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor - 1 \right) \left(\nu_{(i)} - \left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor - 1 \right) - (i(S_{(i)}) - 1) \right) \\ (4.2) \quad &= \mu + \sum_{i=1}^N \left(\left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor \left(\left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor - \nu_{(i)} - 1 \right) + \nu_{(i)} - 1 + p_1(S_{(i)}) \right). \end{aligned}$$

As $\mu = \sum_{i=1}^N \nu_{(i)} (\nu_{(i)} - 1)$, we get the proof. \square

In [4], Dimca and Greuel present an interesting question about the Tjurina number for curves in a given topological class L . More specifically, they ask if $4\tau(S) > 3\mu(S)$ for any curve in L .

As the Tjurina number is semicontinuous in L and we have obtained a formula for the τ_{\min} , we are able to give a lower bound for the Tjurina number in terms of the Milnor number and it answered positively the previous question for the irreducible case.

Corollary 14. *Let S be a singular irreducible plane curve. Then*

$$\tau(S) \geq \frac{3}{4}\mu(S) + \frac{\sqrt{1 + 4\mu(S)} - 1}{8}.$$

In particular, $4\tau(S) > 3\mu(S)$.

Proof. We denote $\mu = \mu(S)$. It is sufficient to show the inequality for the τ_{\min} . By (4.2), the relation below holds

$$4\tau_{\min} - 3\mu = \mu + 4 \sum_{i=1}^N \left(\left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor \left(\left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor - \nu_{(i)} - 1 \right) + \nu_{(i)} - 1 + p_1(\nu_{(i)}) \right).$$

Now, using that $\mu = \sum_{i=1}^N \nu_{(i)} (\nu_{(i)} - 1)$ and $4 \left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor \left(\left\lfloor \frac{\nu_{(i)}}{2} \right\rfloor - \nu_{(i)} - 1 \right) = -\nu_{(i)}^2 - 2\nu_{(i)} + \delta_i$ with $\delta_i = 0$ if $\nu_{(i)}$ is even and $\delta_i = 3$ if $\nu_{(i)}$ is odd, we obtain

$$4\tau_{\min} - 3\mu = \sum_{i=1}^N (\nu_{(i)} + \delta_i + 4(p_1(S_{(i)}) - 1)).$$

Now, by Theorem 12 we have that:

- if $\nu_{(i)}$ is even, then $p_1(S) \geq 1$ and $\nu_{(i)} + 0 + 4(p_1(\nu_{(i)}) - 1) \geq \nu_{(i)}$,
- if $\nu_{(i)}$ is odd, then $p_1(S) \geq 0$ and $\nu_{(i)} + 3 + 4(p_1(\nu_{(i)}) - 1) \geq \nu_{(i)} - 1$.

So, the following inequality follows

$$(4.3) \quad 4\tau_{\min} - 3\mu \geq \sum_{i=1}^N (\nu_{(i)} - 1).$$

As $\mu = \sum_{i=1}^N (\nu_{(i)} - 1)^2 + \sum_{i=1}^N (\nu_{(i)} - 1)$ we get $4\tau_{\min} - 3\mu \geq \mu - \sum_{i=1}^N (\nu_{(i)} - 1)^2$. Using (4.3), that is, $-(4\tau_{\min} - 3\mu)^2 \leq -\left(\sum_{i=1}^N (\nu_{(i)} - 1)\right)^2 \leq -\sum_{i=1}^N (\nu_{(i)} - 1)^2$, we obtain $4\tau_{\min} - 3\mu \geq \mu - (4\tau_{\min} - 3\mu)^2$ and consequently

$$\tau(S) \geq \tau_{\min} \geq \frac{3}{4}\mu + \left(\frac{-1 + \sqrt{1 + 4\mu}}{8}\right).$$

□

Example (6). Let us consider the topological class L determined by the characteristic exponents $(141, 142)$. The Milnor number for any curve in L is $\mu = (141 - 1)(142 - 1) = 19740$. Using the lower bound presented in the last result we obtain $\tau_{\min} \geq 14840$. For this topological class it follows by the Delorme result (*cf.* [3]) that $\tau_{\min} = 14910$.

While we submit the first version of this paper to Arxiv, we discover that, at the same time, a positive answer for the Dimca-Greuel question was obtained by Alberich-Carramiñana *et al.* and published in Arxiv [1] a few days before. Although the methods are a bit different, the key ingredient is still the formula for the generic dimension of the moduli space obtained in [5].

Acknowledgment. The authors are grateful to the anonymous referee for the suggestions that improved this work.

REFERENCES

- [1] M. Alberich-Carraminana, P. Almiron, G. Blanco, and A. Melle-Hernandez. The minimal tjurina number of irreducible germs of plane curve singularities. *arXiv:1904.02652*, 2019.
- [2] C. Camacho, A. Lins Neto, and P. Sad. Topological invariants and equidesingularization for holomorphic vector fields. *J. Differential Geom.*, 20(1):143–174, 1984.
- [3] C. Delorme. Sur les modules des singularités des courbes planes. *Bulletin de la Société Mathématique de France*, 106:417–446, 1978.
- [4] A. Dimca and G.-M. Greuel. On 1-forms on isolated complete intersection curve singularities. *J. Singul.*, 18:114–118, 2018.
- [5] Y. Genzmer. Dimension of the moduli space of a curve in the complex plane. *arXiv:1610.05998v3*, 2017.
- [6] O. A. Laudal, B. Martin, and G. Pfister. Moduli of plane curve singularities with \mathbb{C}^* -action. *Singularities, Banach Cent. Publ.* 20, 255–278., 1988.
- [7] R.I. Michler. On the number of generators of the torsion module of differentials. *Proc. Am. Math. Soc.*, 129(3):639–646, 2001.
- [8] R. Peraire. Moduli of plane curve singularities with a single characteristic exponent. *Proc. Amer. Math. Soc.*, 126(1):25–34, 1998.
- [9] K. Saito. Theory of logarithmic differential forms and logarithmic vector fields. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 27(2):265–291, 1980.
- [10] O. Zariski. Characterization of plane algebroid curves whose module of differentials has maxim torsion. *Proc. Natl. Acad. Sci. USA* 56, 781–786; Erratum. *Ibid.* 1927., 1966.
- [11] O. Zariski. *Le problème des modules pour les branches planes*. University Lecture Series. AMS, 2006.

Genzmer, Y.

yohann.genzmer@math.univ-toulouse.fr

Hernandes, M. E.

mehernandes@uem.br