

Scientists' warning of threats to mountains

Dirk Schmeller, Davnah Urbach, Kieran Bates, Jordi Catalan, Dan Cogălniceanu, Matthew Fisher, Jan Friesen, Leopold Füreder, Veronika Gaube, Marilen Haver, et al.

► To cite this version:

Dirk Schmeller, Davnah Urbach, Kieran Bates, Jordi Catalan, Dan Cogălniceanu, et al.. Scientists' warning of threats to mountains. Science of the Total Environment, 2022, 853, pp.158611. 10.1016/j.scitotenv.2022.158611. hal-03795426

HAL Id: hal-03795426 https://hal.science/hal-03795426v1

Submitted on 7 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Scientists' Warning of Threats to Mountains

3 AUTHORS:

5 6 7	C.Fish Le Ro	Schmeller ^a , DavnahUrbach ^b , KieranBates ^{cde} , JordiCatalan ^{fg} , DanCogălniceanu ^h , Matthew er ^d , JanFriesen ⁱ , LeopoldFüreder ^j , VeronikaGaube ^k , MarilenHaver ^a , DeanJacobsen ¹ , Gael ux ^a , Yu PinLin ^m , Adeline Loyau ^a , Oliver Machate ⁱ , Andreas Mayer ^k , Ignacio Palomo ⁿ ,
8	<u>Christ</u>	oph Plutzar ^k , Hugo Sentenac ^a , Ruben Sommaruga ^j , Rocco Tiberti ^o , William J.Ripple ^p
9 10 11	a b	LEFE, Université de Toulouse, INPT, UPS, Toulouse, France
12 13	C	Global Mountain Biodiversity Assessment, Institute of Plant Sciences, University of Bern, Bern, Switzerland
14 15 16	с	Department of Zoology, University of Oxford, 11a Mansfield Road, Oxford OX1 3SZ, UK
17	d	
18 19	е	MRC Centre for Global Infectious Disease Analysis, Department of Infectious Disease Epidemiology, School of Public Health, Imperial College London, London W2 1PG, UK
20 21 22	f	Institute of Zoology, Zoological Society of London, Regent's Park, London NW1 4RY, UK
23 24 25	g	CREAF Campus UAB, Edifici C, Cerdanyola Del Valles, Spain
26 27	h	CSIC, Campus UAB, Cerdanyola Del Valles, Spain
28 29	i	Ovidius University Constanța, Faculty of Natural Sciences and Agricultural Sciences, Al. Universității 1, 900470 Constanța, Romania
30 31 32	j	Helmholtz Centre for Environmental Research – UFZ, Leipzig, Germany
33 34	ŀr	Department of Ecology, University of Innsbruck, Technikerstr. 25, 6020 Innsbruck, Austria
35 36 37	k	University of Natural Resources and Life Sciences, Vienna, Department of Economics and Social Sciences, Institute of Social Ecology (SEC), Schottenfeldgasse 29, Austria
38	1	and Social Sciences, institute of Social Ecology (SEC), Schouenielugasse 29, Austria
39 40	m	Freshwater Biological Section, Dept. Biology, University of Copenhagen, Denmark
41 42		Department of Bioenvironmental Systems Engineering, National Taiwan University, Taiwan
43	n	
44 45	0	Univ. Grenoble-Alpes, IRD, CNRS, Grenoble INP*, IGE, 38000 Grenoble, France

46 47

- 48
- Department of Earth and Environmental Sciences DSTA, University of Pavia, Via Ferrata 9, 27100 Pavia, Italy
- р Department of Forest Ecosystems and Society, Oregon State University, Corvallis, OR, 49 USA 50
- 51
- 52
- 53

54 Abstract

55 Mountains are an essential component of the global life-support system. They are characterized 56 by a rugged, heterogenous landscape with rapidly changing environmental conditions providing 57 myriad ecological niches over relatively small spatial scales. Although montane species are well 58 adapted to life at extremes, they are highly vulnerable to human derived ecosystem threats. 59 Here we build on the manifesto 'World Scientists' Warning to Humanity', issued by the Alliance 60 of World Scientists, to outline the major threats to mountain ecosystems. We highlight climate 61 change as the greatest threat to mountain ecosystems, which are more impacted than their 62 lowland counterparts. We further discuss the cascade of "knock-on" effects of climate change 63 such as increased UV radiation, altered hydrological cycles, and altered pollution profiles; 64 highlighting the biological and socio-economic consequences. Finally, we present how 65 intensified use of mountains leads to overexploitation and abstraction of water, driving changes 66 in carbon stock, reducing biodiversity, and impacting ecosystem functioning. These 67 perturbations can provide opportunities for invasive species, parasites and pathogens to 68 colonize these fragile habitats, driving further changes and losses of micro- and macro-69 biodiversity, as well further impacting ecosystem services. Ultimately, imbalances in the normal 70 functioning of mountain ecosystems will lead to changes in vital biological, biochemical, and 71 chemical processes, critically reducing ecosystem health with widespread repercussions for 72 animal and human wellbeing. Developing tools in species/habitat conservation and future 73 restoration is therefore essential if we are to effectively mitigate against the declining health of 74 mountains.

75

76 **Keywords:** Pollution, climate change, environmental health, sustainable development 77 goals, policy

79 Main text

80 Following the onset of the industrial revolution and the use of fossil energy, humans can 81 indisputably be seen as major geological agents (Gałuszka et al., 2014) and as global pathogen 82 vectors (Small et al., 2019). Due to the increase in human activities over the last 300 years, 83 human impact has become at least as strong a force as natural processes (Crutzen, 2006), 84 marking the geological era of the Anthropocene (Gałuszka et al., 2014). Already in 1972, 85 scientists lined out that there are limits to human population growth (Meadows et al., 1972). 86 Twenty years later, scientists warned about ozone depletion in the stratosphere, availability of 87 drinking water, climate change, exponential human population growth, and degradation of the 88 environment and biodiversity (Scientists, 1992). This early warning remained largely unheard 89 and unrecognized, and the progress of humanity towards a sustainable lifestyle has been largely 90 insufficient. This insufficiency ushered in a second warning to humanity 25 years later, arguing 91 that our life-support system is on the brink of collapse (Ripple et al., 2017). Scientists are now 92 largely aware of humanity approaching important tipping points (Lenton, 2011) and planetary 93 boundaries (Steffen et al., 2015). A policy-led reaction was the development of 17 Sustainable 94 Development (https://sdgs.un.org/goals) 20 Goals and Aichi targets 95 (https://www.cbd.int/sp/targets/), the creation of the Intergovernmental Panel for Climate Change 96 (IPCC) and the Intergovernmental Science-Policy Platform for Biodiversity and Ecosystem 97 Services (IPBES, Schmeller et al., 2017). However, all recent global reviews of the state of the planet conducted by the United Nations, IPCC and IPBES univocally lay out the dire state of 98 99 Earth (Bridgewater et al., 2019). The direct drivers of the observed changes, as outlined in the 100 global assessment on biodiversity and ecosystem services by IPBES (Intergovernmental 101 Science-Policy Platform on Biodiversity and Ecosystem Services 2019), include climate change, 102 pollution, and increasing demands for energy and materials due to a growing human population. 103 As a response, the UN Convention on Biological Diversity (CBD), developed a global biodiversity 104 framework with 21 targets and 10 milestones to be achieved by 2030 to 'living in harmony with 105 nature' by 2050 (CBD/WG2020/3/3). Goals include reducing threats to biodiversity (8 targets), 106 meeting people's needs through sustainable use and benefit-sharing (5 targets), developing 107 tools and solutions for implementation and mainstreaming (8 targets; CBD/WG2020/3/3). In line 108 with these goals and targets, IPBES, during its 8th plenary, decided to advance with an 109 assessment on the transformative change needed to indicate ways out of the current global 110 crisis (Schmeller and Bridgewater, 2021).

111 Mountains cover a large part of the Earth's terrestrial surface and host a larger proportion 112 of biodiversity than expected by area (Körner et al., 2011). They hold an estimated one-third of

113 terrestrial species diversity (Körner, 2004), and represent 18 of the 36 global biodiversity 114 hotspots (Chape et al., 2005). Nevertheless, even in remote areas human impact is strong, as 115 mountains are part of the global socio-ecological systems which have been shaped by 116 geological forces and by human activities (Turner et al., 1990). In mountains, these impacts and 117 the resulting threats remain largely understudied (Schmeller et al., 2018). Generally, people are 118 unaware of the threats to mountain ecosystems and the services mountains provide to humanity. 119 Mountain ecosystems sequester CO_2 , clean waters and the air, regulate climate, provide 120 biomedical resources, and regulate floods (Martín-López et al., 2019). Mountains also provide 121 for the livelihoods of more than half of humanity (Grêt-Regamey et al., 2012; Grêt-Regamey and 122 Weibel, 2020). All these goods and services are provided by mountain ecosystems through 123 complex processes that are maintained by communities of species interacting with each other 124 and with the abiotic environment (Bestion et al., 2021). These mountain communities are 125 comprised of prokaryotic and eukaryotic microbes, fungi, plankton species, woody and non-126 woody plants, as well as invertebrates and vertebrates. By destructing, rebuilding, changing, and 127 shaping the environment, these species produce organic matter and oxygen as well as bind 128 CO_2 .

129 The mountain environment is characterized by extreme temperature regimes, severe 130 weather events and short growing seasons at high altitudes to which species have adapted 131 (Körner, 2019; Payne et al., 2020). In temperate mountains, comparably few species have been 132 able to widely colonize the diverse habitats with montane conditions during the short time 133 window given by the recent demise of glaciers at the end of the last glacial period about 11,000 134 years ago (McCain and Colwell, 2011; Schabetsberger et al., 2013; Valbuena-Ureña et al., 135 2018). Mountain ecosystems have evolved in partial isolation, separated by a variety of 136 biogeographic barriers. The gradients and dynamics in climate, hydrology, and water chemistry 137 contributed to the formation of a high diversity of microhabitats, harboring numerous species in 138 comparably small areas (McCain and Colwell, 2011), which also explains the high levels of 139 endemism detected in mountains (Rahbek et al., 2019a; b; Swanson et al., 1988). With 140 increasing elevation, montane climates become more extreme, providing habitat for fewer 141 species. System redundancies (i.e. different species with similar functions), available in 142 ecosystems at low altitudes, are increasingly scarce in mountain ecosystems with increasing 143 elevation. Such redundancies usually provide stability to ecosystems (Fonseca and Ganade, 144 2001). The absence of these redundancies renders mountain ecosystem particularly vulnerable 145 to the impacts of global change (Moser et al., 2019). However, multiple threats to mountains are 146 arising from climate change alone. Moreover, interactions with socio-cultural, economic and

political developments, such as the exploitation of mountains, e.g. for timber, food production,
including fish and livestock, tourism, and hydro-electricity, exacerbate these threats, calling for
urgent consideration by policymakers (Figure 1).

150 Here, we highlight the diversity of global threats impacting mountain ecosystems. We 151 focus on the direct drivers climate change, pollution, and land use following earlier science-152 policy reports (IPBES 2019) to compel stakeholders and decision makers of the urgency to act 153 on all of these different threats. We detail how different drivers interact, creating pressures that 154 degrade and destroy valuable mountain ecosystems and their biodiversity. We further outline, 155 how this impacts the services provided by mountains and creates emerging risks for humans. 156 We treat threats to mountains largely equally, as a ranking of threats in regard to their severity 157 appears elusive based on current knowledge. Finally, we provide recommendations for 158 mitigation actions to be taken to preserve mountains, their biodiversity and the ecosystem 159 services they provide to humanity, as well as describing ways of averting detrimental 160 trajectories.

161 Climate Change in Mountains

162 Mountains are defined by rugged terrain and unique climate regimes distinguishing them 163 from lowlands (Körner et al., 2017). The climatic complexity created by mountain topography 164 also influences insolation and air circulation (Dobrowski et al., 2009). Elevation gradients in 165 particular, have a strong impact on many abiotic variables in mountains, and their geographic 166 location is the main control on moisture gradients or seasonality in climate (Körner, 2007). On a 167 regional scale, mountain climate is influenced by large-scale synoptic patterns, proximity to 168 oceans, and the range's longitudinal or latitudinal orientation (Del Barrio et al., 1990). In synergy 169 with climate change impacts, we see important changes in precipitations, temperatures, and 170 frequency of extreme events, such as droughts and floods. Therefore, climate change might be 171 considered the most basic and far-reaching threat to mountains, impacting mountain biodiversity 172 and ecosystems way more intensively as compared to lowland regions (Rangwala and Miller, 173 2012; Scarano, 2019).

Precipitation dynamics are not the same in all mountains, and for example, mountains in tropical regions show precipitation maxima at lower and mid-elevations. Temperate mountains typically have an orographic effect and show increased precipitation towards the top, largely driven by different seasonal dynamics in precipitation intensity (Roe, 2005). These precipitation regimes are also regulated by geology, soils, vegetation, and human land use, leading to a large variety of hydrological behaviors and stream flow regimes in global mountain catchments (Dierauer et al., 2018; Zuecco et al., 2018). Climate change-driven precipitation regimes and

181 their increasing variability, especially in mountains, determine long-term change on soil moisture 182 conditions and are important controls on water levels and on the hydroperiods of shallow lakes. 183 ponds and wetlands (Catalan and Bartumeus, 2006; Stephan et al., 2021). As such they are 184 particularly important variables for mountain habitats. Climate change will increase the 185 unpredictability of precipitation patterns (Myhre et al., 2019). Generally, water availability and its 186 predictability is expected to decrease in the future due to lower water storage capacities in areas 187 with glacier cover and higher outflows of excess water during periods of extreme precipitation 188 and melting events (Rajczak and Schär, 2017). These changes will jeopardize the role of 189 mountains as global water towers and the drinking water supply for billions of people (Viviroli et 190 al., 2007) as well as for plants and animals.

191 Trends in warming of mountain surface air temperatures become more and more 192 apparent (López-Moreno et al., 2008; Niedrist et al., 2018; Niedrist and Füreder, 2021), and 193 high-latitude mountains are projected to warm much faster than temperate and tropical 194 mountains (Negi et al., 2021; Nogues-Bravo et al., 2007). Further, warming is accentuated at 195 higher altitudes (Pepin et al., 2015). The intensity of warming depends on the mountain climatic 196 zone, elevation and season (Pepin and Lundquist, 2008). In the Alps, the mean annual 197 temperature has already increased by 1.5 - 2°C since 1970, and future projections predict an additional rise of 0.25 - 0.36 °C per decade within the next century (Einhorn et al., 2015). Similar 198 199 increases have been projected in other mountain ranges (Urrutia and Vuille, 2009; Valdivia et 200 al., 2013). Increasing temperatures have decreased the annual snow deposition volume with 201 most impacts observed at mid- and lower elevations (Laternser and Schneebeli, 2003), has 202 caused important shifts to earlier snow melt (Kapnick and Hall, 2012), earlier lake-ice melt 203 (Franssen and Scherrer, 2008), and globally accelerated the deglaciation process (Zemp et al., 204 2015). Yearly, world-wide cumulative glacier mass balance data have been showing significant 205 decreasing trends in glacier thickness (Ripple et al., 2021). The shorter period of snow cover will 206 therefore also shorten the time during which the Albedo effect is active and during which 85% to 207 90% of sunlight is reflected. Hence, further acceleration of the temperature rise in alpine 208 mountain regions needs to be expected. Droughts and extreme precipitation events will globally 209 increase in mountains (Gobiet et al., 2014; Urrutia and Vuille, 2009; Valdivia et al., 2013). For 210 the European Alps, climate projections predict more summer droughts and more extreme rainfall 211 events (Rajczak and Schär, 2017). Subtropical mountains tend to experience more frequent 212 summer droughts (McCullough et al., 2016), while for example the Hindu Kush and the Himalaya 213 have been experiencing increased amounts of extreme rainfall events (Hartmann and Andresky, 214 2013; Wester et al., 2019). Droughts in mountains also lead to an increase of the probability of

wildfires in mountain grasslands and forest as the probability of ignition of dry plant material by
lightning increases (Stephens et al., 2018). Such wildfires are difficult to extinguish due to the
difficult terrain and result in the loss of very large areas (i.e. thousands of hectare) of mountain
vegetation with large and long-lasting effects on mountain ecosystems due to low pre-fire
production of seeds due to drought and the generally low recovery speed with increasing altitude
(Werner et al., 2022).

221 Ultraviolet radiation (UVR) both affects and is affected by climate change. These 222 modifications of UV exposure are affecting how people and ecosystems respond to UV, with 223 more pronounced effects in the future (Barnes et al. 2019). UVR reaching mountain ecosystems 224 increases with elevation (Sommaruga et al., 1999). In many high mountain freshwaters, the 225 limited catchment sources of DOC, which results in crystal clear water, facilitates the penetration 226 of radiation far into the water column (Catalan and Donato Rondon, 2016; Laurion et al., 2000; 227 Rose et al., 2009). High levels of UVR in the UV-B spectrum (wavelength range 280-320 nm) 228 have been studied with regard to their effects on aquatic organisms at higher elevations. Colored 229 dissolved organic carbon (DOC) and phytoplankton may attenuate the penetration of UV 230 underwater, shielding freshwater species from negative effects of this radiation (Sommaruga et 231 al., 1999). Variability in UVR reaching mountain freshwaters depends on factors such as cloud 232 cover and air pollution (Brooks et al., 2005; de Oliveira et al., 2021; Diamond et al., 2005; 233 Obertegger and Flaim, 2021; Sommaruga, 2001). Long-term changes in UV-B in turn are due to 234 changes in atmospheric ozone levels, and despite the recovery of atmospheric ozone layers 235 since the Montreal protocol was implemented (Barnes et al., 2019), new reports of ongoing 236 lower stratospheric ozone depletion (Ball et al., 2018) and new ozone depleting agents in the 237 atmosphere have been issued (Fang et al., 2019). We may also see potential opposite effects of 238 global warming in mountains, as for aquatic or semi-aquatic species UVR stress may increase 239 due to earlier snow and ice-cover melt, but may decrease due to higher DOC import from 240 catchment reforestation (Sommaruga et al., 1999). The impact of UVR and stress avoidance 241 behavior remains an understudied field, but may drive further changes in mountain biodiversity 242 and perturb mountain ecosystems (Häder et al., 2011).

Generally, decreasing amounts of annual snow and retreating glaciers have been and will continue to profoundly reshape mountain freshwater habitats and also the terrestrial communities depending on them, threatening mountain species and communities (Jacobsen and Dangles, 2017; Sommaruga, 2015), but also providing new habitats for colonization (Ficetola et al., 2021). Impacts on lowland human populations include difficult to predict water supply (Huss et al., 2017; Viviroli et al., 2007) and destruction of infrastructure through more frequent extreme

floods and/or landslides. Climate change will drive the disappearance of many intermediate and ephemeral habitats in mountains due to droughts, with potential severe consequences for mountain biodiversity and biological processes e.g. carbon storage. Climate change will also impact the water regime of many peatlands. Due to the importance of mountain peat- and wetlands as global carbon sinks, the loss of these habitats may further accentuate climate change (De Jong et al., 2010) and UVR impacts (Barnes et al. 2019).

255 **Pollutants in Mountains**

256 Sources of pollution are manifold in mountain ecosystems (Lei and Wania, 2004; Noyes 257 et al., 2009; Shunthirasingham et al., 2010) and we generally lack a global approach to observe 258 environmental pollution and its impact (Brack et al., 2022). Global atmospheric transport of 259 micropollutants (Hussain et al., 2019; Wania and Mackay, 1993; Yang et al., 2010) and local 260 human activities such as mining, logging, agriculture, pastoralism and tourism are the main 261 pollution sources in mountain environments. Increasing pastoralism and livestock units have 262 been shown to put the health of mountain lakes at risk by introducing highly toxic organic 263 pollutants into mountain lakes (Machate et al., 2022). Further, fish stocking introduced heavy 264 metals such as mercury in mountain ecosystems (Hansson et al., 2017b). Other sources of 265 pollution include extreme weather events releasing legacy pollutants from mining, tourists 266 introducing UV blockers as part of personal care products, and atmospheric transport of 267 pollutants releasing a plethora of different molecules (Gross, 2022; Le Roux et al., 2020; Pozo et 268 al., 2007). Mountains are also at risk of acidification when they are located downwind of Nitrogen 269 or Sulphur emission sources and buffering capacity of freshwater is low. In addition, seasonal 270 events such as snowmelt can change pH up to a full unit (Nodvin et al., 1995). Acidic pulses 271 may co-occur with spikes of toxic element concentrations in water (Havas and Rosseland, 272 1995). Further, higher rates of erosion and weathering, as a result of more frequent events of 273 heavy precipitation, may also lead to a higher import of base cations and increase the pH of 274 mountain freshwater and soil (Kopáček et al., 2017). Finally, global change-driven eutrophication 275 and elevated temperatures lead to an increased growth of cyanobacteria, which can produce 276 toxins (e.g. cyanotoxins), with known negative impacts on human health (Catherine et al., 2013; 277 Funari and Testai, 2008; Zanchett and Oliveira-Filho, 2013).

The scavenging of atmospheric organic and inorganic pollutants is pronounced at high altitudes and can take the form of dry and wet deposition of aerosols to the ground surface (Daly and Wania, 2005; Le Roux et al., 2016; Le Roux et al., 2020). Atmospheric pollutants in mountains include trace elements (Camarero et al., 2009) (Yang et al., 2010), organic and

282 synthetic pollutants (i.e. current-use and legacy pesticides), polycyclic aromatic hydrocarbons 283 (PAHs), endocrine disrupting compounds (EDCs; Rockström et al., 2009; Meire et al., 2012), 284 and polychlorinated biphenyls (PCBs) or microplastics (Allen et al., 2019; Brahney et al., 2020; 285 Reid et al., 2019). These compounds are introduced into the atmosphere via evaporation, or 286 binding to particles light enough to be carried by wind. Volatile compounds tend to evaporate in 287 warmer environments at lower altitudes and condensation progressively takes place with 288 decreasing temperature as air masses travel over mountain slopes (Blais et al., 1998). Less 289 volatile compounds, such as most of the currently used agricultural pesticides, are not prone to 290 directly evaporate into the atmosphere, but are still transported into the mountain environment 291 as they bind to soil particles, which can be uplifted and carried over longer distance during more 292 extreme wind events (Silva et al., 2018). As a consequence, organic and inorganic 293 micropollutants can be introduced to mountains via the atmosphere and over long distances 294 (Camarero et al., 2009; Bradford et al., 2010; 2013; Fig. 2).

295 Transfer of inorganic and organic pollutants to mountain freshwaters can also originate 296 from legacy and recent local human activities. Fish stocking (Hansson et al., 2017b), present 297 and historic mining activities (Hansson et al., 2017a) or forestry, agriculture and tourism (Alpers 298 et al., 2016) introduce pollutants or mobilize their catchment sources such as soils and 299 sediments. Evidence for the introduction of potentially harmful trace elements (PHTE, for 300 example the metals As, Hg, Pb, Se, Sb, Zn, Cu) from distant or local sources is found in 301 mountain lake sediments, peatlands and in snow (Bacardit and Camarero, 2010). Furthermore, 302 glacier melt and snowmelt can mobilize legacy atmospheric pollution and catchment sources of 303 pollutants (Bogdal et al., 2009; Meyer and Wania, 2008), which may increase exposure of 304 mountain species to micropollutants.

305 Despite the accumulated knowledge on global pollution, we still know little about the toxic 306 cocktail accumulating in mountains and its impact on biodiversity (but see Catalan, 2015). There 307 are very few studies that analyzed a broad set of pollutants or even conduct non-target 308 monitoring in a mountain context (Machate et al., 2022), rendering our current knowledge on 309 global mountain pollution patterns largely incomplete. We also know little about each 310 (detectable) compound's physico-chemical characteristics, e.g. the water-air constant (K_{wa}), 311 without which it is difficult to make predictions on future pollution patterns in mountains. 312 Generally, the introduction of new pollutants and changes in pollutant mobilization due to climate 313 change may challenge mountain ecosystem health and increase the vulnerability of species and 314 humans to pathogens, increasing health risks (Brack et al., 2022; Schmeller et al., 2020; 2018).

315 Concerns about adverse toxic effects have especially been raised about, but are not limited to,

316 pollutants introduced from local activities (Machate et al., 2022). In the future, these health risks

- 317 might further increase, by shifting e.g. pastures to higher altitudes (Mayer et al., 2022), where
- ecosystems are already under high pressures due to climate change (Herzog and Seidl, 2018)).
- 319 Tourism and expansion of infrastructure create yet additional pressures on mountain
- 320 biodiversity.

321 Vegetation and land use changes in Mountains

322 Recent mountain vegetation is largely shaped by historical and current land use (Körner 323 et al., 1997; Lavorel et al., 2017), with the degree of influence depending mainly on the 324 accessibility of the area (Tasser and Tappeiner, 2002). Vegetation and changes in rates of 325 carbon sequestration due to shifts in land use can occur through a change in land cover, through 326 intensification or extensification of existing land use practices (Niedertscheider et al., 2017) and 327 acidification (Bowman et al., 2012). Most of these changes are driven by pastoral activities, such 328 as livestock grazing, which is the major agricultural activity in most mountains. Pastoral activities 329 in many mountain regions have a long history. However, especially for European countries it is 330 known that livestock units are increasing, partly as a result of the EU subvention policy, partly as 331 strategy to evade climate change impacts and reduced availability of fodder in lowlands (Mayer 332 et al. 2022). Another (illegal) activity related to pastoralism is slash and burn to avoid expansion 333 of forest areas. The clearing intensity and frequency increases with increasing pastoral pressure. 334 In addition, novel non-native crops have been planted to increase local food availability under 335 optimal environmental conditions. However, plantations of species that are well-adapted to the 336 environment of a particular mountain range, such as cardamom in the Hindu Kush region 337 (Eklabya et al., 2000), provide a larger genetic reservoir and thus a greater buffer against 338 environmental pressures such as climate change (Kelty, 2006). Other land use changes include 339 logging of forest stands (Latty et al., 2004), afforestation (Liu et al., 2021), vegetation regrowth in 340 abandoned lands (Aide et al., 2019). All these changes intervene deeply in the existing 341 ecosystem (Hinojosa et al., 2016), altering and threatening underlying processes and associated 342 ecosystem services (Chiang et al., 2014; Faccioni et al., 2019; Tasser and Tappeiner, 2002). An 343 emerging land-use trend is the growing impact of tourism on ecosystems, where damage to 344 vegetation can occur (Rodway-Dyer and Ellis, 2018) and is playing an increasing role in 345 mountain regions (Niu and Cheng, 2019). In the Dongling Mountains (China), tourism led to a 346 lower species richness, heterogeneity and evenness in impacted subalpine meadows (Zhang et 347 al., 2012). Increasing pressures from land use changes will further accentuate the impacts of 348 climate change and pollution on mountain biodiversity and the health of mountain ecosystems.

349 Introduced species in mountains

350 Biological invasions are increasingly exacerbated by human activities and are 351 responsible for significant biodiversity decline as well as high economic losses to society 352 (Diagne et al., 2021). In addition, they are exacerbated by globalization and climate change 353 (Seebens et al., 2021). The harsh environmental conditions (e.g., high UV-B, low temperatures, 354 variable water availability, poor soil) may limit alien plant invasion and expansion, especially in 355 high mountain areas (Watermann et al., 2020). However, remote mountain areas have been 356 reportedly impacted by the introduction of alien species (Pauchard et al., 2009), partly also 357 through tourism (Hemp, 2008).

358 For example, tourism drives fish stocking of naturally fishless lakes. In such naturally 359 fishless mountain lakes fish introductions have an outsized impact and drive profound ecological 360 change as a highly efficient aquatic predator is introduced in a naïve environment (Miró and 361 Ventura, 2020). Stocking montane lakes with fish for subsistence purposes has been occurring 362 since the Neolithic, through fish translocations from nearby lakes and rivers. However, before 363 1950 such introductions had limited geographic extent and their impacts were rather local 364 (Moser et al., 2019). Since 1950, introductions of fish increased dramatically as a consequence 365 of the increasing popularity of recreational angling, both in large and relatively small lakes as 366 well as in adjacent wetlands (Hansson et al., 2017b). In addition, the use of small fish, mainly 367 minnows *Phoxinus* sp. (Fam. Cyprinidae), as live baits for trout fishing is causing a new and 368 detrimental wave of invasion (Miró and Ventura, 2015). These introduced fish dramatically affect 369 native communities of mountain lakes. Initially considered ecologically harmless and 370 economically beneficial, introductions continued even when their serious ecological 371 consequences became clear (Knapp et al., 2001). Supported by institutional stocking campaigns 372 and non-authorized translocations by anglers, fish spread rapidly in mountain deep-ponds and 373 lakes of all sizes, as well as in all the colonizable downstream habitats (Ventura et al., 2017), 374 with a long list of negative impacts: i) decline/elimination of native species (e.g., invertebrates 375 and amphibians; (Knapp et al., 2001; Tiberti et al., 2014); ii) cascading effects in the trophic 376 network (Schindler et al., 2001), affecting the chemical/microbiological quality of waters, and the 377 ecological linkages with surrounding terrestrial habitats (Epanchin et al., 2010); iii) impacts such 378 as predation, competition, transmission of pathogens and hybridization on native fish inhabiting 379 downstream habitats (Adams et al., 2001), and iv) further collateral introductions of fish used as 380 live bait (Miró and Ventura, 2015). Hence, fish stocking in mountain lakes is particularly 381 detrimental to water quality and biodiversity, especially as now nearly all these ecosystems are

affected, including large lakes, small lakes, ponds, connecting streams and their adjacentmountain wetlands (Ventura et al., 2017).

384 For timber production, fast growing tree species of the genera *Pinus* and *Eucalyptus* 385 have also been introduced in many mountain forests. These exotic tree plantations are subject 386 to serious criticism due to their negative impact on water balance, soil fertility, and native 387 biodiversity (Fahey and Jackson, 1997; Hofstede et al., 2002; Lundgren, 1978). A significant loss 388 of soil carbon and a major reduction in taxonomic and functional diversity of soil invertebrates 389 has been observed in pine tree plantations compared to native forests of very similar soil origins and topographies (Cifuentes-Croquevielle et al., 2020). These impacts may further be 390 391 exaggerated by fast rotation speeds, which would not permit to increase floristic and hence also 392 faunistic biodiversity (Hall et al., 2012). These changes therefore have the potential to aggravate 393 both climate change impacts and biodiversity loss (but see (Balthazar et al., 2015). Further, 394 exotic tree species can become highly invasive under the right environmental conditions, which 395 might also be met by future climate change. Invasibility is also driven by seeds from exotic tree 396 plantations leading to colonisation and replacement of surrounding natural vegetation (van 397 Wilgen, 2012). Reversing impacts associated with those self-sown invasive stands has be 398 proven to be very difficult (van Wilgen and Richardson, 2012), also because waste of e.g. pine 399 harvesting is left at sites or even delivered to nearby rivers, delaying the natural regeneration of 400 indigenous vegetation (Balthazar et al., 2015).

401 Water abstraction from Mountains

402 Most valley bottoms have been heavily altered by human activities that impact freshwater 403 systems (Finlayson and D'Cruz, 2005). These activities include land drainage, dredging, flood 404 protection, water abstraction for hydroelectric powerplants, and inter-basin water transfer, 405 building dams to create reservoirs, and digging new canals for navigation. In mountains, which 406 are increasingly used as recreational area, food and water source, but also as hydroelectric 407 powerplants, water abstraction has been increased to excessive levels. Hydrological 408 interventions include (hydroelectric) dams, pipelines and derivation channels, agricultural ponds, 409 irrigation and snowmaking reservoirs, guarries, water removal, and flow regime alterations. 410 Among their very many consequences, these interventions may lead to the gradual drying up of 411 natural aquatic ecosystems due to excessive water extraction and diversion, as well as changes 412 in the water level of (dammed) lakes, in the flow regime of streams, and in hydrological 413 connectivity. These consequences in turn impact on the structure and function of the unique 414 biodiversity that is characteristic of these habitats, and which includes many endemic and 415 threatened species absent from the lowlands (Fait et al., 2020; Mayerhofer et al., 2021; Schabetsberger et al., 2013). Importantly, ongoing modifications in high-mountain freshwater
ecosystems may also directly and profoundly impact on the wellbeing and livelihoods of peoples
(Schmeller et al., 2018).

419 Recent reports indicate that biodiversity in freshwater ecosystems is declining even 420 faster than in oceans and forests and that the extent of human alteration and impairment of 421 aquatic ecosystems is massive (Tickner et al., 2020). Mountain aquatic systems are no 422 exception, particularly in high-mountain areas (Catalan et al., 2017). Human alterations to alpine 423 aquatic ecosystems are of particular concern given that mountain aquatic habitats provide 424 essential ecosystem services such as drinking water and renewable energy to much of 425 humanity, and that they are of high aesthetic, recreational, and conservation value, particularly in 426 their function as biodiversity reservoirs (Fait et al., 2020). Water abstraction in concert with 427 climate driven changes in hydrological regimes will lead to a gradual drying up of aquatic 428 mountain ecosystems, likely causing massive water shortages in cities that depend on drinking 429 water from mountains (Viviroli et al., 2020; United Nations Environment Programme 2022). The 430 desertification of these ecosystems will also be detrimental to their unique mountain biodiversity, 431 leading to an irreversible degradation of these sensitive ecosystems, if no or too little action for 432 their preservation are put in place immediately (Immerzeel et al., 2020).

433 Threats to and from Mountain Micro-Biodiversity

434 The unseen diversity of micro-organisms and their microbiomes, comprising the 435 community of fungi, yeasts, bacteria, viruses and protists, and the impacts of climate change on 436 them have already been subject to a previous warning (Cavicchioli et al., 2019). In short, micro-437 biodiversity plays a central role and is of global importance in climate change biology, 438 particularly in extreme environments such as mountain ecosystems (Schmeller et al., 2018). 439 Climate change impacts, relevant also for humanity, depend heavily on the responses of 440 microorganisms, which are essential for achieving an environmentally sustainable future 441 (Cavicchioli et al., 2019). Despite their importance, we still know little about the microbial 442 communities or microbiomes, especially in mountain ecosystems (Kammerlander et al., 2015; 443 Schmeller et al., 2018).

Despite their small size, microbial communities drive major processes in and on animals,
plants, as well as in ecosystems (Bates et al., 2022; Bernardo-Cravo et al., 2020; Lin et al.,
2021). In a nutrient poor environment such as mountains, microbial communities likely play an
important role in synthesizing vital nutrients, thereby increasing energy uptake and growth of
plants and animals (Bernardo-Cravo et al., 2020; Schmeller et al., 2020; Sentenac et al., 2022).
Similarly, micro-organisms stabilize whole ecosystems by buffering against change through the

maintenance of biodiversity and ecosystem processes. For example, the interactions between
micro-organisms and plankton constitute the basis of aquatic food webs and determine the
functioning of biogeochemical cycles, accounting for more than half of global carbon fixation
(Cavicchioli et al., 2019; Purcell et al., 2022). Any kind of disturbance to microbial communities
can therefore impact on mountain species and ecosystems.

455 Pathogens, and other microorganisms, can be easily introduced to mountains through 456 pastoralism, tourism or wind drift. However, we remain largely oblivious to how the complexity of 457 the abiotic and biotic environment in mountain ecosystems influences beneficial microbe-species 458 interactions (e.g. microbial loop, mycoloop; (Kagami et al., 2014), host-pathogen interactions 459 (Frenken et al., 2017; Haver et al., 2021; Fisher and Garner, 2020) and health risks for the 460 human population (Schmeller et al., 2018). For example, the transport of microbial pathogens is 461 of special concern for human and livestock health, but also for wildlife and keystone species 462 groups such as amphibians. In particular, fungi and bacteria with resistant aerosolised spores 463 are capable of long-distance transport of e.g. dust (Dadam et al., 2019; Sultan et al., 2005). 464 Global dust dispersion is a natural phenomenon, and occurs when topsoil is transported into the 465 troposphere and carried over long distances by wind currents. However, global warming and 466 changes in land use practices (e.g. deforestation and overgrazing) have accelerated 467 desertification in many areas, resulting in increased dust dispersion even to remote places 468 (Moulin and Chiapello, 2006; Tegen et al., 2004), particularly to high elevation sites (Dong et al., 469 2020). Further anthropogenic impacts via air pollution can intensify both the abundance and 470 community composition of aerial microbes (Yan et al., 2016), but also for vector-borne diseases 471 (Caminade et al., 2019): Malaria has been found at higher altitudes in mountains in Ethiopia and 472 Colombia (Siraj et al., 2014), incidences of Malaria and Dengue are increasing in Nepal's 473 mountains (Dhimal et al., 2015a), altitudinal upward shifts of Dengue and Chikungunya (Dhimal 474 et al., 2015b), and also ticks have been reported, the latter e.g. leading to increased occurrence 475 of Lyme borreliosis in the Alps (Garcia-Vozmediano et al., 2020).

476 Recent data also suggest that we currently see an increase in eutrophication of mountain 477 lakes globally with an upsurge of the diversity of Cyanobacteria (Ho et al., 2019). Cyanobacteria 478 produce a range of toxins (e.g. microcystins, cyanotoxins, Catherine et al., 2013), which have an 479 important impact on the quality of water (Du et al., 2019; Ho et al., 2019), therewith increasing 480 risks of intoxications for humans and livestock. Epilithic biofilms are a highly reactive component 481 in freshwater systems that play a crucial role in the provision of many ecosystem services 482 (Catalan and Donato Rondon, 2016). Especially in smaller mountain lakes, streams and other 483 waterbodies, epilithic biofilms must be considered the major player in carbon cycling and

484 ecosystem productivity (Vadeboncoeur et al., 2008). A better understanding of the risks of 485 proliferation of potentially harmful microbial groups, pathogenic fungi, bacteria and protists due 486 to human-driven input of phosphorus, nitrogen through atmospheric fertilization and microbial 487 pollution is necessary to improve our predictive abilities for human and wildlife risks. For 488 mountain ecosystems, in particular, our predictive abilities are poor for forecasting pathogen 489 proliferation, the dynamics of potentially harmful microorganisms, and for identifying threatened 490 species and habitats. In a mountain context that could mean that resources of clean drinking 491 water will diminish at a much faster rate than currently predicted (Schmeller et al., 2018) and the 492 important ecosystem services such as CO₂ sequestration and nitrogen retention will be 493 suboptimal or absent (Saunders and Kalff, 2001).

494 Threats to Mountain Macro-Biodiversity

495 Mountain areas host many species that live in a delicate balance or at the edge of their 496 distribution and are therefore very susceptible to environmental changes and local extinction. 497 Top predators, such as large carnivores, but also large herbivores play important roles in 498 maintaining mammal, avian, invertebrate, and herpetofauna abundance and richness. Many of 499 these species are threatened with extinction (Ripple et al., 2014) and nearing global collapse 500 (Ripple et al., 2015). Threats to macro-biodiversity in mountains come from chemical pollution, 501 nutrient influx through atmospheric processes and local sources such as livestock (Machate et 502 al., 2022), introduction of non-native taxa, but most importantly from overexploitation and habitat 503 loss (Maxwell et al., 2016). These threats drive the decline of already threatened species 504 (Maxwell et al., 2016) and will change the communities of species, which do not all have the 505 same possibilities of dispersal, recovery and reproduction to avoid disturbances (Kerr and 506 Dequise, 2004; Pimm, 2008). For example, among aquatic organisms, the possibility of dispersal 507 and life traits such as the mode of reproduction are very different among taxonomic groups. 508 Some zooplankton species can reproduce sexually or parthenogenetically, and can produce 509 resting eggs, which can survive for a long time in sediment egg banks (Brendonck and De 510 Meester, 2003; Nielsen et al., 2012). Benthic invertebrates increase their dispersal capacity by 511 producing winged adults. Both strategies may allow a speedy recovery through hatching from 512 resting stages or recolonization by flying. Complete recovery after local extinction, however, is 513 unlikely, and restoration and recovery processes in mountains take a long time (Tiberti et al., 514 2019). For example, recovery from fish impacts took 11-20 years to obtain a similar food web 515 structure. In the same studies, it was evident that recolonization efficiency of species with a 516 parthenogenetic reproduction mode was higher than for sexual reproduction (Knapp and 517 Sarnelle, 2008; Knapp et al., 2001). In any case, for many species dispersal and recolonization

518 can be limited, when the remaining seed pool is not abundant or found too distant or beyond 519 barriers in the land- or waterscape in mountains, due to their relief. Hence, connectivity between 520 different populations of the same mountain species is reduced, hampering recolonization after 521 disturbance (Heino, 2013). Further, upward shifts of distribution areas are not the same for all 522 mountain species, disrupting long-established communities and their interactions. For example, 523 we understand only inadequately, if the observed upward shift of some mountain plant species 524 due to climate change is met by associated microbes and invertebrates (Grabherr et al., 1994; 525 Steinbauer et al., 2018). Other factors leading to unequal dispersal of formerly associated 526 species might be driven by reduced oxygen availability with increasing elevation (Jacobsen, 527 2020), differences in temperature and drought tolerance (Forero-Medina et al., 2011; 528 Schai-Braun et al., 2021), different adaptation abilities through e.g. seasonality or phenology 529 (Parmesan and Yohe, 2003), or different abilities to change depth distribution. Dysfunctional 530 ecosystems, with lower resilience to further impact s, are the likely outcome (Körner, 2019; Pecl 531 et al., 2017). Due to the non-linear loss of biodiversity (Trisos et al., 2020), the expected 532 extinction of endemic plant and animal species after tipping points have been met (Dullinger et 533 al., 2012), may further increase the dysfunctioning of mountain ecosystems. When this will 534 happen and what will be the outcome will be difficult to predict due to the multitude of factors 535 impacting different species in a community.

536 Threats to mountain ecosystem services

537 Negative impacts on mountain biodiversity threatens ecosystem integrity and functioning, 538 and hence also the multiple ecosystem services provided to local communities, populations 539 downstream and local stakeholders, including tourists (Grêt-Regamey et al., 2012; Martín-López 540 et al., 2019; Schirpke et al., 2019). The capacity of mountain ecosystems to provide ecosystem 541 services is deteriorating due to biodiversity loss driven by global change (Palomo, 2017). There 542 is an alarming set of negative consequences from those changes, in stark contrast to the few 543 positive effects that have been reported (Hobbs et al., 2009). These changes will jeopardize 544 water use of at least 1.9 billion people (Immerzeel et al., 2020). Moreover, as a result of glacier 545 decline, water availability will be severely reduced in the dry season, affecting millions of farmers 546 globally (Biemans et al., 2019). Mountains will therefore not remain the reliable and highly 547 important source of water they have been for thousands of years. Even in humid mountain 548 regions, such as the European Alps, droughts have become a problem (Stephan et al., 2021) 549 due to the increasing irregularity of water discharging rates and increasing flood events (Ragettli 550 et al., 2021). The irregularity of water discharging rates in combination with land use and land 551 cover changes can also have synergistic effects on ecosystem functioning, rendering mountains

552 more vulnerable to climate change impacts (Chiang et al., 2014). For the central mountain range 553 of Taiwan, it was shown that these combined effects lead to a relocation or loss of ecosystem 554 services and therefore need to be considered in conservation planning (Lin et al., 2019; Lin et 555 al., 2017).

556 Net primary production (NPP), the amount of biomass or carbon produced by primary 557 producers per unit area and time, is the basis of all ecosystem services and is being altered due 558 to climate change (Haberl et al., 2007; Kastner et al., 2022; Melillo et al., 1993). Overall, there is 559 evidence that increasing temperatures and CO₂ concentrations have increased the NPP of 560 forests when water was not a limiting factor (Boisvenue and Running, 2006). In the Alps, 561 changes in NPP of grasslands on which cattle depend show contrasting regional trends (Jäger 562 et al., 2020). A study combining experimentation and meta-analysis reported stabilization of NPP 563 of grasslands under climate change due to changes in species and increasing allocation towards 564 belowground biomass to resist drought (Liu et al., 2018). Despite a limited body of evidence, 565 local models predict that NPP will increase under climate change in the forests of the mountain 566 region of Changbai in China (Gao et al., 2020).

567 Further, as glaciers retreat and permafrost thaw, the decreased land-surface stability 568 results in increased hazards in the form of landslides and rock fall, increasing risks for wildlife, 569 tourists and livestock (Huss et al., 2017; Temme, 2015). Glacial lake outburst floods may also 570 intensify due to glacier retreat and glacial lake formation, with potentially devastating 571 consequences for populations downstream (Harrison et al., 2018; Milner et al., 2017; Vuille et 572 al., 2018). Cultural ecosystem services are also impacted by climate change. For example, 573 glaciers are considered sacred or have a strong symbolic meaning for several mountain 574 communities, and thus spirituality is being affected (Allison, 2015) and has been documented in 575 various countries in Africa, Asia, and the Americas (Allison, 2015; Mölg et al., 2008; Shijin and 576 Dahe, 2015). Overall, the documented threats to mountain ecosystem services are a major 577 concern worldwide, as they could lead to increased poverty, lower food production, higher health 578 risks and a general decrease of human wellbeing, which may often affect not only mountains but 579 also the populations living downstream.

580 **Conclusions**

581 Mountain ecosystems are complex, dynamic, exceptionally fragile and are highly 582 sensitive to global change. They are therefore considered sentinels of change (Schmeller et al., 583 2018). We are only beginning to understand the functional ecology of mountain ecosystems, but 584 international research already suggests that changing species communities will be detrimental to 585 the environment, to biodiversity and therefore to a critical part of Earth's life-support system. 586 Climate change might be considered the most impactful driver of change in mountain 587 ecosystem, but all the outlined threats to mountains act in synergy. Climate change is modifying 588 and will continue to modify the occurrence of extreme events, the amount of precipitations (rain 589 and snow), as well as freeze and thaw cycles, with impacts on the onset of snow melt (and thus 590 length of growing season) and water temperatures, aggravating impacts from inappropriate land 591 use practices. Global change with all the different pressures outlined above causes imbalances 592 in the functioning of mountain ecosystems, which lead to changes in vital biological, biochemical, 593 and chemical processes, critically reducing ecosystem health with repercussions for animal and 594 human health and wellbeing (Acevedo-Whitehouse and Duffus, 2009; Bradshaw et al., 2021; 595 Lerner and Berg, 2017).

596 Humanity has a wide range of options in its hand to mitigate human-driven impacts on 597 mountains and to change the current trajectory as humanity is at the nexus of it all. All relevant 598 actors need to coordinate their efforts in extensive collaborations to achieve the necessary 599 conservation measures: in mountain areas with a protection status conservation policy needs 600 reinforcement; for mountain areas without a protection status, evaluation of its status, 601 importance and future perspective need to be used to prioritize (i) protective measures, (ii) re-602 evaluations of impacts of touristic and pastoral activities, (iii) evaluation of sustainability 603 management of natural resources, and (iv) development of early-warning systems of ecosystem 604 degradation and biodiversity loss. These measures will then be able to inform about trajectories 605 towards detrimental outcomes (pathogen emergence, ecosystem services; (Huber et al., 2013). 606 As mountain stakeholders are numerous, regional networks and coordination mechanisms must 607 urgently be installed, and a broad communication strategy needs to be developed to raise 608 awareness about the threats to mountains and their complex consequences (Brunner and Grêt-609 Regamey, 2016; Drexler et al., 2016). These consequences may have also an important social 610 component, as people may move out of mountain areas, if the conditions for cultivation and 611 exploitation are unfavourable, not providing for their livelihood. These different aspects need to 612 be included in comprehensive mountain ecosystem management plans, considering the 613 cumulative and hierarchical context of disturbance regimes to prevent reductions in ecological 614 variability and ecosystem resilience (Chiang et al., 2014).

In this light, and in that of the challenging objectives set by global agendas, including the
UN Sustainable Development Agenda, the Convention on Biological Diversity, the recently
launched EU Biodiversity Strategy for 2030 or the UN Decade on Ecosystem Restoration 20212030, investments are needed for the delivery of policy-relevant science on mountain
ecosystems (Körner, 2019), closely following recommendations given in the global biodiversity

- 620 framework (CBD/WG2020/3/3). Only if we maintain a high ecosystem resilience will we be able
- to maintain ecosystem functioning and ecosystem services. Threats to mountains are numerous
- and the repercussions to humanity demand conservation and restoration of mountain
- 623 ecosystems, as they are an essential and highly sensitive part of the global life-support system.
- 624

625 Acknowledgements

D.S.S. holds the AXA Chair for Functional Mountain Ecology funded by the AXA Research Fund
through the project GloMEc. M.C.F. acknowledges funding from the Natural Environment
Research Council (NERC) and the Medical Research Council (MRC) Centre for Global
Infectious Disease Analysis (reference MR/R015600/1) and is a fellow in the Canadian Institute
for Advanced Research (CIFAR) 'Fungal Kingdom' programme. D.C. was partly supported by a
grant of the Ministry of Research, Innovation and Digitization (PN-III-P4-PCE-2021-0818). DSS,
DC, DU, JC, and RS receive funding through the BiodivRestore COFUND Action (BiodivERsA)

- and Water JPI), and the FishME project (ANR-21-BIRE-0002-01, FWF-I-5824, UEFISCDI 276/
- 634 2022). Y-PL was financially supported by the Ministry of Science and Technology (Project
- numbers: MOST 110-2321-B-002-017), and Taiwan Agricultural Research Institute (COA,
 Contract 1113017).
- 637 Contract 11

638 References

- Acevedo-Whitehouse K, Duffus ALJ. Effects of environmental change on wildlife health.
 Philosophical Transactions of the Royal Society B-Biological Sciences 2009; 364: 3429 3438.
- Adams SB, Frissell CA, Rieman BE. Geography of invasion in mountain streams: consequences
 of headwater lake fish introductions. Ecosystems 2001; 4: 296-307.
- Aide TM, Grau HR, Graesser J, Andrade-Nuñez MJ, Aráoz E, Barros AP, et al. Woody
 vegetation dynamics in the tropical and subtropical Andes from 2001 to 2014: Satellite
 image interpretation and expert validation. Global Change Biology 2019; 25: 2112-2126.
- Allen S, Allen D, Phoenix VR, Le Roux G, Jiménez PD, Simonneau A, et al. Atmospheric
 transport and deposition of microplastics in a remote mountain catchment. Nature
 Geoscience 2019; 12: 339-344.
- Allison EA. The spiritual significance of glaciers in an age of climate change. Wiley
 Interdisciplinary Reviews: Climate Change 2015; 6: 493-508.
- Bacardit M, Camarero L. Atmospherically deposited major and trace elements in the winter
 snowpack along a gradient of altitude in the Central Pyrenees: the seasonal record of
 long-range fluxes over SW Europe. Atmospheric Environment 2010; 44: 582-595.
- Ball WT, Alsing J, Mortlock DJ, Staehelin J, Haigh JD, Peter T, et al. Evidence for a continuous
 decline in lower stratospheric ozone offsetting ozone layer recovery. Atmospheric
 Chemistry and Physics 2018; 18: 1379-1394.
- Balthazar V, Vanacker V, Molina A, Lambin EF. Impacts of forest cover change on ecosystem
 services in high Andean mountains. Ecological Indicators 2015; 48: 63-75.
- Barnes PW, Williamson CE, Lucas RM, Robinson SA, Madronich S, Paul ND, et al. Ozone
 depletion, ultraviolet radiation, climate change and prospects for a sustainable future.
 Nature Sustainability 2019; 2: 569-579.
- Bates KA, Sommer U, Hopkins KP, Shelton JMG, Wierzbicki C, Sergeant C, et al. Microbiome
 function predicts amphibian chytridiomycosis disease dynamics. Microbiome 2022; 10:
 44.

- Bernardo-Cravo A, Schmeller DS, Chatzinotas A, Vredenburg VT, Loyau A. Environmental
 Factors and Host Microbiomes Shape Host-Pathogen Dynamics. Trends in Parasitology
 2020; 36: 29-36.
- Bestion E, Haegeman B, Alvarez Codesal S, Garreau A, Huet M, Barton S, et al. Phytoplankton
 biodiversity is more important for ecosystem functioning in highly variable thermal
 environments. Proceedings of the National Academy of Sciences 2021; 118:
 e2019591118.
- Biemans H, Siderius C, Lutz A, Nepal S, Ahmad B, Hassan T, et al. Importance of snow and
 glacier meltwater for agriculture on the Indo-Gangetic Plain. Nature Sustainability 2019;
 2: 594-601.
- Blais JM, Schindler DW, Muir DC, Kimpe LE, Donald DB, Rosenberg B. Accumulation of
 persistent organochlorine compounds in mountains of western Canada. Nature 1998;
 395: 585-588.
- 679Boisvenue C, Running SW. Impacts of climate change on natural forest productivity-evidence680since the middle of the 20th century. Global Change Biology 2006; 12: 862-882.
- Bowman WD, Murgel J, Blett T, Porter E. Nitrogen critical loads for alpine vegetation and soils in
 Rocky Mountain National Park. Journal of Environmental Management 2012; 103: 165 171.
- Brack W, Barcelo Culleres D, Boxall ABA, Budzinski H, Castiglioni S, Covaci A, et al. One
 planet: one health. A call to support the initiative on a global science–policy body on
 chemicals and waste. Environmental Sciences Europe 2022; 34: 21.
- Bradford DF, Stanley K, McConnell LL, Tallent-Halsell NG, Nash MS, Simonich SM. Spatial
 patterns of atmospherically deposited organic contaminants at high elevation in the
 southern Sierra Nevada mountains, California, USA. Environ Toxicol Chem 2010; 29:
 1056-66.
- Bradford DF, Stanley KA, Tallent NG, Sparling DW, Nash MS, Knapp RA, et al. Temporal and
 spatial variation of atmospherically deposited organic contaminants at high elevation in
 Yosemite National Park, California, USA. Environmental toxicology and chemistry 2013;
 32: 517-525.
- Bradshaw CJA, Ehrlich PR, Beattie A, Ceballos G, Crist E, Diamond J, et al. Underestimating
 the Challenges of Avoiding a Ghastly Future. Frontiers in Conservation Science 2021; 1.
- 697 Brahney J, Hallerud M, Heim E, Hahnenberger M, Sukumaran S. Plastic rain in protected areas 698 of the United States. Science 2020; 368: 1257-1260.
- Brendonck L, De Meester L. Egg banks in freshwater zooplankton: evolutionary and ecological
 archives in the sediment. Hydrobiologia 2003; 491: 65-84.
- Bridgewater P, Loyau A, Schmeller DS. The seventh plenary of the intergovernmental platform
 for biodiversity and ecosystem services (IPBES-7): a global assessment and a reshaping
 of IPBES. Biodiversity and Conservation 2019; 28: 2457-2461.
- Brooks PD, O'Reilly CM, Diamond SA, Campbell DH, Knapp R, Bradford D, et al. Spatial and
 temporal variability in the amount and source of dissolved organic carbon: implications
 for ultraviolet exposure in amphibian habitats. Ecosystems 2005; 8: 478-487.
- Brunner SH, Grêt-Regamey A. Policy strategies to foster the resilience of mountain social ecological systems under uncertain global change. Environmental Science & Policy
 2016; 66: 129-139.
- Camarero L, Botev I, Muri G, Psenner R, Rose N, Stuchlík E. Trace elements in alpine and
 arctic lake sediments as a record of diffuse atmospheric contamination across Europe.
 Freshwater Biology 2009; 54: 2518-2532.
- Caminade C, McIntyre KM, Jones AE. Impact of recent and future climate change on
 vector-borne diseases. Annals of the New York Academy of Sciences 2019; 1436: 157.

- Catalan J. Tracking long-range atmospheric transport of trace metals, polycyclic aromatic
 hydrocarbons, and organohalogen compounds using lake sediments of mountain
 regions. Environmental contaminants. Springer, 2015, pp. 263-322.
- Catalan J, Camarero, L., Felip, M., Pla, S., Ventura, M., Buchaca, T., Bartumeus F, Guillermo
 de Mendoza, Miró, A., Casamayor, E.O., Medina-Sánchez, J.M., Bacardit, M., Altuna, M.,
 Bartrons, M., Díaz de Quijano, D. High mountain lakes: extreme habitats and witnesses
 of environmental changes. Limnetica 2006; 25: 551-584.
- Catalan J, Donato Rondon JC. Perspectives for an integrated understanding of tropical and
 temperate high-mountain lakes. Journal of Limnology 2016; 75: 215-234.
- Catalan J, Ninot JM, Aniz MM. High Mountain Conservation in a Changing World. Vol 62. Cham:
 Springer, 2017.
- Catherine Q, Susanna W, Isidora E-S, Mark H, Aurelie V, Jean-François H. A review of current
 knowledge on toxic benthic freshwater cyanobacteria–ecology, toxin production and risk
 management. Water research 2013; 47: 5464-5479.
- Cavicchioli R, Ripple WJ, Timmis KN, Azam F, Bakken LR, Baylis M, et al. Scientists' warning to
 humanity: microorganisms and climate change. Nat Rev Microbiol 2019.
- Chape S, Harrison J, Spalding M, Lysenko I. Measuring the extent and effectiveness of
 protected areas as an indicator for meeting global biodiversity targets. Philosophical
 Transactions of the Royal Society B: Biological Sciences 2005; 360: 443-455.
- Chiang L-C, Lin Y-P, Huang T, Schmeller DS, Verburg PH, Liu Y-L, et al. Simulation of
 ecosystem service responses to multiple disturbances from an earthquake and several
 typhoons. Landscape and urban planning 2014; 122: 41-55.
- Cifuentes-Croquevielle C, Stanton DE, Armesto JJ. Soil invertebrate diversity loss and functional
 changes in temperate forest soils replaced by exotic pine plantations. Scientific Reports
 2020; 10: 7762.
- Crutzen PJ. The "anthropocene". Earth system science in the anthropocene. Springer, 2006, pp.13-18.
- Dadam D, Robinson RA, Clements A, Peach WJ, Bennett M, Rowcliffe JM, et al. Avian malaria mediated population decline of a widespread iconic bird species. Royal Society open
 science 2019; 6: 182197.
- 745 Daly GL, Wania F. Organic contaminants in mountains. Environmental science & technology
 746 2005; 39: 385-398.
- De Jong R, Blaauw M, Chambers FM, Christensen TR, De Vleeschouwer Fo, Finsinger W, et al.
 Climate and peatlands. Changing Climates, Earth Systems and Society. Springer
 Netherlands, 2010, pp. 85-121.
- de Oliveira DG, Schneider G, Itokazu AG, Costa GB, Rörig LR, Simonassi JC, et al. Effects of
 ultraviolet radiation removal on algal communities in three high-elevation Brazilian (ultra)
 oligotrophic lakes. Phycologia 2021; 60: 497-512.
- Del Barrio G, Creus J, Puigdefábregas J. Thermal seasonality of the high mountain belts of the
 Pyrenees. Mountain Research and Development 1990: 227-233.
- Dhimal M, Ahrens B, Kuch U. Climate change and spatiotemporal distributions of vector-borne diseases in Nepal–a systematic synthesis of literature. PloS one 2015a; 10: e0129869.
- Dhimal M, Gautam I, Joshi HD, O'Hara RB, Ahrens B, Kuch U. Risk factors for the presence of
 chikungunya and dengue vectors (Aedes aegypti and Aedes albopictus), their altitudinal
 distribution and climatic determinants of their abundance in central Nepal. PLoS
 Neglected Tropical Diseases 2015b; 9: e0003545.
- Diagne C, Leroy B, Vaissière A-C, Gozlan RE, Roiz D, Jarić I, et al. High and rising economic
 costs of biological invasions worldwide. Nature 2021; 592: 571-576.
- Diamond SA, Trenham PC, Adams MJ, Hossack BR, Knapp RA, Stark SL, et al. Estimated
 ultraviolet radiation doses in wetlands in six national parks. Ecosystems 2005; 8: 462 477.

- Dierauer JR, Whitfield PH, Allen DM. Climate controls on runoff and low flows in mountain
 catchments of Western North America. Water Resources Research 2018; 54: 7495 7510.
- Dobrowski SZ, Abatzoglou JT, Greenberg JA, Schladow S. How much influence does
 landscape-scale physiography have on air temperature in a mountain environment?
 Agricultural and Forest Meteorology 2009; 149: 1751-1758.
- Dong Z, Brahney J, Kang S, Elser J, Wei T, Jiao X, et al. Aeolian dust transport, cycle and
 influences in high-elevation cryosphere of the Tibetan Plateau region: New evidences
 from alpine snow and ice. Earth-Science Reviews 2020; 211: 103408.
- Drexler C, Braun V, Christie D, Claramunt B, Dax T, Jelen I, et al. Mountains for Europe's
 Future–A strategic research agenda. Bern, Switzerland: Mountain Research Initiative,
 Institute of Interdisciplinary Mountain Research 2016.
- Du X, Liu H, Yuan L, Wang Y, Ma Y, Wang R, et al. The Diversity of Cyanobacterial Toxins on Structural Characterization, Distribution and Identification: A Systematic Review. Toxins 2019; 11: 530.
- Dullinger S, Gattringer A, Thuiller W, Moser D, Zimmermann NE, Guisan A, et al. Extinction debt
 of high-mountain plants under twenty-first-century climate change. Nature climate
 change 2012; 2: 619-622.
- Einhorn B, Eckert N, Chaix C, Ravanel L, Deline P, Gardent M, et al. Climate change and
 natural hazards in the Alps. Observed and potential impacts on physical and socio economic systems. Journal of Alpine Research Revue de géographie alpine 2015.
- 787 Eklabya S, Rita S, Singh KK, Sharma G. A Boon for Mountain Populations. Mountain Research
 788 and Development 2000; 20: 108-111.
- Epanchin PN, Knapp RA, Lawler SP. Nonnative trout impact an alpine-nesting bird by altering
 aquatic-insect subsidies. Ecology 2010; 91: 2406-2415.
- Faccioni G, Sturaro E, Ramanzin M, Bernués A. Socio-economic valuation of abandonment and intensification of Alpine agroecosystems and associated ecosystem services. Land Use Policy 2019; 81: 453-462.
- Fahey B, Jackson R. Hydrological impacts of converting native forests and grasslands to pine
 plantations, South Island, New Zealand. Agricultural and Forest Meteorology 1997; 84:
 69-82.
- Fait P, Demierre E, Ilg C, Oertli B. Small mountain reservoirs in the Alps: New habitats for alpine
 freshwater biodiversity? Aquatic Conservation: Marine and Freshwater Ecosystems
 2020; 30: 617-630.
- Fang X, Pyle JA, Chipperfield MP, Daniel JS, Park S, Prinn RG. Challenges for the recovery of
 the ozone layer. Nature Geoscience 2019; 12: 592-596.
- Ficetola GF, Marta S, Guerrieri A, Gobbi M, Ambrosini R, Fontaneto D, et al. Dynamics of
 ecological communities following current retreat of glaciers. Annual Review of Ecology,
 Evolution, and Systematics 2021; 52: 405-426.
- Finlayson C, D'Cruz R. Inland water systems. Ecosystems and human well-being: Current state
 and trends. Island Press, 2005, pp. 551-583.
- Fisher MC, Garner TW. Chytrid fungi and global amphibian declines. Nature Reviews
 Microbiology 2020; 18: 332-343.
- Fonseca CR, Ganade G. Species functional redundancy, random extinctions and the stability of ecosystems. Journal of Ecology 2001; 89: 118-125.
- Forero-Medina G, Joppa L, Pimm SL. Constraints to Species' Elevational Range Shifts as
 Climate Changes. Conservation Biology 2011; 25: 163-171.
- 813 Franssen HH, Scherrer S. Freezing of lakes on the Swiss plateau in the period 1901–2006.
- 814 International Journal of Climatology: A Journal of the Royal Meteorological Society 2008;
 815 28: 421-433.

- Frenken T, Alacid E, Berger SA, Bourne EC, Gerphagnon M, Grossart H-P, et al. Integrating
 chytrid fungal parasites into plankton ecology. Research gaps and needs. Environmental
 Microbiology 2017; 19: 3802–3822.
- Funari E, Testai E. Human health risk assessment related to cyanotoxins exposure. Critical
 reviews in toxicology 2008; 38: 97-125.
- Gałuszka A, Migaszewski ZM, Zalasiewicz J. Assessing the Anthropocene with geochemical
 methods. Geological Society, London, Special Publications 2014; 395: 221-238.
- Gao W-Q, Lei X-D, Fu L-Y. Impacts of climate change on the potential forest productivity based
 on a climate-driven biophysical model in northeastern China. Journal of Forestry
 Research 2020; 31: 2273-2286.
- Barcia-Vozmediano A, Krawczyk AI, Sprong H, Rossi L, Ramassa E, Tomassone L. Ticks climb
 the mountains: ixodid tick infestation and infection by tick-borne pathogens in the
 Western Alps. Ticks and Tick-borne Diseases 2020; 11: 101489.
- Gobiet A, Kotlarski S, Beniston M, Heinrich G, Rajczak J, Stoffel M. 21st century climate change
 in the European Alps—A review. Science of the Total Environment 2014; 493: 1138 1151.
- Grabherr G, Gottfried M, Pauli H. Climate effects on mountain plants. Nature 1994; 369: 448448.
- Grêt-Regamey A, Brunner SH, Kienast F. Mountain ecosystem services: who cares? Mountain
 Research and Development 2012; 32: S23-S34.
- Biggin Grêt-Regamey A, Weibel B. Global assessment of mountain ecosystem services using earth
 observation data. Ecosystem Services 2020; 46: 101213.
- Gross M. Anthropocene at altitude. Current Biology 2022; 32: R441-R444.
- Haberl H, Erb KH, Krausmann F, Gaube V, Bondeau A, Plutzar C, et al. Quantifying and
 mapping the human appropriation of net primary production in earth's terrestrial
 ecosystems. Proceedings of the National Academy of Sciences 2007; 104: 12942-12947.
- Häder DP, Helbling EW, Williamson CE, Worrest RC. Effects of UV radiation on aquatic
 ecosystems and interactions with climate change. Photochemical & Photobiological
 Sciences 2011; 10: 242-260.
- Hall JM, Van Holt T, Daniels AE, Balthazar V, Lambin EF. Trade-offs between tree cover, carbon
 storage and floristic biodiversity in reforesting landscapes. Landscape Ecology 2012; 27:
 1135-1147.
- Hansson SV, Claustres A, Probst A, De Vleeschouwer F, Baron S, Galop D, et al. Atmospheric
 and terrigenous metal accumulation over 3000 years in a French mountain catchment:
 Local vs distal influences. Anthropocene 2017a; 19: 45-54.
- Hansson SV, Sonke J, Galop D, Bareille G, Jean S, Le Roux G. Transfer of marine mercury to
 mountain lakes. Scientific Reports 2017b; 7: 12719.
- Harrison S, Kargel JS, Huggel C, Reynolds J, Shugar DH, Betts RA, et al. Climate change and the global pattern of moraine-dammed glacial lake outburst floods. The Cryosphere 2018; 12: 1195-1209.
- Hartmann H, Andresky L. Flooding in the Indus River basin—a spatiotemporal analysis of
 precipitation records. Global and planetary change 2013; 107: 25-35.
- Havas M, Rosseland BO. Response of zooplankton, benthos, and fish to acidification: an
 overview. Water, Air, and Soil Pollution 1995; 85: 51-62.
- Haver M, Le Roux G, Friesen J, Loyau A, Vredenburg VT, Schmeller DS. The role of abiotic
 variables in an emerging global amphibian fungal disease in mountains. Science of The
 Total Environment 2021: 152735.
- Heino J. The importance of metacommunity ecology for environmental assessment research in
 the freshwater realm. Biological Reviews 2013; 88: 166-178.
- Hemp A. Introduced plants on Kilimanjaro: tourism and its impact. Plant Ecology 2008; 197: 1729.

- Herzog F, Seidl I. Swiss alpine summer farming: current status and future development under
 climate change. The Rangeland Journal 2018; 40: 501-511.
- Hinojosa L, Napoléone C, Moulery M, Lambin EF. The "mountain effect" in the abandonment of grasslands: Insights from the French Southern Alps. Agriculture, Ecosystems & Environment 2016; 221: 115-124.
- Ho JC, Michalak AM, Pahlevan N. Widespread global increase in intense lake phytoplankton
 blooms since the 1980s. Nature 2019; 574: 667-670.
- Hobbs RJ, Higgs E, Harris JA. Novel ecosystems: implications for conservation and restoration.
 Trends in ecology & evolution 2009; 24: 599-605.
- Hofstede RGM, Groenendijk JP, Coppus R, Fehse JC, Sevink J. Impact of Pine Plantations on
 Soils and Vegetation in the Ecuadorian High Andes. Mountain Research and
 Development 2002; 22: 159-167, 9.
- Huber R, Rigling A, Bebi P, Brand FS, Briner S, Buttler A, et al. Sustainable land use in
 mountain regions under global change: synthesis across scales and disciplines. Ecology
 and Society 2013; 18.
- Huss M, Bookhagen B, Huggel C, Jacobsen D, Bradley RS, Clague JJ, et al. Toward mountains
 without permanent snow and ice. Earth's Future 2017; 5: 418-435.
- Hussain BA, Westgate JN, Hayward SJ, Shunthirasingham C, Brown TN, Hung H, et al.
 Polycyclic aromatic hydrocarbons and polychlorinated biphenyls in soils and atmosphere
 of Western Canadian mountains: The role of source proximity, precipitation, forest cover
 and mountain cold-trapping. Atmospheric Environment: X 2019; 1: 100004.
- 888 Immerzeel WW, Lutz A, Andrade M, Bahl A, Biemans H, Bolch T, et al. Importance and 889 vulnerability of the world's water towers. Nature 2020; 577: 364-369.
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services Global
 assessment report on biodiversity and ecosystem services of the Intergovernmental
 Science-Policy Platform on Biodiversity and Ecosystem Services. In: Brondizio ES,
 Settele J, Díaz S, Ngo HT, editors. IPBES Secretariat, Bonn, Germany, 2019, pp. 1148.
- Jacobsen D. The dilemma of altitudinal shifts: caught between high temperature and low
 oxygen. Frontiers in Ecology and the Environment 2020; 18: 211-218.
- Jacobsen D, Dangles O. Ecology of high altitude waters: Oxford University Press, 2017.
- Jäger H, Peratoner G, Tappeiner U, Tasser E. Grassland biomass balance in the European
 Alps: current and future ecosystem service perspectives. Ecosystem Services 2020; 45:
 101163.
- Kagami M, Miki T, Takimoto G. Mycoloop: chytrids in aquatic food webs. Frontiers in
 microbiology 2014; 5: 166.
- Kammerlander B, Breiner H-W, Filker S, Sommaruga R, Sonntag B, Stoeck T. High diversity of
 protistan plankton communities in remote high mountain lakes in the European Alps and
 the Himalayan mountains. FEMS microbiology ecology 2015; 91: fiv010.
- Kapnick S, Hall A. Causes of recent changes in western North American snowpack. Climate
 Dynamics 2012; 38: 1885-1899.
- Kastner T, Matej S, Forrest M, Gingrich S, Haberl H, Hickler T, et al. Land use intensification
 increasingly drives the spatiotemporal patterns of the global human appropriation of net
 primary production in the last century. Global Change Biology 2022; 28: 307-322.
- Kelty MJ. The role of species mixtures in plantation forestry. Forest Ecology and Management
 2006; 233: 195-204.
- Kerr JT, Deguise I. Habitat loss and the limits to endangered species recovery. Ecology Letters
 2004; 7: 1163-1169.
- Knapp R, Sarnelle O. Recovery after local extinction: factors affecting re-establishment of alpine
 lake zooplankton. Ecological Applications 2008; 18: 1850 1859.
- Knapp RA, Matthews KR, Sarnelle O. Resistance and Resilience of Alpine Lake Fauna to Fish
 Introductions. Ecological Monographs 2001; 71: 401-421.

- Kopáček J, Fluksová H, Hejzlar J, Kaňa J, Porcal P, Turek J. Changes in surface water
 chemistry caused by natural forest dieback in an unmanaged mountain catchment.
 Science of the Total Environment 2017; 584: 971-981.
- Körner C. Mountain biodiversity, its causes and function. AMBIO: A Journal of the Human
 Environment 2004; 33: 11-17.
- Körner C. The use of 'altitude'in ecological research. Trends in ecology & evolution 2007; 22:
 569-574.
- Körner C. Mountain biodiversity, its causes and function: an overview. Mountain biodiversity2019: 3-20.
- Körner C, Jetz W, Paulsen J, Payne D, Rudmann-Maurer K, Spehn EM. A global inventory of
 mountains for bio-geographical applications. Alpine Botany 2017; 127: 1-15.
- Körner C, Paulsen J, Spehn EM. A definition of mountains and their bioclimatic belts for global
 comparisons of biodiversity data. Alpine Botany 2011; 121: 73-78.
- Körner W, Dupouey JL, Dambrine E, Benoit M. Influence of Past Land Use on the Vegetation
 and Soils of Present Day Forest in the Vosges Mountains, France. Journal of Ecology
 1997; 85: 351-358.
- Laternser M, Schneebeli M. Long-term snow climate trends of the Swiss Alps (1931–99).
 International Journal of Climatology: A Journal of the Royal Meteorological Society 2003;
 23: 733-750.
- Latty EF, Canham CD, Marks PL. The effects of land-use history on soil properties and nutrient dynamics in northern hardwood forests of the Adirondack Mountains. Ecosystems 2004; 7: 193-207.
- Laurion I, Ventura M, Catalan J, Psenner R, Sommaruga R. Attenuation of ultraviolet radiation in mountain lakes: Factors controlling the among-and within-lake variability. Limnology and Oceanography 2000; 45: 1274-1288.
- Lavorel S, Grigulis K, Leitinger G, Kohler M, Schirpke U, Tappeiner U. Historical trajectories in
 land use pattern and grassland ecosystem services in two European alpine landscapes.
 Regional Environmental Change 2017; 17: 2251-2264.
- Le Roux G, Hansson SV, Claustres A. Inorganic chemistry in the mountain critical zone: are the
 mountain water towers of contemporary society under threat by trace contaminants?
 Developments in Earth Surface Processes. 21. Elsevier, 2016, pp. 131-154.
- Le Roux G, Hansson SV, Claustres A, Binet S, De Vleeschouwer F, Gandois L, et al. Trace
 Metal Legacy in Mountain Environments. Biogeochemical Cycles, 2020, pp. 191-206.
- Lei YD, Wania F. Is rain or snow a more efficient scavenger of organic chemicals? Atmospheric
 Environment 2004; 38: 3557-3571.
- Lenton TM. Early warning of climate tipping points. Nature Climate Change 2011; 1: 201-209.
- Lerner H, Berg C. A Comparison of Three Holistic Approaches to Health: One Health,
 EcoHealth, and Planetary Health. Frontiers in Veterinary Science 2017; 4.
- Lin Y-P, Chen C-J, Lien W-Y, Chang W-H, Petway J, Chiang L. Landscape Conservation
 Planning to Sustain Ecosystem Services under Climate Change. Sustainability 2019; 11:
 1393.
- Lin Y-P, Wunderlich RF, Lin C-M, Uphoff N, Schmeller DS, Shipin OV, et al. Topsoil microbial
 community structure responds to land cover type and environmental zone in the Western
 Pacific region. Science of The Total Environment 2021; 764: 144349.
- Lin YP, Lin W-C, Wang Y-C, Lien W-Y, Huang TH, Hsu CC, et al. Systematically designating
 conservation areas for protecting multiple ecosystem services. Environmental Modelling
 & Software 2017; 90: 126-146.
- Liu H, Mi Z, Lin L, Wang Y, Zhang Z, Zhang F, et al. Shifting plant species composition in
 response to climate change stabilizes grassland primary production. Proceedings of the
 National Academy of Sciences 2018; 115: 4051-4056.

- Liu X, Dou L, Ding X, Sun T, Zhang H. Influences of different afforestation systems on the soil
 properties of limestone mountains in the mid-eastern region of China. Catena 2021; 201:
 105198.
- 971 López-Moreno JI, Beniston M, García-Ruiz JM. Environmental change and water management
 972 in the Pyrenees: Facts and future perspectives for Mediterranean mountains. Global and
 973 Planetary Change 2008; 61: 300-312.
- Lundgren B. Soil conditions and nutrient cycling under natural and plantation forests in
 Tanzanian highlands: Swedish University of Agricultural Sciences, 1978.
- Machate O, Schmeller DS, Loyau A, Paschke A, Krauss M, Carmona E, et al. Complex chemical
 cocktail, containing insecticides diazinon and permethrin, drives acute toxicity to
 crustaceans in mountain lakes. Science of The Total Environment 2022: 154456.
- Martín-López B, Leister I, Lorenzo Cruz P, Palomo I, Grêt-Regamey A, Harrison PA, et al.
 Nature's contributions to people in mountains: a review. PloS one 2019; 14: e0217847.
- Maxwell SL, Fuller RA, Brooks TM, Watson JEM. Biodiversity: The ravages of guns, nets and
 bulldozers. Nature 2016; 536: 143-145.
- Mayer A, Egger C, Loyau A, Plutzar C, Schmeller DS, Gaube V. Mountain pastures increase the
 resilience of livestock farming to extreme events in the Ariège department, France.
 Agronomy for Sustainable Development 2022; 42: 49.
- Mayerhofer J, Wächter D, Calanca P, Kohli L, Roth T, Meuli RG, et al. Environmental and
 Anthropogenic Factors Shape Major Bacterial Community Types Across the Complex
 Mountain Landscape of Switzerland. Frontiers in Microbiology 2021; 12: 500.
- McCain CM, Colwell RK. Assessing the threat to montane biodiversity from discordant shifts in temperature and precipitation in a changing climate. Ecology letters 2011; 14: 1236-1245.
- McCullough IM, Davis FW, Dingman JR, Flint LE, Flint AL, Serra-Diaz JM, et al. High and dry:
 high elevations disproportionately exposed to regional climate change in Mediterranean climate landscapes. Landscape Ecology 2016; 31: 1063-1075.
- Meadows DH, Meadows DL, Randers J, Behrens III WW. The limits to growth: a report to the
 club of Rome (1972). New York, Washington, DC (USA): Universe Books, Potomac
 Associates, 1972.
- Meire RO, Lee SC, Yao Y, Targino AC, Torres JPM, Harner T. Seasonal and altitudinal
 variations of legacy and current-use pesticides in the Brazilian tropical and subtropical
 mountains. Atmospheric Environment 2012; 59: 108-116.
- Melillo JM, McGuire AD, Kicklighter DW, Moore B, Vorosmarty CJ, Schloss AL. Global climate
 change and terrestrial net primary production. Nature 1993; 363: 234-240.
- Milner AM, Khamis K, Battin TJ, Brittain JE, Barrand NE, Füreder L, et al. Glacier shrinkage
 driving global changes in downstream systems. Proceedings of the National Academy of
 Sciences 2017; 114: 9770-9778.
- 1006Miró A, Ventura M. Evidence of exotic trout mediated minnow invasion in Pyrenean high1007mountain lakes. Biological Invasions 2015; 17: 791-803.
- 1008 Miró A, Ventura M. Introduced fish in Pyrenean high mountain lakes: impact on amphibians and 1009 other organisms, and conservation implications. Limnetica 2020; 39: 283-297.
- Mölg T, Hardy DR, Cullen NJ, Kaser G. Tropical glaciers, climate change, and society.
 Darkening peaks: Glacier retreat, science, and society 2008: 168-182.
- Moser KA, Baron JS, Brahney J, Oleksy IA, Saros JE, Hundey EJ, et al. Mountain lakes: Eyes
 on global environmental change. Global and Planetary Change 2019; 178: 77-95.
- Moulin C, Chiapello I. Impact of human-induced desertification on the intensification of Sahel
 dust emission and export over the last decades. Geophysical Research Letters 2006; 33:
 L18808.

- Myhre G, Alterskjær K, Stjern CW, Hodnebrog Ø, Marelle L, Samset BH, et al. Frequency of
 extreme precipitation increases extensively with event rareness under global warming.
 Scientific Reports 2019; 9: 16063.
- Negi VS, Tiwari DC, Singh L, Thakur S, Bhatt ID. Review and synthesis of climate change
 studies in the Himalayan region. Environment, Development and Sustainability 2021.
- Niedertscheider M, Tasser E, Patek M, Rüdisser J, Tappeiner U, Erb K-H. Influence of Land-Use
 Intensification on Vegetation C-Stocks in an Alpine Valley from 1865 to 2003.
 Ecosystems 2017; 20: 1391-1406.
- Niedrist G, Psenner R, Sommaruga R. Climate warming increases vertical and seasonal water
 temperature differences and inter-annual variability in a mountain lake. Climatic Change
 2018; 151: 473-490.
- 1028 Niedrist GH, Füreder L. Real-time warming of alpine streams:(Re) defining invertebrates' 1029 temperature preferences. River Research and Applications 2021; 37: 283-293.
- Nielsen DL, Smith D, Petrie R. Resting egg banks can facilitate recovery of zooplankton
 communities after extended exposure to saline conditions. Freshwater Biology 2012; 57:
 1306-1314.
- Niu L, Cheng Z. Impact of tourism disturbance on forest vegetation in Wutai Mountain, China.
 Environmental Monitoring and Assessment 2019; 191: 81.
- Nodvin S, Van Miegroet H, Lindberg S, Nicholas N, Johnson D. Acidic deposition, ecosystem
 processes, and nitrogen saturation in a high elevation southern Appalachian watershed.
 Water, Air, and Soil Pollution 1995; 85: 1647-1652.
- Nogues-Bravo D, Araújo MB, M.P. E, Martínez-Rica JP. Exposure of global mountain systems to
 climate warming during the 21st Century. Global Environmental Change 2007; 17: 420 428.
- 1041 Noyes PD, McElwee MK, Miller HD, Clark BW, Van Tiem LA, Walcott KC, et al. The toxicology
 1042 of climate change: environmental contaminants in a warming world. Environ Int 2009; 35:
 1043 971-86.
- 1044Obertegger U, Flaim G. A 40-year perspective of an alpine lake: Is everything the same?1045Limnologica 2021; 91: 125929.
- 1046Palomo I. Climate change impacts on ecosystem services in high mountain areas: a literature1047review. Mountain Research and Development 2017; 37: 179-187.
- 1048Parmesan C, Yohe G. A globally coherent fingerprint of climate change impacts across natural1049systems. Nature 2003; 421: 37-42.
- Pauchard A, Kueffer C, Dietz H, Daehler CC, Alexander J, Edwards PJ, et al. Ain't no mountain
 high enough: plant invasions reaching new elevations. Frontiers in Ecology and the
 Environment 2009; 7: 479-486.
- Payne D, Spehn EM, Prescott GW, Geschke J, Snethlage MA, Fischer M. Mountain Biodiversity
 Is Central to Sustainable Development in Mountains and Beyond. One earth 2020; 3:
 530-533.
- Pecl GT, Araújo MB, Bell JD, Blanchard J, Bonebrake TC, Chen I-C, et al. Biodiversity
 redistribution under climate change: Impacts on ecosystems and human well-being.
 Science 2017; 355: eaai9214.
- Pepin N, Bradley RS, Diaz H, Baraër M, Caceres E, Forsythe N, et al. Elevation-dependent
 warming in mountain regions of the world. Nature climate change 2015; 5: 424-430.
- Pepin N, Lundquist J. Temperature trends at high elevations: patterns across the globe.
 Geophysical Research Letters 2008; 35.
- Pimm SL. Biodiversity: Climate change or habitat loss Which will kill more species? Current
 Biology 2008; 18: R117-R119.
- Pozo K, Urrutia R, Barra R, Mariottini M, Treutler H-C, Araneda A, et al. Records of
 polychlorinated biphenyls (PCBs) in sediments of four remote Chilean Andean Lakes.
 Chemosphere 2007; 66: 1911-1921.

- Purcell AM, Hayer M, Koch BJ, Mau RL, Blazewicz SJ, Dijkstra P, et al. Decreased growth of
 wild soil microbes after 15 years of transplant-induced warming in a montane meadow.
 Global Change Biology 2022; 28: 128-139.
- 1071 Ragettli S, Tong X, Zhang G, Wang H, Zhang P, Stähli M. Climate change impacts on summer
 1072 flood frequencies in two mountainous catchments in China and Switzerland. Hydrology
 1073 Research 2021; 52: 4-25.
- 1074 Rahbek C, Borregaard MK, Antonelli A, Colwell RK, Holt BG, Nogues-Bravo D, et al. Building
 1075 mountain biodiversity: Geological and evolutionary processes. Science 2019a; 365:
 1076 1114-1119.
- 1077 Rahbek C, Borregaard MK, Colwell RK, Dalsgaard B, Holt BG, Morueta-Holme N, et al.
 1078 Humboldt's enigma: What causes global patterns of mountain biodiversity? Science
 1079 2019b; 365: 1108-1113.
- 1080 Rajczak J, Schär C. Projections of future precipitation extremes over Europe: A multimodel
 1081 assessment of climate simulations. Journal of Geophysical Research: Atmospheres
 2017; 122: 10,773-10,800.
- 1083Rangwala I, Miller JR. Climate change in mountains: a review of elevation-dependent warming1084and its possible causes. Climatic change 2012; 114: 527-547.
- Reid AJ, Carlson AK, Creed IF, Eliason EJ, Gell PA, Johnson PT, et al. Emerging threats and
 persistent conservation challenges for freshwater biodiversity. Biological Reviews 2019;
 94: 849-873.
- 1088 Ripple W, J., Estes J, A., Beschta R, L., Wilmers C, C., Ritchie E, G., Hebblewhite M, et al.
 1089 Status and Ecological Effects of the World's Largest Carnivores. Science 2014; 343:
 1090 1241484.
- 1091 Ripple WJ, Newsome TM, Wolf C, Dirzo R, Everatt KT, Galetti M, et al. Collapse of the world's
 1092 largest herbivores. Science advances 2015; 1: e1400103.

1093Ripple WJ, Wolf C, Newsome TM, Galetti M, Alamgir M, Crist E, et al. World Scientists' Warning1094to Humanity: A Second Notice. BioScience 2017: bix125-bix125.

- 1095 Ripple WJ, Wolf C, Newsome TM, Gregg JW, Lenton TM, Palomo I, et al. World Scientists'
 1096 Warning of a Climate Emergency 2021. BioScience 2021.
- Rockström J, Steffen W, Noone K, Persson A, Chapin FS, Lambin EF, et al. A safe operating
 space for humanity. Nature 2009; 461: 472-475.
- Rodway-Dyer S, Ellis N. Combining remote sensing and on-site monitoring methods to
 investigate footpath erosion within a popular recreational heathland environment. Journal
 of Environmental Management 2018; 215: 68-78.
- 1102 Roe GH. Orographic precipitation. Annu. Rev. Earth Planet. Sci. 2005; 33: 645-671.
- Rose KC, Williamson CE, Saros JE, Sommaruga R, Fischer JM. Differences in UV transparency
 and thermal structure between alpine and subalpine lakes: implications for organisms.
 Photochemical & Photobiological Sciences 2009; 8: 1244-1256.
- Saunders DL, Kalff J. Nitrogen retention in wetlands, lakes and rivers. Hydrobiologia 2001; 443:
 205-212.
- Scarano FR. Biodiversity Sector: Risks of Temperature Increase to Biodiversity and
 Ecosystems. In: Nobre CA, Marengo JA, Soares WR, editors. Climate Change Risks in
 Brazil. Springer International Publishing, Cham, 2019, pp. 131-141.
- Schabetsberger R, Kaiser R, Rott E, Lenzenweger R, Traunspurger W, Kotov AA, et al. On the
 brink investigating biodiversity in endangered crater lakes of the Amber Mountains
 National Park (Madagascar). Aquatic Conservation: Marine and Freshwater Ecosystems
 2013; 23: 316-331.

Schai-Braun SC, Jenny H, Ruf T, Hackländer K. Temperature increase and frost decrease driving upslope elevational range shifts in Alpine grouse and hares. Global Change Biology 2021; 27: 6602-6614.

1118 Schindler DE, Knapp RA, Leavitt PR. Alteration of Nutrient Cycles and Algal Production 1119 Resulting from Fish Introductions intoMountain Lakes. Ecosystems 2001: 4: 308-321. 1120 Schirpke U, Tappeiner U, Tasser E. A transnational perspective of global and regional 1121 ecosystem service flows from and to mountain regions. Scientific reports 2019; 9: 1-11. 1122 Schmeller DS, Bridgewater P. The eighth plenary of the Intergovernmental Science-Policy 1123 Platform for Biodiversity and Ecosystem Services (IPBES-8): online, nexus, and 1124 transformative change. Biodiversity and Conservation 2021; 33: 2857-2862. 1125 Schmeller DS, Courchamp F, Killeen G. Biodiversity loss, emerging pathogens and human 1126 health risks. Biodiversity and Conservation 2020: 3095–3102. 1127 Schmeller DS, Loyau A, Bao K, Brack W, Chatzinotas A, De Vleeschouwer F, et al. People, 1128 pollution and pathogens – Global change impacts in mountain freshwater ecosystems. 1129 Science of The Total Environment 2018; 622-623: 756-763. 1130 Schmeller DS, Niemelä J, Bridgewater P. The Intergovernmental Science-Policy Platform on 1131 Biodiversity and Ecosystem Services (IPBES): getting involved. Biodiversity and 1132 Conservation 2017; 26: 2271-2275. 1133 Scientists UoC. World scientists' Warning to Humanity. Union of Concerned Scientists, 1134 http://www.ucsusa.org/sites/default/files/attach/2017/11/World%20Scientists%27%20War ning%20to%20Humanity%201992, 1992. 1135 Seebens H, Bacher S, Blackburn TM, Capinha C, Dawson W, Dullinger S, et al. Projecting the 1136 1137 continental accumulation of alien species through to 2050. Global Change Biology 2021; 1138 27: 970-982. 1139 Sentenac H, Loyau A, Leflaive J, Schmeller DS. The significance of biofilms to human, animal, 1140 plant and ecosystem health. Functional Ecology 2022; 33: 294 - 313. 1141 Shijin W, Dahe Q. Mountain inhabitants' perspectives on climate change, and its impacts and 1142 adaptation based on temporal and spatial characteristics analysis: a case study of Mt. 1143 Yulong Snow, Southeastern Tibetan Plateau. Environmental Hazards 2015; 14: 122-136. 1144 Shunthirasingham C, Oyiliagu CE, Cao X, Gouin T, Wania F, Lee S-C, et al. Spatial and 1145 temporal pattern of pesticides in the global atmosphere. Journal of Environmental 1146 Monitoring 2010; 12: 1650-1657. Silva V, Montanarella L, Jones A, Fernández-Ugalde O, Mol HG, Ritsema CJ, et al. Distribution 1147 1148 of glyphosate and aminomethylphosphonic acid (AMPA) in agricultural topsoils of the 1149 European Union. Science of the Total Environment 2018; 621: 1352-1359. 1150 Siraj A, Santos-Vega M, Bouma M, Yadeta D, Carrascal DR, Pascual M. Altitudinal changes in 1151 malaria incidence in highlands of Ethiopia and Colombia. Science 2014; 343: 1154-1158. 1152 Small ST, Labbé F, Coulibaly YI, Nutman TB, King CL, Serre D, et al. Human migration and the 1153 spread of the nematode parasite Wuchereria bancrofti. Molecular biology and evolution 1154 2019; 36: 1931-1941. 1155 Sommaruga R. The role of solar UV radiation in the ecology of alpine lakes. Journal of 1156 Photochemistry and Photobiology B: Biology 2001; 62: 35-42. 1157 Sommaruga R. When glaciers and ice sheets melt: consequences for planktonic organisms. 1158 Journal of plankton research 2015; 37: 509-518. 1159 Sommaruga R, Sattler B, Oberleiter A, Wille A, Wögrath-Sommaruga S, Psenner R, et al. An in situ enclosure experiment to test the solar UVB impact on plankton in a high-altitude 1160 1161 mountain lake. II. Effects on the microbial food web. Journal of Plankton Research 1999; 1162 21. 1163 Steffen W, Richardson K, Rockström J, Cornell SE, Fetzer I, Bennett EM, et al. Planetary 1164 boundaries: Guiding human development on a changing planet. Science 2015; 347. 1165 Steinbauer MJ, Grytnes J-A, Jurasinski G, Kulonen A, Lenoir J, Pauli H, et al. Accelerated 1166 increase in plant species richness on mountain summits is linked to warming. Nature 1167 2018; 556: 231-234.

- Stephan R, Erfurt M, Terzi S, Žun M, Kristan B, Haslinger K, et al. An inventory of Alpine drought impact reports to explore past droughts in a mountain region. Nat. Hazards Earth Syst.
 Sci. 2021; 21: 2485-2501.
- 1171 Stephens SL, Collins BM, Fettig CJ, Finney MA, Hoffman CM, Knapp EE, et al. Drought, Tree 1172 Mortality, and Wildfire in Forests Adapted to Frequent Fire. BioScience 2018; 68: 77-88.
- Sultan B, Labadi K, Guégan J-F, Janicot S. Climate drives the meningitis epidemics onset in
 West Africa. PLoS medicine 2005; 2: e6.
- Swanson F, Kratz T, Caine N, Woodmansee R. Landform effects on ecosystem patterns and processes. BioScience 1988; 38: 92-98.
- Tasser E, Tappeiner U. Impact of land use changes on mountain vegetation. Applied Vegetation
 Science 2002; 5: 173-184.
- Tegen I, Werner M, Harrison S, Kohfeld K. Relative importance of climate and land use in
 determining present and future global soil dust emission. Geophysical Research Letters
 2004; 31: L05105.
- Temme AJ. Using climber's guidebooks to assess rock fall patterns over large spatial and
 decadal temporal scales: an example from the swiss alps. Geografiska Annaler: Series
 A, Physical Geography 2015; 97: 793-807.
- Tiberti R, Bogliani G, Brighenti S, Iacobuzio R, Liautaud K, Rolla M, et al. Recovery of high
 mountain Alpine lakes after the eradication of introduced brook trout Salvelinus fontinalis
 using non-chemical methods. Biological invasions 2019; 21: 875-894.
- 1188Tiberti R, von Hardenberg A, Bogliani G. Ecological impact of introduced fish in high altitude1189lakes: a case of study from the European Alps. Hydrobiologia 2014; 724: 1-19.
- Tickner D, Opperman JJ, Abell R, Acreman M, Arthington AH, Bunn SE, et al. Bending the curve of global freshwater biodiversity loss: an emergency recovery plan. BioScience 2020; 70: 330-342.
- 1193 Trisos CH, Merow C, Pigot AL. The projected timing of abrupt ecological disruption from climate 1194 change. Nature 2020; 580: 496-501.
- 1195Turner B, Clark W, Kates R, Richards J, Mathews J. The earth as transformed by human action:1196global and regional changes in the biosphere over the past 300 years.1990.
- United Nations Environment Programme A Scientific Assessment of the Third Pole Environment,
 Nairobi, 2022.
- Urrutia R, Vuille M. Climate change projections for the tropical Andes using a regional climate
 model: Temperature and precipitation simulations for the end of the 21st century. Journal
 of Geophysical Research: Atmospheres 2009; 114.
- Vadeboncoeur Y, Peterson G, Vander Zanden MJ, Kalff J. Benthic algal production across lake
 size gradients: interactions among morphometry, nutrients, and light. Ecology 2008; 89:
 2542-2552.
- Valbuena-Ureña E, Oromi N, Soler-Membrives A, Carranza S, Amat F, Camarasa S, et al. Jailed
 in the mountains: Genetic diversity and structure of an endemic newt species across the
 Pyrenees. PLOS ONE 2018; 13: e0200214.
- 1208 Valdivia Ć, Thibeault J, Gilles JL, García M, Seth A. Climate trends and projections for the 1209 Andean Altiplano and strategies for adaptation. Adv. Geosci. 2013; 33: 69-77.
- van Wilgen BW. Evidence, Perceptions, and Trade-offs Associated with Invasive Alien Plant
 Control in the Table Mountain National Park, South Africa. Ecology and Society 2012; 17.
- van Wilgen BW, Richardson DM. Three centuries of managing introduced conifers in South
 Africa: Benefits, impacts, changing perceptions and conflict resolution. Journal of
 Environmental Management 2012; 106: 56-68.
- Ventura M, Tiberti R, Buchaca T, Buñay D, Sabás I, Miró A. Why should we preserve fishless
 high mountain lakes? In: Catalan J, Ninot JM, Aniz M, editors. High mountain
 conservation in a changing world. Springer, Cham, 2017, pp. 181-205.

- Viviroli D, Dürr HH, Messerli B, Meybeck M, Weingartner R. Mountains of the world, water
 towers for humanity: Typology, mapping, and global significance. Water Resources
 Research 2007; 43.
- 1221 Viviroli D, Kummu M, Meybeck M, Kallio M, Wada Y. Increasing dependence of lowland 1222 populations on mountain water resources. Nature Sustainability 2020; 3: 917-928.
- Vuille M, Carey M, Huggel C, Buytaert W, Rabatel A, Jacobsen D, et al. Rapid decline of snow
 and ice in the tropical Andes–Impacts, uncertainties and challenges ahead. Earth Science Reviews 2018; 176: 195-213.
- Wania F, Mackay D. Global fractionation and cold condensation of low volatility organochlorine
 compounds in polar regions. Ambio 1993; 22: 10-18.
- Watermann LY, Hock M, Blake C, Erfmeier A. Plant invasion into high elevations implies
 adaptation to high UV-B environments: a multi-species experiment. Biological Invasions
 2020; 22: 1203-1218.
- Werner CM, Harrison SP, Safford HD, Bohlman GN, Serata R. Extreme pre-fire drought
 decreases shrub regeneration on fertile soils. Ecological Applications 2022; 32: e02464.
- 1233 Wester P, Mishra A, Mukherji A, Shrestha AB. The Hindu Kush Himalaya assessment: 1234 mountains, climate change, sustainability and people: Springer Nature, 2019.
- Yan D, Zhang T, Su J, Zhao L-L, Wang H, Fang X-M, et al. Diversity and composition of airborne
 fungal community associated with particulate matters in Beijing during haze and non haze days. Frontiers in microbiology 2016; 7: 487.
- Yang H, Engstrom DR, Rose NL. Recent changes in atmospheric mercury deposition recorded
 in the sediments of remote equatorial lakes in the Rwenzori Mountains, Uganda.
 Environmental science & technology 2010; 44: 6570-6575.
- Zanchett G, Oliveira-Filho EC. Cyanobacteria and cyanotoxins: from impacts on aquatic
 ecosystems and human health to anticarcinogenic effects. Toxins 2013; 5: 1896-1917.
- Zemp M, Frey H, Gärtner-Roer I, Nussbaumer SU, Hoelzle M, Paul F, et al. Historically
 unprecedented global glacier decline in the early 21st century. Journal of glaciology
 2015; 61: 745-762.
- Zhang J-T, Xiang C, Li M. Effects of Tourism and Topography on Vegetation Diversity in the
 Subalpine Meadows of the Dongling Mountains of Beijing, China. Environmental
 Management 2012; 49: 403-411.
- Zuecco G, Penna D, Borga M. Runoff generation in mountain catchments: long-term
 hydrological monitoring in the Rio Vauz Catchment, Italy. Cuadernos de investigación
 geográfica/Geographical Research Letters 2018: 397-428.
- 1252
- 1253
- 1254
- 1255
- 1255
- 1256