

Fate of polychlorobiphenyls in the insect Tenebrio molitor: consequences for further use as food and feed.

Christelle Planche, Frédéric Mercier, Johanna Rivas, Magaly Angénieux, Hanli Wang, Benjamin Calmont, Sébastien Crépieux, jérémy ratel, Erwan Engel

▶ To cite this version:

Christelle Planche, Frédéric Mercier, Johanna Rivas, Magaly Angénieux, Hanli Wang, et al.. Fate of polychlorobiphenyls in the insect Tenebrio molitor: consequences for further use as food and feed.. 10. International Symposium on Recent Advances in Food Analysis, Sep 2022, Prague, Czech Republic. 81 (OCE2), pp.E71; PII S0029665122000945, 10.1017/S0029665122000945. hal-03795507

HAL Id: hal-03795507 https://hal.science/hal-03795507v1

Submitted on 4 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.







September 6–9, 2022 • Prague, Czech Republic

Liberté Égalité Fraternité

C. Planche¹, F. Mercier¹, J. Rivas¹, M. Angénieux¹, H. Wang¹, B. Calmont², S. Crépieux³, J. Ratel¹, E. Engel¹

¹MASS group, UR QuaPA, INRAE, F-63122 Saint-Genès-Champanelle, France ²SHNAO, Société d'histoire naturelle Alcide-d'Orbigny, F-63170 Aubière, France ³Invers, F-63720 Saint-Ignat, France

Fate of polychlorobiphenyls in the insect Tenebrio molitor: consequences for further use as food and feed

Background



There is a growing demand for animal protein while the resources used for animal feeding are limited or may raise environmental issues. Thus, there is a need to find alternative and sustainable protein sources. Edible insects may be an interesting solution with a good nutritional value [1]. Learning form the BSE prion crisis, it is essential to ensure the safety of insect products and the chemical contamination is a crucial issue [2]. Among priority contaminants, persistent organic pollutants (POPs) such as polychlorobiphenyls (PCBs), dioxins (PCDD/Fs) or polycyclic aromatic hydrocarbon (PAHs) are of serious concern in food supply [3]. In this context, this study explored the ability of yellow mealworm (Tenebrio molitor), the most widely bred and traded insect species in Europe, to bioaccumulate PCBs from their feeding substrate.

Methods



Results



Analytical method set up









➢ 100°C / 1500psi Internal standard

Clean-up (GPC) S-X3 bioBeads column ➤ DCM ≽ 5mL.min⁻¹





Concentration Redissolution Quantification (GC-µECD) ➢ Hexane Low pressure steam heat ➤ Internal transfer standard ➢ 40°C / 1h

Recovery rates: 98% - 129% ➢ RSD: 10% - 18%

Bioaccumulation factors

Bioaccumulation factor (BAF) = PCB concentration in larvae / PCB concentration in wheat bran



> No significant difference between spiking levels for low-molecular weight PCB congeners

BAF increases with the contamination level for high-molecular weight PCB congeners

Thermo Scientific)



Conclusions

- No significant impact of PCBs on larval growth
- Bioaccumulation factors of up to almost 3
- Importance of monitoring the quality of the substrates used for farming insects as food and feed



Centre

Clermont-Auvergne-Rhône-Alpes

UR370 QuaPA Route de Theix 63122 Saint-Genès-Champanelle Tél. : + 33 (0)4 73 62 45 14 christelle.planche@inrae.fr www6.clermont.inrae.fr/quapa

References

[1] van Huis, A. (2021). Prospects of insects as food and feed. Organic Agriculture, *11(2)*, 301-308.

[2] Meyer, A. M., Meijer, N., et al. (2021). Chemical food safety hazards of insects reared for food and feed. Journal of Insects as Food and Feed, 7(5), 823-831. [3] Guo, W., Pan, B., et al. (2019). Persistent organic pollutants in food: contamination sources, health effects and detection methods. International Journal of Environmental Research and Public Health, 16(22), 4361.