



HAL
open science

Large-cell transformation is an independent poor prognostic factor in Sézary syndrome: analysis of 117 cases

Christophe Bontoux, Adèle de Masson, Nicolas Thonnart, Caroline Ram-Wolff, Flavien Caraguel, Luciana Batista, Sabrina Carpentier, Hélène Moins-Teisserenc, Jacqueline Rivet, Marie-dominique Vignon-Pennamen, et al.

► **To cite this version:**

Christophe Bontoux, Adèle de Masson, Nicolas Thonnart, Caroline Ram-Wolff, Flavien Caraguel, et al.. Large-cell transformation is an independent poor prognostic factor in Sézary syndrome: analysis of 117 cases. *British Journal of Dermatology*, 2022, 10.1111/bjd.21738 . hal-03795726

HAL Id: hal-03795726

<https://hal.science/hal-03795726>

Submitted on 22 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

DERMATOLOGY IS CHANGING

By investigating critical pathways
and exploring innovative solutions
we aspire to change dermatology.
And patients' lives.





Incyte Dermatology



SOLVE
ON.

roflumilast is <7% that of apremilast in Denmark, with generic versions already available in some countries. Furthermore, experimental studies have found roflumilast to be up to 90 times more potent in inhibiting PDE4 isoforms compared with apremilast.⁸ In contrast to biologic therapy, oral roflumilast treatment does not require routine laboratory monitoring, and with proven efficacy in COPD and weight loss commonly reported during treatment, roflumilast may also work directly on associated HS comorbidities.⁵

In the present case, we observed considerable improvements in clinical HS presentation and quality-of-life measures with oral roflumilast therapy. In addition, the patient achieved a 10% weight loss, which may have contributed to the reduction of disease burden.¹ Roflumilast could represent a novel and convenient treatment option for all severity stages of HS as well as associated comorbidities. To expand upon the rather limited treatment options in HS, larger studies investigating the long-term efficacy and safety of oral PDE4 inhibitors are warranted.

Hans Christian Ring ^{1,2}, Alexander Egeberg ²,
Claus Zachariae,¹ Simon F. Thomsen ^{2,3} and
Mette Gyldenløve ¹

¹Department of Dermatology and Allergy, Herlev and Gentofte Hospital, University of Copenhagen, Denmark; ²Department of Dermato-Venereology & Wound Healing Centre, Bispebjerg and Frederiksberg Hospital, University of Copenhagen, Denmark; and ³Department of Biomedical Sciences, University of Copenhagen, Denmark

Email: hans.christian.ring@regionh.dk

Funding sources: none.

Conflicts of interest: the authors declare they have no conflicts of interest.

Data availability: the data that support the findings of this study are available on request from the corresponding author.

References

- 1 Jemec GBE. Clinical practice. Hidradenitis suppurativa. *N Engl J Med* 2012; **366**:158–64.
- 2 Miller IM, Ellervik C, Vinding GR et al. Association of metabolic syndrome and hidradenitis suppurativa. *JAMA Dermatol* 2014; **150**:1273–80.
- 3 Kimball AB, Okun MM, Williams DA et al. Two phase 3 trials of adalimumab for hidradenitis suppurativa. *N Engl J Med* 2016; **375**:422–34.
- 4 Ring HC, Maul JT, Yao Y et al. Drug survival of biologics in patients with hidradenitis suppurativa. *JAMA Dermatol* 2022; **158**:184–8.
- 5 European Medicines Agency. Daxas [roflumilast]. Summary of product characteristics. Available at: https://www.ema.europa.eu/documents/product-information/daxas-epar-product-information_en.pdf (last accessed 6 July 2022).
- 6 Egeberg A, Meteran H, Gyldenløve M, Zachariae C. Complete clearance of severe plaque psoriasis with 24 weeks of oral roflumilast therapy. *Br J Dermatol* 2021; **185**:1251–2.
- 7 Vossen ARJV, van Doorn MBA, van der Zee HH, Prens EP. Apremilast for moderate hidradenitis suppurativa: results of a randomized controlled trial. *J Am Acad Dermatol* 2019; **80**:80–8.

- 8 Walsh N. Comparison of the modes of action of apremilast and roflumilast. PhD thesis, University of Glasgow, 2015. Available at: <http://theses.gla.ac.uk/7427> (last accessed 6 July 2022).

Large-cell transformation is an independent poor prognostic factor in Sézary syndrome: analysis of 117 cases

DOI: 10.1111/bjd.21738

DEAR EDITOR, Sézary syndrome (SS) is a rare cutaneous T-cell lymphoma (CTCL) grouped with mycosis fungoides (MF) in the international classification and staging criteria of CTCL.^{1,2} Large-cell transformation (LCT) has been widely described in MF and associated with reduced overall survival, suggesting the importance of early and sequential histological screening of LCT in MF.³ However, LCT has never been studied and characterized in a large cohort of SS. Another concern is that LCT in MF was defined in the 1980s using diagnostic criteria for LCT in follicular lymphoma.⁴ Although widely used since, the reliability of these criteria has never been specifically studied in SS. Additionally, the presence of large circulating Sézary cells (SCs) based on cytomorphological and flow cytometry analysis was independently associated with poor outcome and might predict LCT occurrence in skin.^{5,6} Nevertheless, the prognostic impact of structure parameters [forward scatter (FSC) and side scatter (SSC)] of circulating cells in cytometry and correlation with LCT remains to be determined in SS.

The main objective of our study was to characterize LCT in SS. All patients with SS diagnosed at Saint-Louis hospital (Paris, France) between 1998 and 2020 according to European Organisation for Research and Treatment of Cancer–World Health Organization criteria were included. The gating strategy of KIR3DL2+ SC among lymphocytes was previously described.⁷ Circulating KIR3DL2+ SC $\geq 200 \text{ mm}^{-3}$ was used to define KIR3DL2-positive status.⁸ For each patient, all skin biopsy samples performed were included. LCT was histologically defined by the presence in the lymphocytes' infiltrate of > 25% or aggregates/nodules of large cells (more than four times the diameter of a small lymphocyte).^{3,4} Haematoxylin-, eosin- and safran-stained slides were then digitized and an analysis using HALO software was performed. All blood samples from patients with flow cytometry data between 2015 and 2020 were included for FSC/SSC analysis. This study received the Institutional Review Committee agreement (LYMPHOTEQ reference: CPP 2019-AO1158-49).

In total, 117 patients were included with a median follow-up of 41 months (interquartile range 1–81). Overall, 6% (six of 100) and 16% (18 of 112) of patients were diagnosed with LCT on skin biopsy samples at diagnosis and during follow-up, respectively. Interobserver reliability between two independent pathologists was excellent [$k = 0.88$; 95% confidence interval (CI) 0.78–0.98]. Considering all skin biopsy

samples at diagnosis, CD30 > 10% and Ki67 > 20% was more frequent in LCT+ samples than in LCT- samples (67% vs. 9%; $P = 0.003$ and 100% vs. 22%; $P = 0.016$, respectively).

We then compared visual histopathological analysis with digital pathology analysis of skin biopsy samples ($n = 189$). Mean cell surface was significantly higher in LCT+ than in LCT- skin biopsy sample images [$27 \mu\text{m}^2$ (SD 3.3) vs. $22 \mu\text{m}^2$ (SD 2.3); $P < 0.001$].

Subsequently, we compared cell size between blood and skin compartments, on 231 blood samples from 112 patients with SS. The maximal mean FSC value of circulating tumour cells in patients who were LCT+ prior to LCT occurrence was not significantly higher than that found in patients who were LCT- [561 273 (SD 99 359) vs. 531 839 (SD 68 040)] ($P = 0.37$). Among patients with a maximum mean FSC value < 600 000 during follow-up, 17% (10 of 60) subsequently presented LCT vs. 14% (one of seven) of patients with a

maximum mean FSC value $\geq 600\,000$ ($P > 0.99$). We obtained similar results for SSC.

Finally, median overall survival was shorter in patients who were LCT+ than in those who were LCT- at diagnosis (35 months vs. 80 months) (HR 9.5, 95% CI 1.9–47.1; $P = 0.006$). In multivariate analysis, age > 60 years (HR 4.46, 95% CI 1.14–17.41; $P = 0.031$), elevated LDH level (HR 2.63, 95% CI 1.03–6.72; $P = 0.044$), CD4 + CD26- circulating cell $\geq 10\,000 \text{ mm}^{-3}$ (HR 3.71, 95% CI 1.23–11.19; $P = 0.020$) and LCT at diagnosis (HR 4.77, 95% CI 1.11–20.4; $P = 0.035$) were independently associated with shorter overall survival (Table 1). During follow-up, median overall survival after LCT occurrence was 21 months, with a 5-year survival of 12% (95% CI 1–38).

In conclusion, our study characterized for the first time LCT in a large homogeneous cohort of patients with SS using histological, digital pathology and flow cytometry

Table 1 Univariate and multivariate survival analysis





Variables	No. of patients with complete data	No. of patients (%)	Median survival, months	Univariate analysis		Multivariate analysis	
				HR (95% CI)	P-values	HR (95% CI)	P-values
Sex	117			1.3 (0.67–2.4)	0.470		
Male		53 (45)	77			–	–
Female		64 (56)	150				
Age, years	117			3.3 (1.7–6.3)	< 0.001	4.46 (1.14–17.41)	0.031
> 60		38 (32)	62				
≤ 60		79 (68)	258				
Stage	103			7.4 (1.3–41.3)	0.022	0.66 (0.16–2.7)	0.562
IVA2		10 (9)	39				
IVA1		93 (91)	80				
LDH	97			2.4 (1.2–5.1)	0.018	2.63 (1.03–6.72)	0.044
Elevated		43 (44)	42				
Normal		54 (56)	92				
Circulating KIR3DL2+ SC mm^{-3}	87			5.9 (1.3–26.2)	0.036	–	–
≥ 10 000		7 (8)	39				
< 10 000		80 (92)	75				
Circulating CD4 + CD26- cells mm^{-3}	83			6.8 (1.7–27.7)	0.007	3.71 (1.23–11.19)	0.020
≥ 10 000		12 (14)	35				
< 10 000		71 (86)	75				
FSC baseline	96			1.6 (0.4–6.4)	0.537	–	–
≥ 600 000		9 (9)	NR				
< 600 000		87 (91)	109				
Large-cell transformation at diagnosis	100			9.5 (1.9–47.1)	0.006	4.77 (1.11–20.4)	0.035
Yes		6 (6)	35				
No		94 (94)	80				
CD30 expression in skin at diagnosis	63			0.8 (0.3–2.2)	0.649	–	–
> 10%		9 (14)	45				
≤ 10%		54 (86)	43				
Ki67 in skin at diagnosis	52			1.7 (0.5–6.3)	0.323	–	–
> 20%		14 (27)	41				
≤ 20%		38 (73)	55				

HR, hazard ratio; NR, not reached; LDH, lactate dehydrogenase; CI, confidence interval; FSC, forward scatter; SC, Sézary cell. P-values < 0.05 are provided in bold [log-rank (univariate) and Cox regression (multivariate) statistical test].

approaches. LCT incidence in SS is lower than for MF. Studies specifically investigating LCT in advanced-stage MF reported variable cumulative incidences ranging from 20% to 55% and large international cohorts have reported a variable incidence of LCT from 4.7% (all stages) to 20% (advanced stages) of MF/SS but without distinguishing the two entities.²

Moreover, we have shown that LCT at diagnosis was an independent unfavourable prognostic factor that could not be predicted by the presence of large circulating tumour cells. In parallel, patients with visually diagnosed LCT on skin biopsy had lymphoid cells with a significantly higher mean surface in digital pathology analysis. The criteria used in MF for LCT diagnosis proved reliable and reproducible.

Thus, histological evaluation of LCT in SS, assisted by digital pathology, is a major means of prognostic staging at diagnosis and during follow-up, allowing the early selection of patients that might benefit from therapeutic changes.

Christophe Bontoux ^{1,3,4}, Adèle de Masson ^{2,3,4},
Nicolas Thonnart,^{3,4} Caroline Ram-Wolff,^{2,3,4}
Flavien Caraguel,⁵ Luciana Batista,⁵ Sabrina Carpentier,⁵
Hélène Moins-Teisserenc,^{4,6,7} Jacqueline Rivet,¹
Marie-Dominique Vignon-Pennamen,¹ Anne Marie-Cardine,^{3,4}
Martine Bagot ^{2,3,4} and Maxime Battistella ^{1,3,4}

¹Department of Pathology, Hôpital Saint-Louis, AP-HP, Paris, France;

²Department of Dermatology, Hôpital Saint-Louis, AP-HP, Paris, France;

³INSERM U976, HIPI, Paris, France; ⁴Université de Paris, Institut de Recherche Saint-Louis, Paris, France; ⁵Innate Pharma, Marseille, France;

⁶Clinical Haematology Laboratory, Hôpital Saint-Louis, AP-HP, Paris, France; and ⁷INSERM U1160, Paris, France

Correspondence: Maxime Battistella.

Email: maxime.battistella@aphp.fr

Funding sources: none.

Conflicts of interest: the authors declare they have no conflicts of interest.

Data availability: the data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

Ethics statement: this study received Institutional Review Committee agreement (LYMPHOTEQ reference: CPP 2019-AO1158-49).

References

- Olsen E, Whittaker S, Willemze R et al. Primary cutaneous lymphoma: recommendations for clinical trial design and staging update from the ISCL, USCLC, and EORTC. *Blood* 2021; DOI: <https://doi.org/10.1182/blood.2021012057>.
- Scarlsbrick J, Prince H, Vermeer M et al. Cutaneous lymphoma international consortium study of outcome in advanced stages of mycosis fungoides and Sézary syndrome: effect of specific

prognostic markers on survival and development of a prognostic model. *J Clin Oncol* 2015; **33**:3766–73.

- Vergier B, de Muret A, Beylot-Barry M et al. Transformation of mycosis fungoides: clinicopathological and prognostic features of 45 cases. *Blood* 2000; **95**:2212–18.
- Salhany KE, Cousar JB, Greer JP et al. Transformation of cutaneous T cell lymphoma to large cell lymphoma. A clinicopathologic and immunologic study. *Am J Pathol* 1988; **132**:265–77.
- Clark RA, Shackelton JB, Watanabe R et al. High-scatter T cells: a reliable biomarker for malignant T cells in cutaneous T-cell lymphoma. *Blood* 2011; **117**:1966–76.
- Vonderheid EC, Jakubowski J, Hou JS. High-scatter lymphocytes in the blood of erythrodermic cutaneous T-cell lymphoma: evidence for large-cell transformation? *Clin Lymphoma Myeloma Leuk* 2020; **20**:624–31.e2.
- Moins-Teisserenc H, Daubord M, Clave E et al. CD158k is a reliable marker for diagnosis of Sézary syndrome and reveals an unprecedented heterogeneity of circulating malignant cells. *J Invest Dermatol* 2015; **135**:247–57.
- Roelens M, de Masson A, Ram-Wolff C et al. Revisiting the initial diagnosis and blood staging of mycosis fungoides and Sézary syndrome with the KIR3DL2 marker. *Br J Dermatol* 2020; **182**:1415–22.

Differences in epidemiology, comorbidities and treatment choice between plaque psoriasis and pustular psoriasis: results from the BIOBADADERM registry

DOI: 10.1111/bjd.21763

DEAR EDITOR, Pustular psoriasis is a group of inflammatory skin conditions characterized by clinically visible sterile pustules. It has been considered as a form of psoriasis vulgaris, but they are phenotypically different, respond differently to treatments and are genetically distinct. Variations in IL36RN, CARD14, APIS3, MPO and SERPINA3 genes have been linked to generalized pustular psoriasis.^{1,2} The European Rare and Severe Psoriasis Expert Network (ERASPEN) recently presented a consensus classification of clinical phenotypes of pustular psoriasis.³ There are limited data on the differences between pustular and plaque psoriasis. Therefore, we aimed to compare the demographic characteristics, comorbidities and prescriptions between these clinical variants in clinical practice.

We used data from the BIOBADADERM registry, a previously described prospective multicentre cohort registry of patients with psoriasis treated with systemic drugs in Spain intended to detect adverse events related to systemic therapy.⁴ Participants enter the cohort when they start a therapy that they have never used before. This study contains analysis of the data extracted from BIOBADADERM from October 2008 to December 2021. Therapy in the first 5 years of disease was described in a subset of patients with disease onset close to cohort entry. Plaque psoriasis (PP) was used as the reference group for all comparisons.

There were 3864 patients with PP, 41 patients with generalized pustular psoriasis (GPP) and 294 patients with