



**HAL**  
open science

## **Sensitive 1,1-dicyanovinyl push-pull dye for primary amine sensing in solution by fluorescence**

Lulu Bu, Maxime Rémond, Pauline Colinet, Erwann Jeanneau, Tangui Le Bahers, Frédéric Chaput, Chantal Andraud, Yann Bretonnière

### ► **To cite this version:**

Lulu Bu, Maxime Rémond, Pauline Colinet, Erwann Jeanneau, Tangui Le Bahers, et al.. Sensitive 1,1-dicyanovinyl push-pull dye for primary amine sensing in solution by fluorescence. *Dyes and Pigments*, 2022, 202, pp.110258. <10.1016/j.dyepig.2022.110258>. <hal-03795976>

**HAL Id: hal-03795976**

**<https://hal.science/hal-03795976v1>**

Submitted on 4 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# Sensitive 1,1-dicyanovinyl push-pull dye for primary amine sensing in solution by fluorescence

Lulu Bu,<sup>a,b</sup> Maxime Rémond,<sup>a</sup> Pauline Colinet,<sup>a</sup> Erwan Jeanneau,<sup>c</sup> Tangui Le Bahers,<sup>a</sup> Frédéric Chaput,<sup>a</sup> Chantal Andraud,<sup>a</sup> Yann Bretonnière<sup>a,\*</sup>

<sup>a</sup> Université de Lyon, École Normale Supérieure de Lyon, Université Claude Bernard Lyon 1, CNRS UMR 5182, Laboratoire de Chimie, 46 allée d'Italie, F69364 Lyon, France

<sup>b</sup> School of Chemistry and Chemical Engineering, Huainan Normal University, Huainan, China

<sup>c</sup> Université de Lyon, Centre de Diffractométrie Henri Longchambon, Université Lyon I, 43 boulevard du 11 Novembre 1918, 69622 Villeurbanne Cedex (France)

\* corresponding author: [yann.bretonniere@ens-lyon.fr](mailto:yann.bretonniere@ens-lyon.fr)

## Abstract

1,1-dicyanovinyl push-pull dye **PTTCN** undergoes rapid and reversible aza-Michael addition with volatile primary aliphatic amines in solution in DMSO, yielding a highly fluorescent imine derivative. This reaction is accompanied by a dramatic colour change as well as a strong increase in fluorescence (1200-fold increase) allowing sensitive turn-on detection of primary amine in solution. In the solid state, **PTTCN** is emissive and reaction with primary amines vapours results in a shift in fluorescence and a possible ratiometric detection of primary amine. The selectivity is excellent for primary amines over secondary, tertiary and aromatic amines as well as over thiols and fluorescence spectroscopy allow to discriminate between primary and other nitrogen nucleophiles such as hydrazine and ammonia. Finally, we show as proof-of-concept that the fluorescence turn-on response can be used in response to the vapours of volatile amines released during food spoilage.

**Keywords:** Primary amine; Fluorescence turn-on; Ratiometric; Fluorescent Thin Film

## 1. Introduction

Aliphatic primary amines are organic compounds omnipresent in the pharmaceutical, cosmetics or agrochemical industry, as well as in biology and it is often necessary to be able to accurately detect trace levels in real time and

28 in different matrices. Some natural primary amines of low-molecular weight, like histamine for instance,  
29 modulate essential metabolic and physiological functions in living organisms, but may also be of major health  
30 concern when present in food. They thus have been identified as important markers for evaluation of food  
31 quality or specific indicators for various diseases or health issues.[1-3]

32 For these reasons, many techniques have been developed so far for the qualitative and quantitative detection of  
33 amines, as diverse as high performance liquid chromatography, capillary electrophoresis, gas chromatography,  
34 or biosensor, to cite a few, each with its pro and cons.[4, 5] Of these, those based on chemical optical probes in  
35 general and photoluminescence (PL) in particular offer definite advantages in terms of rapidity, sensitivity with  
36 demonstrated limits of detection to nM, ease of handling and possibility to be used in solution or solid  
37 supports.[6, 7]

38 As recently summarized in excellent reviews, various strategies exploiting the basic or nucleophilic properties of  
39 amines have been reported to design chemical probes whose optical properties, whether absorption or PL,  
40 change upon exposure to amines.[6] [5, 7, 8]

41 For instance, direct addition of a nucleophilic amine to a chromophore's electron-withdrawing group or  
42 1,2-addition to C=C double bond conjugated with this group heavily modifies the electron delocalisation within  
43 the dye molecule, therefore triggering consequent spectral changes (absorption and/or emission) and directly  
44 inducing a signalling response. Thus, reversible conversion of the trifluoroacetyl group into a hemiaminal  
45 function [9-11] or irreversible conversion of the tricyanovinyl group into the 1-amino-2,2-dicyanovinyl group  
46 [12] results in a decrease of the electron-withdrawing strength and consequently in a blue-shift in the absorbance  
47 or PL spectra. On the other hand, reversible addition of amine on aldehyde- [13-19] or ketone-containing [20]  
48 chromophores gave highly fluorescent imine or iminium derivatives with usually red-shifted optical properties.

49 Dicyanovinyl group is probably one of the simplest, yet powerful, electron-withdrawing groups for this effect.  
50 Aza-Michael nucleophilic addition of amine to the dicyanovinyl group is followed by the loss of a malononitrile  
51 molecule to yield an imine, leading to a large spectral-shift and most often a PL turn-on response. This was  
52 shown by Cheng and He who developed high sensitivity probes for primary amine vapour with polymers  
53 encompassing benzylidenemalononitrile units [21] and by Shi, Dong and co-workers who described the  
54 detection of amine vapours through formation of a bis-imine derivatives with a bis- dicyanovinyl based  
55 probes.[22] Previously, turn-on fluorogenic probe for detecting MDMA from ecstasy tablets [23] and dual-mode  
56 detection of aliphatic amines and hydrazine by PL [24] were also reported built upon a bis-diarylurea-

57 dicyanovinyl push-pull fluorophore and a dicyanovinyl-substituted oligothiophene derivative associated with  
58 zinc tetraphenylporphyrin, respectively. Mohr *et al.* also described a colorimetric read-out sensor based on  
59 push-pull azo dye and therefore working at wavelengths compatible with low-cost light sources and detectors.[25,  
60 26] However in all these works, the end product was not characterized or simply postulated as being the 1,2  
61 adduct.[26] Sathiyarayanan *et al.* reported the design of a ratiometric PL sensor for primary amine based on a  
62 dicyanovinyl substituted phenanthridine derivatives and they clearly identified the end product as being the  
63 corresponding imine.[27] Finally, in 2020, Burn, Show and coworkers reported two dicyanovinyl-  
64 fluorene-benzothiadiazole-based fluorescent compounds, which showed rapid response to a wide range of  
65 amines. Primary amines react with the dicyanovinyl by aza-Michael (1,2) addition to give the corresponding  
66 imine, while PL quenching *via* photoinduced hole transfer was observed in the solid-state with other amines.[28]

67 It should be noted, however, that nucleophilic addition on dicyanovinyl based chromophores is not specific to  
68 primary amines. Hydrazine indeed yields hydrazone, a much weaker electron-withdrawing group than  
69 dicyanovinyl,[29] whereas thiols and cyanide give the 1,2-addition product. Here also the modification or the  
70 breaking of the dye  $\pi$ -conjugated system leads to considerable changes in the optical properties (blue shift and  
71 generally PL enhancement). Extremely sensitive and “selective” colorimetric and/or fluorescent sensors to  
72 hydrazine,[30-34] cyanide,[30, 35-42] and thiols [43, 44] [36, 45] are reported. In most cases, no mention is  
73 made of possible interferences from competitive amines.[25]

74 We have identified **2-((5'-(piperidin-1-yl)-[2,2'-bithiophen]-5-yl)methylene)malononitrile (PTTCN,**  
75 Scheme 1), a 1,1-dicyanovinyl push-pull dye, that undergoes rapid reaction with primary amine to yield  
76 fluorescent imine with emission intensities as high as 1200 times greater than the starting materials. We want to  
77 present here preliminary results about its use as fluorescent probe for primary amine detection in solution and in  
78 the solid state.

79 Compound **PTTCN** and other analogues, in which the 1,1-dicyanovinyl group is connected to a strong donor  
80 2-(*N*-dialkyl)-amino- or 2-(*N*-diaryl)-amino- across a 2,2'-bithiophene  $\pi$ -conjugated bridge conferring moderate  
81 aromatic stabilization to the donor-acceptor system and therefore a high degree of conjugation between the donor  
82 and acceptor end groups, were originally developed as highly polarizable chromophores for 2<sup>nd</sup> order nonlinear  
83 optics.[46, 47] [48, 49] The spectroscopic properties were later studied in detail and are characteristic of such  
84 donor-acceptor substituted heterocyclic compounds with a strong intramolecular charge transfer (ICT)  
85 absorption band in the visible region of the spectrum, and a weak red PL that decreases in intensity as the solvent

86 polarity increases.[50, 51] The PL also increased with the viscosity, making this compound and other analogues  
87 potential viscosity probes [52] and explaining the solid-state emission we observed. On the other hand, very few  
88 examples of imine derivatives built on the 2-(*N*-dialkylamino or diarylamino)-2,2'-bithiophene pattern are  
89 described,[53, 54] mostly chiral bichromophoric systems developed for CD exciton chirality studies or for  
90 second-order NLO applications. Bereva and coworkers, nevertheless, highlighted the intense blue-green PL of  
91 *N*-[[5'-(Dimethylamino)[2,2'-bithiophen]-5-yl]methylene]-cyclohexylamine and  
92 *N*-[[5'-(1-Pyrrolidinyl)[2,2'-bithiophen]-5-yl]methylene]-cyclohexylamine.[53]

93

## 94 **2 Experimental**

### 95 **2.1 Materials and Methods**

96 All solvents and reagents were commercially available and used without further purification. Column  
97 chromatography was carried out on silica gel (35-70  $\mu\text{m}$ ).  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded at ambient  
98 temperature on a Bruker Advance 400 spectrometer operating at 400.0 MHz for proton and 101.00 MHz for  
99 carbon, respectively. Chemical shifts are reported in parts per million ( $\delta/\text{ppm}$ ) relative to tetramethylsilane with  
100 the residual solvent peaks as internal standard (2.50 and 39.5 ppm for DMSO). High-resolution mass  
101 spectrometry (HRMS) were measured by ESI-TOF (Bruker Daltonics Micro TOF-Q II) at the *Centre Commun*  
102 *de Spectrométrie de Masse* (UCBL, Villeurbanne, France). UV/Vis absorption spectra were measured using a  
103 dual beam Jasco 670 spectrometer. Fluorescence spectra in diluted solution and in thin film were recorded on a  
104 Horiba Jobin-Yvon Fluorolog-3® spectrofluorimeter equipped with red-sensitive Hamamatsu R928 PMT.  
105 Spectra were reference corrected for both the excitation source light intensity variation (lamp and grating) and  
106 the emission spectral response (detector and grating).

107

### 108 **2.2 Synthesis**

#### 109 **2.2.1 Synthesis of 2-((5'-(piperidin-1-yl)-[2,2'-bithiophen]-5-yl)methylene)malononitrile (PTTCN)**

110 In a 20 mL microwave tube, 5-formyl-5'-piperidino-2,2'-bithiophene (**1**) (300 mg, 1.08 mmol) and  
111 malononitrile (85 mg, 1.28 mmol) were dissolved in anhydrous EtOH (5 mL), followed by addition of one drop  
112 of piperidine. The tube was sealed and heated at 100°C by microwave irradiation for 1 h. After cooling down to  
113 room temperature, the solid was filtrated and purified by column chromatography on silica (eluent:  $\text{CH}_2\text{Cl}_2$ ) to

114 afford **PTTCN** as a dark purple solid (244 mg, yield: 69%). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>, ppm) δ 8.38 (s, 1H),  
115 7.77 (d, <sup>3</sup>*J* = 4.3 Hz, 1H), 7.50 (d, <sup>3</sup>*J* = 4.3 Hz, 1H), 7.29 (d, <sup>3</sup>*J* = 4.3 Hz, 1H), 6.27 (d, <sup>3</sup>*J* = 4.3 Hz, 1H), 3.29 (m,  
116 4H), 1.61 (m, 4H), 1.82-1.43 (m, 6H). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>, ppm) δ 162.9, 151.4, 150.7, 143.6, 131.0,  
117 129.8, 121.5, 117.5, 115.7, 114.9, 105.6, 68.6, 50.9, 24.5, 23.0. HR-MS (ESI-QTOF) *m/z*: [M+H]<sup>+</sup> calcd for  
118 C<sub>17</sub>H<sub>16</sub>N<sub>3</sub>S<sub>2</sub>, 326.0786; found, 326.0770.

### 119 **2.2.2 Synthesis of (E)-1-(5'-(piperidin-1-yl)-[2,2'-bithiophen]-5-yl)-N-propylmethanimine (PTT-imine)**

120 Under an inert atmosphere, **1** (50.0 mg, 0.18 mmol) and n-propylamine (10.6 mg, 0.18 mmol) were dissolved  
121 in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) and 4 Å molecular sieve (MS) was added. The mixture was heated to reflux for 2 h. MS was  
122 filtered off. Evaporation under reduced pressure to dryness afforded **PTT-imine** as a yellow solid sufficiently  
123 pure to be used without further purification (57 mg, yield: 99%). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>, ppm) δ 8.33 (s,  
124 1H), 7.28 (d, *J* = 3.8 Hz, 1H), 7.06 (d, *J* = 4.0 Hz, 1H), 6.98 (d, *J* = 3.8 Hz, 1H), 6.08 (d, *J* = 4.0 Hz, 1H), 3.44 (t,  
125 *J* = 6.7 Hz, 2H), 3.23-2.99 (m, 4H), 1.80-1.35 (m, 8H), 0.87 (t, *J* = 7.4 Hz, 3H). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>,  
126 ppm) δ 159.5, 154.1, 140.7, 138.4, 132.1, 125.2, 120.8, 120.6, 104.6, 61.8, 51.3, 24.6, 23.7, 23.2, 11.7. HR-MS  
127 (ESI-QTOF) *m/z*: [M+H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>22</sub>N<sub>2</sub>S<sub>2</sub>, 319.1297; found, 319.1295.

### 128 **2.2.3 Preparation of sol-gel film**

129 Silica-based sol preparation from methyltriethoxysilane (MTEOS): In a 2 L flask, citric acid (3.96 g, 0.02  
130 mole) was dissolved in DI water (396 mL, 22 moles). Under vigorous stirring, MTEOS (400 mL, 2 moles) was  
131 added to the aqueous solution. With the formation of an emulsion, the medium quickly became opaque and then  
132 increasingly transparent as the ethanol was released following the hydrolysis of MTEOS. The resulting mixture  
133 was heated to 45°C overnight under continuous slow stirring. The ethanol produced by the hydrolysis reaction of  
134 the alkoxy silane was subsequently removed from the reaction solution at reduced pressure. The silica-based sol  
135 was then extracted by adding ether which caused phase separation. The aqueous phase containing the citric acid  
136 was then separated. In order to completely remove citric acid from the sol, the ethereal phase was washed several  
137 times by adding water. The ether is then removed in vacuo and the resulting sol is dissolved in THF while  
138 controlling the solid content (54% by mass).

139 Preparation of doped sol-gel film: **PTTCN** (2.0 mg) dissolved in THF (1 mL) was added to the previously  
140 prepared MTEOS sol (2 g, 54 wt% in THF). Once the mixing has been carried out, part of the THF (1 mL) is  
141 removed under vacuum in order to obtain a viscous doped sol. The latter was spread evenly on a glass substrate  
142 by centrifugation at 2000 rpm. The sol-gel film was further dried by heating at 90°C for 1 h in an oven.

143

#### 144 **2.2.4 Preparation of PMMA thin film**

145 A solution of **PTTCN** and **PMMA** in dichloromethane (1 mg dye for 100 mg PMMA in 1 mL) was spread  
146 evenly on a glass surface by spin-coating at 2000 rpm using a Karl Suss CT 62 spin-coater.

147

### 148 **2.3 Computational method**


149 All calculations were performed using Gaussian package.[55] At the ground state, the structure of all the  
150 compounds was optimized at the DFT/6-311G(d) level of theory using hybrid functionals PBE0.[56] Bulk  
151 solvent effects (DMSO) were considered using continuum solvation model (C-PCM).[57, 58] For each  
152 compound, 10 vertical excitation energies and associated excited-state densities of interest were computed at the  
153 TD-DFT level with the same basis set and CAM-B3LYP functional.[59] The charge transfer transitions were  
154 characterized thanks to the DCT index developed by Le Bahers et al. using the cube files generated by the  
155 cubegen utility provided by the Gaussian package.[60] To simulate the colour, the trichromatic components were  
156 obtained from the transmission spectra through a code developed in our group and converted into RGB ones.

157

## 158 **3 Results and discussion**

### 159 **3.1 PTTCN and PTT-imine synthesis and characterization**

160



161

162 **Scheme 1.** Synthesis of **PTTCN** and **PTT-imine**.

163

164 **PTTCN** was obtained as a dark violet powder from malononitrile and aldehyde **1** by Knoevenagel condensation.

165 **1** was prepared in five steps from thiophene, succinic anhydride and piperidine by construction of the second

166 thiophene ring by a combination of Friedel-Craft and Lawesson reactions,[61] followed by formylation by

167  $\alpha$ -lithiation and reaction with DMF (Scheme 1). This approach was preferred to classical bithiophene synthesis

168 using palladium catalysed coupling reaction as it gives higher yield. **PTT-imine** was in turn easily obtained from

169 **1** by reaction with n-propylamine in presence of 4 Å MS.

170 Crystals of **PTTCN**, suitable for X-ray diffraction, were grown by slow diffusion of EtOH in concentrated dye

171 solution in  $\text{CHCl}_3$ . Crystallographic data, basic structural parameters and the X-ray crystal structure are given in

172 SI. The latter showed a quasi-planar  $\pi$ -delocalized system with twenty-two atoms forming the molecular mean

173 plane (maximum deviation of 0.114 Å). Only the dicyanovinylene group is slightly twisted forming a 5.7° angle

174 with the bis-thiophene entity. This is indicative of a full conjugation between the donor and the acceptor end of

175 the push-pull molecule.

176 In solution, the electronic absorption spectrum of **PTTCN** is characteristic of intramolecular charge transfer

177 (ICT) transitions in push-pull dipolar molecules with a broad absorption in the visible ( $\lambda_{\text{abs}} = 575 \text{ nm}$  in DMSO,


178 in good agreement with the calculated value of 509 nm, cam-B3LYP/6-311G(d)/PCM(DMSO), large molar

179 absorption coefficient values, and a small bathochromic shift in the maximum as the solvent polarity increases  
180 (Figure SI 2). Upon excitation in the main absorption band PL is observed, whose peak maximum is gradually  
181 red-shifted and decreases in intensity as the solvent polarity increases from toluene ( $\lambda_{em} = 640$  nm,  $\phi = 18$  %) to  
182 DMSO where it almost vanishes ( $\lambda_{em} = 695$  nm,  $\phi < 1$  %). **PTTCN** also showed interesting far-red emission in  
183 the solid state ( $\lambda_{em} = 665$  nm,  $\phi = 24$  %) easily visualized under illumination of a handheld UV lamp at 365 nm.

### 184 **3.2 Sensing properties in solution**

185 The reaction **PTTCN** with primary amine was first investigated by  $^1\text{H-NMR}$ . Stepwise addition of  
186 n-propylamine to a solution of **PTTCN** in  $\text{-DMSO-}d_6$  immediately yielded the formation of a single new product  
187 that concurs with colour change from violet to yellow along with appearing of an intense green PL that can be  
188 seen under UV light. The  $^1\text{H-NMR}$  spectrum (Figure 1) of the new product displayed similar splitting patterns  
189 but significant upfield shifts for aromatic protons and the persistence of a vinylidene singlet integrating for one  
190 proton at 8.33 ppm. This is in agreement with an imine structure in which the push-pull character is preserved  
191 but a weaker electron-acceptor group is present. A simple nucleophilic reaction at the  $\text{C}\beta$  position, as assumed by  
192 Finšgar *et al.* for similar dicyanovinyl azobenzene dyes [26] or as is observed with thiols [25] and the cyanide  
193 anion,[35, 42] would lead to a drastic upfield shift of the proton originally at the  $\beta$ -position as well as those of  
194 the closest thiophene ring due to a disruption of the  $\pi$ -conjugation from the donor to the acceptor end. The imine  
195 structure was further confirmed by mass spectrometry and the independent synthesis of **PTT-imine** from  
196 aldehyde **1** and n-propylamine (Scheme 1), whose  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra (Figure SI 15 and SI 16) perfectly  
197 match those of the new product.

198




199  
200 **Figure 1.** Aromatic region of the  $^1\text{H}$ -NMR spectra (300 MHz, DMSO- $d_6$ ) of **PTTCN** prior (a) and immediately  
201 after addition of 0.6 equiv. (b), 1.2 equiv. (c), 2.5 equiv. (d), and 4 equiv. (e) of n-propylamine.

202  
203 More interestingly, in the case of a volatile amine having low boiling point such as n-propylamine,  $^1\text{H}$  NMR also  
204 evidences the reverse transformation of imine to the initial dicyanovinyl (Figure SI 4). Removal of the excess  
205 amine by evaporation under vacuum induced a colour change back to the original purple of dye **PTTCN** as well  
206 as a disappearance of the PL and the reappearing of the NMR signal of **PTTCN**. Note, however, that addition of  
207 excess malononitrile to a solution of **PTT-imine** in DMSO- $d_6$  immediately resulted in a colour change from  
208 yellow-green to dark red then purple, but the signals obtained are broad and the reformation of **PTTCN** could  
209 not be unambiguously demonstrated.

210 Spectroscopic studies also evidenced the interest of **PTTCN** as a primary amine fluorescent probe. The changes  
 211 in the absorption and PL spectra on addition of n-octylamine (or n-propylamine) to solution of **PTTCN** in  
 212 DMSO (10  $\mu\text{M}$ ) are shown in Figure 2. With increasing amount of amine added, the characteristic ICT band at  
 213 575 nm disappeared while the colour of the dye solution turns yellow with the concomitant appearing of a new  
 214 peak with a maximum absorbance at 412 nm. The isobestic point at 462 nm indicates that a single chemical  
 215 reaction takes place. Kinetic studies revealed that the reaction is first order with respect to both reactants primary  
 216 amine and **PTTCN** (Figures SI 5 and SI 6).

217



218

219 **Figure 2.** Spectral changes for a solution of **PTTCN** (10  $\mu\text{M}$ ) in DMSO immediately after addition of increasing  
 220 amount of n-octylamine (0 to 140  $\mu\text{M}$ ): (a) absorption and (b) emission spectra ( $\lambda_{\text{ex}} = 462$  nm, isobestic point of  
 221 the absorption spectra). Inset: solution of **PTTCN** without (left) and with (right) excess octylamine (140  $\mu\text{M}$ ),  
 222 daylight and under UV-light (365 nm).


223

224 At the same time, an intense emission appeared at 535 nm that steadily increased while adding amine. At this  
 225 high dye concentration (10  $\mu\text{M}$ ) the reaction is fast and the fluorescence is therefore seen immediately. The  
 226 quantum yield of the newly formed product reaches unity ( $\phi = 0.96$ ) whereas **PTTCN** itself is barely emissive in  
 227 DMSO, a “turn-on” PL response (1160-fold emission intensity enhancement, Figure 2b) is then achieved. As  
 228 shown is Figure SI 7, a theoretical limit of detection of n-octylamine was estimated to be as low as 60 nM for a  
 229 dye concentration of 5  $\mu\text{M}$ . Note however that for such low concentration in amine, the reaction is much slower  
 230 and the measure has to be performed 15 hours after amine addition.

231 The dramatic colour change observed when switching from dicyonovinyl to imine is clearly explained by a

232 decrease in intramolecular charge transfer and in the push-pull character. This is confirmed by DFT and TD-DFT  
233 calculations on **PTTCN** and **PTT-imine**.

234 Calculations reproduced the modification of the absorption properties towards the blue range could also be  
235 reproduced computationally (Figure 3-a) with a shift of the wavelength of maximum absorption from 509 nm  
236 (0.28 eV difference compared to experimental value) for **PTTCN** to 383 nm (0.15 eV difference compared to  
237 experimental value) for the **PTT-imine**. A fluorescence maximum at 505 nm is computed in good agreement  
238 with what is observed (0.14 eV difference). The computed distance characteristic of the charge transfers under  
239 absorption decreases from 3.41 Å to 2.95 Å from **PTTCN** to **PTT-imine** as illustrated on Figure 3-b, in good  
240 agreement with a less efficient charge transfer.



241  
242 **Figure 3.** b) Simulated absorption spectra (at CAM-B3LYP/6-311G(d) level of theory) of the two molecules in  
243 DMSO along with their simulated colour (top insert). b) Representation of the difference between excited- and  
244 ground-state electron densities (cyan, density increase upon transition; yellow, density decrease; isovalue of .0005)  
245 as well as the CT distance (in grey) for PTTCN (top) and PTT-imine (bottom) molecules. Carbon, hydrogen, sulfur  
246 and nitrogen atoms are depicted respectively by brown, white, yellow and blue spheres.


247

### 248 3.3 Selectivity

249 Qualitative PL-sensing studies were performed with various aliphatic amines (ammonia, putrescine, tyramine,  
250 cadaverine, propylamine, histamine, tryptamine, spermidine, 1,3-diamine, diethylamine, diisopropylamine,  
251 trimethylamine), aromatic amine (aniline, p-anisidine, *N,N*-dimethyl-p-phenylenediamine) and other nucleophile  
252 (ethanethiol, hydrazine). Addition of secondary amine (diethylamine, diisopropylamine), tertiary amine


253 (trimethylamine) or thiol (ethanethiol) did not induce any colour change nor any significant change in the PL  
 254 spectra of **PTTCN** in DMSO solution (Figure 4-a). The same applies when an aromatic amine is added,  
 255 including electron enriched p-anisidine and N,N-dimethyl-p-phenylenediamine. However, when a primary amine  
 256 (here n-propylamine, 50 equiv.) was added after addition of 50 equivalent of diethylamine, trimethylamine, or  
 257 thiol, the colour immediately changed and the strong PL characteristic of the imine formation is immediately  
 258 observed highlighting the selectivity for the primary amine and the absence of interference from secondary  
 259 analytes (Figure 4-b).

260 On the other hands, **PTTCN** has fast PL-response to ammonia and hydrazine compounds (hydrazine or  
 261 dimethylhydrazine). However, the new products obtained in these cases display slightly blue-shifted absorption  
 262 and emission compared to the imine for hydrazine and red-shifted for other analytes, and are less emissive  
 263 (Figure 5, Table 1), allowing to distinguish them.



264  
 265 **Figure 4.** PL intensity at 535 nm ( $\lambda_{exc} = 460$  nm) of 10  $\mu$ M **PTTCN** solution in DMSO with: a) different amines  
 266 and nucleophiles (50 equiv); b) different amines and nucleophiles (50 equiv) then n-octylamine (50 equiv), in  
 267 black the initial PL intensity before adding n-octylamine.

268



269

270 **Figure 5.** Absorption (dotted lines) and emission spectra (solid lines) of PTTCN upon addition of excess of  
 271 ammonia, hydrazine or dimethylhydrazine (10  $\mu$ M in DMSO,  $\lambda_{exc} = 400$  nm).

272

273 **Table 1.** Spectroscopic data for PTTCN and for PTTCN with addition of various N nucleophiles.

	PTTCN			+ n-propylamine			+ Ammonia			+ Hydrazine			+ N,N-dimethylhydr.		
	$\lambda_{abs}$ (nm)	$\lambda_{em}$ (nm)	$\Phi$	$\lambda_{abs}$ (nm)	$\lambda_{em}$ (nm)	$\Phi$	$\lambda_{abs}$ (nm)	$\lambda_{em}$ (nm)	$\Phi$	$\lambda_{abs}$ (nm)	$\lambda_{em}$ (nm)	$\Phi$	$\lambda_{abs}$ (nm)	$\lambda_{em}$ (nm)	$\Phi$
DMSO	575	695	< 0.01	412	535	0.96	413	543	0.58	400	516	0.30	419	541	0.03
Film (sol-gel)	566	662	23	403	506	0.43	410	543	0.26	398	514	0.15	418	535	0.01
Film (PMMA)	553	656	13	404	499	0.57	nd	nd	nd	nd	nd	nd	nd	nd	nd

274

### 275 3.4 Primary amine vapour sensing in thin-film


276 Having established PTTCN as a robust PL turn-on and reversible probe to detect primary amines in solution, we  
 277 then investigated primary amine vapour detection by thin films of the sensing material. For this purpose,  
 278 optically thin films of PTTCN in sol/gel or PMMA were obtained on glass substrate by spin-coating. The violet  
 279 films displayed large absorption band at 566 nm and 553 nm respectively and showed strong far-red PL upon  
 280 excitation at the absorption peak ( $\lambda_{em} = 661$  nm /  $\phi = 23\%$  and  $\lambda_{em} = 656$  nm /  $\phi = 13\%$ , for sol-gel and PMMA  
 281 films resp., Table 1). We then choose n-propylamine as an example for its high volatility, to test the film upon  
 282 exposure to amine vapours. Exposing the sol-gel films to an atmosphere saturated with n-propylamine vapours

283 induced colour change from violet to almost colourless ( $\lambda_{\text{max}} = 402 \text{ nm}$ ) with complete transformation in less  
284 than 5 minutes. For PMMA films, the same colour change is observed ( $\lambda_{\text{max}} = 403 \text{ nm}$ ) upon exposure to amine  
285 vapour, but total conversion takes more than 20 min. The faster kinetic observed in sol-gel is probably due to the  
286 more porous material in this case and therefore the greater accessibility of the dye. In the same time, for both  
287 kind of films, a significant variation of emission from red to green was observed by naked eye when the films are  
288 illuminated under 365 nm UV-light (Figure 6, see also SI). Upon excitation in the absorption peak strong green  
289 PL signals were observed for the sol-gel and the PMMA films after exposure with amine, with maxima at  
290 508 nm and 497 nm respectively. Good emission quantum yields were measured, respectively  $\Phi = 57 \%$  and  $\Phi =$   
291 43 %. A ratiometric PL response to primary amine vapour is therefore possible when choosing the excitation  
292 wavelength at 455 nm, wavelength at which the absorption of both **PTTCN** and imine are identical.

293 Similar colour changes were observed with exposure to ammonia or dimethylhydrazine vapours (Figure SI 8).  
294 PLQY quantum yields in solid reflect what was observed in solution with weaker values for ammonia and  
295 dimethylhydrazine than for n-propylamine. Always in agreement to what was observed in solution, no  
296 fluorometric nor colourmetric responses were observed, even after 1 h, when the films were exposed to vapours  
297 of secondary amine, tertiary amine, ether, ester, amide, alcohol, or thiol.

298 Unfortunately, the reversibility highlighted in solution could not easily be reproduced. Vacuum pumping of the  
299 film exposed to propylamine vapour partially converts the film to its original state, but pumping for too long  
300 results in severe flaking-off the films. The same observation was made when fuming malononitrile vapours after  
301 amine exposure. The colour of the films reverted back to purple and the original emission partially recovered,  
302 but the sample were damaged.

303




304

305 **Figure 6.** Normalized absorption (dotted line) and PL spectra (solid line) of **PTTCN** sol-gel film before (red) and  
 306 after (green) exposure to n-propylamine vapour. Inset: the photography showed the film mounted in the support  
 307 before (right) and after (left) exposure to n-propylamine of **PTTCN** in the absence (black) and presence (red) of  
 308 saturated propylamine vapor under daylight or UV-light (365 nm).

309

310 **3.5 Tentative of monitoring seafood freshness**



311  
 312 **Figure 7.** Attempt to control shrimp freshness, followed by a week: a) initial situation, b) after 24 hours, c) after 72  
 313 hours, d) after 150 hours. For each time: picture of a solution (20  $\mu\text{M}$  in DMSO in the small crystallizer) and of a  
 314 sol-gel film on a glass substrate, placed inside the box with the whole animal, under daylight (left) and under  
 315 handheld UV light (right); picture of the spectroscopic cell used for recording the fluorescence under daylight (left)  
 316 and under 365nm UV-light of a portable lamp (right); and emission spectra recorded in the spectrofluorimeter –  
 317  $\lambda_{\text{exc}} = 430 \text{ nm}$ .

318  
 319 Various volatile amine/ammonia vapours are generated during food spoilage. Using shrimps as an example,  
 320 we tested whether **PTTCN** and the prepared materials can be used as fluorescent probe for monitoring seafood  
 321 freshness. To do so, sol-gel spin-coated film as well as a DMSO solution of **PTTCN** (20  $\mu\text{M}$ ) were placed inside  
 322 a packing box in which a shrimp was stored at room temperature for 1 week. A picture of the whole was taken  
 323 every 24 hours and the emission spectra of the solution and films ( $\lambda_{\text{exc}} = 430 \text{ nm}$ ) were also recorded. These data  
 324 are summarised in Figure 7. Initially, only **PTTCN** emission was observed both in film and in solution (weak  
 325 emission in concentrated solution, Figure 7a). Changes can be directly observed by naked eye in the solution  
 326 sample. After 24 h, while the solution was still purple, indicating that **PTTCN** was still the main species in  
 327 solution, a blue fluorescence emanated from the solution when viewed under 365 nm UV-light. The  
 328 corresponding emission spectrum (Figure 7b) showed the appearing of a new emission band at 540 nm, in  
 329 addition to the main **PTTCN** emission. This new band strongly increased in intensity after 48 and then 72 hours

330 (Figure 7b and 7c), as the solution lost its purple colour to grey, and then became almost the only detectable  
331 emission after one week when the solution turned yellow (Figure 7d).

332 Unfortunately, during the same period of time, the changes to the film are not as visually spectacular. The  
333 overall colour does not change and remains purple, albeit slightly faded, as does the fluorescence when viewed  
334 under 365 nm UV-light, dominated by the intense red emission of **PTTCN**. It is only when the emission spectra  
335 are recorded with a spectrofluorimeter, that a change is clearly seen with the appearing of a new band at 540 nm  
336 in addition to the (intense) fluorescence of **PTTCN** (Fig. SI-12) when exciting at a wavelength close to the imine  
337 absorption maximum ( $\lambda_{\text{exc}} = 430$  nm). When PMMA films are used instead, no change both in absorption and  
338 fluorescence spectra are observed. This could originate from the weak concentration of amine in the atmosphere  
339 of the box reducing the kinetic of the reaction that is already slower in PMMA than in sol-gel film (see above).

340 While these film results confirm the previous observations and the formation of new emissive species, they  
341 also reveal the limitation of the system. The detection of volatile amines during food spoilage could not be easily  
342 followed with the naked eye, although it could be possible with the use of an appropriate measuring instrument.

## 343 **Conclusion**

344 In solution in DMSO or embedded in PMMA or sol-gel films, the dicyanovinyl compound **PTTCN** reacts with  
345 strong nucleophilic nitrogen-containing analytes and especially primary aliphatic amine in a transformation  
346 dicyanovinyl  $\rightarrow$  imine, that is reversible in nature. This transformation results in significant changes in the  
347 optical properties, particularly fluorescence. **PTTCN** is only slightly emissive but displays a large emission in  
348 the far-red in the solid-state, while the imine compounds formed by reaction with primary amines, on the other  
349 hand, are highly emissive in both solution and in the solid state. In consequence, **PTTCN** can serve as a very  
350 sensitive turn-on probe to detect primary amines in solution with a 1200-fold enhancement of the fluorescence  
351 intensity during the reaction of **PTTCN** with n-propylamine. Sensitivity in the nanomolar range by fluorescence  
352 monitoring was achieved and an excellent selectivity for primary amines over other amine nucleophiles was  
353 evidenced both in solution and in the solid-state. We have also shown that monitoring of seafood spoilage is  
354 possible with colour changes in response to the vapours released during shrimp spoilage, visible to naked-eye  
355 accompanied by spectacular fluorescence enhancement over time in solution. Spin-coated films can also serve as  
356 probe, highlighting the potential of **PTTCN** for low-cost amine probes, even though the response is less obvious  
357 and only visible by instrumental recording.

358

359 **Declaration of Competing Interest**

360 The authors declare that they have no known competing financial interests or personal relationships that could  
361 have appeared to influence the work reported in this paper.

362 **Acknowledgements**

363 L.B. thank the thanks the Chinese scholarship council and ENS de Lyon for a PhD grant.

364 **Appendix A. Supplementary data**

365 The following is Supplementary data to this article: Experimental details, additional figures, films and  
366 characterizations.

367 **References**

- 368 [1] Y. Özogul, F. Özogul, Chapter 1: Biogenic amines formation, toxicity, regulations in food, in Biogenic  
369 amines in food: Analysis, occurrence and toxicity, The Royal Society of Chemistry, 2020, pp. 1-17,  
370 <https://doi.org/10.1039/9781788015813-00001>.
- 371 [2] C. Ruiz-Capillas, A. M. Herrero, Impact of biogenic amines on food quality and safety, *Foods* 8 (2019) 62,  
372 <https://doi.org/10.3390/foods8020062>
- 373 [3] D. Doeun, M. Davaatseren, M.-S. Chung, Biogenic amines in foods, *Food Sci Biotechnol* 26 (2017)  
374 1463-1474, <https://doi.org/10.1007/s10068-017-0239-3>.
- 375 [4] A. Önal, A review: Current analytical methods for the determination of biogenic amines in foods, *Food*  
376 *Chem.* 103 (2007) 1475-1486, <https://doi.org/10.1016/j.foodchem.2006.08.028>.
- 377 [5] S. K. Kannan, B. Ambrose, S. Sudalaimani, M. Pandiaraj, K. Giribabu, M. Kathiresan, A review on  
378 chemical and electrochemical methodologies for the sensing of biogenic amines, *Anal. Methods* 12 (2020)  
379 3438-3453, <https://doi.org/10.1039/d0ay00358a>.
- 380 [6] J. L. Klockow, K. S. Hettie, T. E. Glass, Fluorescent sensors for amines, in: Atwood JL, *Comprehensive*  
381 *supramolecular chemistry II*, Elsevier, Oxford, 2017, pp. 447-467,  
382 <https://doi.org/10.1016/b978-0-12-409547-2.12630-7>.
- 383 [7] A. I. Danchuk, N. S. Komova, S. N. Mobarez, S. Y. Doronin, N. A. Burmistrova, A. V. Markin, A.  
384 Duerkop, Optical sensors for determination of biogenic amines in food, *Anal. Bioanal. Chem.* 412 (2020)  
385 4023-4036, <https://doi.org/10.1007/s00216-020-02675-9>.
- 386 [8] N. Kaur, S. Chopra, G. Singh, P. Raj, A. Bhasin, S. K. Sahoo, A. Kuwar, N. Singh, Chemosensors for  
387 biogenic amines and biothiols, *J. Mater. Chem. B* 6 (2018) 4872-4902, <https://doi.org/10.1039/c8tb00732b>.
- 388 [9] G. J. Mohr, C. Demuth, U. E. Spichiger-Keller, Application of chromogenic and fluorogenic reactands in  
389 the optical sensing of dissolved aliphatic amines, *Anal. Chem.* 70 (1998) 3868-3873,  
390 <https://doi.org/10.1021/ac980279q>.
- 391 [10] S. Reinert, G. J. Mohr, Chemosensor for the optical detection of aliphatic amines and diamines, *Chem*  
392 *Commun* (2008) 2272-2274, <https://doi.org/10.1039/b717796h>.
- 393 [11] S. Körsten, G. J. Mohr, Star-shaped tripodal chemosensors for the detection of aliphatic amines, *Chem. Eur.*  
394 *J.* 17 (2011) 969-975, <https://doi.org/10.1002/chem.201000787>.
- 395 [12] G. J. Mohr, A tricyanovinyl azobenzene dye used for the optical detection of amines via a chemical

396 reaction in polymer layers, *Dyes Pigm.* 62 (2004) 77-81, <https://doi.org/10.1016/j.dyepig.2003.11.011>.

397 [13] K. Secor, J. Plante, C. Avetta, T. Glass, Fluorescent sensors for diamines, *J. Mater. Chem.* 15 (2005)

398 4073-4077, <https://doi.org/10.1039/b503269e>.

399 [14] E. K. Feuster, T. E. Glass, Detection of amines and unprotected amino acids in aqueous conditions by

400 formation of highly fluorescent iminium ions, *J. Am. Chem. Soc.* 125 (2003) 16174-16175,

401 <https://doi.org/10.1021/ja036434m>.

402 [15] B. Lee, R. Scopelliti, K. Severin, A molecular probe for the optical detection of biogenic amines, *Chem*

403 *Commun* 47 (2011) 9639-9641, <https://doi.org/10.1039/c1cc13604f>.

404 [16] C. Patze, K. Broedner, F. Rominger, O. Trapp, U. H. F. Bunz, Aldehyde cruciforms: Dosimeters for

405 primary and secondary amines, *Chem. Eur. J.* 17 (2011) 13720-13725, <https://doi.org/10.1002/chem.201101871>.

406 [17] J. Freudenberg, J. Kumpf, V. Schäfer, E. Sauter, S. J. Wörner, K. Brödner, A. Dreuw, U. H. F. Bunz,

407 Water-soluble cruciforms and distyrylbenzenes: Synthesis, characterization, and pH-dependent amine-sensing

408 properties, *J. Org. Chem.* 78 (2013) 4949-4959, <https://doi.org/10.1021/jo400576y>.

409 [18] J. Kumpf, J. Freudenberg, S. T. Schwaebel, U. H. F. Bunz, Amine sensing with distyrylbenzenes and their

410 hexamethylene-linked polymers: Spraying them on, *Macromolecules* 47 (2014) 2569-2573,

411 <https://doi.org/10.1021/ma500486u>.

412 [19] J. L. Klockow, T. E. Glass, Development of a fluorescent chemosensor for the detection of kynurenine, *Org.*

413 *Lett.* 15 (2013) 235-237, <https://doi.org/10.1021/ol303025m>.

414 [20] V. Valderrey, A. Bonasera, S. Fredrich, S. Hecht, Light-activated sensitive probes for amine detection,

415 *Angew. Chem. Int. Ed.* 56 (2017) 1914-1918, <https://doi.org/10.1002/anie.201609989>.

416 [21] L. Shi, Y. Fu, C. He, D. Zhu, Y. Gao, Y. Wang, Q. He, H. Cao, J. Cheng, A mild and catalyst-free

417 conversion of solid phase benzylidenemalononitrile/benzylidenemalonate to N-benzylidene-amine and its

418 application for fluorescence detection of primary alkyl amine vapor, *Chem Commun* 50 (2014) 872-874,

419 <https://doi.org/10.1039/c3cc48299e>.

420 [22] L. Kong, Y. Zhang, H. Mao, X. Pan, Y. Tian, Z. Tian, X. Zeng, J. Shi, B. Tong, Y. Dong,

421 Dimalononitrile-containing probe based on aggregation-enhanced emission features for the multi-mode

422 fluorescence detection of volatile amines, *Faraday Discuss.* 196 (2017) 101-111,

423 <https://doi.org/10.1039/c6fd00178e>.

424 [23] D. Moreno, B. D. d. Greñu, B. García, S. Ibeas, T. Torroba, A turn-on fluorogenic probe for detection of

425 MDMA from ecstasy tablets, *Chem Commun* 48 (2012) 2994-2996, <https://doi.org/10.1039/c2cc17823k>.

426 [24] T. Liu, L. Yang, W. Feng, K. Liu, Q. Ran, W. Wang, Q. Liu, H. Peng, L. Ding, Y. Fang, Dual-mode

427 photonic sensor array for detecting and discriminating hydrazine and aliphatic amines, *ACS Appl. Mater. Inter.*

428 12 (2020) 11084-11093, <https://doi.org/10.1021/acsami.0c00568>.

429 [25] T. Mastnak, A. Lobnik, G. J. Mohr, M. Turel, Design and characterization of dicyanovinyl reactive dyes for

430 the colorimetric detection of thiols and biogenic amines, *Sensors* 18 (2018) 814,

431 <https://doi.org/10.3390/s18030814>.

432 [26] T. Mastnak, A. Lobnik, G. J. Mohr, M. Finšgar, Indicator layers based on ethylene-vinyl acetate copolymer

433 (EVA) and dicyanovinyl azobenzene dyes for fast and selective evaluation of vaporous biogenic amines, *Sensors*

434 18 (2018) 4361, <https://doi.org/10.3390/s18124361>.

435 [27] M. Saravanakumar, B. Umamahesh, R. Selvakumar, J. Dhanapal, S. K. Ashok kumar, K. I.

436 Sathiyarayanan, A colorimetric and ratiometric fluorescent sensor for biogenic primary amines based on

437 dicyanovinyl substituted phenanthridine conjugated probe, *Dyes Pigm.* 178 (2020) 108346,

438 <https://doi.org/https://doi.org/10.1016/j.dyepig.2020.108346>.

439 [28] G. Zhang, A. S. Loch, J. C. M. Kistemaker, P. L. Burn, P. E. Shaw, Dicyanovinyl-based fluorescent sensors

440 for dual mechanism amine sensing, *J. Mater. Chem. C* 8 (2020) 13723-13732,

441 <https://doi.org/http://doi.org/10.1039/d0tc03974h>.

442 [29] B. Roy, S. Bandyopadhyay, The design strategies and mechanisms of fluorogenic and chromogenic probes  
443 for the detection of hydrazine, *Anal. Methods* 10 (2018) 1117-1139, <https://doi.org/10.1039/c7ay02866k>.

444 [30] Y. Tan, J. Yu, J. Gao, Y. Cui, Y. Yang, G. Qian, A new fluorescent and colorimetric probe for trace  
445 hydrazine with a wide detection range in aqueous solution, *Dyes Pigm.* 99 (2013) 966-971,  
446 <https://doi.org/10.1016/j.dyepig.2013.08.008>.

447 [31] M. Sun, J. Guo, Q. Yang, N. Xiao, Y. Li, A new fluorescent and colorimetric sensor for hydrazine and its  
448 application in biological systems, *J. Mater. Chem. B* 2 (2014) 1846-1851, <https://doi.org/10.1039/c3tb21753a>.

449 [32] J. Fan, W. Sun, M. Hu, J. Cao, G. Cheng, H. Dong, K. Song, Y. Liu, S. Sun, X. Peng, An ICT-based  
450 ratiometric probe for hydrazine and its application in live cells, *Chem Commun* 48 (2012) 8117-8119,  
451 <https://doi.org/10.1039/c2cc34168a>.

452 [33] Y. Zhang, L. Yuan, S. Jia, X. Liu, J. Zhao, G. Yin, Dicyanovinyl substituted push-pull chromophores:  
453 effects of central C=C/phenyl spacers, crystal structures and application in hydrazine sensing, *Phys. Chem.*  
454 *Chem. Phys.* 21 (2019) 3218-3226, <https://doi.org/10.1039/c8cp07127f>.

455 [34] B. Chen, X. Sun, X. Li, H. Ågren, Y. Xie, TICT based fluorescence “turn-on” hydrazine probes, *Sens.*  
456 *Actuator B-Chem.* 199 (2014) 93-100, <https://doi.org/10.1016/j.snb.2014.03.087>.

457 [35] X. Cheng, Y. Zhou, J. Qin, Z. Li, Reaction-based colorimetric cyanide chemosensors: Rapid naked-eye  
458 detection and high selectivity, *ACS Appl. Mater. Inter.* 4 (2012) 2133-2138, <https://doi.org/10.1021/am3001083>.

459 [36] J. w. Jeong, S. Angupillai, I. J. Kim, J. Jeong, H.-S. Kim, H.-S. So, Y.-A. Son, Michael addition-based  
460 colorimetric and fluorescence chemodosimeters for the nanomolar-level tracking of cyanide ions in  
461 aqueous-organic media, *Sens. Actuator B-Chem.* 237 (2016) 341-349, <https://doi.org/10.1016/j.snb.2016.06.107>.

462 [37] Q. Zhang, J. Zhang, H. Zuo, C. Wang, Y. Shen, A novel colorimetric and fluorescent sensor for cyanide  
463 anions detection based on triphenylamine and benzothiadiazole, *Tetrahedron* 72 (2016) 1244-1248,  
464 <https://doi.org/10.1016/j.tet.2016.01.019>.

465 [38] A. Panja, K. Ghosh, Pyridylazo derivatives with dicyanovinyl appendage in selective Sensing of CN<sup>-</sup> in  
466 sol-gel medium, *ChemistrySelect* 3 (2018) 1809-1814, <https://doi.org/10.1002/slct.201702839>.

467 [39] L. Wang, L. Li, D. Cao, A BODIPY-based dye with red fluorescence in solid state and used as a fluorescent  
468 and colorimetric probe for highly selective detection of cyanide, *Sens. Actuator B-Chem.* 239 (2017) 1307-1317,  
469 <https://doi.org/10.1016/j.snb.2016.09.112>.

470 [40] Y.-D. Lin, Y.-S. Pen, W. Su, K.-L. Liao, Y.-S. Wen, C.-H. Tu, C.-H. Sun, T. J. Chow, Reaction-based  
471 colorimetric and ratiometric fluorescence sensor for detection of cyanide in aqueous media, *Chem. Asian J.* 7  
472 (2012) 2864-2871, <https://doi.org/10.1002/asia.201200578>.

473 [41] J. Orrego-Hernández, J. Portilla, Synthesis of dicyanovinyl-substituted 1-(2-pyridyl)pyrazoles: Design of a  
474 fluorescent chemosensor for selective recognition of cyanide, *J. Org. Chem.* 82 (2017) 13376-13385,  
475 <https://doi.org/10.1021/acs.joc.7b02460>.

476 [42] E. Thanayupong, K. Suttisintong, M. Sukwattanasinitt, N. Niamnont, Turn-on fluorescent sensor for the  
477 detection of cyanide based on a novel dicyanovinyl phenylacetylene, *New J. Chem.* 41 (2017) 4058-4064,  
478 <https://doi.org/10.1039/c6nj03794a>.

479 [43] H. Kwon, K. Lee, H.-J. Kim, Coumarin-malonitrile conjugate as a fluorescence turn-on probe for biothiols  
480 and its cellular expression, *Chem Commun* 47 (2011) 1773-1775, <https://doi.org/10.1039/c0cc04092d>.

481 [44] J. Mei, J. Z. Sun, A. Qin, B. Z. Tang, Comparative study of the dicyanovinyl-functionalized  
482 1,1-dimethyl-2,3,4,5-tetraphenylsilole derivatives on their structures, properties, and applications in thiol  
483 detection, *Dyes Pigm.* 141 (2017) 366-378, <https://doi.org/10.1016/j.dyepig.2017.02.039>.

484 [45] X. Lou, Y. Hong, S. Chen, C. W. T. Leung, N. Zhao, B. Situ, J. W. Y. Lam, B. Z. Tang, A selective  
485 glutathione probe based on AIE fluorogen and its application in enzymatic activity assay, *Sci. Rep.* 4 (2014)

486 4272-4272, <https://doi.org/10.1038/srep04272>.

487 [46] P. V. Bedworth, Y. Cai, A. Jen, S. R. Marder, Synthesis and relative thermal stabilities of diphenylamino-  
488 vs piperidiny-substituted bithiophene chromophores for nonlinear optical materials, *J. Org. Chem.* 61 (1996)  
489 2242-2246, <https://doi.org/10.1021/jo9517555>.

490 [47] F. Steybe, F. Effenberger, S. Beckmann, P. Krämer, C. Glania, R. Wortmann, Enhanced nonlinear optical  
491 properties and thermal stability of donor-acceptor substituted oligothiophenes, *Chem. Phys.* 219 (1997) 317-331,  
492 [https://doi.org/10.1016/S0301-0104\(97\)00103-1](https://doi.org/10.1016/S0301-0104(97)00103-1).

493 [48] F. Steybe, F. Effenberger, U. Gubler, C. Bosshard, P. Günter, Highly polarizable chromophores for  
494 nonlinear optics: syntheses, structures and properties of donor-acceptor substituted thiophenes and  
495 oligothiophenes, *Tetrahedron* 54 (1998) 8469-8480, [https://doi.org/10.1016/S0040-4020\(98\)00450-5](https://doi.org/10.1016/S0040-4020(98)00450-5).

496 [49] M. M. M. Raposo, G. Kirsch, Formylation, dicyanovinyl and tricyanovinyl of 5-alkoxy- and  
497 5-amino- substituted 2,2'-bithiophenes, *Tetrahedron* 59 (2003) 4891-4899,  
498 [https://doi.org/10.1016/S0040-4020\(03\)00705-1](https://doi.org/10.1016/S0040-4020(03)00705-1).

499 [50] F. Effenberger, F. Wuerthner, F. Steybe, Synthesis and solvatochromic properties of  
500 donor-acceptor-substituted oligothiophenes, *J. Org. Chem.* 60 (1995) 2082-2091,  
501 <https://doi.org/10.1021/jo00112a032>.

502 [51] K. Eckert, A. Schröder, H. Hartmann, Preparation and characterization of cyanovinyl-substituted  
503 2-aminothiophenes and 2-aminothiazoles and some of their heterooligomers, *Eur. J. Org. Chem.* 2000 (2000)  
504 1327-1334, [https://doi.org/10.1002/1099-0690\(200004\)2000:7<1327::aid-ejoc1327>3.0.co;2-t](https://doi.org/10.1002/1099-0690(200004)2000:7<1327::aid-ejoc1327>3.0.co;2-t).

505 [52] M. M. Oliva, J. Casado, M. M. M. Raposo, A. M. C. Fonseca, H. Hartmann, V. Hernández, J. T. López  
506 Navarrete, Structure-property relationships in push-pull amino/cyanovinyl end-capped oligothiophenes:  
507 Quantum chemical and experimental studies, *J. Org. Chem.* 71 (2006) 7509-7520,  
508 <https://doi.org/10.1021/jo060318v>.

509 [53] N. Ikemoto, I. Estevez, K. Nakanishi, N. Berova, Fluorescent bithiophene chromophores: Synthesis and  
510 application in CD exciton chirality studies, *Heterocycles* 46 (1997) 489-501, <https://doi.org/10.3987/com-97-S57>.

511 [54] S. Zrig, G. Koeckelberghs, T. Verbiest, B. Andrioletti, E. Rose, A. Persoons, I. Asselberghs, K. Clays,  
512  $\Lambda$ -type regioregular oligothiophenes: Synthesis and second-order NLO properties, *J. Org. Chem.* 72 (2007)  
513 5855-5858, <https://doi.org/10.1021/jo070888a>.

514 [55] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V.  
515 Barone, G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato, A. V. Marenich, J. Bloino, B. G. Janesko, R.  
516 Gomperts, B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, Williams, F. Ding, F.  
517 Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N.  
518 Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y.  
519 Honda, O. Kitao, H. Nakai, T. Vreven, K. Throssell, J. A. Montgomery Jr., J. E. Peralta, F. Ogliaro, M. J.  
520 Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K.  
521 Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C.  
522 Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman, D. J. Fox. *Gaussian*  
523 16 Rev. C.01. Wallingford, CT2016.

524 [56] C. Adamo, V. Barone, Toward reliable density functional methods without adjustable parameters: The  
525 PBE0 model, *The Journal of Chemical Physics* 110 (1999) 6158-6170, <https://doi.org/10.1063/1.478522>.

526 [57] V. Barone, M. Cossi, Quantum Calculation of Molecular Energies and Energy Gradients in Solution by a  
527 Conductor Solvent Model, *The Journal of Physical Chemistry A* 102 (1998) 1995-2001,  
528 <https://doi.org/10.1021/jp9716997>.

529 [58] M. Cossi, N. Rega, G. Scalmani, V. Barone, Energies, structures, and electronic properties of molecules in  
530 solution with the C-PCM solvation model, *Journal of Computational Chemistry* 24 (2003) 669-681,

531 <https://doi.org/https://doi.org/10.1002/jcc.10189>.  
532 [59] T. Yanai, D. P. Tew, N. C. Handy, A new hybrid exchange–correlation functional using the  
533 Coulomb-attenuating method (CAM-B3LYP), *Chemical Physics Letters* 393 (2004) 51-57,  
534 <https://doi.org/https://doi.org/10.1016/j.cplett.2004.06.011>.  
535 [60] T. Le Bahers, C. Adamo, I. Ciofini, A Qualitative Index of Spatial Extent in Charge-Transfer Excitations,  
536 *Journal of Chemical Theory and Computation* 7 (2011) 2498-2506, <https://doi.org/10.1021/ct200308m>.  
537 [61] M. M. M. Raposo, G. Kirsch, A combination of Friedel-Crafts and Lawesson reactions to 5-substituted  
538 2,2'-bithiophenes, *Heterocycles* 55 (2001) 1487-1498, <https://doi.org/10.3987/com-01-9249>.