



HAL
open science

Cardiac alterations with preserved left ventricular ejection fraction association with hepatic steatosis in female prediabetic rats submitted to high-fat-high-sucrose diet

Alexis Jouene, Kenza Hamici, Isabelle Varlet, Carole Lan, Julien Astier, Charlène Couturier, Frank Kober, Jean-Francois Landrier, Monique Bernard, Martine Desrois

► To cite this version:

Alexis Jouene, Kenza Hamici, Isabelle Varlet, Carole Lan, Julien Astier, et al.. Cardiac alterations with preserved left ventricular ejection fraction association with hepatic steatosis in female prediabetic rats submitted to high-fat-high-sucrose diet. Printemps de la Cardiologie, May 2021, Virtual Meeting, France. pp.172, 10.1016/j.acvdsp.2022.04.039 . hal-03796432

HAL Id: hal-03796432

<https://hal.science/hal-03796432>

Submitted on 27 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Cardiac alterations with preserved left ventricular ejection fraction association with hepatic steatosis in female prediabetic rats submitted to high-fat-high-sucrose diet

A. JOUENNE (1) ; K. Hamici (1) ; I. Varlet (1) ; C. Lan (1) ; J. Astier (2) ; C. Couturier (2) ; F. Kober (1) ; JF. Landrier (2) ; M. Bernard (1) ; M. Desrois (1)

(1) *Système cardiovasculaire, Centre de Résonance Magnétique Biologique et Médicale - UMR 7339, Marseille, France;* (2) *Nort, C2VN, Marseille, France*

Introduction

Cardiovascular disease (CVD) is known to be linked with nonalcoholic fatty liver disease (NAFLD) in type 2 diabetes, but few studies are interested in this relationship in prediabetes. This association needs to be explored especially among prediabetic women, who are at greater risk of CVD than men.

Objective

We aimed to evaluate cardiac alterations and its association with hepatic inflammation and lipid metabolism in prediabetic female rats submitted to 5 months of high-fat-high-sucrose diet (HFS).

Method

Wistar female rats were divided in 2 groups fed for 5 months with standard or HFS diet (CTRL/HFS). We analyzed cardiac function, morphology, perfusion and fibrosis by Magnetic Resonance Imaging, then rats underwent an IPGTT. The expression of genes involved in lipid metabolism and inflammation was afterwards studied in liver samples, as well as total lipid mass and triglyceride (TG) contents.

Results

5 months of HFS diet induced glucose intolerance ($p < 0.05$), and cardiac alterations with increased heart perfusion ($p < 0.01$), diastolic and systolic myocardial wall thickness ($p < 0.05$ and $p < 0.001$, respectively) but also increased diastolic left ventricular mass ($p < 0.05$). No significant differences were found in left ventricular ejection fraction, cardiac fibrosis, cardiac output and index between groups. HFS diet induced hepatic lipid accumulation with increased total lipid mass ($p < 0.001$) and TG contents, but also lipid metabolism adaptations with increased acyl CoA oxidase ($p < 0.0001$), CPT1a ($p < 0.0001$) and LCAD ($p < 0.0001$) expressions. No significant differences were found for FAS, MCAD and inflammatory cytokines (IL-6, TNF- α and MCP1) expressions.

Conclusion

HFS diet in female rats induced cardiac alterations associated with hepatic lipid metabolism adaptations characterized by increased lipid peroxidation and β oxidation. Further analyses must be performed to study the relation between cardiac and hepatic alterations observed in female prediabetic rats.