

Present state of global wetland extent and wetland methane modelling: methodology of a model intercomparison project (WETCHIMP)

R. Wania, Joe R. Melton, E. L. Hodson, Ben Poulter, Bruno Ringeval, Renato Spahni, T. Bohn, C. A. Avis, G. Chen, A. V. Eliseev, et al.

▶ To cite this version:

R. Wania, Joe R. Melton, E. L. Hodson, Ben Poulter, Bruno Ringeval, et al.. Present state of global wetland extent and wetland methane modelling: methodology of a model intercomparison project (WETCHIMP). 2022. hal-03796920

HAL Id: hal-03796920 https://hal.science/hal-03796920

Preprint submitted on 5 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Geosci. Model Dev. Discuss., 5, 4071–4136, 2012 www.geosci-model-dev-discuss.net/5/4071/2012/ doi:10.5194/gmdd-5-4071-2012 © Author(s) 2012. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal Geoscientific Model Development (GMD). Please refer to the corresponding final paper in GMD if available.

Present state of global wetland extent and wetland methane modelling: methodology of a model intercomparison project (WETCHIMP)

R. Wania^{1,*}, J. R. Melton^{2,**}, E. L. Hodson^{3,***}, B. Poulter⁴, B. Ringeval^{4,5,6}, R. Spahni⁷, T. Bohn⁸, C. A. Avis^{9,****}, G. Chen¹⁰, A. V. Eliseev^{11,12}, P. O. Hopcroft⁵, W. J. Riley¹³, Z. M. Subin^{13,*****}, H. Tian¹⁰, V. Brovkin¹⁴, P. M. van Bodegom¹⁵, T. Kleinen¹⁴, Z. C. Yu¹⁶, J. S. Singarayer⁵, S. Zürcher⁷, D. P. Lettenmaier⁸, D. J. Beerling¹⁷, S. N. Denisov¹¹, C. Prigent¹⁸, F. Papa¹⁹, and J. O. Kaplan²

 ¹Institut des Sciences de l'Evolution (UMR5554, CNRS), Université Montpellier 2, Place Eugène Bataillon, 34090 Montpellier, France
 ²ARVE Group, École Polytechnique Fédérale de Lausanne, Switzerland
 ³Swiss Federal Research Institute WSL, Switzerland
 ⁴Laboratoire des Sciences du Climat et de L'Environment, CNRS–CEA, UVSQ, Gif-sur Yvette, France
 ⁵BRIDGE, School of Geographical Sciences, University of Bristol, UK

⁶Department of Earth Sciences, VU University, Amsterdam, The Netherlands



GMDD

 ⁷Climate and Environmental Physics, Physics Institute & Oeschger Centre for Climate Change Research, University of Bern, Switzerland ⁸Dept. of Civil and Environmental Engineering, University of Washington, USA ⁹School of Earth and Ocean Sciences, University of Victoria, Canada ¹⁰International Center for Climate and Global Change Research and School of Ecrestry and 	Discussion Pa	GN 5, 4071–4	IDD 136, 2012
Wildlife Sciences, Auburn University, Auburn, AL 36849, USA ¹¹ A.M. Obukhov Institute of Atmospheric Physics, Russian Academy of Sciences,	iper	WET(metho	CHIMP dology
Russia ¹² Kazan (Volga region) Federal University, Russia ¹³ Earth Sciences Division (ESD), Lawrence Berkeley National Lab, USA ¹⁴ May Planek Institute für Metaarologia, Hamburg, Cormony	Discus	R. War	nia et al.
¹⁵ Department of Ecological Sciences, VU University, Amsterdam, The Netherlands ¹⁶ Department of Earth and Environmental Sciences, Lehigh University, USA	sion Pap	Title	Page
¹⁹ LEGOS, IRD, 18 Ave. Edouard Belin, 31400 Toulouse, France	ber	Conclusions	Reference
now at: Lanser Strasse 30, 6080 Igls, Austria ** now at: Canadian Centre for Climate Modelling and Analysis, Environment Canada, Victoria, BC, V8W 2Y2, Canada	Discuss	Tables	Figures
¹ now at: AAAS Science and Technology Policy Fellow, Office of Climate Change Policy and Technology, US Department of Energy, USA ¹ now at: Physics and Astronomy department, Camosun College, Victoria, BC, Canada	sion Pap		•
now at: Princeton Environmental Institute, Princeton University, Princeton, New Jersey, USA	er	Back Full Scr	Close een / Esc
Received: 30 October 2012 – Accepted: 26 November 2012 – Published: 10 December 2012	Dis		
Correspondence to: J. R. Melton (joe.melton.sci@gmail.com)	cuss	Printer-frier	ndly Version
Published by Copernicus Publications on behalf of the European Geosciences Union.	sion	Interactive	Discussion
	Paper	œ	BY



Abstract

The Wetland and Wetland CH_4 Intercomparison of Models Project (WETCHIMP) was created to evaluate our present ability to simulate large-scale wetland characteristics and corresponding methane (CH_4) emissions. A multi-model comparison is essential to

- ⁵ evaluate the key uncertainties in the mechanisms and parameters leading to methane emissions. Ten modelling groups joined WETCHIMP to run eight global and two regional models with a common experimental protocol using the same climate and atmospheric carbon dioxide (CO₂) forcing datasets. We reported the main conclusions from the intercomparison effort in a companion paper (Melton et al., 2012). Here we provide
- ¹⁰ technical details for the six experiments, which included an equilibrium, a transient, and an optimized run plus three sensitivity experiments (temperature, precipitation, and atmospheric CO_2 concentration). The diversity of approaches used by the models is summarized through a series of conceptual figures, and is used to evaluate the wide range of wetland extents and CH_4 fluxes predicted by the models in the equilibrium run.
- ¹⁵ We discuss relationships among the various approaches and patterns in consistencies of these model predictions. Within this group of models, there are three broad classes of methods used to estimate wetland extent: prescribed based on wetland distribution maps, prognostic relationships between hydrological states based on satellite observations, and explicit hydrological mass balances. A larger variety of approaches was
- ²⁰ used to estimate the net CH_4 fluxes from wetland systems. Even though modelling of wetland extents and CH_4 emissions has progressed significantly over recent decades, large uncertainties still exist when estimating CH_4 emissions: there is little consensus on model structure or complexity due to knowledge gaps, different aims of the models, and the range of temporal and spatial resolutions of the models.



1 Introduction

In order to study the importance of wetlands in the global water and carbon cycle a variety of hydrological and biogeochemical models have been developed over the last three decades. The first studies of global-scale wetland CH_4 modelling appeared twenty-five

- $_{5}$ years ago (Matthews and Fung, 1987). Matthews and Fung (1987) combined vegetation, soil and fractional inundation maps along with estimates of CH₄ flux intensity to generate a map of global wetland distribution and an annual wetland emissions estimate of ~ 100 Tg CH₄ yr⁻¹. Aselman and Crutzen (1989) soon followed developing their own wetland distribution datasets, and assumed CH₄ emission flux rates, yielding
- ¹⁰ a wetland emissions estimated range of 40–160 Tg CH_4 yr⁻¹. These early approaches are limited by uncertainties inherent in up-scaling point measurements to large regions, and an inability to predict changes to wetland systems due to changes in climate and hydrology because of the use of static wetland extents and simple scaling-based estimates of CH_4 emissions.
- As an attempt to circumvent these limitations, process-based modelling of global CH_4 emissions from wetland systems was first pioneered by Fung et al. (1991) followed by Christensen and Cox (1995), Christensen et al. (1996), and Cao et al. (1996). While those early global studies used the static wetland maps of Matthews and Fung (1987), they differed in their approach to simulate the CH_4 emissions. Christensen and Cox
- ²⁰ (1995) was the first study to introduce a formulation for oxidation and a soil vertical discretization in a one-dimensional, single-column model. The simple approach of Christensen et al. (1996) estimates net CH_4 emissions as a fraction of heterotrophic respiration calculated by an equilibrium vegetation model (BIOME2) giving a climate sensitive, but perhaps simplistic CH_4 emissions estimate. A more mechanistic approach
- ²⁵ was adopted by Cao et al. (1996) whose CH₄ emission model assumes substrate supply to methanogens is controlled by plant primary productivity and soil organic matter decomposition. Methane production is then modelled as a function of soil temperature, soil organic matter decomposition, water table position, and a fixed ratio of CH₄



production to decomposed organic carbon. Oxidation of produced CH_4 is assumed to scale with gross primary productivity (GPP) and to increase to a fixed fraction under non-inundated conditions (Cao et al., 1996).

- None of these initial modelling studies performed transient simulations and the models originally accounted for hydrologic variability only in the vertical dimension, not spatially within model gridcells. Other process-based models soon followed but were not applied on a global-scale, at least initially (Walter et al., 1996; Potter, 1997; Walter and Heimann, 2000). These initial papers included mechanistic modelling of such processes as diffusive, aerenchymal, and ebullition gas and oxygen transport. More recent
 work has devoted much effort to improving modelling of these processes (Segers and Leffelaar, 2001; van Bodegom et al., 2001a,b; Zhuang et al., 2006) and other controls on CH₄ production such as pH (Zhuang et al., 2004). Oxidation in the oxic portion of
- the soil, water column, and rhizosphere has also been parameterized (Ridgwell et al., 1999; Segers and Leffelaar, 2001; Zhuang et al., 2006; Curry, 2007, 2009). Model simulations have also moved on from equilibrium-only simulations to transient simulations
- (Walter et al., 2001a,b; Shindell et al., 2004; Gedney et al., 2004; Zhuang et al., 2006). Regional- to global-scale models have now been applied for the recent past (Ringeval et al., 2010; Hodson et al., 2011; Spahni et al., 2011; Riley et al., 2011), more distant past climates (Kaplan, 2002; Valdes et al., 2005; Hopcroft et al., 2011; Singarayer
- et al., 2011; Beerling et al., 2011), and to project responses to future climate change (Shindell et al., 2004; Gedney et al., 2004; Bohn et al., 2007; Bohn and Lettenmaier, 2010; Ringeval, 2011). Wetland and wetland CH_4 models are now becoming included in intermediate complexity (Shindell et al., 2004; Gedney et al., 2004; Avis et al., 2011) and comprehensive (Riley et al., 2011) global climate and earth system models.
- ²⁵ A number of models have also integrated approaches allowing for dynamic wetland response to climate changes. Approaches to simulate wetland distribution in order to study the interaction between climate and free water bodies were developed by Coe (1997, 1998) and Krinner (2003). The earliest attempt at wetland modelling for the purpose of estimating wetland CH₄ emissions was designed to estimate wetland



emissions during the Last Glacial Maximum (LGM, ~ 21 ka before present Kaplan, 2002). The simple scheme of Kaplan (2002) used threshold values of slope and soil moisture content to define wetland areas, with the soil moisture calculated by an equilibrium vegetation model (BIOME4); an approach adopted by other models (Shindell et al., 2004; Weber et al., 2010; Avis et al., 2011). Later schemes used land cover datasets to outline peatland regions (Wania et al., 2009a, 2010; Spahni et al., 2011),

and/or satellite-derived inundation datasets to prescribe wetlands either directly (Hodson et al., 2011; Ringeval et al., 2010), or indirectly (Ringeval et al., 2011; Riley et al., 2011). Other wetland distribution schemes use internally-calculated water table positions or soil moisture thresholds to locate wetlands (Chen et al., 2012).

In this context, the WETland and wetland CH_4 Inter-comparison of Models Project (WETCHIMP) was designed to offer the first multi-model comparison highlighting similarities and differences between modelling approaches and results. The advantage of using a multi-model comparison is that the range of the current state-of-the-art model

- estimates for wetland extents and CH₄ emissions can be studied in parallel. This approach allows us to study the sources of uncertainties and spatial and temporal differences in model behaviour. The results of this multi-model comparison are presented in Melton et al. (2012). In this paper, we provide the technical background for WETCHIMP, presenting details of the modelling protocol (Sect. 2), descriptions of the models as
 used for WETCHIMP (Sect. 3), conceptual comparisons of the models involved and results from the model default simulation to illustrate the differences between models
- are presented in Sect. 4.

2 Modelling protocol

The models participating in WETCHIMP followed a common modelling protocol for six experiments (Table 1) and adhered to it as closely as possible; divergences from the modelling protocol are described in the individual model description section.



As briefly described in Melton et al. (2012), WETCHIMP consisted of six experiments, including both a transient simulation and several equilibrium state simulations with step-changes to climate forcing. The first experiment ("Experiment 1-equil") was an equilibrium simulation under repeating 1901–1931 climate and a carbon dioxide concentration ([CO₂]) of 303 ppmv. The second experiment ("Experiment 2-trans") was a transient historical simulation from 1901–2009, using observed climate and atmospheric CO₂ concentration ([CO₂]), with the final (equilibrium) state of Experiment 1-equil as its initial state. Some models require observed fractional inundation values as an input; these were provided and cover the period 1993–2004 by the Global Inundation Extent from Multi-Satellites (GIEMS) dataset (Prigent et al., 2007; Papa et al., 2010). Thus, the period 1993–2004 was selected from Experiment 2-trans for comparison of model results. A third experiment ("Experiment 3-opt") was run for the same time period as Experiment 2-trans, but allowed the models to run under user-defined optimal configurations (e.g. running coupled into an earth system model or using differ-

¹⁵ ent meteorological forcing or remotely-sensed inundation datasets than those common to Experiment 2-trans).

The remaining three experiments applied step-changes to each model's equilibrium state from Experiment 1-equil. The fourth experiment ("Experiment 4- CO_2 ") applied an instantaneous increase in atmospheric [CO_2] to 857 ppmv (SRES A2 year 2100 levels from LDCC (2000)) while helding the other state state state instantaneous increase in the other state state state state state.

- from IPCC (2000)) while holding the other meteorological inputs identical to Experiment 1-equil; this perturbed simulation was then run until each model had reached a new equilibrium state. Experiment 5 ("Experiment 5-T") investigated the effect of an instantaneous increase of +3.4 °C in surface air temperature (SAT). The magnitude of this increase was chosen from the SRES A2 year 2100 multi-model mean SAT warming
- for the period 2080–2099 relative to 1980–1999 (Meehl et al., 2007). However, since this increase was applied to the mean climate of 1901–1931, it represented a slightly smaller departure from the 1901–1931 equilibrium than from the climate of 1980–1999. The final experiment ("Experiment 6-P") examined model responses to changes in precipitation with an instantaneous increase of +3.9 % (SRES A2 2100 level; 30-yr global



average for 2071–2100 relative to 1961–1990) (Prentice et al., 2001). In all cases, the step increases were applied to all months and gridcells uniformly. While actual changes in climate are projected to vary in both space and time, these uniform changes are suitable for the purpose of sensitivity tests (Melton et al., 2012). An overview of which groups conducted which simulations is shown in Table 2.

All data are freely available for download on http://arve.epfl.ch/pub/wetchimp, please send request for a username and password to joe.melton.sci@gmail.com.

2.1 Data sets

5

2.1.1 Climate data

- The CRU TS3.1 time series (Mitchell and Jones, 2005; Jones and Harris, 2008) was used for monthly climate forcing data and – dependent on the model – precipitation, 2 m air temperature, percentage cloud cover, number of wet days, and vapour pressure were used from this data set. Models that required data with a higher temporal resolution used the CRUNCEP data, which is the correction of the 6-hourly NCEP reanalysis
- ¹⁵ by the CRU TS3.1 data (Viovy and Ciais, 2011). CRUNCEP provides incoming longand short-wave radiation, air specific humidity (used to compute the relative humidity), pressure, total precipitation, temperature, and the zonal and meridional components of the wind. UVic-ESCM used surface winds and diurnal temperature range from the NCEP reanalysis directly.

20 2.1.2 Soil and wetland distribution data

The soil data used in WETCHIMP are given in Table 3 and are allocated to each model in Table 4. The model requirements for soil data are too broad to accommodate a uniform soil data set easily. Soil data sets used, and model treatment of soil textural information, is thus considered part of the wetland model itself.



There are several wetland distribution maps that were used for our simulations: (i) remotely sensed inundation area from GIEMS (Prigent et al., 2007; Papa et al., 2010) (Fig. 1), (ii) northern peatland distribution from NCSCD (Tarnocai et al., 2007, 2009) (Fig. 2), and (iii) peatland distribution for the West Siberian Lowlands (Sheng et al., 2004) (Fig. 3). In addition, some groups made use of the rice distribution data set by Leff et al. (2004) (Fig. 4) and the GICEW waterbodies and land ice data set (Fig. 5) to exclude areas from their wetland distribution map.

2.1.3 Global Inundation Extent from Multi-Satellites (GIEMS)

As the GIEMS dataset is used extensively by several models, and forms a comparison for the model outputs in Melton et al. (2012), it will be described in more detail here. The GIEMS dataset (Fig. 1) is a global, multi-year product quantifying the monthly variations of the distribution and extent of episodic and seasonal inundations, wetlands, rivers, lakes and irrigated agriculture at 0.25° resolution at the equator. GIEMS is derived from a complementary suite of satellite observations including passive microwave ob-

- ¹⁵ servations (SSM/I emissivities), active microwave observations (ERS scatterometer), along with AVHRR-NDVI. The complete methodology is described in detail in Prigent et al. (2007) and Papa et al. (2010) and is briefly summarized here. First, an unsupervised classification of the three sources of satellite data is performed, and pixels with satellite signatures likely related to inundation are retained. For each inundated
- pixel, the monthly fractional coverage by open water is obtained using the passive microwave signal and a linear mixture model with end-members calibrated with scatterometer observations to account for the effects of vegetation cover. We use here the dataset available at a monthly time scale for the period 1993–2004. More detailed information concerning the seasonal and inter-annual behaviour of GIEMS dataset can
- ²⁵ be found in Prigent et al. (2012) for the global scale analysis and in Papa et al. (2006) and Papa et al. (2008) for for the tropical and boreal regions, respectively.



3 Participating models

In this section, we describe each model briefly and refer to published papers where more detailed information can be found. Modifications to models that were made after recent publications or specifically for WETCHIMP are described in the respective model

5 section. An overview of which models performed which experiments is given in Table 2 and a list of forcing data for each model is provided in Table 4. The models follow the prescribed modelling protocol listed in Table 1 unless otherwise stated in the respective model description.

3.1 CLM4Me

- ¹⁰ The version of CLM4Me used for this project is described in Riley et al. (2011), and is incorporated into the Community Land Model 4 (CLM4; Lawrence et al., 2011), the land-surface component of CESM1 (Community Earth System Model 1). Using the hydrology, soil carbon cycling, and soil thermal physics predicted in CLM4, net CH₄ fluxes are computed separately in inundated and non-inundated areas in each gridcell, including untake of streaghboring Cline. The meeting and transport equations for Cline and
- ¹⁵ including uptake of atmospheric CH₄. The reaction and transport equations for CH₄ and oxygen (where applicable) include production, consumption, aerenchyma transport, ebullition, and diffusion.

3.1.1 WETCHIMP setup

The CH₄ model code deviates slightly from that described in Riley et al. (2011); these changes resulted in less than a 5 % difference from the global budget presented in Riley et al. (2011). The changes in the code include: (i) the calculation of below-ground root mass for determining aerenchyma area now uses the time-lagged (1-yr decay time) belowground-to-aboveground NPP ratio, instead of the instantaneous one, and (ii) in calculating the water availability for permafrost vegetation, root fraction is weighted



over all soil layers down to last year's maximum active layer depth, rather than the instantaneous active layer depth, thereby causing a slight delay in growth in the spring.

The CLM model requires a number of forcings in addition to the lower atmospheric boundary conditions and fluxes specified in the CRUNCEP forcing. For all the experi-

ments except Experiment 3-opt, the standard CLM4 year 2000 conditions are used for atmospheric [CO₂] experienced by plant stomata (except Experiment 4-CO₂), atmospheric nitrogen deposition, atmospheric aerosol deposition, and vegetation distributions. For Experiment 3-opt, the same configuration as in Riley et al. (2011) is used, namely a spin up to "1850" conditions using the 1850 [CO₂], nitrogen, aerosols, and vegetation distributions, and then repeated 1948–1972 (Qian et al., 2006) corrected-NCEP forcing. A transient simulation from 1850–2004 is run using transient data for [CO₂], nitrogen, aerosols, and vegetation, using repeated 1948–1972 forcing through 1972, at which point the model is switched to actual-year forcing through 2004.

The model is run at 1.9 × 2.5° resolution and the standard CLM4 datasets are used,
except that the default CLM4 1 × 10⁶ km² of inland non-vegetated wetland area that were used in Riley et al. (2011), were eliminated. As described in that paper, the CLM4Me model requires three parameters at each gridcell to calculate the inundated fraction as a function of the modelled water table and lagged surface runoff, based on an inversion to Prigent et al. (2007) satellite observations for 1994–1998. For
WETCHIMP, the parameters generated in a previous model described in Riley et al. (2011) were used (similar to Experiment 3-opt); however, the parameters were not reoptimised with the CRUNCEP forcing, hence the CRUNCEP 1990s inundated area (e.g. Experiment 2-trans) may differ from that simulated in Experiment 3-opt.

3.2 DLEM

²⁵ The Dynamic Land Ecosystem Model (DLEM) is a process-based model that simulates daily carbon, water and nitrogen fluxes and pool sizes for land and riverine ecosystems. These pools and fluxes are influenced by changes in atmospheric chemistry (CO₂, ozone concentration and nitrogen deposition), climate, land-cover and land-use



change, management practices (e.g. irrigation, fertilization, rotation, and technology improvement), and other disturbances (e.g. fire, hurricane, insects, disease, and forest harvest) (Ren et al., 2007, 2011; Tian et al., 2010a,b, 2011a,b, 2012; Chen et al., 2012; Lu et al., 2012; Liu et al., 2012a,b). For WETCHIMP, the disturbance submodel and the influence of ozone chemistry were not used due to a lack of spatially explicit driving data.

The soil biogeochemistry module simulates CH_4 uptakes in upland ecosystem and emissions in wetland ecosystems. The mechanisms and algorithms for simulating CH_4 fluxes have been described in Tian et al. (2010b, 2011b); Xu et al. (2010). DLEM requires input datasets for daily climate (average, maximum, and minimum air temperature, precipitation, gross radiation, and relative air humidity), atmospheric composition ([CO_2], nitrogen deposition and ozone), annual land use information, soil condition information (soil texture, pH, and soil depth), and topographic data (elevation, slope, and aspect).

10

- Wetlands are defined as those areas that are inundated or saturated by surface water at a frequency and duration sufficient to support vegetation growth, which leads to five wetland types: (i) rice paddy, (ii) permanent herbaceous wetland, (iii) permanent woody wetland, (iv) seasonal herbaceous wetland and (v) seasonal woody wetland. The distribution map for different wetland types are determined based on the data from Stillwell-
- Soller et al. (1995); Aselman and Crutzen (1989) and Lehner and Döll (2004). DLEM simulates water transport to rivers based upon catchments, but does not explicitly move water through gridcells and thereby does not influence conditions in neighbouring gridcells. The version of DLEM used to simulate CH₄ fluxes for WETCHIMP has been described by Tian et al. (2010b, 2011a); Xu et al. (2010). The CH₄ exchanges between
- ecosystems and the atmosphere are a combination of CH₄ production, oxidation, and transportation from soil/water to the atmosphere. DLEM only considers CH₄ production from dissolved organic carbon (DOC), which is indirectly controlled by environmental factors including soil pH, temperature, soil texture and soil moisture content.



3.2.1 WETCHIMP setup

In an earlier version of DLEM, wetland extent was prescribed by external input datasets, which means that wetland extent could not be influenced by environmental factors (Tian et al., 2010a; Liu et al., 2008, 2012a; Chen et al., 2012). For WETCHIMP, the DLEM water cycling processes were improved to dynamically simulate natural wet-

- land extent through the coupling of TOPMODEL and other models with DLEM. This model setup has been described in Liu et al. (2012a) and Yang et al. (2012). Major improvements include: (i) more soil layers (ten), (ii) fractional vegetation structure, and (iii) a river routing system. After these improvements, DLEM can be used to dynam-
- ically identify daily patterns of wetland distribution extent. This new version of DLEM uses both water table position and soil moisture to determine the wetland percentage. In order to integrate the provided GIEMS and rice paddy (Leff et al., 2004) datasets with the DLEM-simulated wetland distribution, some of the DLEM parameterizations were adapted for a semi-prognostic approach for wetland areal determination. DLEM
- separately simulated the extent of two major natural wetland types: permanent and seasonal wetlands. For the permanent natural wetland distribution, areas of the GIEMS dataset that were continually inundated during the growing season (May to October) of 1993–2004, were regarded as permanent wetlands. In these areas the soil moisture was prescribed at saturation. Excluding areas of permanent wetlands, seasonal
- wetlands were determined using the DLEM prognostic parameterizations as discussed previously. Thus minimum annual wetland area is controlled by the GIEMS dataset, but daily and seasonal wetland areal dynamics above this were determined by internal DLEM model dynamics. Areal extent of rice paddies was used directly from Leff et al. (2004).

25 3.3 IAP-RAS model

The present version of the IAP-RAS wetland CH_4 emission module is described by Mokhov et al. (2007). The module consists of two parts; one for soil temperature



calculations, and a second for calculations of CH_4 emissions. Soil temperature is calculated based on the model by Kudryavtsev et al. (1977). This model represents a generalized solution of Stephan's problem with annual temperature variations at the lower boundary of the atmosphere, while accounting for the influence of snow

- ⁵ and moss cover. Moss cover is prescribed as a layer of 10 cm thickness in presence of boreal and tundra vegetation. The depth of seasonal thaw/freeze depends on the annual variation of the near-surface temperature and precipitation. The influence of the effect of snow metamorphism is ignored. The seasonal thaw depth was assessed based on the thickness of the active layer and temperature of the soil sur-
- face. Only soil layers to a certain limit depth were included in the calculations. In the standard version, the depths of 15 and 60 cm were used for tropical and extratropical zones, respectively. Similar depth values are obtained for the organic carbon content in soil at the characteristic peat density of 200 kgm⁻³ based on data from http://soils.usda.gov/use/worldsoils/mapindex/soc.html. Deeper layers were ignored in calculations of CH₄ emissions by wetlands. The amount of water in wetlands is con-
- sidered to be always sufficient for inundation. Methane emissions are calculated based on the empirical model of Christensen and Cox (1995).

3.3.1 WETCHIMP setup

For WETCHIMP simulations, the model was run at $0.5^{\circ} \times 0.5^{\circ}$ resolution using CRU

- TS3.1 data set as climate forcing (Mitchell and Jones, 2005). Wetland areal extent was prescribed according to CDIAC NDP017 dataset, also known as the Olson database (http://cdiac.esd.ornl.gov/ndps/ndp017.html). In this data set, only areas with ecosystem codes 44 ("bog/mire of cool or cold climates"), 45 + 72 ("warm and hot wetlands"), 64 ("heath and moorland"), and 53 ("tundra") are considered as wetlands. The inclusion
- of tundra regions as methane–producing area was specifically for WETCHIMP; earlier applications of IAP-RAS model neglected their contribution. Outside of wetlands, soil thermophysical parameters are homogeneously prescribed as loam everywhere.



3.4 LPJ-Bern

5

LPJ-Bern is a subsequent development of the Lund-Potsdam-Jena dynamic global vegetation model (Sitch et al., 2003; Joos et al., 2004; Gerber et al., 2003) that combines process-based, large-scale representations of terrestrial vegetation dynamics, soil hydrology (Gerten et al., 2004; Wania et al., 2009a), human induced land use changes (Strassmann et al., 2008; Stocker et al., 2011), permafrost and peatland establishment (Wania et al., 2009a,b) and simulation of biogeochemical trace gas emissions, such as

CH₄ (Wania et al., 2010; Spahni et al., 2011; Zürcher et al., 2012).

3.4.1 WETCHIMP setup

¹⁰ The CH₄ model within the LPJ-Bern version differs slightly from the LPJ-WHyMe CH₄ model that was used in Wania et al. (2010) and Spahni et al. (2011). The main differences with respect to CH₄ emissions concern peatland modelling, global carbon cycle parameters and input data. The differences between the model as used in this study and Spahni et al. (2011) (and thus to LPJ-WHyMe version 1.3.1, Wania et al., 2010) are described below ordered by CH₄ source and sink category.

LPJ-Bern uses a different ebullition mechanism for CH_4 emissions from peatlands, which includes variations in partial pressure of CO_2 (Zürcher et al., 2012). The carbon balance over all layers is now preserved after every gas diffusion time step, whereas in LPJ-WHyMe a correction factor for carbon balance is applied at the end of the year. The

- ²⁰ possible plant functional types in peatlands are limited to flood-tolerant graminoids and *Sphagnum* mosses. Additionally, the prescribed fractional peatland cover per gridcell is taken from NCSCD (Tarnocai et al., 2007, 2009). NCSCD covers histels and histosols in the northern high-latitudes with a total area of 2.7×10^6 km², which is larger than the the extent (2.06×10^6 km²) used in Spahni et al. (2011). The global scaling factor used by Wania et al. (2010) to account for the lack of microtopography in the model is
- thus reduced from 75 % to 26 % to constrain CH_4 emissions from peatlands in 2004 to 28.2 Tg CH_4 yr⁻¹ (Spahni et al., 2011).



For CH₄ emissions in inundated areas the GIEMS monthly fractional inundation data set for 1993–2004 was averaged by month (e.g. mean January, mean February, etc.). The fractional area of inundation per gridcell is further limited by the fraction of land available, i.e. land not covered by permanent water or ice (GICEW, http://luh.sr.unh. edu/). In gridcells containing peatlands (35–90° N), the inundated fraction was assumed

- edu/). In gridcells containing peatlands (35–90° N), the inundated fraction was assumed to include peatlands. If the inundated fraction exceeds the constant peatland fraction, the difference is interpreted as the inundated fraction of mineral soils. This is different to the treatment in Spahni et al. (2011), where the inundated fraction was explicitly set to zero north of 45° N. The fraction of inundated areas was further divided into natural
 wetlands and rice agriculture using the scheme as described in Spahni et al. (2011).
- For these two categories the CH_4 to CO_2 conversion tuning parameter was adjusted to obtain total CH_4 emissions in 2004 of 81.3 Tg CH_4 yr⁻¹ for natural wetlands and of 43.1 Tg CH_4 yr⁻¹ rice agriculture (Spahni et al., 2011).
- For CH_4 emissions in wet mineral soils, the above changes were included and the ¹⁵ remaining non-inundated and non-peatland land cover was taken as fractional area of mineral soils. These mineral soils can either function as a CH_4 source or sink, depending on their soil moisture (Spahni et al., 2011). For this study the CH_4 -to- CO_2 conversion factor – a global scaling factor – for CH_4 emissions from wet mineral soils was scaled to obtain emissions of 63.1 Tg CH_4 yr⁻¹ for 2004. For the CH_4 uptake ²⁰ the concentration-to-flux tuning factor was reduced to reach a global consumption of 25.8 Tg CH_4 yr⁻¹ (Spahni et al., 2011).

While the peatland fraction is a separate tile in each gridcell with its own carbon and soil water pools, the other three CH₄ source types and the sink share the same tile. So, for the non-peatland areas, there is no interaction between water table position and vegetation growth. Compared to Spahni et al. (2011), an updated soil type map based on the World Harmonized Soil Database (FAO/IIASA/ISRIC/ISSCAS/JRC, 2009) was used by selecting the dominant soil type in each 0.5° × 0.5° gridcell. However, soil properties for the corresponding 9 soil types were not changed to previous simulations (Spahni et al., 2011).



When model results were compared to other WETCHIMP models a unique feature was observed in CH_{4} emissions from northern peatlands as simulated by LPJ-Bern. Emissions for the years 1998, 1999 and 2001 are extremely high for some sites compared to average peatland emissions (Melton et al., 2012). Investigating the model output showed the high emission peaks in those years originates from CH_4 gas reser-5 voirs within and below frozen peat layers. Two processes are responsible for the establishment of these CH₄ gas reservoirs. First, in LPJ-Bern, frozen peat layers act as a barrier for gas diffusion thus allowing CH₄ to accumulate beneath. Second, if environmental conditions are suitable methanogenesis can occur in unfrozen layers, regardless whether layers above are frozen. Normally, CH_{4} production in deeper layers 10 is negligible, but for the WETCHIMP simulations a considerable amount of CH₄ gas accumulated beneath a frozen layer during the model spin-up procedure (1000 yr). This stored CH₄ could not escape to the atmosphere until the year 1998 in the transient run, the first year showing an exceptional boreal warming that lead to thawing of peat layers and burst emissions of CH_4 . While the process as such could be questioned 15

the magnitude of CH_4 emissions is certainly too large, as can be concluded from the model-data comparison in the Hudson's Bay Lowlands (Melton et al., 2012).

3.5 LPJ-WHyMe

Methane emissions for peatlands north of 35° N were simulated using LPJ-WHyMe (Wania et al., 2009a,b, 2010; Spahni et al., 2011). Location and fractional cover of peatlands are taken from NCSCD (Tarnocai et al., 2007, 2009). Monthly air temperature, precipitation, percentage cloud cover and number of wet days are taken from CRU TS3.1 to force all simulations. The data from 1901–1930 are used repeatedly to spin up the model for 1000 yr before running it transiently, either for 1901–1930 or 1901–2009.



3.5.1 WETCHIMP setup

Instead of running LPJ-WHyMe only once for an average topographical microform, two parallel runs were executed for each experiment. One run represents the wetter microform, lawns, which emit more CH_4 and the other one represents the drier microform,

- ⁵ hummocks. The model modifications to approximate these two microforms include setting the daily drainage rate to 0.2 mm (lawns) and to 0.6 mm (hummocks), whereas it was 0 mm in Wania et al. (2010). These modifications lower the water table position in hummocks compared to lawns. The vegetation for hummocks is restricted to *Sphagnum* mosses, whereas lawns are able to grow any plant functional type depending on
 the water table position (Wania et al., 2009b). Methane emissions from the two parallel runs are averaged under the assumption that hummocks and hollows cover approxi
 - mately the same surface area.

3.6 LPJ-WSL

The LPJ-WSL CH₄ model used in this analysis is the same as presented in Hodson ¹⁵ et al. (2011), but has been recalibrated to a new set of regional CH₄ fluxes as noted below. The wetland CH₄ flux (*E*) at each $0.5^{\circ} \times 0.5^{\circ}$ gridcell (*x*) and monthly time step (*t*) is calculated as a linear function of two scaling factors ($r_{CH_4:C}$ and f_{ecosys}), wetland extent (*A*) and heterotrophic respiration (R_{hetr}) according to the following equation:

$$E(x,t) = r_{\text{CH4:C}} \cdot f_{\text{ecosys}}(x) \cdot A(x,t) \cdot R_{\text{h}}(x,t)$$

²⁰ The notation in Eq. (1) has been modified from Hodson et al. (2011) to follow Table 5 (to convert between the notation in Eq. (1) and Hodson et al. (2011): $r_{CH4:C} = \beta$; $f_{ecosys} = F$; A = S)

Together, $r_{CH4:C}$ and $f_{ecosys}(x)$ comprise the scaling ratio F(x), which converts C to CH₄ fluxes and is a function of two weighted-regional scaling factors, one representing tropical (T) and another representing boreal (B) wetland climates (with temperate conditions represented as a combination of tropical and boreal). This approach allows the



(1)

model to account for broad ecosystem differences in CH_4 -emitting capacity between wetland types (Eq. (2)). The weighting of wetland type (i.e. tropical vs. boreal) in each gridcell is calculated based on surface temperature (Eq. 3).

$$F(x) = r_{\text{CH4:C}} \cdot f_{\text{ecosvs}}(x) = \sigma(x)F_{\text{T}} + (1 - \sigma(x))F_{\text{B}}$$
⁽²⁾

where T(x) is the mean near-surface temperature between 1960 and 1990, and $T_{\text{max}} = 303.35$ °K. Equations 2 and 3 correct unintentional omissions in both equations as written in Hodson et al. (2011).

10 3.6.1 WETCHIMP setup

For WETCHIMP, we constrained the scaling ratios, F_T and F_B , by minimizing the error between our model fit, inverse modelling results (Spahni et al., 2011) and a regional flux estimate from the Hudson Bay Lowlands (Pickett-Heaps et al., 2011), yielding $F_T = 0.152$ and $F_B = 0.049$. Total global wetland and rice fluxes were con-

¹⁵ strained at 215.8 TgCH₄ yr⁻¹, wetland and rice fluxes north of 45° N at 39.6 TgCH₄ yr⁻¹ (Spahni et al., 2011), and wetland and rice fluxes from 50–60° N and 75–96° W at 2.3 TgCH₄ yr⁻¹ (Pickett-Heaps et al., 2011). These are different constraints than were used in Hodson et al. (2011).

R_{hetr} was calculated using the LPJ-WSL dynamic global vegetation model (DGVM),
 based on the LPJv3.1 DGVM (Sitch et al., 2003; Gerten et al., 2004). The monthly climatology inputs (precipitation, mean temperature, cloud cover, wet days) were taken from CRU TS3.1 and the non-gridded annual CO₂ concentration inputs to LPJ-WSL are described in Hodson et al. (2011). In addition, as in Hodson et al. (2011), soil texture was prescribed from the Food and Agriculture Organization (Zobler, 1986), using a 2 apil lawer budralaginal model with a total apil denth of 1.5 m. For according 2.

²⁵ using a 2-soil layer hydrological model with a total soil depth of 1.5 m. For scenarios 2 and 3, a 1000-yr spin up was implemented by recycling the first 30 yr of climate data (1901–1930) with pre-industrial CO₂ concentrations to equilibrate soil and vegetation

(3)

carbon pools, followed by a transient climate and CO₂ simulation running from 1901–2005. For scenarios 1, 4, 5, and 6, instead of using a repeating 1901–1930 climate, first a 12-monthly mean annual data set was created and repeated until equilibrium is reached. The default soil respiration equation in LPJ was used for all scenarios except s experiment 3.

We used the same temperature and moisture dependent equation as in Hodson et al. (2011), which allows the soil respiration to drop to zero when soil moisture is zero. For experiment 3, we used the soil moisture-respiration equation from Sitch et al. (2003) that fixes the soil respiration to a minimum of 0.25 in the upper one meter of soil, even when soil moisture drops to values below 0.25.

For all experiments except experiment 3, wetland extent (*A*) represents natural wetlands and lakes and is the original, monthly varying GIEMS inundation dataset processed to $0.5^{\circ} \times 0.5^{\circ}$ spatial resolution with rice growing regions removed (Leff et al., 2004). For experiments 1, 4, 5, and 6, this wetland area satellite product was averaged across all years from 1993–2004 to create a 12-month mean wetland area product. For experiment 3, a combined satellite and model product was used, which is described in Hodson et al. (2011).

3.7 ORCHIDEE

10

15

The ORCHIDEE model (Krinner et al., 2005) has been implemented with a wetland CH₄ emissions scheme. This version of ORCHIDEE has been previously used to simulate the evolution of wetland CH₄ emissions under future climate change (Koven et al., 2011) and to study the feedback between climate, atmospheric CH₄, and CO₂ (Ringeval et al., 2011). Simulations of ORCHIDEE for the current time period have also been performed and evaluated against top-down simulations to investigate the role of wetlands in the current atmospheric CH₄ concentration growth rate (I. Pison, personal communication, 2012).



4091

The wetland CH₄ emissions, $E_{CH_4}(g,t)$, are computed for each gridcell, g, and for each time step, t, through the following equation:

$$E_{CH_4}(g,t) = \sum_{WTP_i} S_{WTP_i}(g,t) D_{WTP_i}(g,t)$$

where S_{WTP_i} is the fraction of g covered by a wetland whose the water table position

- ⁵ is equal to WTP_i . D_{WTP_i} is the CH₄ flux (i.e. gCH_4 (m⁻² ofwetland) time⁻¹) for a wetland whose water table position is equal to WTP_i . S_{WTP_i} and D_{WTP_i} are respectively computed by (i) the coupling between a TOPMODEL (Beven and Kirkby, 1979) approach and ORCHIDEE, and (ii) the coupling between a (slightly modified) version of the Walter et al. (2001a) model and ORCHIDEE.
- ¹⁰ The main modification to the Walter et al. (2001a) model, as described in Ringeval et al. (2010), concerns the methanogenesis substrate. A fraction, α , of the natural labile carbon pool computed by ORCHIDEE is used to estimate the methanogenesis substrate. α has been optimized against three sites then extrapolated on all grid-cells sharing the same vegetation type (boreal, temperate or tropical).
- ¹⁵ In contrast to LPJ-WHyMe and its derivatives, ORCHIDEE did not implement wetland-specific PFTs. Instead, a fraction of the mean natural labile carbon pool over the gridcell is used to estimate the substrate supply.

For the computation of S_{WTP_i} in each gridcell, TOPMODEL allows distribution of the mean water deficit computed by ORCHIDEE according to the sub-grid topographic in-

- dex distribution. This leads to the diagnostic of the gridcell fraction with a null deficit. The mean deficit over the gridcell is computed from a gap to the field capacity (and not to the saturation which cannot be reached in ORCHIDEE). The saturated wetland extents are computed from these "field capacity extents" using a shift of the topographic index distribution into each gridcell. The value of this shift is the same for all gridcells
- and has been chosen to obtain a global coverage by wetlands close to 4%. The simulated wetland extent have been evaluated both through the induced modification on the simulated riverflows and against the GIEMS data (Ringeval et al., 2012a). The

(4)

TOPMODEL approach is used to simulate not only the saturated wetland extent but also the wetland extent with a WTP below the soil surface.

3.7.1 WETCHIMP setup

In the simulations performed for the WETCHIMP inter-comparison, the WTP_i values chosen for each gridcell are: 0, -3 and -9cm. A value of water table position equal to 0 means that the water is at the soil surface while negative values corresponds to water table position below the soil surface. Thus, for each time-step, three fractions of each gridcell are given: (i) a fraction covered by a saturated wetland, (ii) a fraction covered by a wetland with a mean WTP equal to -3cm (i.e. where the deficit is between 0 and -6cm) and (iii) a fraction covered by a wetland with a mean WTP equal to -9cm (i.e.

where the deficit is between -6 and -12 cm).

As in Ringeval et al. (2011) the wetland extent are corrected to subtract the systematic biases of the model by normalizing the mean yearly wetland extent to the GIEMS data (i.e. both the seasonal and year-to-year variability come from TOPMODEL). In the WETCHIMP simulations, a Q_{10} equal to 3 (close to the mean value in Ringeval et al., 2010) has been chosen for all grid-cells. As in Ringeval et al. (2012b), the reference temperature for methanogenesis is defined as the mean surface temperature computed by ORCHIDEE when forced by the 1960–1991 CRUNCEP climatology.

3.8 SDGVM

- The SDGVM (Sheffield Dynamic Global Vegetation Model: Woodward et al., 1995; Beerling and Woodward, 2001) was used in conjunction with a modified version of the Cao et al. (1996) wetland emissions model to perform the WETCHIMP simulations. The modelling setup follows the approach of Singarayer et al. (2011), however in that study an equilibrium approach was taken wherein the vegetation and CH₄ models were
- ²⁵ forced with averaged (30-yr) climatologies from a series of general circulation model simulations. For WETCHIMP, a transient approach was required whereby the models



were forced sequentially with monthly climatic inputs. The SDGVM and CH_4 models were therefore run in a similar manner as described by Hopcroft et al. (2011) in which a transient setup of the model is described.

SDGVM requires monthly mean inputs of surface air temperature, precipitation and relative humidity, as well as a map of soil texture and global annual mean atmospheric CO₂ concentrations. The CRU vapour pressure was converted to relative humidity using standard formulae, whilst the temperatures and precipitation were used without modification.

In the CH₄ model, the supply of carbon substrate for methanogenesis is taken to scale with 1 m soil heterotrophic respiration as simulated in SDGVM. 1 m is the is the model's soil depth. CH₄ production, M_{prod} , is then given by

 $M_{\text{prod}} = R_{\text{hetr}} P_0 f_{\text{w}}(\text{WTP}) f_T(T)$

where R_{hetr} is the soil heterotrophic respiration rate (gCm⁻²month⁻¹) from SDGVM, P_0 represents the fraction of decomposed matter converted to CH₄ under optimal conditions (0.47, see Cao et al., 1996). f_w and f_T are dimensionless scaling functions which parametrise the effects of water table position (WTP, in cm) and temperature (T in °C) on emission rates. These are given by:

$$f_{w}(WTP) = 0.383e^{0.096 \times WTP}, \quad WTP \le 10 \text{ cm}$$
$$f_{T}(T) = \frac{e^{0.0405 \times T}}{3.375}, \quad 5^{\circ}C < T \le 30^{\circ}C$$

20

where f_w is 1.0 for WTP > 10 cm and f_T is 1.0 for $T > 30^{\circ}$ C and 0.0 for $T \le 5.0^{\circ}$ C. f_w follows observations from Roulet et al. (1992) and Eq. (7) implies a Q_{10} value of 1.5. 90% of CH₄ produced is assumed to be oxidised. The water table position in each gridcell is calculated from the SDGVM simulated 1 m total soil moisture content using the relations from Cao et al. (1996) for tundra (their Eqs. 15 and 16) and a constant global soil porosity.

(5)

(6)

(7)

SDGVM also includes the biogeochemical coupling between the above- and below-ground carbon and nitrogen cycles. This is an important feature of modelling realistic changes in land surface primary production which depend on, and should be in line with, realistic biological and anthropogenic nitrogen fixation rates (Hungate et al., 2003). In SDGVM, litter production influences soil carbon and nitrogen pools via the Century soil nutrient cycling model (Parton et al., 1993), which in turn feedback to influence the primary production of vegetation; details are provided elsewhere (Beerling and Woodward, 2001).

3.8.1 WETCHIMP setup

- ¹⁰ The model output saved from SDGVM simulations includes annual NPP, soil carbon content, and monthly heterotrophic respiration, soil moisture content, and GPP. The monthly outputs of CH₄ emissions and water table position were saved from the CH₄ model. The experiment protocol called for monthly and annual maximum wetland area (mmax_weta, amax_weta). Wetland area in this model is not used by the emissions
- ¹⁵ model, which is instead a function of water table position (see Eq. 5). CH₄ emitting area could be used as a proxy for wetland area, but this would include gridcells with water-table far from the surface and with very small CH₄ fluxes. Since most of the other models in the inter-comparison were parameterized using inundated area from satellite observations, the SDGVM output was tailored to be somewhat comparable.
- The wetland area was here taken as simulated inundated area. Since there is no subgrid hydrology in the model, in each grid cell this area will either be 0 or the area of the gridcell. However, the model includes a correction for sub-grid orography based on ETOPO5 dataset which is applied to CH₄ emissions. This correction was also applied to the calculated inundation area. Two climate-dependent conditions on CH₄ emissions
- ²⁵ are also currently used within the model: (i) the monthly air temperature must be above 5° C, and (ii) if the temperature in a given gridcell during the current year is always > 0°C, then in a given month, the evapo-transpiration must not exceed precipitation.



These two conditions were both also used to correct the simulated inundated area, so that it is more directly relevant to the simulated CH_4 emissions.

3.9 UVic-ESCM

The UVic Earth System Climate Model (Weaver et al., 2001; Meissner et al., 2003) is an intermediate complexity climate model consisting of several coupled components: an energy-moisture balance atmospheric model, a three-dimensional ocean general circulation model, a dynamic-thermodynamic sea ice model and a land surface scheme which includes a representation of vegetation dynamics. The model was recently modified to include a representation of permafrost and global wetlands (Avis et al., 2011).

- Wetlands in the UVic model are determined using empirically derived threshold values for unfrozen soil moisture content and terrain slope so that wetlands are present where ground is sufficiently wet and flat (Kaplan, 2002). Wetlands are either "on or off" in a particular gridcell. If they satisfy the soil moisture criterion they are "on" and they occupy the fraction of the gridcell with the requisite terrain slope. Organic and mineral soil
- properties were specified by using the ISLSCP-II data sets as the model is presently incapable of generating the observed high soil carbon values in northern high latitudes (none of the participating models is able to couple soil biogeochemical with thermal characteristics yet).

3.9.1 WETCHIMP setup

- The UVic model is nominally run in a fully coupled configuration with coupling between atmospheric, ocean, land surface and other model components. For the purpose of participating in WETCHIMP, the land surface scheme was decoupled from the other model components and run in an offline configuration. This offline configuration uses the monthly CRU data to drive the land surface scheme. The model smoothly interpo-
- ²⁵ lates between these CRU fields to obtain data for a particular timestep. Some of the atmospheric code from the coupled model's energy moisture balance model (Weaver



et al., 2001) was retained to calculate factors such as incoming solar radiation, which need to be specified to the land surface scheme and was not specified in the monthly CRU datasets.

Surface air temperature, precipitation and relative humidity were specified from the CRU data, while surface winds and the diurnal temperature range were drawn from the NCEP reanalysis. Incoming longwave and shortwave radiation are calculated within the model's atmospheric module.

The land surface scheme also has the capacity to simulate vegetation dynamics using the TRIFFID dynamic vegetation model. As vegetation parameters were derived based on the coupled model climate, for the purposes of the WETCHIMP simulations, the vegetation distribution and characteristics were fixed and set equal to their mean year-1900 values from an equilibrated version of the coupled model, rather than adjusting vegetation parameters to fit the CRU climate data. Monthly mean fields of vegetation fraction, leaf area index, vegetation height and litterfall are then obtained from the

mean year-1900 model output. The vegetation dynamics are consequently switched off for the offline run, and vegetation parameters for a given model timestep are specified by smoothly interpolating between these monthly fields. Non-plant competition based vegetation parameters do remain interactive in the model. For example, plant stomata still open and close in response to factors like changing soil moisture or atmospheric
 CO₂ concentration.

For the equilibrium run, the model was forced repeatedly with 1901-1931 data for 2000 yr. This time period was found to be sufficient to allow for equilibration of soil moisture and temperature variables. When applying the CO₂, temperature and moisture runs, the 1901-1931 spinup period was repeated for an additional 2000 yr to allow a new equilibrium to be established.

3.10 UW-VIC

25

The University of Washington team used the Variable Infiltration Capacity (VIC) model, version 4.1.2, with some extensions specifically tailored for the modelling of boreal



peatlands described in Bohn et al. (2007) and Bohn and Lettenmaier (2010). UW-VIC is a large-scale hydrologic model that balances the water and energy budgets of the land surface at an hourly time step and spatial resolutions ranging from 1° to 5 km. Most of the model physics are described in Liang et al. (1994). Land cover is repre-

- ⁵ sented as a collection of "tiles", each containing a different plant functional type, overlaying a single soil column divided into 3 hydrologic layers, down to varying depths, but generally no deeper than 3 m. While UW-VIC does not track the storage of carbon in biomass, it computes NPP via a scheme taken from the BETHY model (Knorr, 2000). The seasonal cycle of LAI is prescribed at each gridcell based on the MODIS
- LAI product (Myneni et al., 2002). Stomatal resistance is a function of day length, temperature, soil moisture, and vapor pressure deficit. UW-VIC models permafrost and the soil temperature profile via the finite difference scheme of Cherkauer and Lettenmaier (1999) with an exponential node distribution down to 50 m depth and a no-flux bottom boundary condition. Thermal properties of organic soil are also taken into account
- (Farouki, 1981). To account for dynamic surface water storage (lakes and seasonally-flooded wetlands) UW-VIC's lake/wetland model was employed (Bowling and Letten-maier, 2010). This feature allocates one land cover tile to contain a combination of a lake (representing all lakes in the gridcell) and its surrounding catchment. Within the tile, the inundated area fraction is dynamic, changing as a function of storage and
- ²⁰ bathymetry. Thus, while the maximum possible extent of inundation within a gridcell is prescribed, the actual inundated extent is a dynamic function of environmental conditions. In the exposed portion of the tile, the water table position is assumed to have a distribution based on peatland microtopography: the peatland consists of a mix of hummocks (or ridges) and hollows (or pools), with the peat underneath hummocks up
- to 70 cm thicker than under the deepest points of the hollows. The fraction of peatland covered by hummocks is a calibrated parameter. Local water table position under any given point is computed as a function of soil moisture via the formulation described in Frolking et al. (2002). Methane emissions were computed for the lakes, inundated wetlands, and each point in the water table position distribution in the exposed wetlands



as a function of water table position, soil temperature, and NPP via the model of Walter and Heimann (2000). A lake emission rate of $375 \text{ mgCH}_4 \text{ m}^{-2} \text{ d}^{-1}$ is assumed for during the ice-free season and half that rate during ice-covered season.

3.10.1 WETCHIMP setup

- 5 For these simulations, each gridcell was separated into two parts: an upland fraction, underlain by mineral soils, with soil textures supplied by the FAO Digital Soil Map of the World (Batjes, 1997); and a lake/wetland fraction, underlain by peat soils, with peat depths given by the database of Sheng et al. (2004) and other characteristics taken from Letts et al. (2000). Simulations were run separately for each portion of the gridcell. The lake/wetland portion of each gridcell was determined as the superset of the Sheng et al. (2004) peatland map; wetlands, wet tundra, and croplands (so that nearby lakes could have a surrounding catchment) given by the Bartalev et al. (2003) land cover classification; and lakes given by the Global Lake and Wetland Database (GLWD; Lehner and Döll, 2004. Bathymetries for the lake/wetlands were estimated by
- combining lake size distributions from the GLWD; average lake depths from literature for bog pools, Arctic thaw lakes, and other boreal lakes; and topography of surrounding wetlands from the ASTER (Hayakawa et al., 2008) and STRM (Farr and Kobrick, 2000) DEMs. When lake storage increased beyond the bounds of the "permanent" lake, it was allowed to flood the surrounding wetlands, with drainage rate controlled by a calibrated
- parameter. Both this parameter and the area fraction of hummocks within the peatland 20 were calibrated to optimize the match with global inundation datasets. For the optimized runs (Experiment 3), the global daily AMSR-E/QuickSCAT-based dataset of Schroeder et al. (2010) was used; for all other runs, the GIEMS dataset was used (Prigent et al., 2007; Papa et al., 2010). Parameters for the CH₄ emissions model were calibrated to
- optimize the in-situ observations of Glagolev et al. (2010) across West Siberia.

)iscussion Pa	GMDD 5, 4071–4136, 2012		
per	WETC metho	CHIMP dology	
Discus	R. Wan	ia et al.	
sion F	Title	Page	
aper	Abstract	Introduction	
_	Conclusions	References	
Discu	Tables	Figures	
Issio	I	►I	
n Pap	•	•	
ber	Back	Close	
_	Full Scre	en / Esc	
Discuss	Printer-frier	dly Version	
ion F	Interactive	Discussion	
aper	œ	BY	

4 Results and discussion

We provide two conceptual overviews of the participating models highlighting similarities as well as differences between the chosen approaches. These overviews are

deigned to assist discussions of the differences in modelling results (Fig. 8 and in
 Melton et al., 2012), but they also represent the first attempt at conceptually describing the state-of-the-art approaches used in wetland extent and wetland CH₄ modelling. For the conceptual figures describing the modelling approaches, we defined two variables of interest: the CH₄ producing area (Fig. 6) and CH₄ flux (Fig. 7). We used these metrics to explore the dominant processes responsible for differences between the models.

10 4.1 CH₄ producing area

15

We use the term " CH_4 producing areas" (MPAs) to include all terrestrial areas that may produce CH_4 biogenically. We include wet mineral soils, presently only simulated by LPJ-Bern, that are proposed to function as a CH_4 source or sink depending on the soil moisture level. The participating models use a large diversity of methods to determine MPAs (Fig. 6). We identified the features of the models that we found most strongly controlled the MPAs and visualized the concepts of the models.

The starting points to locate MPAs are either "Prescribed constant wetland extents", "Remotely-sensed inundation" or a "Hydrological model" (Fig. 6). The simplest case of estimating MPAs is where "Prescribed constant wetland extents" are taken from annu-

- ally non-varying distribution maps, and are used without modifications. This approach is applied by LPJ-WHyMe and LPJ-Bern (peatlands) that use the northern peatland map from NCSCD (Tarnocai et al., 2007, 2009), and IAP-RAS that uses the Olson data set for global MPA location. A similar approach takes seasonally varying, "Remotely-sensed inundation" to prescribe MPAs. LPJ-Bern wetlands uses an averaged monthly
 mean extent from GIEMS, while LPJ-WSL (all experiments except 3) uses the GIEMS
- dataset without modification. A further step up the complexity ladder is LPJ-Bern wetsoils, the most basic wetland extent which uses model output. LPJ-Bern wetsoils uses a



"Hydrological model" to derive "Soil moisture" and "Unsaturated" MPAs. "Unsaturated" means that the pore-space in the soil is not completely filled with water. This could be the case when – even though a water table position is calculated – it is below the surface or when the soil moisture is estimated as a homogeneous average over the

- ⁵ soil depth and its values do not reach saturation. Next, we include approaches that comprise of "Topography" in addition to "Hydrological model" as an additional factor to locate "Unsaturated" areas (UVic-ESCM). SDGVM uses a similar approach to UVic-ESCM but simulates "Water table position" before determining "Unsaturated" as well as "Saturated/inundated" MPAs. CLM4Me, DLEM, ORCHIDEE, UW-VIC and LPJ-WSL
- (Exp. 3) all use "Remotely sensed inundation" (GIEMS) data in their approaches, but they use these data in different ways: e.g. ORCHIDEE guides the mean simulated wetland extent over the 1993–2004 period and CLM4Me uses the GIEMS data set to invert for parameters that allow the hydrological state (i.e. water table depth and runoff) to be used to determine wetland extent. More details on the use of GIEMS can be found in
- the description of each model (Sect. 3). Once the "Water table position" is determined, CLM4Me, DLEM, ORCHIDEE, and UW-VIC identify the MPAs that are either "Unsaturated" or "Saturated/inundated" while LPJ-WSL (Experiment 3-opt) determines MPAs which are "Saturated/inundated" only. The UW-VIC model is the most complex model and takes advantage of all of the features included in Fig. 6, using the fractional peatland cover by Sheng et al. (2004) only as maximal boundaries, rather than as a fixed
- Iand cover by Sheng et al. (2004) only as maximal boundaries, rather than as a fixed map.

From the conceptual overview (Fig. 6) one can see that the only two models that simulate MPAs without the guidance of other wetland-relevant observations are the UVic-ESCM and the SDGVM. The difference between those two models is that the

²⁵ UVic-ESCM uses only soil moisture (as well as topography) to find "Unsaturated" areas, whereas the SDGVM also calculates the water table position to find "Unsaturated" as well as "Saturated/inundated" areas. As the UVic-ESCM model was designed to identify wetland areas, not specifically MPAs (it presently has no CH₄ model), the model



uses gridcell mean *unsaturated* soil moisture values and terrain slope as a mean of approximating *saturated* areas.

4.2 CH₄ flux

The second variable we analyze in detail is the "CH₄ flux" calculation by nine out of the
ten participating models – the UVic-ESCM does not yet include CH₄ fluxes. Figure 7 shows which pools and processes models consider to determine CH₄ flux. All models but IAP-RAS base their CH₄ production on some kind of carbon flux, where two groups can be distinguished – one that uses "Wetland PFTs" and one that uses "Upland PFTs" to simulate vegetation net primary production (NPP); only DLEM utilizes NPP (and also GPP) simulated by both types of PFTs for CH₄ production. The UW-VIC model uses NPP in the algorithm for CH₄ production, ORCHIDEE uses a fraction of the most labile of the "Litter + soil C" pool and all remaining models use "Heterotrophic respiration" as the basis for their "CH₄ production" (see also Table 5). LPJ-Bern, LPJ-WHyMe and

- DLEM add "Exudates"-derived carbon to the "Heterotrophic respiration" calculation. All models calculate CH_4 production and half of the models consider "Transport" mechanisms such as ebullition, plant-mediated transport, and diffusion to derive " CH_4 fluxes" (Fig. 7, Table 5). This table also gives insight into which models include oxidation of soil-derived CH_4 and how they combine production and oxidation rates to simulate the final net CH_4 flux. Only three of the models include atmospheric CH_4 ox-
- ²⁰ idation (CLM4Me, DLEM, LPJ-Bern wetsoils). Thus, we include atmospheric oxidation in Table 5 for completeness, but we excluded atmospheric oxidation from model results as far as possible in order to compare gross CH_4 fluxes. The separation of gross CH_4 fluxes and atmospheric CH_4 uptake fluxes was not completely feasible in CLM4Me as the CH_4 uptake occurs implicitly in the reaction-transport solution, although for Melton
- et al. (2012) an estimate was determined to allow better comparison between models. Across the models, the complexity of equations for CH_4 production covers a wide range. The IAP-RAS model is the simplest model relating CH_4 production only to temperature, whereas all other models use some estimate of the available carbon flux rate.



All but two models (IAP-RAS and LPJ-WSL) include some kind of soil-derived CH_4 oxidation, which can be as simple as reducing production by a fixed fraction (SDGVM) or by including up to five different terms in the equation (Table 5).

4.3 Methane producing area and methane flux

- ⁵ Mean annual maximum extent of wetland area and mean annual CH₄ fluxes for Experiment 1-equil are shown in Fig. 8. Total MPAs and CH₄ emissions for each model are listed in the bottom left corner of each sub-figure. Two of the models are regional models (LPJ-WHyMe and UW-VIC), the rest are global models of which the UVic-ESCM does not simulate CH₄ fluxes. CLM4Me, DLEM, LPJ-Bern (without mineral soils), LPJ-WSL, and ORCHIDEE share similar wetland distributions due to their varying degrees of reliance on remotely-sensed inundation data (see Fig. 6). The similarity of approaches is also reflected in the total MPA of these models (CLM4Me: 6.8 × 10⁶, DLEM: 7.9 × 10⁶, LPJ-Bern without mineral soils: 7.9 × 10⁶, LPJ-WSL: 7.4 × 10⁶, and ORCHIDEE: 9.2 × 10⁶ km²). Two models (IAP-RAS and LPJ-Bern with wet mineral soils)
- stand out visually because of their large areas of 80–100 % MPA per gridcell. The IAP-RAS model uses a binary approach – either a gridcell is a wetland or it is not – resulting in a total MPA of 20.3×10^6 km², which is an entirely prescribed amount. Given the definition of the wet mineral soils CH₄ source, the LPJ-Bern wet mineral soils map should be interpreted as a map of "potential CH₄ emissions in at least one month per year".
- Since LPJ-Bern does not use a sub-gridcell hydrology for wet mineral soils to estimate the CH₄ production capacity, the wet mineral soils component of LPJ-Bern is also a binary approach. However, the extent of wet mineral soils in a given gridcell can be reduced by peatland area and inundated area so that they jointly sum to 100 %, but as soon as a gridcell qualifies as wet mineral soils, the MPA of that gridcell is 100 %. This approach leads to the largest total MPA of 76.6 × 10⁶ km² of the WETCHIMP models.

The only two models that use an explicit water balance scheme to simulate wetland extent without relying on wetland or inundation data sets are the SDGVM and the UVic-ESCM (Fig. 8). They show a similar spatial distribution but differ notably in Eastern



Siberia, the western United States and northern Canada. SDGVM uses soil moisture content to first diagnose water table position and then MPAs, whereas UVic-ESCM uses soil moisture directly to derive MPAs. The differences between these two models could be related to parameterization of permafrost (present in UVic-ESCM, but absent in SDGVM) and other soil physics or hydrology parameters, i.e. hydraulic conductivity,

⁵ In SDGVM) and other soil physics or hydrology parameters, i.e. hydraulic conductivity, porosity, etc. Further, the percentages of gridcells covered by wetlands are generally higher for SDGVM than for the UVic-ESCM leading to a higher overall wetland area of 34.9×10^6 km² vs. 14.9×10^6 km², respectively.

Of the two regional models, LPJ-WHyMe uses a fixed peatland distribution (Fig. 8e) whereas the UW-VIC model uses the most sophisticated method of all participating models to simulate saturated and unsaturated wetland areas in the West Siberian Lowlands (Fig. 8j). A comparison focused on the West Siberian Lowlands is planned to evaluate the differences between a highly regionalized model like the UW-VIC model and the rest of the WETCHIMP models (personal communication Ted Bohn, June 2012).

- ¹⁵ Simulated CH₄ fluxes of nine of the participating models are shown on the right hand side in Fig. 8. Methane fluxes ranged from 0 to over 250 gCH₄ m⁻² of wetland yr⁻¹ with CLM4Me, DLEM, LPJ-WSL, ORCHIDEE, and UW-VIC showing widespread high fluxes (Fig. 8a, b, g, h and j) and IAP-RAS, LPJ-Bern, LPJ-WHyMe, and SDGVM showing low fluxes (Fig. 8c, d, e and i). Of the five models that show widespread high fluxes, three
 ²⁰ base their CH₄ flux on upland PFTs (CLM4Me, LPJ-WSL, ORCHIDEE), one on wetland PFTs (UW-VIC) and one on both (DLEM) (Fig. 7). Of the four models that show low
- CH_4 fluxes, two rely on wetland PFTs (LPJ-Bern, LPJ-WHyMe), one on upland PFTs (SDGVM) and one does not rely on PFTs at all (IAP-RAS). This could indicate a general tendency to higher CH_4 fluxes when upland PFTs instead of wetland PFTs are used
- to simulate NPP. Some of the models show higher fluxes in the tropics than in the extra-tropics (CLM4Me, IAP-RAS, SDGVM), whereas others show equally high fluxes (DLEM, LPJ-Bern, LPJ-WSL, ORCHIDEE), which may be linked to the model-inherent temperature sensitivities of e.g. NPP, heterotrophic respiration or CH₄ production, but



without specific parameter sensitivity studies it is impossible to evaluate where the differences arise from.

The patterns of CH₄ fluxes do not always match the pattern of wetland distribution (e.g. compare wetland area and CH₄ fluxes for the Amazon for CLM4Me). Total
CH₄ emissions for a gridcell are calculated as the product of fluxes and wetland area (except for CLM4Me, which also considers production in upland soils. Therefore, models may simulate similar global CH₄ emissions with completely different MPAs and CH₄ fluxes (e.g. CLM4Me: 186 TgCH₄yr⁻¹ vs. SDGVM: 183 TgCH₄yr⁻¹ or IAP-RAS: 154.2 TgCH₄yr⁻¹ vs. LPJ-Bern: 156.6 TgCH₄yr⁻¹ vs. LPJ-WSL: 151.5 TgCH₄yr⁻¹)
(Fig. 8). The comparability of these simulated global CH₄ emissions emphasizes the fact that most models are tuned to some degree towards a global total global CH₄ emissions value, which allows the MPAs to vary more between the models than global CH₄ emissions. As highlighted in Melton et al. (2012), the fact that the models agree fairly well on global CH₄ emissions with very different MPAs and CH₄ fluxes underlines

the importance of regional-scale observational estimates to constrain this dichotomy.

5 Summary and conclusions

WETCHIMP is the first multi-model comparison of wetland extents and wetland CH_4 emissions. Our analysis demonstrates how diverse modelling approaches, wetland definitions, and wetland extents can be, while still leading to comparable values of global

- ²⁰ CH₄ emissions. In terms of modelling CH₄ producing areas (MPAs), there are three main approaches (i) the fixed MPA (IAP-RAS, LPJ-Bern (peatlands and wetlands), LPJ-WHyMe), (ii) the guided MPA (CLM4Me, DLEM, LPJ-WSL, ORCHIDEE, UW-VIC) and (iii) the fully simulated MPA (UVic-ESCM, SDGVM, LPJ-Bern wetsoils). Total MPA can vary significantly between models depending on their definitions, which also influences
- $_{25}$ CH₄ fluxes, but does not have as much impact on the global CH₄ emissions. Achieving similar global CH₄ emissions with very different MPA distributions also means that the CH₄ fluxes between the models differ greatly. A wide range of parameterization



complexity is used to simulate CH_4 fluxes in the participating models, which influences not just the present day flux but also its sensitivity to climate factors.

Each individual model's approach needs to be considered carefully when interpreting results, especially past and future climate change experiments or sensitivity ex-

- ⁵ periments such as those that were conducted as part of WETCHIMP (Melton et al., 2012). There are several factors that need to be kept in mind: A fixed wetland distribution as used by some models or a seasonally varying distribution based on presently observed patterns is unlikely to be representative of past or future conditions. Another limitation is the absence of wetland specific PFTs in most models. Models that lack wetland energing PFTs (i.e. CLMAME, LPL, WELL, SPC) Matching and an action of the presentation.
- wetland specific PFTs (i.e. CLM4Me, LPJ-WSL, SDGVM, LPJ-Bern non-peatland, OR-CHIDEE, IAP-RAS) may overestimate NPP due to an unrealistic lack of plant stress that would be caused by inundation or nutrient limitation. We expect these models to show different responses to changes in temperature, precipitation and CO₂ fertilization than the models that include wetland specific PFTs (i.e. DLEM, LPJ-WHyMe,
- LPJ-Bern peatlands, UW-VIC). For example, changes in precipitation will affect wetland specific PFTs that grow under inundated conditions differently than upland plants. Also, the effect of CO₂ fertilization on wetland plants is still unclear (Berendse et al., 2001; Heijmans et al., 2001, 2002a,b; Boardman et al., 2011) and therefore wetland NPP under CO₂ fertilisation calculated by models that include wetland specific PFTs
 remains highly uncertain.

There are features that are still missing, or are crudely represented, in almost all of the models, partially due to the difficulties of simulating small-scale processes in large-scale models. Such features include (i) lateral transport of water and groundwater dy-namics, and explicit treatments of floodplains and mangroves; (ii) plant nutrients (nitro-

gen, phosphorus and sulphur) and their interactions (presently only SDGVM, DLEM, and CLM4Me include carbon-nitrogen interactions); (iii) microtopographical features such as lawns, hollows or hummocks and their impacts upon overall CH₄ dynamics; (iv) vertically-resolved carbon pools and soil organic matter remineralization modelling; (v) permafrost-preserved carbon; (vi) feedbacks between peat or carbon dynamics and



thermal and hydrological processes in soil; (vii) hydrology affected by thawing permafrost; and (viii) wetland specific vegetation (improvements for boreal peatlands, introduction of tropical wetland PFTs).

- WETCHIMP provides a the first multi-model platform to explore the current knowl-5 edge, recent improvements, and necessary future developments of models simulating wetland extents and wetland CH_4 emissions. The design of future iterations of WETCHIMP will be focused on analyzing and understanding the different uncertainties and sensitivities of participating models with the goal of greatly improving the performance of the models for both wetland and wetland CH₄ modelling. The simulations conducted in WETCHIMP are available (http://arve.epfl.ch/pub/wetchimp, please contact J. R. Melton for immediate access) and their use is encouraged to advance re-
- search in this area.

Acknowledgements. The authors are grateful to the COST Action ES0805 TERRABITES for providing support for the WETCHIMP workshop. We thank Sylvia Houston and Olga Petrikova for assistance with organizing the project meeting and Kristen Krumhardt for assistance producing the figures. J. R. M. was supported by the Swiss Ministry for Research and Education

- (grant C09.0054). J. O. K. acknowledges support from the Swiss National Science Foundation (grant PP0022_119049 and PP00P2_139193). The contributions of W. J. R. and Z. M. S. were supported by the Director, Office of Science, Office of Biological and Environmental Research,
- Climate and Environmental Science Division, of the US Department of Energy under Contract 20 No. DE-AC02-05CH11231 to Berkeley Lab (IMPACTS and C-Climate Uncertainties projects). The contributions of T. J. B. and D. P. L. were supported by NASA's ROSES program, grant NNX08AH97G. The contribution of Z. C. Y. was supported by a US NSF grant (ARC-1107981) and a grant from The Max Planck Society (Germany). A. V. E. and S. N. D. were supported
- by the the President of Russia grant 5467.2012.5, by the Russian Foundation for Basic Re-25 search, and by the programs of the Russian Academy of Sciences. The contributions of R. S. and S. Z. were supported by the Swiss National Science Foundation and by the European Research Council advanced grant MATRICs (ERC grant agreement no. 226172) under the European Community's Seventh Framework Programme. P. O. H. is supported by a NERC UK grant NE/I010912/1 and previously through NERC UK/INSU France QUEST-DESIRE project.
- **Discussion** Paper GMDD 5, 4071-4136, 2012 **WETCHIMP** methodology **Discussion** Paper R. Wania et al. **Title Page** Abstract Introduction Conclusions References **Discussion** Paper Tables **Figures** 14 Back Full Screen / Esc Discussion **Printer-friendly Version** Interactive Discussion Paper

15

E. L. H. was supported by ETH Competence Center Environment and Sustainability's grant MAIOLICA.

References

15

Aselman, I. and Crutzen, P. J.: Global distribution of natural freshwater wetlands and rice pad-

- dies, and their net primary productivity, seasonality and possible methane emissions, J. Atmos. Chem., 8, 307–358, 1989. 4074, 4082
 - Avis, C. A., Weaver, A. J., and Meissner, K. J.: Reduction in areal extent of high-latitude wetlands in response to permafrost thaw, Nat. Geosci., 4, 444–448, 2011. 4075, 4076, 4095
- Bartalev, S. A., Belward, A. S., Erchov, D. V., and Isaev, A. S.: A new SPOT4-VEGETATION derived land cover map of Northern Eurasia, Int. J. Remote Sens., 24, 1977–1982, 2003. 4098, 4122
 - Batjes, N. H.: A world dataset of derived soil properties by FAO-UNESCO soil unit for global modelling, Soil Use Manage., 13, 9–16, 1997. 4098, 4122
 - Beerling, D. J. and Woodward, F. I.: Vegetation and the Terrestrial Carbon Cycle: Modelling the first 400 Million Years, Cambridge University Press, Cambridge, 2001. 4092, 4094
 - Beerling, D. J., Fox, A., Stevenson, D. S., and Valdes, P. J.: Enhanced chemistryclimate feedbacks in past greenhouse worlds, P. Natl. Acad. Sci. USA, 108, 9770–9775, doi:10.1073/pnas.1102409108, 2011. 4075

Berendse, F., Breemen, N. V., Rydin, H., Buttler, A., Heijmans, M., Hoosbeek, M. R., Lee, J. A.,

Mitchell, E., Saarinen, T., Vasander, H., and Wallen, B.: Raised atmospheric CO₂ levels and increased N deposition cause shifts in plant species composition and production in *Sphagnum* bogs, Glob. Change Biol., 7, 591–598, 2001. 4105

Beven, K. J. and Kirkby, M. J.: A physically based, variable contributing area model of basin hydrology, Hydrol. Sci. J., 24, 43–69, 1979. 4091

²⁵ Boardman, C. P., Gauci, V., Watson, J. S., Blake, S., and Beerling, D. J.: Contrasting wetland CH₄ emission responses to simulated glacial atmospheric CO₂ in temperate bogs and fens, New Phytol., 192, 898–911, 2011. 4105

Bohn, T. J. and Lettenmaier, D. P.: Systematic biases in large-scale estimates of wetland methane emissions arising from water table formulations, Geophys. Res. Lett., 37, L22401,

³⁰ doi:10.1029/2010GL045450 2010. 4075, 4097

2						
	GMDD					
	5, 4071–4	5, 4071–4136, 2012				
)						
	WETC metho	CHIMP dology				
!	R. Wan	ia et al.				
		_				
,	Title	Page				
	Abstract	Introduction				
	Conclusions	References				
2	Tables	Figures				
	14	۶I				
)		•				
	Back	Close				
	Full Scre	en / Esc				
!						
	Printer-frier	dly Version				
	Interactive	Discussion				
	œ	D BY				

Jiscussion Pape

Discussion Pape

Discussion Pape

Discussion Paper

4108

- Bohn, T. J., Lettenmaier, D. P., Sathulur, K., Bowling, L. C., Podest, E., McDonald, K. C., and Friborg, T.: Methane emissions from western Siberian wetlands: heterogeneity and sensitivity to climate change, Environm. Res. Lett., 2, 259–268, 2007. 4075, 4097
- Bowling, L. C. and Lettenmaier, D. P.: Modeling the effects of lakes and wetlands on the water balance of Arctic environments, J. Hydrometeorol., 11, 276–295, 2010. 4097
- balance of Arctic environments, J. Hydrometeorol., 11, 276–295, 2010. 4097
 Cao, M. K., Marshall, S., and Gregson, K.: Global carbon exchange and methane emissions from natural wetlands: application of a process-based model, J. Geophys. Res., 101, 14399– 14414, 1996. 4074, 4075, 4092, 4093

Chen, G., Tian, H., Zhang, C., Liu, M., Ren, W., Zhu, W., Chappelka, A., Prior, S. A., and

- Lockaby, G.: Drought in the southern United States over the last century: variability and its impacts on terrestrial ecosystem productivity and carbon storage, Clim. Change, 114, 379–397, doi:10.1007/s10584-012-0410-z, 2012. 4076, 4082, 4083
 - Cherkauer, K. A. and Lettenmaier, D. P.: Hydrologic effects of frozen soils in the upper Mississippi River basin, J. Geophys. Res.-Atmos., 104, 19599–19610, 1999. 4097
- ¹⁵ Christensen, T. R. and Cox, P.: Response of methane emission from arctic tundra to climatic change: results from a model simulation, Tellus B, 47, 301–309, 1995. 4074, 4084 Christensen, T. R., Prentice, I. C., Kaplan, J., Haxeltine, A., and Sitch, S.: Methane flux from northern wetlands and tundra – an ecosystem source modelling approach, Tellus B, 48,
 - 652–661, 1996. 4074 Coe. M. T.: Simulating continental surface water
- ²⁰ Coe, M. T.: Simulating continental surface waters: an application to Holocene northern Africa, J. Clim., 10, 1680–1689, doi:10.1175/1520-0442(1997)010%3C1680:SCSWAA%3E2.0.CO;2, 1997. 4075
 - Coe, M. T.: A linked global model of terrestrial hydrologic processes: simulation of modern rivers, lakes, and wetlands, J. Geophys. Res., 103, 8885–8899, 1998. 4075
- ²⁵ Curry, C. L.: Modelling the soil consumption of atmospheric methane at the global scale, Global Biogeochem. Cy., 21, gB4012, doi:10.1029/2006GB002818, 2007. 4075
 - Curry, C. L.: The consumption of atmospheric methane by soil in a simulated future climate, Biogeosciences, 6, 2355–2367, doi:10.5194/bg-6-2355-2009, 2009. 4075
 - Dentener, F., Drevet, J., Lamarque, J., Bey, I., Eickhout, B., Fiore, A., Hauglustaine, D.,
- ³⁰ Horowitz, L., Krol, M., and Kulshrestha, U.: Nitrogen and sulfur deposition on regional and global scales: a multimodel evaluation, Global Biogeochem. Cy., 20, gB4003, doi:10.1029/2005GB002672, 2006. 4122



- ETOPO: 2-minute gridded global relief data (ETOPO2v2), US Department of Commerce, National Oceanic and Atmospheric Administration, National Geophysical Data Center, available at: http://www.ngdc.noaa.gov/mgg/fliers/06mgg01.html (last access: 14 August 2012), 2006. 4122
- 5 FAO/IIASA/ISRIC/ISSCAS/JRC: Harmonized World Soil Database (version 1.1), Tech. rep., FAO, Rome, Italy and IIASA, Laxenburg, Austria, 2009. 4086, 4122
 - Farouki, O. T.: The thermal-properties of soils in cold regions, Cold Reg. Sci. Technol., 5, 67–75, 1981. 4097
 - Farr, T. G. and Kobrick, M.: Shuttle radar topography mission produces a wealth of data, EOS Trans. Am. Geophys. Union, 81, 583–585, 2000. 4098

10

Fries, R. D., Hansen, M., Townshend, J., and Sohlberg, R.: Global land cover classifications at 8 km spatial resolution: the use of training data derived from Landsat imagery in decision tree classifiers, Int. J. Remote Sens., 19, 3141–3168, 1998. 4122

Frolking, S., Roulet, N. T., Moore, T. R., Lafleur, P. M., Bubier, J. L., and Crill, P. M.: Modeling

- 15 seasonal to annual carbon balance of Mer Bleue Bog, Ontario, Canada, Global Biogeochem. Cy., 16, doi:10.1029/2001GB001457 2002. 4097
 - Fung, I. Y., John, J., Lerner, J., Matthews, E., Prather, M., Steele, L. P., and Fraser, P. J.: Threedimensional model synthesis of the global methane cycle, J. Geophys. Res., 96, 13033– 13065, 1991. 4074
- Gedney, N., Cox, P. M., and Huntingford, C.: Climate feedback from wetland methane emissions, Geophys. Res. Lett., 31, L20503, doi:10.1029/2004GL020919, 2004. 4075
 Gerber, S., Joos, F., Brugger, P., Stocker, T., Mann, M., Sitch, S., and Scholze, M.: Constraining temperature variations over the last millennium by comparing simulated and observed atmospheric CO₂, Clim. Dynam., 20, 281–299, doi:10.1007/s00382-002-0270-8, 2003. 4085
- Gerten, D., Schaphoff, S., Haberlandt, U., Lucht, W., and Sitch, S.: Terrestrial vegetation and water balance – hydrological evaluation of a dynamic global vegetation model, J. Hydrol., 286, 249–270, 2004. 4085, 4089
 - Glagolev, M. V., Kleptsova, I. E., Filippov, I. V., Kazantsev, V. S., Machida, T., and Maksyutov, S.: Methane emissions from subtaiga mires of Western Siberia: the Standard Model Bc5, Moscow Univ. Soil Sci. Bull., 65, 86–93, 2010. 4098
- Bc5, Moscow Univ. Soil Sci. Bull., 65, 86–93, 2010. 4098
 Global Soil Data Task Group: Global gridded surfaces of selected soil characteristics (IGBP-DIS), Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge,



Tennessee, USA, available at: http://www.daac.ornl.gov (last access: 14 August 2012), 2000. 4122

- Hayakawa, Y. S., Oguchi, T., and Lin, Z.: Comparison of new and existing global digital elevation models: ASTER G-DEM and SRTM-3, Geophys. Res. Lett., 35, 19599–19610, 2008. 4098
- ⁵ Heijmans, M. M. P. D., Arp, W. J., and Berendse, F.: Effects of elevated CO₂ and vascular plants on evapotranspiration in bog vegetation, Glob. Change Biol., 7, 817–827, 2001. 4105
 - Heijmans, M. M. P. D., Klees, H., and Berendse, F.: Competition between Sphagnum magellanicum and Eriophorum angustifolium as affected by raised CO₂ and increased N deposition, Oikos, 97, 415–425, 2002a. 4105
- Heijmans, M. M. P. D., Klees, H., de Visser, W., and Berendse, F.: Response of a *Sphagnum* bog plant community to elevated CO₂ and N supply, Plant Ecol., 162, 123–134, 2002b. 4105
 Hodson, E. L., Poulter, B., Zimmermann, N. E., Prigent, C., and Kaplan, J. O.: The El Niño-Southern Oscillation and wetland methane interannual variability, Geophys. Res. Lett., 38, L08810, doi:10.1029/2011GL046861, 2011. 4075, 4076, 4088, 4089, 4090
- Hopcroft, P. O., Valdes, P. J., and Beerling, D. J.: Simulating idealized Dansgaard-Oeschger events and their potential impacts on the global methane cycle, Quaternary Sci. Rev., 30, 3258–3268, 2011. 4075, 4093
 - Hungate, B. A., Dukes, J. S., Shaw, M. R., Luo, Y. Q., and Field, C. B.: Nitrogen and climate change, Science, 302, 1512–1513, 2003. 4094
- ²⁰ IPCC: Emissions Scenarios, Special Report of the Intergovernmental Panel on Climate Change, Tech. Rep., Cambridge University Press, Cambridge UK, 2000. 4077
- ISLSCP-II: Global Gridded Surfaces of Selected Soil Characteristics for the International Satellite Land Surface Climatology Project (ISLSCP) Initiative II Data Collection, Global Soil Data Task Group, Oak Ridge, Tennessee, USAm available at: https://daac.ornl.gov (last access: 14 August 2012), 2009. 4122
- Jones, P. D. and Harris, I.: CRU Time Series (TS) high resolution gridded datasets, University of East Anglia Climatic Research Unit (CRU), available at: http://badc.nerc.ac.uk/view/badc. nerc.ac.uk_ATOM_dataent_1256223773328276 (last access: 28 August 2012), 2008. 4078, 4122
- Joos, F., Gerber, S., Prentice, I. C., Otto-Bliesner, B. L., and Valdes, P. J.: Transient simulations of Holocene atmospheric carbon dioxide and terrestrial carbon since the Last Glacial Maximum, Global Biogeochem. Cy., 18, GB2002, doi:10.1029/2003GB002156, 2004. 4085

Discussion Paper	GM 5, 4071–4 WETC metho	GMDD 5, 4071–4136, 2012 WETCHIMP methodology			
Discu	R. Wan	ia et al.			
ission Pa	Title	Page			
aper	Abstract	Introduction			
_	Conclusions	References			
Discu	Tables	Figures			
oissi	14	►I			
n Pa	•	•			
per	Back	Close			
—	Full Scre	en / Esc			
Discussi	Printer-frier	Printer-friendly Version			
on F	Interactive	Discussion			
aper	$\overline{\mathbf{c}}$	BY			

- Discussion Paper GMDD 5, 4071-4136, 2012 **WETCHIMP** methodology **Discussion** Paper R. Wania et al. Title Page Abstract Introduction Conclusions References **Discussion** Paper Tables **Figures** Back Full Screen / Esc **Discussion** Paper **Printer-friendly Version** Interactive Discussion
- Kaplan, J. O.: Wetlands at the Last Glacial Maximum: Distribution and methane emissions, Geophys. Res. Lett., 29, 1079, doi:10.1029/2001GL013366, 2002. 4075, 4076, 4095
- Knorr, W.: Annual and interannual CO₂ exchanges of the terrestrial biosphere: process-based simulations and uncertainties, Global Ecol. Biogeogr., 9, 225–252, 2000. 4097
- Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G., and Tarnocai, C.: Permafrost carbon-climate feedbacks accelerate global warming, P. Natl. Acad. Sci., 108, 14769–14774, 2011. 4090
 - Krinner, G.: Impact of lakes and wetlands on boreal climate, J. Geophys. Res., 108, 4520,doi:10.1029/2002JD002597, 2003. 4075
- Krinner, G., Viovy, N., de Noblet-Ducoudre, N., Ogee, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I. C.: A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system, Global Biogeochem. Cy., 19, 941–962, 2005. 4090
 - Kudryavtsev, V. A., Garagula, L. S., Kondratiyeva, K. A., and Melamed, V. E.: Frost Forecasting in Geological Engineering Investigations, CRREL, Hanover, NH, 1977. 4084
- ¹⁵ Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slate, A. G.: Parameterization improvements and functional and structural advances in version 4 of the Community Land Model, J. Adv. Model. Earth. Sys., 3, M03001, doi:10.1029/2011MS000045, 2011. 4080
- ²⁰ Lawrence, P. J. and Chase, T. N.: Representing a new MODIS consistent land surface in the Community Land Model (CLM 3.0), J. Geophys. Res., 112, G01023, doi:10.1029/2006JG000168, 2007. 4122
 - Leff, B., Ramankutty, N., and Foley, J.: Geographic distribution of major crops across the world, Global. Biogeochem. Cy., 18, GB1009, doi:10.1029/2003GB002108, 2004. 4079, 4083, 4090, 4122, 4129
 - Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, J. Hydrol., 296, 1–22, 2004. 4082, 4098, 4122
 - Letts, M. G., Roulet, N. T., Comer, N. T., Skarupa, M. R., and Verseghy, D. L.: Parametrization of peatland hydraulic properties for the Canadian Land Surface Scheme, Atmos. Ocean, 38, 141–160, 2000, 4000
- ³⁰ 141–160, 2000. 4098

25

Liang, X., Lettenmaier, D. P., Wood, E. F., and Burges, S. J.: A simple hydrologically based model of land-surface water and energy fluxes for general-circulation models, J. Geophys. Res., 99, 14415–14428, 1994. 4097

Liu, M. L., Tian, H. Q., Chen, G. S., Ren, W., Zhang, C., and Liu, J. Y.: Effects of land-use and land-cover change on evapotranspiration and water yield in China during 1900–2000, J. Am. Water Resour. As., 44, 1193–1207, 2008. 4083

Liu, M., Tian, H., Yang, Q., Yang, J., Song, X., Lohrenz, S. E., and Cai, W.: Variations of evapo-

- transpiration and runoff over the drainage basins of the Gulf of Mexico driven by climate and land use change during 1901–2008, Water Resour. Res., in revision, 2012a. 4082, 4083
 - Liu, M. L., Tian, H. Q., Lu, C. Q., Xu, X. F., Chen, G. S., and Ren, W.: Effects of multiple environment stresses on evapotranspiration and runoff over eastern China, J. Hydrol., 426, 39–54, 2012b. 4082
- ¹⁰ Lu, C. Q., Tian, H. Q., Liu, M. L., Ren, W., Xu, X. F., Chen, G. S., and Zhang, C.: Effect of nitrogen deposition on China's terrestrial carbon uptake in the context of multifactor environmental changes, Ecol. Appl., 22, 53–75, 2012. 4082
 - Matthews, E. and Fung, I. Y.: Methane emission from natural wetlands: Global distribution, area, and environmental characteristics of sources, Global Biogeochem. Cy., 1, 61–86, 1987. 4074
- ¹⁵ Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J. M., Noda, A., Raper, S. C. B., Watterson, I. G., Weaver, A. J., and Zhao, Z.-C.: Global climate projections, in: Climate change 2007: The physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z.,
- ²⁰ Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., chap. 10, Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 747–846, 2007. 4077
 - Meissner, K. J., Weaver, A. J., Matthews, H. D., and Cox, P. M.: The role of land surface dynamics in glacial inception: a study with the UVic Earth System Model, Clim. Dynam., 21, 515–537, 2003. 4095
- Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen, G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: conclusions from a model intercomparison project
- ³⁰ (WETCHIMP), Biogeosciences Discuss., 9, 11577–11654, doi:10.5194/bgd-9-11577-2012, 2012. 4073, 4076, 4077, 4078, 4079, 4087, 4099, 4101, 4104, 4105

Discussion Pap	GMDD 5, 4071–4136, 2012					
oer	WET(metho	CHIMP dology				
Discu	R. Wan	R. Wania et al.				
ission Pa	Title	Page				
aper	Abstract	Introduction				
—	Conclusions	References				
Discu	Tables	Figures				
oissi	14	►I.				
n Pa	•	•				
per	Back	Close				
—	Full Scre	en / Esc				
Discussion	Printer-frier Interactive	ndly Version				
Paper	C	BY				

Mitchell, T. D. and Jones, P. D.: An improved method of constructing a database of monthly climate observations and associated high-resolution grids, Int. J. Climatol., 25, 693–712, 2005. 4078, 4084, 4122

Mokhov, I. I., Eliseev, A. V., and Denisov, S. N.: Model diagnostics of variations in methane

- emissions by wetlands in the second half of the 20th century based on reanalysis data, Doklady Earth Sci., 417, 1293–1297, doi:10.1134/S1028334X07080375, 2007. 4083
 - Myneni, R. B., Hoffman, S., Knyazikhin, Y., Privette, J. L., Glassy, J., Tian, Y., Wang, Y., Song, X., Zhang, Y., Smith, G. R., Lotsch, A., Friedl, M., Morisette, J. T., Votava, P., Nemani, R. R., and Running, S. W.: Global products of vegetation leaf area and fraction absorbed PAR from year one of MODIS data, Remote Sens. Environ., 83, 214–231, 2002. 4097
- one of MODIS data, Remote Sens. Environ., 83, 214–231, 2002. 4097
 Papa, F., Prigent, C., Durand, F., and Rossow, W. B.: Wetland dynamics using a suite of satellite observations: a case study of application and evaluation for the Indian Subcontinent, Geophys. Res. Lett., 33, L08401, doi:10.1029/2006GL025767, 2006. 4079
- Papa, F., Prigent, C., and Rossow, W. B.: Monitoring flood and discharge variations in the large Siberian rivers from a multi-satellite technique, Surv. Geophys., 29, 297–317,
 - doi:10.1007/s10712-008-9036-0, 2008. 4079
 Papa, F., Prigent, C., Aires, F., Jimenez, C., Rossow, W. B., and Matthews, E.: Interannual variability of surface water extent at the global scale, 1993–2004, J. Geophys. Res., 115, D12111, doi:10.1029/2009JD012674, 2010. 4077, 4079, 4098, 4122, 4126
- Parton, W. J., Scurlock, J. M. O., Ojima, D. S., Gilmanov, T. G., Scholes, R. J., Schimel, D. S., Kirchner, T., Menaut, J. C., Seastedt, T., Moya, E. G., Kamnalrut, A., and Kinyamario, J. I.: Observations and modeling of biomass and soil organic-matter dynamics for the grassland biome worldwide, Global Biogeochem. Cy., 7, 785–809, 1993. 4094

Pickett-Heaps, C. A., Jacob, D. J., Wecht, K. J., Kort, E. A., Wofsy, S. C., Diskin, G. S., Wor-

thy, D. E. J., Kaplan, J. O., Bey, I., and Drevet, J.: Magnitude and seasonality of wetland methane emissions from the Hudson Bay Lowlands (Canada), Atmos. Chem. Phys., 11, 3773–3779, doi:10.5194/acp-11-3773-2011, 2011. 4089

Potter, C. S.: An ecosystem simulation model for methane production and emission from wetlands, Global Biogeochem. Cy., 11, 495–506, 1997. 4075

Prentice, I. C., Farquhar, G. D., Fasham, M. J. R., Goulden, M., Heimann, M., Jaramillo, V., Kheshgi, H., Quéré, C.-L., Scholes, R., and Wallace, D.: The Carbon Cycle and Atmospheric Carbon Dioxide, in: Climate Change 2001: The scientific basis, Contributions of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change,



edited by: Pitelka, L. F. and Rojas, A. R., Cambridge University Press, Cambridge, 183–237, 2001. 4078

- Prigent, C., Papa, F., Aires, F., Rossow, W. B., and Matthews, E.: Global inundation dynamics inferred from multiple satellite observations, 1993–2000, J. Geophys. Res., 112, 305–317, doi:10.1029/2006JD007847, 2007. 4077, 4079, 4081, 4098, 4122
- doi:10.1029/2006JD007847, 2007. 4077, 4079, 4081, 4098, 4122
 Prigent, C., Papa, F., Aires, F., Jimenez, C., Rossow, W. B., and Matthews, E.: Changes in land surface water dynamics since the 1990s and relation to population pressure, Geophys. Res. Lett., 39, L08403, doi:10.1029/2012GL051276, 2012. 4079
 - Qian, T. T., Dai, A., Trenberth, K. E., and Oleson, K. W.: Simulation of global land surface
- conditions from 1948 to 2004, part I: forcing data and evaluations, J. Hydrometeorol., 7, 953–975, 2006. 4081

15

- Ren, W., Tian, H., Chen, G., Liu, M., Zhang, C., Chappelka, A. H., and Pan, S.: Influence of ozone pollution and climate variability on net primary productivit and carbon storage in China's grassland ecosystems from 1961 to 2000, Environ. Pollut., 149, 327–335, 2007. 4082
- Ren, W., Tian, H. Q., Xu, X. F., Liu, M. L., Lu, C. Q., Chen, G. S., Melillo, J., Reilly, J., and Liu, J. Y.: Spatial and temporal patterns of CO₂ and CH₄ fluxes in China's croplands in response to multifactor environmental changes, Tellus B, 63, 222–240, 2011. 4082
 Ridgwell, A., Marshall, S. J., and Gregson, K.: Consumption of atmospheric methane by soils:
- a process-based model, Global Biogeochem. Cy., 13, 59–70, 1999. 4075
 - Riley, W. J., Subin, Z. M., Lawrence, D. M., Swenson, S. C., Torn, M. S., Meng, L., Mahowald, N. M., and Hess, P.: Barriers to predicting changes in global terrestrial methane fluxes: analyses using CLM4Me, a methane biogeochemistry model integrated in CESM, Biogeosciences, 8, 1925–1953, doi:10.5194/bg-8-1925-2011, 2011. 4075, 4076, 4080, 4081
- Ringeval, B.: Interactions entre climat et émissions de méthane par les zones humides à l'échelle global, Ph. D. thesis, Université Pierre & Marie Curie, Paris, France, 2011. 4075
- Ringeval, B., de Noblet-Ducoudre, N., Ciais, P., Bousquet, P., Prigent, C., Papa, F., and Rossow, W. B.: An attempt to quantify the impact of changes in wetland extent on methane emissions on the seasonal and interannual time scales, Global Biogeochem. Cy., 24, 611–617, 2010, 4075, 4076, 4091, 4092
 - Ringeval, B., Friedlingstein, P., Koven, C., Ciais, P., de Noblet-Ducoudré, N., Decharme, B., and Cadule, P.: Climate-CH4 feedback from wetlands and its interaction with the climate-CO₂ feedback, Biogeosciences, 8, 2137–2157, doi:10.5194/bg-8-2137-2011, 2011. 4076, 4090



- Ringeval, B., Decharme, B., Piao, S. L., Ciais, P., Papa, F., de Noblet-Ducoudré, N., Prigent, C., Friedlingstein, P., Gouttevin, I., Koven, C., and Ducharne, A.: Modelling sub-grid wetland in the ORCHIDEE global land surface model: evaluation against river discharges and remotely sensed data, Geosci. Model Dev., 5, 941–962, doi:10.5194/gmd-5-941-2012, 2012a. 4091
- ⁵ Ringeval, B., Hopcroft, P. O., Valdes, P. J., Ciais, P., Ramstein, G., Dolman, A. J., and Kageyama, M.: Response of methane emissions from wetlands to the Last Glacial Maximum and an idealized Dansgaard-Oeschger climate event: insights from two models of different complexity, Clim. Past Discuss., 8, 3093–3142, doi:10.5194/cpd-8-3093-2012, 2012b. 4092 Roulet, N. T., Moore, T. R., Bubier, J., and Lafleur, P.: Northern fens: methane flux and climate change, Tellus, 44B, 100–106, 1992. 4093
- Schroeder, R., Rawlins, M. A., McDonald, K. C., Podest, E., Zimmermann, R., and Kueppers, M.: Satellite microwave remote sensing of North Eurasian inundation dynamics: development of coarse-resolution products and comparison with high-resolution synthetic aperture radar data, Environ. Res. Lett., 5, 14415–14428, 2010. 4098, 4122
- ¹⁵ Segers, R. and Leffelaar, P. A.: Modeling methane fluxes in wetlands with gas-transporting plants 3, plot scale, J. Geophys. Res., 106, 3541–3558, 2001. 4075
 - Sellers, P. J., Meeson, B. W., Closs, J., Corprew, F., Dazlich, D., Hall, F. G., Kerr, Y., Koster, R., Los, S., Mitchell, K., McManus, J., Myers, D., Sun, K.-J., and National Aeronautics and Space Administration and Goddard Space Flight Center, P. J. T.: International Satellite Land Surface
- 20 Climatology Project Initiative I data collection (ISLSCP I), available at: http://badc.nerc.ac. uk/view/badc.nerc.ac.uk__ATOM__dataent_ISLSCP (last access: 1 September 2012), 1996. 4122
 - Sheng, Y. W., Smith, L. C., MacDonald, G. M., Kremenetski, K. V., Frey, K. E., elichko, A. A., Lee, M., Beilman, D. W., and Dubinin, P.: A high-resolution GIS-based inventory of the west
- ²⁵ Siberian peat carbon pool, Global Biogeochem. Cy., 18, 14415–14428, 2004. 4079, 4098, 4100, 4122, 4128
 - Shindell, D. T., Walter, B. P., and Faluvegi, G.: Impacts of climate change on methane emissions from wetlands, Geophys. Res. Lett., 31, L21202, doi:10.1029/2004GL021009, 2004. 4075, 4076
- Singarayer, J. S., Valdes, P. J., Friedlingstein, P., Nelson, S., and Beerling, D. J.: Late Holocene methane rise caused by orbitally controlled increase in tropical sources, Nature, 470, 82–91, 2011. 4075, 4092



- Sitch, S., Smith, B., Prentice, I. C., Arneth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, Glob. Change Biol., 9, 161–185, 2003. 4085, 4089, 4090
- ⁵ Spahni, R., Wania, R., Neef, L., van Weele, M., Pison, I., Bousquet, P., Frankenberg, C., Foster, P. N., Joos, F., Prentice, I. C., and van Velthoven, P.: Constraining global methane emissions and uptake by ecosystems, Biogeosciences, 8, 1643–1665, doi:10.5194/bg-8-1643-2011, 2011. 4075, 4076, 4085, 4086, 4087, 4089, 4122

Stillwell-Soller, L. M., Klinger, L. F., Pollard, D., and Thompson, S. L.: The Global Distribution of

- ¹⁰ Freshwater Wetlands, Tech. Rep. NCAR/TN-416+STR, NCAR, available at: http://www.cisl. ucar.edu/ess/services/dbst/wetlands/ (last access: 11 September 2012), 1995. 4082
 - Stocker, B. D., Strassmann, K., and Joos, F.: Sensitivity of Holocene atmospheric CO₂ and the modern carbon budget to early human land use: analyses with a process-based model, Biogeosciences, 8, 69–88, doi:10.5194/bg-8-69-2011, 2011. 4085
- Strassmann, K. M., Joos, F., and Fischer, G.: Simulating effects of land use changes on carbon fluxes: past contributions to atmospheric CO₂ increases and future commitments due to losses of terrestrial sink capacity, Tellus B, 60, 583–603, doi:10.1111/j.1600-0889.2008.00340.x, 2008. 4085

Tarnocai, C., Swanson, D., Kimble, J., and Broll, J.: Northern Circumpolar Soil Carbon

- Database, Tech. Rep. Version 1, Research Branch, Agriculture and Agri-Food Canada,, available at: http://wms1.agr.gc.ca/NortherCircumpolar/northercircumpolar.zip (last access: 1 October 2012), 2007. 4079, 4085, 4087, 4099, 4122, 4127
 - Tarnocai, C., Canadell, J. G., Schuur, E. A. G., Kuhry, P., Mazhitova, G., and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region, Global Biogeochem.
- Cy., 23, GB2023, doi:10.1029/2008GB003327, 2009. 4079, 4085, 4087, 4099, 4122, 4127
 Tian, H., Xu, X., Liu, M., Ren, W., Zhang, C., Chen, G., and Lu, C.: Spatial and temporal patterns of CH₄ and N₂O fluxes in terrestrial ecosystems of North America during 1979–2008: application of a global biogeochemistry model, Biogeosciences, 7, 2673–2694, doi:10.5194/bg-7-2673-2010, 2010. 4082, 4083
- Tian, H. Q., Chen, G. S., Liu, M. L., Zhang, C., Sun, G., Lu, C. Q., Xu, X. F., Ren, W., Pan, S. F., and Chappelka, A.: Model estimates of net primary productivity, evapotranspiration, and water use efficiency in the terrestrial ecosystems of the southern United States during 1895– 2007, Forest Ecol. Manag., 259, 1311–1327, 2010b. 4082



- Tian, H. Q., Melillo, J., Lu, C. Q., Kicklighter, D., Liu, M. L., Ren, W., Xu, X. F., Chen, G. S., Zhang, C., Pan, S. F., Liu, J. Y., and Running, S.: China's terrestrial carbon balance: contributions from multiple global change factors, Global Biogeochem. Cy., 25, 222–240, 2011a. 4082
- ⁵ Tian, H. Q., Xu, X. F., Lu, C. Q., Liu, M. L., Ren, W., Chen, G. S., Melillo, J., and Liu, J. Y.: Net exchanges of CO₂, CH₄, and N₂O between China's terrestrial ecosystems and the atmosphere and their contributions to global climate warming, J. Geophys. Res., 116, G02011, doi:10.1029/2010JG001393, 2011b. 4082

Tian, H. Q., Chen, G. S., Zhang, C., Liu, M. L., Sun, G., Chappelka, A., Ren, W., Xu, X. F.,

¹⁰ Lu, C. Q., Pan, S. F., Chen, H., Hui, D. F., McNulty, S., Lockaby, G., and Vance, E.: Centuryscale responses of ecosystem carbon storage and flux to multiple environmental changes in the southern United States, Ecosystems, 15, 674–694, 2012. 4082

Valdes, P. J., Beerling, D. J., and Johnson, C. E.: The ice age methane budget, Geophys. Res. Lett., 32, 941–962, 2005. 4075

- van Bodegom, P., Goudriaan, J., and Leffelaar, P.: A mechanistic model on methane oxidation in a rice rhizosphere, Biogeochem., 55, 145–177, 2001a. 4075
 - van Bodegom, P., Wassmann, R., and Metra-Corton, T.: A process-based model for methane emission predictions from flooded rice paddies, Global Biogeochem. Cy., 15, 247–263, 2001b. 4075
- Viovy, N. and Ciais, P.: CRUNCEP data set for 1901–2008, Tech. Rep. Version 4, Laboratoire des Sciences du Climat et de l'Environnement, available at: http://dods.extra.cea.fr/ data/p529viov/cruncep/, last access: 1 September, 2011. 4078, 4122
 - Walter, B. P. and Heimann, M.: A process-based, climate-sensitive model to derive methane emissions from natural wetlands: application to five wetland sites, sensitivity to model pa-
- rameters, and climate, Global Biogeochem. Cycles, 14, 745–765, 2000. 4075, 4098
 Walter, B. P., Heimann, M., Shannon, R. D., and White, J. R.: A process-based model to derive methane emissions from natural wetlands, Geophys. Res. Lett., 23, 3731–3734, 1996. 4075
 Walter, B. P., Heimann, M., and Matthews, E.: Modeling modern methane emissions from natural wetlands 1, model description and results, J. Geophys. Res., 106, 34189–34206, 2001a.
 4075, 4091
 - Walter, B. P., Heimann, M., and Matthews, E.: Modeling modern methane emissions from natural wetlands 2, interannual variations 1982–1993, J. Geophys. Res., 106, 34207–34219, 2001b. 4075

	GMDD 5, 4071–4136, 2012						
	WETC	WETCHIMP methodology					
	R. Wani Title F	a et al.					
5	Abstract	Introduction					
_	Conclusions	References					
7	Tables	Figures					
5	I	۶I					
	•	•					
5	Back	Close					
-	Full Scree	en / Esc					
	Printer-friend	dly Version					
5	Interactive E	Discussion					
5	<u></u>	D BY					

- Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model, I: evaluation and sensitivity of physical land surface processes, Global Biogeochem. Cy., 23, GB3014, doi:10.1029/2008GB003412, 2009a. 4076, 4085, 4087
- ⁵ Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model, II: evaluation and sensitivity of vegetation and carbon cycle processes, Global Biogeochem. Cy., 23, GB3015, doi:10.1029/2008GB003413, 2009b. 4085, 4087, 4088

Wania, R., Ross, I., and Prentice, I. C.: Implementation and evaluation of a new methane model

- within a dynamic global vegetation model: LPJ-WHyMe v1.3.1, Geosci. Model Dev., 3, 565–584, doi:10.5194/gmd-3-565-2010, 2010. 4076, 4085, 4087, 4088
 - Weaver, A. J., Eby, M., Wiebe, E. C., Bitz, C. M., Duffy, P. B., Ewen, T. L., Fanning, A. F., Holland, M. M., MacFadyen, A., Matthews, H. D., Meissner, K. J., Saenko, O., Schmittner, A., Wang, H. X., and Yoshimori, M.: The UVic Earth System Climate Model: model description,
- climatology, and applications to past, present and future climates, Atmos. Ocean, 39, 361– 428, 2001. 4095
 - Weber, S. L., Drury, A. J., Toonen, W. H. J., and van Weele, M.: Wetland methane emissions during the Last Glacial Maximum estimated from PMIP2 simulations: Climate, vegetation, and geographic controls, J. Geophys. Res., 115, D06111, doi:10.1029/2009JD012110, 2010. 4076

20

25

- Woodward, F. I., Smith, T. M., and Emanuel, W. R.: A global land primary productivity and phytogeography model, Global Biogeochem. Cy., 9, 471–490, 1995. 4092
- Xu, X. F., Tian, H. Q., Zhang, C., Liu, M. L., Ren, W., Chen, G. S., Lu, C. Q., and Bruhwiler, L.: Attribution of spatial and temporal variations in terrestrial methane flux over North America, Biogeosciences, 7, 3637–3655, doi:10.5194/bg-7-3637-2010, 2010. 4082
- Yang, Q., Tian, H., Liu, M., and Yang, J.: Impacts of land use/cover change and climate change on land surface runoff and water export in the US East Coast area during 1901–2008, In prep., 2012.4083

Zhuang, Q., Melillo, J. M., Kicklighter, D. W., Prinn, R. G., McGuire, A. D., Steudler, P. A.,

³⁰ Felzer, B. S., and Hu, S.: Methane fluxes between terrestrial ecosystems and the atmosphere at northern high latitudes during the past century: A retrospective analysis with a process-based biogeochemistry model, Global Biogeochem. Cy., 18, GB3010, doi:10.1029/2004GB002239, 2004. 4075



- Zhuang, Q., Melillo, J. M., Sarofim, M. C., Kicklighter, D. W., McGuire, A. D., Felzer, B. S., Sokolov, A., Prinn, R. G., Steudler, P. A., and Hu, S.: CO₂ and CH₄ exchanges between land ecosystems and the atmosphere in northern high latitudes over the 21st century, Geophys. Res. Lett., 33, L17403, doi:10.1029/2006GL026972, 2006. 4075
- ⁵ Zobler, L.: A world soil file for global climate modelling, Technical Memorandum 87802, 32, NASA, 1986. 4089
 - Zürcher, S., Spahni, R., Joos, F., Steinacher, M., and Fischer, H.: Impact of an 8.2-kyr-like event on methane emissions in northern peatlands, Biogeosciences Discuss., 9, 13243–13286, doi:10.5194/bgd-9-13243-2012, 2012. 4085

_	GMDD 5, 4071–4136, 2012				
	WETCHIMP methodology R. Wania et al.				
-	Title Page				
	Abstract	Introduction			
	Conclusions	References			
	Tables Figures				
	I	۶I			
	•	•			
	Back	Close			
	Full Screen / Esc				
	Printer-friendly Version				
	Interactive	Discussion			

Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper



Table 1. Description of the WETCHIMP modelling protocol.

Name	Experiment	Climate years	Description
Experiment 1-equil	Equilibrium	1901–1931	Spin up of models with 1901–1931 climate until equilibrium.
Experiment 2-trans	Transient	1932–2004	Continuing from equilibrium, models are run tran- siently. Comparison phase is 1993–2004.
Experiment 3-opt	Optimized	User-defined	Model run with user selected "optimal" configura- tion. Comparison phase is 1993–2004.
Experiment 4-CO ₂	Atmospheric [CO ₂] sensitivity	1901–1931	From the model state at end of Experiment 1-equil simulation, apply a globally uniform step increase in $[CO_2]$ to reach SRES A2 2100 levels (857 ppmv). Run model until equilibrium ^a is re-established.
Experiment 5-T	Temperature sensitivity	1901–1931	From the model state at end of equilibrium run, apply a step increase in air temperature reflecting mean SRES A2 2100 increase (multi-model mean SAT warming for 2090 to 2099 relative to 1980 to 1999: \approx +3.4°C). Run model until equilibrium ^a reestablished.
Experiment 6-P	Moisture sensitivity	1901–1931	From the model state at end of transient equilibrium run, a step increase in precipitation to reflecting mean SRES A2 2100 increase (30-yr average 2071 to 2100 relative to 1961 to 1990: $\approx +3.9\%^{b}$). Run model until equilibrium ^a re-established.

^a Each modelling group used their own criteria for what equilibrium meant: LPJ-WHyMe, LPJ-Bern, SDGVM used the stability of the soil C pool; UVic-ESCM used soil moisture and temperature variables; DLEM specified an upper limit for inter-annual changes in total ecosystem C storage (< 0.1 gCm^{-2}), soil moisture (< 0.1), and nitrogen storage (< 0.1 gNm^{-2}). LPJ-WSL used soil and vegetation carbon. IAP-RAS is an equilibrium model and thus does not require spin-up.

^b As the IPCC AR4 report does not contain a globally averaged number for the mean precipitation change, this value is from the IPCC TAR Report of 2001.

Table 2. List of experiments (described in Table 1) performed by each of the participating modelling groups. "Prognostic" indicates that a model located at least part of its wetlands based either on an inversion to GIEMS and modelled hydrological state or used GIEMS as a spatial or temporal constraint. "Prescribed" means that the model used a distribution map for wetlands and "simulated" means that the model did not input any kind of wetland observational data to locate wetlands (see Sect. 4.1).

Model	Areal extent	Wetlands	CH ₄ fluxes	Experiments performed	Contact
CLM4Me	global	prognostic	simulated	1, 2, 3, 4, 5, 6	W. J. Riley
DLEM	global	prognostic	simulated	1, 2, 3, 4, 5, 6	H. Tian
IAP-RAS	global	prescribed	simulated	1, 2, 3, , 5, 6	A. V. Eliseev
LPJ-Bern	global	prognostic	simulated	1, 2, 3, 4, 5, 6	R. Spahni
LPJ-WHyMe	35–90° N	prescribed	simulated	1, 2, , 4, 5, 6	R. Wania
LPJ-WSL*	global	prescribed	simulated	1, 2, 3, 4, 5, 6	E. L. Hodson
ORCHIDEE	global	prognostic	simulated	1, 2, 3, 4, 5, 6	B. Ringeval
SDGVM	global	simulated	simulated	1, 2, , 4, 5, 6	P. O. Hopcroft
UVic-ESCM	global	simulated	n/a	1, 2, 3, 4, 5, 6	C. A. Avis
UW-VIC	W-Siberia	prognostic	simulated	1, 2, , , ,	T. Bohn

* LPJ-WSL uses the "prognostic" approach for Experiment 3-opt, using GIEMS as guidance for the wetland distribution.

GN 5, 4071–4	GMDD 5, 4071–4136, 2012		
WETCHIMP methodology R. Wania et al.			
Title	Page		
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
14	►I		
•	•		
Back	Close		
Full Scr	een / Esc		
Printer-friendly Version			
Interactive Discussion			

Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper

Table 3. Names and types of input datasets together with references and a list of models that used the data.

Name of dataset	Description	Used by	Reference
CRU TS3.1	Climate	All models ^a	Mitchell and Jones (2005); Jones and Harris (2008)
CRUNCEP	Climate	All models ^a	Viovy and Ciais (2011)
Law Dome Composite	CO ₂	All models	http://grkapweb1.epfl.ch/pub/ARVE_tech_report2_co2spline.pdf
Dentener et al. (2006)	Nitrogen deposition	DLEM	Dentener et al. (2006)
FAO	Soil texture classes	ORCHIDEE	Batjes (1997)
HWSD	Soil texture classes	LPJ-Bern	FAO/IIASA/ISRIC/ISSCAS/JRC (2009)
IGBP-DIS	Soil texture classes	CLM4Me, DLEM	Global Soil Data Task Group (2000)
ISLSCP I	Soil texture classes	SDGVM	Sellers et al. (1996)
ISLSCP II	Soil texture classes,	UVic-ESCM	ISLSCP-II (2009)
	soil carbon density		
MODIS	Distribution of plant	UW-VIC	Bartalev et al. (2003)
	functional types (PFTs)		
ETOPO 2v2	Topography	SDGVM, UVic-ESCM	ETOPO (2006)
HYDRO1k	Topography	ORCHIDEE	http://webgis.wr.usgs.gov/globalgis/metadata_qr/metadata/hydro1k.htm
CLM soil colours	Soil colours	CLM4Me	Lawrence and Chase (2007)
GIEMS	Monthly inundated	CLM4Me, DLEM,	Prigent et al. (2007); Papa et al. (2010)
	wetland area	LPJ-Bern, LPJ-WSL,	
		ORCHIDEE, UW-VIC	
	1993–2004 (Fig. 1)		
Schroeder et al. (2010)	Remotely-sensed	UW-VIC ^D	Schroeder et al. (2010)
	inundation dataset		
CDIAC NDP017	Wetland area	IAP-RAS	http://cdiac.esd.ornl.gov/ndps/ndp017.html
GLWD	Global land cover	DLEM	Lehner and Döll (2004)
NCSCD	Annual fractional	LPJ-Bern,	Tarnocai et al. (2007, 2009)
	cover of northern	LPJ-WHyMe	
	peatlands (Fig. 2)		
Sheng	Peatland fraction (Fig. 3)	UW-VIC	Sheng et al. (2004)
	and peat depths		
Leff	Annual fractional cover of	DLEM,	Leff et al. (2004), Spahni et al. (2011)
	rice fields scaled by	LPJ-Bern,	
	monthly inundation (Fig. 4)	LPJ-WSL	
Fries et al. (1998)	Global land cover	DLEM	Fries et al. (1998)
GICEW	waterbodies and land	LPJ-Bern	http://luh.sr.unh.edu/
	ice excluding ice sheets (Fig. 5)		

^a These datasets were required for use in Experiments 1, 2, 4, 5, and 6.

^b Used in experiment 3.



Table 4. A list of the models that contributed simulations to WETCHIMP. The "Wetland types" gives a quick overview of what kind of wetlands are used or simulated by each model. The explanations for the wetland types and soil data sets, as well as the full references are given in Table 3.

Model	Resolution (lon × lat)	Wetland types	Climate forcing	Soil data
CLM4Me	2.5° × 1.9°	Simulated inundated area based on predicted water table and runoff and an inversion to GIEMS	CRUNCEP	IGBP-DIS
DLEM	$0.5^{\circ} \times 0.5^{\circ}$	Mixture between prescribed and simulated, rice mask by Leff	CRUNCEP	IGBP-DIS
IAP-RAS	0.5° × 0.5°	Olson data for wetlands (bogs/mires, swamps, heaths/ moorlands, tundra)	CRU3.1	Peat in peatlands, loam elsewhere
LPJ-Bern	0.5° × 0.5°	Peatlands from NCSCD, inundated wetlands from GIEMS, rice mask by Leff, permanent water or ice from GICEW, simulated wet soils	CRU3.1	HWSD
LPJ-WHvMe	$0.5^{\circ} \times 0.5^{\circ}$	Peatlands from NCSCD	CRU3.1	n/a ^a
LPJ-WSL	0.5° × 0.5°	Inundated area from GIEMS, rice mask by Leff for all experiments except 3 ^b	CRU3.1	FAO
ORCHIDEE	1° × 1°	Simulated, but annual mean over 1993–2004 adjusted to mean of GIEMS	CRUNCEP	FAO for mineral
SDGVM	$0.5^{\circ} \times 0.5^{\circ}$	All simulated	CRU3.1	ISLSCP I
UVic-ESCM	3.6° × 1.8°	All simulated	CRU3.1+NCEP ^c	ISLSCP II
UW-VIC	100 km ^d	Simulated lakes and peatlands	CRUNCEP	FAO for mineral soils, Sheng for peatland fraction

^a LPJ-WHyMe is a peatland only model, thus no "soil" data is required.

^b LPJ-WSL exp 3 is a mix between prescribed (GIEMS) and simulated inundation area based upon an empirical relationship between simulated water runoff and GIEMS.

- $^{\rm c}$ Surface winds and diurnal temperature are taken from the NCEP reanalysis.
- d 100 km polar azimuthal equal area grid (EASE grid), resampled to $0.5^\circ \times 0.5^\circ.$

GM 5, 4071–4	GMDD 5, 4071–4136, 2012		
WETCHIMP methodology			
R. Wania et al.			
Title	Title Page		
Abstract	Introduction		
Conclusions	References		
Tables	Figures		
I	۶I		
•	×.		
Back	Close		
Full Scre	Full Screen / Esc		
Printer-frier	Printer-friendly Version		
Interactive Discussion			

Discussion Paper

Discussion Paper

Discussion Paper

Discussion Paper

Table 5. A general description of the dependencies of CH_4 production, CH_4 oxidation (does not include atmospheric CH_4 oxidation) and CH_4 flux. All of the fluxes are modulated by a CH_4 -producing area (see Fig. 6). Parameters and variables used in the models were harmonized where possible, but identical names do not indicate identical values in the different models.

Model	CH ₄ production (P)	CH ₄ oxidation (O)	Atmospheric CH_4 oxidation (O_{atm})	Net CH ₄ flux (F)
CLM4Me	$P = R_{\rm het} r_{\rm CH4:C} f_{\rm pH} f_{\rm pE} Q_{10}$	$O = O_{\max} f_{[O_2]} f_{[CH_4]} f_{\Theta} Q_{10}$	$O_{\text{atm}} = R_{\text{o},\text{max}} f_{[O_2]} f_{\text{atm}[CH_4]} f_{[CH_4]} f_{\odot} Q_{10}$	$F = (P - O)f_{\text{transport}} - O_{\text{atm}}$
DLEM	$P = P_{\max} C_{\text{labile}} f_T f_{\text{pH}} f_{\Theta}$	$\begin{aligned} O_{\text{trans}} &= O_{\text{trans,max}} f_{\text{planttrans}} f_{\mathcal{T}} \\ O_{\text{soil}} &= O_{\text{soil,max}} f_{[\text{CH}_4]} f_{\mathcal{T}_{\text{soil}}} f_{\text{pH}} f_{\text{oxid,}\Theta} \end{aligned}$	$O_{\text{atm}} = O_{\text{atm,max}} f_{\text{atm[CH_4]}} f_{\mathcal{T}_{\text{air}}} f_{\text{pH}} f_{\text{oxid},\Theta}$	$F = (P - O_{\text{trans}} - O_{\text{soil}})f_{\text{transport}} - O_{\text{atm}}$
IAP-RAS	$P = f_T$	-	-	$F = P f_{\Theta} Q_{10}$
LPJ-Bern peat	$P = R_{\rm het} r_{\rm CH4:C} f_{\rm root} f_{\rm WTP}$	$O = f_{[O_2]} f_{[CH_4]} r_{O_2}$	-	$F = (P - O)f_{\text{transport}}$
LPJ-Bern wetlands	$P = R_{\rm het} r_{\rm CH4:C}$	-	-	F = P
LPJ-Bern rice	$P = R_{\rm het} r_{\rm CH4:C}$	-	-	F = P
LPJ-Bern wetsoils	$P = R_{\rm het} r_{\rm CH4:C} f_{\Theta}$	-	$O_{\rm atm} = f_{\rm atm[CH_4]} f_7 f_{\Theta} f_{\rm soil}$	$F = P - O_{\text{atm}}$
LPJ-WHyMe	$P = R_{\rm het} r_{\rm CH4:C} f_{\rm root} f_{\rm WTP}$	$O = f_{[O_2]} f_{[CH_4]} r_{O_2}$	-	$F = (P - O)f_{\text{transport}}$
LPJ-WSL	$P = R_{\rm het} r_{\rm CH4:C} f_{\rm ecosys}$	-	-	F = P
ORCHIDEE	$P = R_0 C_{labile} f_{WTP} f_T Q_{10}$	$O = f_{WTP} f_{[CH_4]} Q_{10}$	-	$F = (P - O)f_{\text{transport}}$
SDGVM	$P = R_{\rm het} r_{\rm CH4:C} f_{\rm WTP} f_T Q_{10}$	<i>O</i> = 0.9 <i>P</i>	-	F = P - O
UW-VIC	$P = R_0 f_{\rm NPP} f_{\rm root} f_T Q_{10}$	$O = f_{[CH_4]}Q_{10}$	-	$F = (P - O)f_{\text{transport}}$



Table 6. Explanation of variable names used in Table 5.

Variable name	Description
C _{labile}	labile carbon pool
$O_{\rm air}$	atmospheric oxidation, i.e. CH ₄ uptake
$O_{\rm air,max}$	maximum atmospheric oxidation rate
O _{soil}	oxidation in the soil pore water
$O_{\rm soil.max}$	maximum oxidation rate in the soil pore water
$O_{\rm trans}$	oxidation associated with transport through plants
$O_{ m trans,max}$	maximum oxidation associated with transport through plants
O_{\max}	maximum oxidation rate
P _{max}	maximum CH ₄ production
Q_{10}	factor describing dependence on temperature
R _{hetr}	heterotrophic respiration
R_0	CH_4 production rate
f _[CH4]	function of pore water CH ₄ concentration
$f_{\text{atm[CH}_4]}$	function of atmospheric CH ₄ concentration
f _{ecosys}	function of ecosystem type
f _{GPP}	function of the ratio of monthly to annual net primary production (NPP)
f _[O2]	function of pore water O_2 concentration, determined by rate of O_2
	diffusion through soil water and aerenchyma
f _{pE}	function of alternative electron acceptors
f _{pH}	function of pH value
f _{planttrans}	function of plant-mediated CH ₄ transport
froot	function of vertical root distribution
t _{soil}	function of soil type
t _T	function of temperature
t _Θ	function of soil moisture
T _{transport}	function of CH ₄ transport
I _{WTP}	runction of water table position
r _{CH4:C}	fraction of C converted to CH ₄
<i>r</i> _{O₂}	traction of O_2 used for OH_4 oxidation





Fig. 1. Mean annual maximum fraction of inundated land between 1993 and 2004 identified by a multi-satellite approach (Papa et al., 2010).





Fig. 2. Fraction of land covered by northern peatlands at present (Tarnocai et al., 2007, 2009).











Fig. 4. Annual maximum fraction of land covered by rice fields (Leff et al., 2004).





Fig. 5. Fractional gridcell covered by permanent water bodies or ice not considered to be wetlands (GICEW, http://luh.sr.unh.edu/). The Greenland ice sheet is masked out.





Fig. 6. Caption on next page.



Fig. 6. Conceptual overview of how the participating models derive their "CH₄ producing area" (MPA). Each model (and in some cases, version) is represented by a different colour. The flow of a particular model starts with the model's name and ends at the "CH₄ producing area". The simplest models rely on "Prescribed constant wetland extents" to define their MPAs (LPJ-WHyMe, LPJ-Bern peatlands, IAP-RAS), whereas UW-VIC uses "Prescribed constant wetland extents" only as maximal boundaries. LPJ-WSL and LPJ-Bern wetlands use remotely-sensed inundation directly. "Soil moisture" is exclusively simulated by a "Hydrological model" and is used to either derive "Unsaturated" MPAs directly (LPJ-Bern wetsoils) or in combination with "Topography" (UVic-ESCM). Of the remaining models that use "Topography", all but SDGVM depend on "Remotely-sensed inundation" data to arrive at the "Water table position", which CLM4Me, DLEM, ORCHIDEE, UW-VIC and SDGVM use in combination with the other factors (e.g. CLM4Me also uses runoff) to determine "Unsaturated" and "Saturated/inundated" MPAs. LPJ-WSL (Exp. 3) uses "Water table position" to obtain only the "Saturated/inundated" MPA. The order in which processes are calculated do not always strictly follow the path used in this schematic representation.





Fig. 7. Caption on next page.



Fig. 7. Conceptual overview of the pathways from the carbon source to "CH₄ flux" (CH₄ flux per m² of wetland) in the participating models. Each model (and in some cases, version) is represented by a different colour. The flow of a particular model starts with the model's name. The hatching of the lines indicates that the CH_4 model is not influenced by wetland hydrology (beyond changes in extent). "Wetland PFTs" means that the model uses wetland-specific PFTs, whereas "Upland PFTs" indicates that the model uses the already existing PFTs used for upland ecosystems. "NPP" stands for net primary production, "Exudates" are root exudates carbon pool. All models but the IAP-RAS model use NPP as a precursor of the carbon used directly in CH₄ production or indirectly in CH₄ production by estimating "Litter and Soil C", "Exudates", and "Heterotrophic respiration". The models then calculate " CH_4 production" and the oxidation based on the equations given in Table 5. Some models include the effect of "Transport" mechanisms explicitly, whereas others include transport only implicitly by either producing less CH_4 or oxidizing it before emitting it to the atmosphere. All models use some sort of temperature dependence when calculating NPP, heterotrophic respiration, and/or CH₄ production. In this figure, LPJ-Bern "nonpeat" includes both wetlands and wetsoils, which also incorporate plant exudates (graphical simplification).





Fig. 8. Mean annual maximum extent of prescribed or simulated wetland area and mean annual CH_4 flux for Experiment 1-equil over the 1901–1931 period. Global total wetland area (Mkm² = Million km²) and CH_4 emissions (Tg = Tg CH₄ per year) have been added to each plot.







Fig. 8. Continued.