



**HAL**  
open science

# Microtubule remodelling as a driving force of axon guidance and pruning

Melody Atkins, Xavier Nicol, Coralie Fassier

► **To cite this version:**

Melody Atkins, Xavier Nicol, Coralie Fassier. Microtubule remodelling as a driving force of axon guidance and pruning. *Seminars in Cell and Developmental Biology*, In press, 10.1016/j.semcdb.2022.05.030 . hal-03799094

**HAL Id: hal-03799094**

**<https://hal.science/hal-03799094v1>**

Submitted on 6 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# 1 **Microtubule remodelling as a driving force of axon guidance and pruning**

2 Melody Atkins<sup>1</sup>, Xavier Nicol<sup>2</sup>, Coralie Fassier<sup>2</sup>

3 1: INSERM, UMR-S 1270, Institut du Fer à Moulin, Sorbonne Université, F-75005 Paris,  
4 France

5 2: Institut de la Vision, Sorbonne Université, INSERM, CNRS, F-75012 Paris, France

6 \*Correspondence should be addressed to CF (e-mail: coralie.fassier@inserm.fr).

## 7 **Abstract**

8 The establishment of neuronal connectivity relies on the microtubule (MT) cytoskeleton,  
9 which provides mechanical support, roads for axonal transport and mediates signalling events.  
10 Fine-tuned spatiotemporal regulation of MT functions by tubulin post-translational  
11 modifications and MT-associated proteins is critical for the coarse wiring and subsequent  
12 refinement of neuronal connectivity. The defective regulation of these processes causes a  
13 wide range of neurodevelopmental disorders associated with connectivity defects. This review  
14 focuses on recent studies unravelling how MT composition, post-translational modifications  
15 and associated proteins influence MT functions in axon guidance and/or pruning to build  
16 functional neuronal circuits. We here summarise experimental evidence supporting the key  
17 role of this network as a driving force for growth cone steering and branch-specific axon  
18 elimination. We further provide a global overview of the MT-interactors that tune developing  
19 axon behaviours, with a special emphasis on their emerging versatility in the regulation of MT  
20 dynamics/structure. Recent studies establishing the key and highly selective role of the tubulin  
21 code in the regulation of MT functions in axon pathfinding are also reported. Finally, our  
22 review highlights the emerging molecular links between these MT regulation processes and  
23 guidance signals that wire the nervous system.

## 24 **Key words**

25 Microtubules; Axon guidance; Axon pruning; Microtubule-associated proteins; tubulin  
26 isotypes; tubulin posttranslational modifications.

27

## 28 **Introduction**

29 The nervous system governs vital functions as well as complex behaviours enabling  
30 animals to interact with their environment. The mature nervous system is composed of  
31 billions of neurons that are interconnected according to a highly precise map (*i.e.* the  
32 connectome). Fine-tuned regulation of these connections is crucial for the accurate  
33 functioning of the network. Neuronal connectivity is established during development by  
34 linked processes of axon specification, outgrowth/navigation, synapse formation and pruning  
35 of exuberant connections and undergoes limited remodelling in the mature central nervous  
36 system [1,2]. Consequently, ectopic, exuberant or imprecise connections can lead to major  
37 neurodevelopmental disorders [3–6]. A crucial step in the assembly of neuronal circuits lies in  
38 the ability of axons to accurately navigate towards their appropriate targets, which can be  
39 located over very long distances (*i.e.* more than one meter). During their road trip in the  
40 developing embryo, the distal tips of growing axons (*i.e.* the growth cone) undergo cycles of  
41 growth, pausing, turning or retraction behaviours, which are dictated by a myriad of physical,

42 mechanical and chemical cues present in their extracellular environment. Different classes of  
43 chemical cues are encountered by the growth cone along its journey such as permissive (*e.g.*  
44 Netrin-1) or adhesive molecules (*e.g.* CAMs), extra-cellular matrix components (*e.g.*  
45 laminin), anti-adhesive substrate-bound cues (*e.g.* Slits) and diffusible chemotropic cues (*e.g.*  
46 Semaphorins, morphogens, neurotrophic factors), which will elicit growth cone attraction or  
47 repulsion [2,7,8]. Importantly, several studies have demonstrated that the response of  
48 attraction versus repulsion is not due to the intrinsic property of a particular cue. Rather, it is  
49 due to the specific growth cone receptors and downstream effectors engaged, including  
50 cytoskeleton-associated proteins [9,10; Figure 1 and Figure 2].

51 While the repertoire of guidance cues and associated receptors wiring the nervous system  
52 has mostly been identified, the intracellular pathways that modulate axon responsiveness to  
53 guidance signals are far from being deciphered. Notably, how guidance signals are integrated  
54 and translated into the cytoskeleton remodelling that underlies growth cone mechanical  
55 behaviours remains largely unknown. Pioneer studies aiming at linking guidance signals to  
56 the cytoskeleton remodelling first focused on the actin cytoskeleton [11,12]. Since then,  
57 several cues have been shown to drive growth cone attraction, repulsion or collapse through  
58 RhoGTPase-mediated actin remodelling [for review, see 13–15]. By contrast, molecular links  
59 between guidance signals and the MT cytoskeleton have remained largely unstudied until  
60 recently. This is mostly due to the fact that unlike F-actin, MTs have only recently emerged as  
61 key driving forces and direct targets of guidance cues in axon pathfinding and pruning  
62 [16,17]. Indeed, several MT-interacting proteins have been identified as decisive players in  
63 these developmental processes *in vitro* and/or *in vivo* [18–27]. Furthermore, tubulin isoforms  
64 and posttranslational modifications have lately come out as key regulators of MT functions in  
65 the axon navigation processes influencing MT properties as well as the activity/MT binding  
66 affinity of MT-interacting proteins [28–31]. This review summarises the main experiments  
67 that have established MT remodelling as a key driving force in axon guidance and pruning  
68 and provides a detailed overview of the molecular mechanisms that tune MT properties and  
69 functionalities to steer developing axons or prune exuberant connections.

70

## 71 **I. Neuronal microtubules: Key cytoskeletal elements with various properties and** 72 **functionalities**

73 Microtubules (MTs) are one of the major cytoskeletal components of eukaryotic cells and are  
74 essential for cell morphology, movement and intracellular trafficking. MTs are hollow 25 nm  
75 diameter tubes composed of polarised linear protofilaments built of head-to-tail assembled  $\alpha$ -  
76 and  $\beta$ -tubulin heterodimers [32]. MTs are highly dynamic polymers that stochastically switch  
77 between polymerisation and depolymerisation, a process known as dynamic instability. This  
78 dynamic property is driven by GTP binding to the  $\alpha\beta$ -tubulin dimer and its hydrolysis into  
79 GDP. The transition from MT growth to shrinkage is called catastrophe, while the reverse  
80 reaction is termed rescue [33]. Since their discovery in 1950, MTs have emerged as critical  
81 structures in neuronal development and function, providing mechanical support and rails for  
82 axonal transport, as well as mediating key signalling pathways. The fine-tuned spatiotemporal  
83 regulation of MT structure, mechanical properties, dynamics and crosstalk with the actin  
84 cytoskeleton (F-actin), which is mediated by tubulin isotypes, posttranslational modifications  
85 and MT-associated proteins (MAPs), is decisive to build functional neuronal circuits [34].  
86 This is highlighted by the increasing number of neurodevelopmental disorders (*e.g.* ASD,  
87 schizophrenia, lissencephaly...) caused by mutations in genes encoding the MT building  
88 blocks (*i.e.* tubulins), tubulin-modifying enzymes, or MT-regulatory proteins [35–37].

89 Over the past decades, the emergence of novel imaging technologies with high spatiotemporal  
90 resolution (correlation EM microscopy, TIRF, super-resolution, expansion microscopy, motor

91 painting approach, cryoelectron tomography) has gradually added to the complexity of the  
92 neuronal MT architecture and function [38–44]. Unlike what was initially thought, the MT  
93 cytoskeleton is not a homogeneous network, but rather a highly mosaic array composed of  
94 several subpopulations of MTs. These subpopulations (i) are decorated by different  
95 posttranslational modifications (PTMs) and MT-associated proteins [MAPs; 34,38,45,46] and  
96 (ii) exhibit distinct structural/dynamic properties and functionalities. In axons, unlike  
97 dendrites, MTs are organised in a highly polarised manner with a fast-growing end (plus end)  
98 pointed outwards, towards the synapse, and a slow-growing end (minus end) pointed inwards,  
99 towards the soma. This typical polarity pattern of axonal MTs has recently been shown to rely  
100 on non-centrosomal MT nucleation, mediated by the augmin- $\gamma$ TuRC complex [47].  
101 Furthermore, the axonal MT network is organised into longitudinal parallel bundles and  
102 consists of a mixed population of stable (*i.e.* long persistence) MTs decorated with distinct  
103 posttranslational modifications, including acetylation, detyrosination, glutamylation. While  
104 some of these PTMs influence the binding affinity and/or activity of several MAPs *in vitro*  
105 [46,48,49], the functional specificity of these different subtypes of stable axonal MTs remains  
106 mostly unknown under physiological conditions [50]. The distal tip of developing axons (*i.e.*  
107 the growth cone) is divided into three compartments, which are characterised by their specific  
108 composition and arrangements of cytoskeletal elements [51; Figure 1 and Figure 2]. The  
109 central domain (C-domain) exclusively contains stable and acetylated MTs. These polymers  
110 extend from the axon shaft and are locally restricted and bundled by an actin barrier (*i.e.* actin  
111 contractile arcs) present at the transition zone [T-zone; 52,53]. In this T-zone, the local  
112 coupling between MT growing plus ends and the actin retrograde flow drags MTs back into  
113 the C-domain, thereby inducing MT looping and bending. The peripheral domain (P-domain)  
114 of the growth cone is composed of an F-actin branched network called lamellipodia-like veils  
115 and long bundles of F-actin forming the filopodia. These F-actin-based structures are locally  
116 intermingled with a few individual MTs (also called pioneer MTs) that have escaped from the  
117 C-domain. In contrast to MTs that are constrained to the C-domain, MTs that penetrate the  
118 actin-rich P-domain are tyrosinated and highly labile [*i.e.* dynamic; 52,53]. Notably, this  
119 subpopulation of MTs are often found in close apposition with F-actin bundles in filopodia,  
120 where their growing plus tips sometimes reach the growth cone margin. Their local exposition  
121 to extracellular cues makes them relevant candidates to integrate axon guidance signals [53–  
122 55].

123 The following sections will emphasise how the characteristic organisation and heterogeneity  
124 of the neuronal MT cytoskeleton are critical for its functions in axonal navigation.

## 125 **II. Microtubules in axon guidance: the outside players**

### 126 **II.1 Instructive role of microtubules in growth cone steering**

127 MTs, together with the actin cytoskeleton, are major determinants of the mechanical  
128 properties of growing axons. Despite this evidence, the role of MTs as driving forces of  
129 growth cone mechanical turning has long been disregarded compared to that of the actin  
130 cytoskeleton. Indeed, due to their highly dynamic and protrusive behaviours at the leading  
131 edge of the growth cone and their well-established role in cell migration, actin filaments (F-  
132 actin) have long held the leading (if not exclusive) role in growth cone steering processes. By  
133 contrast, MT remodelling was thought to occur in response to F-actin changes and to be  
134 required to consolidate the novel portion of the axon shaft and stabilise growth in the  
135 appropriate direction. MTs were first postulated to play an instructive role in growth cone  
136 steering decisions in the late 1990s. This hypothesis was based on live imaging studies from  
137 the Kirschner lab revealing that in advancing growth cones, a few individual MTs

138 dynamically explore the P-domain and that their growth orientation often precedes growth  
139 cone turning and predicts the steering direction [56,57]. Several groups subsequently showed  
140 that the asymmetric invasion and stabilisation of these dynamic MTs in filopodia facing an  
141 adhesive/attractive cue or located on the opposite side of a repulsive cue is decisive to drive  
142 growth cone turning towards or away from the cue, respectively [58–61,62; Figure 1 and  
143 Figure 2]. The first experimental evidence establishing the instructive role of MTs in growth  
144 cone decision-making behaviours comes from the elegant work of Buck and Zheng in 2002.  
145 Their study shows that modulation of MT dynamics on one side of the growth cone by photo-  
146 uncaging a MT-stabilising (*i.e.* Taxol) or destabilising agent (*i.e.* Nocodazole) is sufficient to  
147 induce growth cone turning towards or away from the treated side, respectively [16]. These  
148 data have been strengthened by the work of several groups revealing that (i) asymmetric  
149 manipulation of MT-associated protein activity with MICRO-CALI techniques is sufficient to  
150 drive growth cone turning [63–65] and that (ii) MTs (more precisely, brain specific tubulin  
151 isoforms) can be direct targets of guidance cues in axon navigation processes [29]. However,  
152 it is worth noting that the instructive role of MTs in growth cone steering critically relies on  
153 their fine-tuned coupling to the actin network. The latter either acts as a barrier preventing  
154 premature MT invasion of the P-domain or as a guide facilitating MT growth and stabilisation  
155 within filopodia [62,66,67]. Notably, the myosin II-based actin retrograde flow, which is  
156 regulated by the RhoA GTPase, has been shown to play a significant role in restricting and  
157 bundling stable MTs within the C-domain [68]. It has further been shown to regulate dynamic  
158 MT growth alongside filopodia [53,55,69–71].

159 Despite the now well-established role of MTs as driving forces of axon guidance, the  
160 numerous actors that control MT remodelling and interplay with F-actin in navigating axons,  
161 as well as their specific mode of action and regulation by guidance cues are only starting to be  
162 deciphered. This is mostly explained by the technical challenge associated with the  
163 monitoring of these nanoscale structures with high spatiotemporal resolution in growth cones  
164 while locally applying guidance cue *in vitro* and above all in single-labelled axons navigating  
165 in a complex *in vivo* environment. However, over the past decade, the emergence of novel  
166 microscopy approaches such as super resolution microscopy (e.g. SIM, STED) and live TIRF  
167 imaging as well as of optically transparent vertebrate models with straightforward genetic  
168 manipulations (*e.g.* zebrafish and *Xenopus* models) have allowed several groups to overcome  
169 the above-mentioned limitations. This has contributed to provide novel pieces of information  
170 regarding the regulation of MT functions in axon guidance and targeting [22,24–26,72–74].

## 171 **II.2 Microtubule-interacting proteins: key effectors of guidance signals in growth cone** 172 **steering**

173 In neurons, MT organisation, dynamics and cross-talk with F-actin are tightly regulated by a  
174 myriad of MT-interacting proteins. The vast majority of them, including MT plus-end  
175 tracking proteins (+TIPs), structural MAPs, MT-depolymerising and severing enzymes, or  
176 molecular motors, have been shown to play critical roles in axon extension. However, only  
177 few of them have been reported to influence growth cone steering *in vitro* or to ensure the  
178 fidelity of axon targeting *in vivo*. We will here focus on the different classes of MT-  
179 interacting proteins whose regulation of MT functions in neurons is crucial to evoke growth  
180 cone behaviours (Table 1). It is worth noting that even if these MT-interactors remain  
181 classified in different functional categories, data collected over the past years have gradually  
182 revealed their functional versatility in neuronal circuit wiring and homeostasis.

### 183 ***II.2.1 Plus-end tracking proteins as key signalling integrators in axon navigation***

184 MT plus-end tracking proteins (+TIPs) bind the growing plus end of MTs, which can  
185 extend to the growth cone periphery where guidance receptors reside. This specific  
186 subcellular distribution combined with their ability to regulate MT dynamics, anchoring at the  
187 cell cortex and interplay with F-actin makes them ideal signalling integrators of axon  
188 guidance processes. CLASP proteins, for cytoplasmic linker protein (CLIP)-associated  
189 protein, were the first +TIPs to be implicated in axon navigation. In 2004, the *Drosophila*  
190 CLASP orthologue was identified as a downstream effector of the Slit/Robo repellent  
191 pathway and the Abelson (Abl) tyrosine kinase during midline crossing [19]. This new role in  
192 axon repulsion came as a surprise given the previously identified role of CLASP in MT rescue  
193 and stabilisation [75–78]. A later study, conducted during axon elongation, has nevertheless  
194 offered possible keys to the problem, reporting two distinct MT-binding activities for CLASP  
195 in nerve growth cones: both at MT plus ends and on the MT lattice [79]. While GSK3 $\beta$   
196 inhibition induces CLASP binding to the MT lattice and impedes growth cone advance,  
197 CLASP plus-end activity enables axon elongation by stabilising MTs. Further supporting a  
198 dual CLASP binding activity, elevation of Abl activity in spinal cord growth cones has also  
199 been shown to delocalise CLASP away from MT plus ends, in association with slower growth  
200 cone advance [80]. Abl has moreover been found to bind and phosphorylate CLASP in  
201 response to extracellular signals, such as serum or PDGF. Finally, using biochemical assays  
202 as well as culture experiments, the authors suggest a role for the Abl-induced CLASP  
203 phosphorylation in CLASP-mediated MT-F-actin crosslinking. While such studies  
204 highlight CLASP as a modulator of MT dynamics downstream of guidance signals, the role of  
205 its plus-end binding activity – rather than its MT lattice activity – needs further  
206 characterisation, especially in a context of axon navigation (Figure 2C).

207 The adenomatous polyposis coli (APC) protein is another +TIP involved in axon navigation.  
208 Originally identified as a tumour suppressor mutated in colorectal cancers [81], APC has been  
209 reported in many studies to act in synergy with EBs (End-Binding proteins) to promote MT  
210 stabilisation, in addition to its role as a regulator of the Wnt/ $\beta$ -catenin canonical pathway [82–  
211 84]. Zhou and colleagues were the first to report a role for APC at MT plus ends in axon  
212 elongation, downstream of NGF and its GSK3 $\beta$  effector [69]. Since then, studies have  
213 involved APC in growth cone steering *per se*. Indeed, local MICRO-CALI inactivation of  
214 APC in one half of retinal ganglion cell growth cones is sufficient to regulate axon turning  
215 [64]. Wnt3a has also been shown to delocalise APC from MT plus ends in DRG sensory  
216 neurons, resulting in a MT growth directionality and lopping phenotype that resembles APC  
217 loss of function [85]. Similarly, APC2, an APC-like protein [86], is also important in the  
218 ephrin-A2-induced repulsive guidance of chick retinal axons [87; Figure 2D], and to promote  
219 *Drosophila* axon growth across the midline [27]. Of note, studies have implicated APC in the  
220 crosslinking of the MT and actin cytoskeleton during cell migration [88–91]. Given the  
221 importance of this crosstalk during axon navigation, future investigations will be of interest to  
222 assess its influence on axon pathfinding.

223 Importantly, a study in *Drosophila* has shown that Msps (minispindles), the fly ortholog of  
224 the +TIP and MT polymerase XMAP215, physically interacts with CLASP, and functions  
225 during axon guidance by antagonising both CLASP and Abl activity [92]. The authors  
226 propose a model in which CLASP and Msps converge in an antagonistic manner on the Abl  
227 signalling pathway to regulate midline crossing. A study in *Xenopus laevis* embryos further  
228 confirms this role for XMAP215 in axon guidance. Indeed, using SIM super-resolution  
229 microscopy and live imaging in cultured spinal cord explants, the Lowery lab reveals that  
230 XMAP215 promotes MT/F-actin interactions and is required to regulate growth cone  
231 morphology and dynamic behaviour. They further show that XMAP215 accurately mediates  
232 repulsion to the ephrinA5 guidance cue in an *in vitro* stripe assay [25]. Importantly, as for

233 CLASP, a dual mode of XMAP215 binding to MTs has been identified, on the MT lattice and  
234 at the MT plus ends in the growth cone [93]. Both modes of MT binding appear to regulate  
235 MT-F-actin coupling, in addition to the known XMAP215 MT polymerase activity at plus  
236 ends (Figure 2D). Understanding how these different XMAP215 functions are coordinated  
237 and synchronised by intracellular signalling pathways to regulate different aspects of axon  
238 outgrowth and/or pathfinding requires further investigations. It is likely that a  
239 phosphorylation-based regulation of XMAP215 by kinases acting downstream specific  
240 guidance cues such as Abelson may regulate its subcellular targeting and coupling with MTs  
241 and/or F-actin, as shown for CLASP and APC [53].

242 A known binding partner of XMAP215 is TACC3, a member of the family of proteins  
243 carrying a transforming acidic coiled-coil (TACC) domain. TACC3 was initially known as a  
244 centrosome-associated protein regulating MT growth and stability during mitosis, at least in  
245 part via its interaction with XMAP215 [94–97]. However, TACC3 has also been identified as  
246 a +TIP that colocalises with XMAP215 in vertebrate growth cones and is required to enhance  
247 the MT growth dynamics influencing axon outgrowth [98]. Since then, the same group has  
248 explored the role of TACC3 in axon steering. Erdogan and colleagues report that TACC3  
249 depletion induces spinal motor axon misrouting in whole-mount *Xenopus* spinal cords.  
250 Consistently, TACC3 overexpression leads to an increased resistance to repulsive signal-  
251 induced growth cone collapse, such as Slit2 or ephrin-A5 [21,99; Figure 2C]. Interestingly,  
252 TACC3 phosphorylation by the Abl tyrosine kinase has also been involved in axon outgrowth  
253 [99]. Altogether, these results suggest that TACC3 could be acting in concert with XMAP215  
254 to drive the MT remodelling required for growth cone responsiveness to specific repellent  
255 signals such as ephrin-A5. However, further investigations will be needed to confirm the  
256 physiological relevance of the TACC3-XMAP215 complex in a context of axon guidance.

257 Another +TIP, navigator-1 (NAV1), has recently been reported to play a role in axon  
258 navigation via its MT/actin crosslinking activity. NAV proteins were initially described as  
259 +TIPs regulating MT bundling and stabilisation [100,101]. Despite a few studies reporting  
260 misguided neuronal tracts in *C. elegans* or mouse mutant animals for Nav proteins [102,103],  
261 most studies have so far focused on a role for these +TIPs in axon outgrowth, and the precise  
262 cellular and molecular mechanisms involving Nav proteins in axon navigation have remained  
263 unexplored until recently. Sanchez-Huertas et al. have recently identified NAV1 as a MT-  
264 actin crosslinker required to promote MT persistence in the P-domain of exploring cortical  
265 growth cones and to elicit axon attractive turning in response to Netrin-1 gradients in Dunn  
266 chamber assays [26; Figure 1B]. More studies will be required to explore the role of other  
267 NAV proteins in axon pathfinding, and to decipher the signalling cascades that could regulate  
268 their activities.

269 Similarly to NAV proteins, the spectraplakin +TIPs have mostly been studied for their role in  
270 axon extension, despite a few reports mentioning guidance defects in loss of function models  
271 [104–107]. Interestingly, during axon outgrowth, as for many other +TIPs, a role for  
272 spectraplakins in MT-actin crosstalk has been described [108–110], as well as a dual MT  
273 lattice and plus-end binding activity [111]. Exploring the mechanisms that regulate this  
274 functional switch therefore appears of particular interest in a context of axon guidance.  
275 Finally, the *Drosophila* Shot spectraplakin has also been shown to support midline axon  
276 repulsion via its binding to the Krasavietz (Kra) translation inhibition factor [106], suggesting  
277 an additional role for +TIPs in mediating local translation during growth cone steering.

278 Surprisingly, while the core proteins of MT plus-end complexes (EBs) act as hubs locally  
279 recruiting many of the above mentioned +TIPs, functional evidence demonstrating their role  
280 in growth cone steering are still lacking. Nevertheless, three MT-interacting proteins critical

281 for the guidance of different populations of axons (MAP1B, Tau and Fidgetin-like 1) have  
282 been reported to regulate EB binding at MT plus ends through distinct mechanisms [22,112–  
283 114], suggesting a key role for these MT plus-end scaffolding proteins in axon guidance.  
284 Altogether, data from the past decade reveal the diversity of processes that +TIPs may  
285 coordinate at the cell cortex to promote steering events. They further highlight the importance  
286 of their fine-tuned subcellular targeting and activity for proper neuronal circuit development.  
287

## 288 ***II.2.2 Structural MAPs in axon guidance: more than just MT stabilisers***

289 Structural MAPs form a family of MT-interacting proteins whose characteristics have  
290 considerably evolved since their discovery in the 1970s [115,116]. Early on, structural MAPs  
291 have been defined as proteins that can be co-purified with MTs and that have the ability to  
292 promote the assembly and stabilisation of these polymers. However, during the past 20 years,  
293 the extensive characterisation of neuronal MAPs has gradually expanded their functional  
294 diversity, unravelling among others their roles as key MT/actin crosslinkers [24,117] and as  
295 signalling molecules in developing neurons [118].

296 MAP1B was the first MT stabilising protein involved in axon guidance *in vivo*. Its loss of  
297 function leads to white matter deficits in mice and humans, including corpus callosum  
298 agenesis [119,120]. MAP1B is highly expressed in growth cones where it is required for the  
299 Netrin-1-induced outgrowth of cortical axons [20], the guidance of DRG axons facing a non-  
300 permissive substrate [121], as well as for nitric oxide- or Draxin-evoked growth cone collapse  
301 [122,123]. Notably, MAP1B functions in axon guidance are critically dependent on its  
302 posttranslational modifications [124]. Asymmetric blockage of phospho-MAP1B activity in  
303 growth cones using MICRO-CALI approaches is sufficient to induce growth cone turning *in*  
304 *vitro* [63]. Consistently, GSK3 $\beta$ -mediated phosphorylation of MAP1B regulates the dynamics  
305 and spatial distribution of tyrosinated MTs in growth cones [125] and is necessary for both  
306 Netrin-1- and Draxin-evoked axon outgrowth and repulsion, respectively [20,122; Figure 1B  
307 and Figure 2F]. By contrast, MAP1B S-nitrosylation mediates nitric oxide-induced axon  
308 retraction [122; Figure 2E]. Furthermore, in developing axons, MAP1B was reported on the  
309 one hand to enhance the assembly rate of tyrosinated MTs [125] and on the other to  
310 negatively regulate EB binding at MT plus ends via EB sequestration [112]. While these  
311 independent studies might at first glance appear contradictory, they both converge on the  
312 same conclusion that physiological levels of MAP1B are critical to maintain MT dynamic  
313 growth and proper axon extension. Further challenging its role as a structural MAP, MAP1B  
314 has been suggested to influence MT/actin crosstalk [126]. However, whether MAP1B directly  
315 or indirectly (*i.e.* via its influence on EBs) modulates MT/actin interplay in growth cones and  
316 whether this emerging function of MAP1B is required for axon responsiveness to the Netrin-  
317 1/GSK3 $\beta$  or Draxin/GSK3 $\beta$  pathways remains to be assessed.

318 During the past decades, particular attention has been paid to the Tau protein due to its  
319 implication in several neurodegenerative disorders. Tau is a structural MAP, which was first  
320 shown to crosslink protofilaments within MTs. Tau has then been shown to increase MT  
321 polymerisation, prevent MT shrinkage [127] and regulate MT network organisation [128].  
322 Interestingly, as for MAP1B, Tau has been shown to reduce EB binding at MT plus ends  
323 [113,114], supporting an emerging key role for structural MAPs in fine tuning MT dynamics  
324 through the regulation of EB levels at MT plus ends. Although mainly studied for its role in  
325 neuronal homeostasis, Tau is highly expressed in the developing brain where it is required for  
326 Wnt5a-evoked cortical axon outgrowth and repulsive turning [72]. In this context,  
327 phosphorylation of the Tau MT binding domain at Ser262 by CAMKII (*i.e.* phosphorylation  
328 that impacts Tau binding to MTs *in vitro*) is required for the reorientation and bundling of



329 dynamic MTs that drive Wnt5a-induced axon extension and repulsive steering [72; Figure  
330 2G]. Elie et al. have moreover revealed that Tau directly binds to F-actin, promotes MT/F-  
331 actin crosslinking and regulates the co-polymerisation of both filaments in cell-free  
332 reconstructed systems [117]. Consistently, using super-resolution STED microscopy, Biswas  
333 and colleagues have shown that Tau localises to growth cone regions characterised by an  
334 extensive crosstalk between these two networks (*i.e.* T-zone and filopodia). Its loss of  
335 function further impairs MT growth alongside F-actin bundles in growth cone filopodia [24].  
336 Notably, by combining Dunn chamber guidance assays with live TIRF imaging of  
337 cytoskeletal dynamics, they show that Tau-mediated regulation of MT invasion in filopodia is  
338 critical for both Wnt5a-induced axon extension and directional turning [24; Figure 2G].  
339 Interestingly, Tau has recently been shown to act in concert with the above mentioned +TIPs,  
340 XMAP215/Msps and EB1 to regulate MT polymerisation and bundling in developing axons,  
341 both *in vitro* and *in vivo* [129]. While these proteins have independently been involved in  
342 axon guidance, whether their cooperative role in the regulation of MT dynamics/organisation  
343 is required to drive the growth cone turn in response to specific extracellular cues remains to  
344 be confirmed.

345 MAP6 (also known as STOP) is a versatile regulator of neuronal connectivity that was first  
346 identified as a MT stabilising protein required for MT protection against cold- and  
347 nocodazole-induced depolymerisation [130,131]. Neuroanatomical analysis of MAP6  
348 knockout brains reveals substantial hypoplasia of several commissures and axonal tracts,  
349 including the absence of the post-commissural part of the fornix [132–134]. Regarding this  
350 latter phenotype, MAP6 is required for the Sema3E-induced axon outgrowth of subicular  
351 neurons (*i.e.* axons that form the fornix). Notably, Deloulme and colleagues have  
352 unexpectedly revealed that the function of MAP6 as a downstream effector of Sema6E is  
353 independent of its MT-binding domains. Rather, via its N-terminal prolin-rich domain, MAP6  
354 acts as a signalling hub promoting interactions between components of the Sema6E receptor  
355 complex and key downstream effectors during axon guidance [e.g. PI3K, Src; 130; Figure  
356 1C]. However, they identify the MAP6-interacting MAP CRMP4 as the final effector of this  
357 Sema6E signalling pathways directly connecting the Sema6E receptor complex to the MT-  
358 cytoskeleton in extending subicular neurons [134; Figure 1C]. Interestingly, the same group  
359 has recently revealed another functional facet of MAP6 in neuronal connectivity. Indeed,  
360 neuronal activity-induced phosphorylation of MAP6 by CAMKII relocalises MAP6 to  
361 dendritic spines, where it locally binds to F-actin and stabilises this network to adapt dendritic  
362 spine morphology [135]. Since MAP6 also binds to F-actin in growth cones, it will be of  
363 particular interest to determine whether its ability to switch from one cytoskeleton to another  
364 in response to specific signalling pathways influences the targeting of specific axonal  
365 populations. Another important issue regards the relevance of its emerging role as a MT  
366 intraluminal protein (MIP) in a context of neuronal circuit wiring. It will be interesting to  
367 investigate whether the suggested ability of luminal MAP6 to relieve MT mechanical tension  
368 by promoting MT lattice torsion [136] increases the flexibility and resistance of navigating  
369 axons to compressive forces generated by the surrounding developing tissues.

370 Collapsin response mediator proteins (CRMPs) were first identified as signalling molecules,  
371 before being classified as structural MAPs based on the ability of a subset of their members to  
372 stimulate the MT assembly and stabilisation [118,137–139]. CRMP2 has long been described  
373 as a key effector of Sema-3A repulsive cues [140,141]. In its non-phosphorylated state,  
374 CRMP2 binds to tubulin dimers and promotes their polymerisation [137]. However, upon  
375 phosphorylation by the Sema-3A downstream kinases Cdk5 and GSK3 $\beta$ , CRMP2 dissociates  
376 from MTs, thereby allowing the MT remodelling underlying Sema3A-induced axon repulsion  
377 [142,143; Figure 2B]. Supporting these findings, CRMP2 loss of function has been shown to

378 mostly affect the pathfinding of Sema3A-sensitive axonal populations both in invertebrate  
379 [144,145] and vertebrate models, including peripheral nerves [140,146] and callosal axons  
380 [146]. By contrast, CRMP4 is required for Sema6E-induced axon outgrowth of subicular  
381 neurons [134; Figure 1C]. Supporting the non-overlapping role of CRMP2 and CRMP4 in  
382 axon guidance, knockdown approaches in the zebrafish have revealed that while CRMP2  
383 regulates retinal axon outgrowth, CRMP4 promotes midline crossing, most likely downstream  
384 of Neuropilin 1 [147; Figure 1D]. Whether CRMP4-mediated regulation of growth cone  
385 decision-making behaviours at the zebrafish optic chiasm relies on its dual regulation of MT  
386 and actin dynamics in growth cones [148] needs to be clarified. Still in the zebrafish, CRMP2  
387 has been identified as an effector of the Rac-GAP  $\alpha 2$ -chimaerin in ocular motor axon wiring  
388 and has been suggested to act in concert with the MT depolymerising enzymes stathmin 1/2 in  
389 this process [149; Figure 2B]. Interestingly, CRMPs have been identified as direct binding  
390 partners and potential regulators of the MT-severing spastin [150–152], supporting possible  
391 cooperative roles for CRMPs and MT destabilising proteins in neuronal circuit wiring.

392 Doublecortin (DCX) is a MT-associated protein enriched at the distal tip of developing  
393 neurites, which mutations cause lissencephaly, a neurodevelopmental disorder characterised  
394 by severe cortical malformations [153]. DCX has been classified among structural MAPs  
395 based on its ability to promote MT assembly through catastrophe inhibition and to induce MT  
396 bundling [154–156]. DCX is critical for cortical migration (Kappeler et al., 2006; Friocourt et  
397 al., 2007). However, its MT-related functions also seem to be required for later stages of  
398 neuronal development, including axon guidance. A recent study from the Winckler lab has  
399 revealed its role as a downstream effector of the intermediate filament protein Nestin in the  
400 regulation of growth cone morphology and sensitivity to Sema3A [157]. Indeed, Nestin  
401 increases growth cone responsiveness to Sema3A by facilitating DCX phosphorylation by the  
402 Cdk5 kinase through scaffolding activity. However, the molecular mechanisms involved  
403 remain to be clarified. The authors suggest that Nestin-dependent phosphorylation of DCX by  
404 Cdk5/p35 might release MTs from DCX stabilisation/protection by promoting DCX  
405 detachment from MTs as shown for CRMP2 downstream of a Sema3A/Cdk5/Nestin-  
406 independent pathway. This mechanism might thereby promote the MT remodelling required  
407 for Sema3A-evoked axon repulsion [157; Figure 2B].

408 Interestingly, among their emerging functions, some of the above-mentioned structural MAPs  
409 (*e.g.* Tau and CRMPs) influence the activity of MT severing or depolymerising enzymes in  
410 developing axons [150,151,157], which paves the way for future investigations of their  
411 cooperative or antagonistic roles in axon navigation processes.

### 412 ***II.2.3 MT severing/depolymerising enzymes: Engines that power the growth cone turn***

413 MT severing enzymes including spastin, p60-katanin and fidgetin are nanomachines from the  
414 AAA+ superfamily [ATPases Associated with diverse cellular Activities; 157] that combine  
415 chemical (*i.e.* ATP hydrolysis) and mechanical forces to promote the extraction of tubulin  
416 dimers from MTs through tubulin unfolding [160]. Propagation of the nanodamages generated  
417 by these enzymes leads to MT severing. Based on this functional property, MT severers are  
418 critical regulators of MT mass, length and organisation in cells [161,162] and thereby of axon  
419 elongation and/or branching [158,163–166]. Nevertheless, their ability to rapidly breakdown  
420 taxol-stabilised MTs in cell-free systems [160,167] and to dismantle the MT network when  
421 overexpressed in non-neuronal cells [168–170] positioned them as relevant candidates to  
422 promote the asymmetric disassembly of MTs that drives the growth cone turn [52]. Using  
423 several genetic approaches combined with *in vivo* live imaging in zebrafish larvae, the Hazan  
424 team has recently uncovered the decisive and non-overlapping roles of p60-katanin [31] and

425 spastin main isoforms [M1 versus M61; 23] in zebrafish motor circuit wiring and larval  
426 locomotion. They show that loss of function of spastin main isoforms leads to highly specific  
427 pathfinding errors of rostral motor nerves, while p60-katanin deficiency mainly impairs dorsal  
428 motor nerve targeting (*i.e.* abnormal splitting). This group further reveals that the functional  
429 specificity of these severers in zebrafish axon guidance relies on their preference for specific  
430 populations of MTs and on their selective involvement in guidance signalling pathways.  
431 Indeed, they identify TLL6-mediated MT-polyglutamylation as a selective regulator of p60-  
432 katanin activity in motor axon guidance [31; Figure 3B]. Moreover, they uncover a selective  
433 role for M1 and M61 spastin as a BMP signalling inhibitor and a Neuropilin-1 downstream  
434 effector, respectively [23; Figure 2B]. Although not investigated in these studies, the selective  
435 regulation of the p60-katanin and spastin MT-severing activity by structural MAPs [*e.g.* Tau  
436 or CRMP proteins, respectively; 143–145,155] might also underlie their specificity of action  
437 in axon pathfinding, as suggested by the overlapping motor neuron phenotypes associated  
438 with Sema3A [171], CRMP2 [172] and spastin M61 knockdown [23] in zebrafish embryos.  
439 Nevertheless, an important unsolved issue concerns the molecular mechanisms by which MT  
440 severers drive growth cone turning. Spastin and p60-katanin knockdown drastically reduce  
441 the number of dynamic growing plus ends in zebrafish motor neuron axons, a phenotype  
442 consistent with a lack of MT severing activity [165]. However, depending on whether newly  
443 generated ends are stabilised or not, MT-severing enzymes can either promote MT breakdown  
444 or on the contrary, increase MT mass [161,173]. Furthermore, the *Drosophila* orthologue of  
445 p60-katanin has been shown to have a MT plus-end depolymerising activity *in vitro*, in  
446 addition to its MT severing activity [174,175]. This dual function seems to be dependent on  
447 its concentration and might thereby underlie the dose-dependent effect of p60-katanin on  
448 motor axon targeting [31]. Finally, an elegant work from the Roll-Mecak lab - based on *in*  
449 *vitro* reconstructed systems - reveals that nanodammages generated by MT severers can be  
450 repaired through the incorporation of novel GTP-tubulin dimers. This suggests an additional  
451 role for these AAA+ machines in MT lattice renewal [176]. *In vivo* investigations in the  
452 zebrafish model, which offers the possibility to monitor cytoskeleton dynamics in navigating  
453 growth cones, should provide crucial information regarding the basic mechanisms governing  
454 this “Cut and Turn process”.

455 Fidgetin-like 1 (*fign1*) belongs to the same subfamily of AAA+ engines as spastin and p60-  
456 katanin. However, unlike its extensively studied homologues, *fign1* functions in the nervous  
457 system had never been investigated until the pioneer study from Fassier et al. in 2018. Their  
458 work identifies this MT-associated ATPase as a critical player in zebrafish motor axon  
459 pathfinding [22,73]. The authors show that *fign1* controls growth cone morphology and  
460 steering behaviours at guidance choice points through the regulation of MT plus-end  
461 dynamics. Using *in vitro* systems, they have unexpectedly discovered that unlike its related  
462 proteins, *fign1* does not sever MTs but rather regulates MT plus-end composition and  
463 dynamics. Indeed, alternative translation of the *fign1* transcript generates two isoforms with  
464 distinct functionalities. The full-length protein displaces EBs from MT plus ends in an ATP-  
465 dependent manner. By contrast, the N-terminal truncated variant exhibits an F-actin-  
466 dependent cortical MT depolymerising activity [22]. Through this functional versatility,  
467 *fign1* may concomitantly tune +TIP-dependent MT behaviours (*i.e.* MT growth  
468 directionality, MT/F-actin crosstalk and MT anchoring at the cell cortex) and promote MT  
469 dynamics in the P-domain to drive the growth cone turn. By taking advantage of the zebrafish  
470 model, the authors were able to monitor MT dynamics in navigating axons of live *fign1*-  
471 deficient larvae. The *in vivo* MT phenotype they describe, including the sustained stability of  
472 MTs, their aberrant invasion and growth directionality in the P domain, as well as the  
473 increased MT growth duration in filopodia, fits with the above-mentioned hypothesis [22].  
474 Deciphering how the actin network influences the MT-depolymerising activity of *fign1* and

475 connecting this process to guidance signals should shed new light on the regulation of the  
476 MT/actin crosstalk underlying axon guidance.

477 Another family of MT-destabilising enzymes that has recently been involved in axon  
478 guidance is the stathmin family. Stathmins promote MT depolymerisation by decreasing the  
479 concentration of free-tubulin through tubulin dimer sequestration [177]. While stathmins have  
480 long been identified as critical regulators of axon extension [178,179], Carretero-Rodriguez et  
481 al. have unravelled a role for stathmin1/2 as downstream effectors of the Rac-GAP alpha2-  
482 Chimaerin required for zebrafish oculo-motor axon fasciculation and guidance [149]. Further  
483 work will be required to define the role and regulation modes of stathmins downstream of this  
484 signalling protein.

485 Altogether, these studies emphasise the key role of alternative translation in multiplying the  
486 functional diversity of MT-destabilising enzymes in directed axon outgrowth. This conclusion  
487 is strengthened by the recent work of Atkins and colleagues, uncovering an additional role for  
488 the MT AAA+ machine *figl1* in the regulation of molecular motor motility and its decisive  
489 contribution to axon guidance [73; *See § I.2.4*].

490

#### 491 ***II.2.4 Molecular motors: Microtubule sliding and bidirectional cargo transport***

492 Navigating growth cones need to be provided with a diverse set of building and navigation  
493 material (lipids, membrane receptors, mRNAs, mitochondria...) in order to accurately  
494 respond to their environment in a spatiotemporally regulated manner. It is therefore generally  
495 assumed that molecular motors are required for axon navigation. Confirming this idea,  
496 various animal models carrying mutations affecting molecular motors or altering kinesin  
497 binding sites on specific tubulin isoforms have been associated with guidance defects [180–  
498 184]. However, the precise mode of action and molecular mechanisms through which  
499 molecular motors may control growth cone turning responses remain poorly characterised.

500 A first set of studies reports a role for molecular motors in MT sliding [for review, see 185]  
501 and asymmetric invasion of the growth cone during turning events [186–189]. In a model  
502 proposed by Baas and colleagues, extracellular guidance cues signal to the growth cone  
503 molecular motors in order to regulate this MT sliding process. In this model, dynein counters  
504 myosinII-driven forces to instruct MT invasion of the P-domain on the turning side of the  
505 growth cone. On the opposite side, kinesins 5 and 12 antagonise dynein-driven forces, thereby  
506 locally restricting MTs to the C domain [190]. These synchronised force balances between  
507 dynein and myosin II on the one hand, and dynein and kinesins on the other, could ensure the  
508 polarised MT sliding required for the growth cone to turn.

509 Importantly, other forms of axonal transport, such as receptor-containing vesicle trafficking,  
510 are also starting to emerge in a context of axon navigation. First, it has been shown that  
511 axonal transport of different cargoes can be activated by extracellular guidance cues, such as  
512 TrkB anterograde axonal transport downstream of NGF [191,192] or retrograde transport of a  
513 Plexin A4-TrkA downstream of Sema3A [193]. Selective links between specific molecular  
514 motors and cargo complexes are also just starting to emerge in a context of axon guidance.  
515 For example, in mouse embryonic cortical axons, Netrin-1 regulates axon initiation,  
516 branching and targeting by promoting anterograde transport of the Myosin X molecular motor  
517 in a KIF13B-dependent manner [194; Figure 1B]. Similarly, Saez and colleagues have linked  
518 the kinesin-1 motor to the axonal transport of cannabinoid type-1 receptors, a process required  
519 for mouse corticofugal axon pathfinding [195]. Alternatively, some studies have focused on  
520 adaptor/regulatory proteins of known molecular motors. The ATPase *figl1* has been shown  
521 in this context to restrict dynein-based retrograde transport velocity through dynein coupling  
522 to the opposite polarity-directed motor Kif1b $\beta$  thereby fine tuning bidirectional vesicular

523 trafficking *in vivo* in navigating axons of zebrafish embryos. Importantly, pharmacological  
524 inhibition of the dynein motor rescues the pathfinding defects associated with *figl1* loss of  
525 function, confirming the key role of *figl1* as a dynein speed limiter required for accurate  
526 axon targeting [73]. The kinesin-1 adaptor proteins FEZ1 (fasciculation and elongation  
527 protein zeta 1) and Calsyntenin 1 have also been involved in axon pathfinding events.  
528 Calsyntenin1 regulates Robo1 and Frizzled3 trafficking during midline crossing and  
529 longitudinal axon guidance, respectively [196]. By contrast, FEZ1 binds to Neuropilin1  
530 (Nrp1) and DCC and regulates the response of rat hippocampal axons to Sema3A and Netrin-  
531 1, respectively [197].  
532 Importantly, such studies highlight the variety of cargoes that can be loaded via different  
533 adaptor complexes onto distinct sets of molecular motors. Further studies will be required to  
534 increase our knowledge on the possible combinations and how their precise synchronisation  
535 in space and time is required to instruct accurate growth cone turning.

### 536 **II.3 Emerging key role of the tubulin code in axon navigation**

537 As mentioned above (§ I.), our vision of the MT network has considerably evolved across the  
538 past decades. It is now fully accepted that the MT cytoskeleton is a highly mosaic network  
539 composed of mixed populations of MTs that differ from one another based on their  
540 composition (*i.e.* the nature of the tubulin isoforms that compose the polymer) and the  
541 posttranslational modifications (PTMs) that decorate the tubulin dimers [48,49]. Both  
542 elements establish the tubulin code (*i.e.* by analogy to the histone code), which is starting to  
543 emerge as a highly selective regulator of MT functions in axon guidance.

#### 544 ***II.3.1 Tubulin isoforms as direct targets of guidance cues in growth cone steering***

545 Tubulin dimers are the building blocks of the eukaryotic MT cytoskeleton and are formed  
546 through chaperone-assisted folding and assembly of one  $\alpha$ - and one  $\beta$ -tubulin isoform. The  
547 tubulin gene family has gradually expanded across evolution, from 2 $\alpha$ - and 1 $\beta$ - tubulin genes  
548 in yeast to 9  $\alpha$ - and 9  $\beta$ -tubulin genes in humans [198]. Functional specification of tubulin  
549 isotypes is supported by (i) their different spatiotemporal expression patterns, (ii) their  
550 selective incorporation into specialised populations of MTs (*e.g.* axonemal, axonal...), (iii)  
551 their critical and non-overlapping roles in the development/homeostasis of particular  
552 tissues/cells, and above all (iv), the broad spectrum of human disorders caused by mutations  
553 in tubulin genes [*i.e.* Tubulinopathies; for review: 48,49,198,199]. Notably, mutations in  
554 genes encoding the neuronal-specific  $\beta$ -tubulin isoforms, TUBB2 and TUBB3, are  
555 responsible for brain malformations in humans associated with axon guidance defects  
556 including ocular nerve hypoplasia, dysgenesis of the corpus callosum, anterior commissure  
557 and corticospinal tracts [181,184,201,202; Figure 3]. Direct binding of TUBB3 to DCC or  
558 UNC5C receptors has been shown to couple Netrin-1 signalling to MT dynamics in axon  
559 guidance [28,29]. In this context, Netrin-1 stimulation promotes DCC or reduces UNC5C  
560 interaction and colocalisation with polymerised TUBB3 in primary neuron growth cones and  
561 locally influences MT dynamics (Figure 3A). Importantly, TUBB3 knockdown or pathogenic  
562 mutations impair these processes and thereby affect Netrin-evoked growth cone attraction or  
563 repulsion *in vitro* (Figure 1B & Figure 2H), as well as the pathfinding of spinal axon  
564 populations sensitive to Netrin-1/DCC or Netrin-1/UNC5C signalling *in vivo* [28,29,202,203;  
565 Figure 3B]. Interestingly, the TUBB2 E421K pathogenic mutation (*i.e.*; located in the C-ter  
566 tubulin tail), responsible for commissural axon pathfinding errors in mice and humans, affects  
567 the recruitment of the MT depolymerising motor Kip3p at MT plus ends and increases MT  
568 stability in yeast [184]. These effects strikingly contrast with other TUBB2 pathogenic  
569 variants, which drastically impair tubulin heterodimerisation and cause brain malformations

570 without guidance defects [205]. These data demonstrate the critical role of residues lying in  
571 the C-terminal tail of tubulin isoforms for their functional specification. Further investigations  
572 will be required to clarify how these mutations impact growth cone MT behaviours and  
573 responsiveness to guidance cues. An attractive hypothesis is that through their sequence  
574 specificity, tubulin isoforms generate specific signalling hubs on MTs selectively recruiting  
575 guidance receptors, downstream kinases and MT-regulatory proteins that will act in concert to  
576 locally tune MT dynamics and thereby axon behaviours.

577 Altogether, these studies emphasise how the genetic diversity of MT structural subunits  
578 influences MT functions in axon guidance. The functional specification of tubulin isoforms  
579 has been shown to arise at transcriptional, translational and posttranslational levels [200].  
580 Nevertheless, despite their striking degree of homology, tubulin isoforms exhibit a significant  
581 divergence in their C-terminal tail, which is subjected to extensive PTMs, suggesting a  
582 prominent role for PTMs in the functional specification of tubulins. This hypothesis is  
583 strengthened by the above-mentioned work of Jaglin et al. and the recent work of Zheng and  
584 colleagues, uncovering a novel  $\alpha$ -tubulin isotype in *C.elegans* which, unlike other  $\alpha$ -tubulin  
585 isoforms, destabilises MTs and lacks potential sites for polyamination and polyglutamylation  
586 [206].

### 587 ***II.3.2 Tubulin posttranslational modifications: a selective code that tunes critical MAPs in*** 588 ***axon guidance***

589 With the exception of acetylation, which occurs on the luminal Lysine 40 of  $\alpha$ -tubulin, PTMs  
590 consist in the covalent addition of amino acids with different physio-chemical properties (*e.g.*  
591 glutamate, glycine, tyrosine) on the outer surface of MTs, and more specifically on the acidic  
592 C-terminal tail of tubulins. Tubulin PTMs modulate MT properties and/or tune the activity of  
593 MT-interacting proteins at molecular and cellular scales [48,49]. However, the physiological  
594 role of these modifications remains largely unknown, especially in a context of neural circuit  
595 wiring where MTs are key driving forces for axon guidance. Since excellent reviews  
596 describing the tubulin PTM observed in neurons and their influence on neuronal MT functions  
597 have recently been published [45,46], we will here focus on the very few studies that directly  
598 involve tubulin PTMs in axon pathfinding.

599 Tubulin tyrosination was the first PTM to be involved in axon guidance. Following the  
600 incorporation of tubulin dimers in protofilaments, the C-terminal tyrosine residue of  $\alpha$ -  
601 tubulins is removed by the tubulin carboxypeptidases vasoinhibins (VASH1/2), assisted by its  
602 regulatory subunit, the small vasoinhibin binding protein [SVBP; 207,208]. This tyrosine  
603 residue is subsequently re-added on free tubulin dimers by the tubulin-tyrosine ligase [TTL;  
604 209,210]. This tubulin detyrosination/tyrosination cycle has been shown to be critical for axon  
605 specification and outgrowth *in vitro* [207,211,212], as well as for neuronal connectivity *in*  
606 *vivo*. Indeed, TTL knockout mice exhibit several axonal tract defects including a disrupted  
607 cortico-thalamic loop [211] and aberrant navigational behaviours of olivary fibres at the floor  
608 plate [30]. Moreover, SVBP knockout mice (*i.e.* equivalent of VASH1/2 knockout mice)  
609 show axonal tract agenesis associated with connectivity defects [213]. At the cellular and  
610 molecular levels, TTL-deficient neurons show abnormal growth and branching behaviours at  
611 guidance choice points or on specific substrates. They also exhibit enlarged growth cones  
612 with defective polarisation of F-actin protrusions and mislocalisation of several cytoskeleton  
613 regulators, including the CLIP-170 +TIP and the Myosin IIb motor [30,211]. Importantly,  
614 TTL loss of function does not directly impact MT dynamics [214]. Thus, tubulin tyrosination  
615 may act as a local signal within growth cones for the recruitment of key mediators of the  
616 MT/actin crosstalk underlying axon steering (Figure 3B). Investigating the role of the  $\alpha$ 4A-

617 tubulin isotype, a natural source of detyrosinated tubulin in neurons [214,215], during nervous  
618 system development should provide additional key insight into the contribution of the tubulin  
619 tyrosination/detyrosination cycle to axon pathfinding.

620  
621 An additional tubulin PTM that has recently been involved in neuronal circuit wiring is  
622 polyglutamylolation. Tubulin polyglutamylolation is the most abundant tubulin PTM in neurons.  
623 It consists in the enzymatic addition of glutamate side chains on tubulin C-terminal tail  
624 glutamate residues. This modification is catalysed by enzymes from the Tubulin Tyrosine  
625 Ligase Like family [TTLL; 217,218] and removed by enzymes from the Cytosolic  
626 Carboxypeptidase family [CCP; 219,220]. Each TTLL has an enzymatic specificity (*i.e.*  
627 generation of short or long glutamate side chains) and a substrate preference for  $\alpha$ - versus  $\beta$ -  
628 tubulin [221]. Tubulin polyglutamylolation has been shown to tune the binding affinity and/or  
629 activity of molecular motors [222–224], structural MAPs [225,226] and MT-severing  
630 enzymes [227–230] *in vitro*. It has therefore emerged as a potential regulator of axon  
631 guidance. A recent study from Ten Martin and colleagues provides the first *in vivo* evidence  
632 that tubulin glutamylases act in a highly selective manner to tune the activity of MT severers  
633 during neuronal circuit wiring [31]. TTLL6 and TTLL11 are two tubulin glutamylases  
634 enriched in the zebrafish developing spinal cord [231], which catalyse the addition of long  
635 glutamate side chains on the C-terminal tail of  $\alpha$ -tubulins [218]. Using loss of function and  
636 rescue experiments, Ten Martin et al. show that despite their strong similarities, both  
637 glutamylases have non-overlapping key roles in zebrafish motor axon pathfinding. Notably,  
638 the authors reveal that TTLL6-mediated polyglutamylolation specifically boosts p60-katanin  
639 activity in navigating axons to control their targeting [31]. However, TTLL6 and TTLL11-  
640 deficient larvae also exhibit motor neuron migratory and axon guidance defects that resemble  
641 those described in spastin mutants [23,31]. These data suggest a concerted action of TTLL6  
642 and TTLL11 in the fine tuning of spastin activity in motor circuit wiring (Figure 3B).

643 In conclusion, the emerging key roles of tubulin isoforms and PTMs in the establishment of  
644 neuronal connectivity emphasise the need to use multi-scale and system approaches in order  
645 to crack the tubulin code that regulates MT functions in developing axons. Notably,  
646 investigating the role of other PTMs present in neurons such as tubulin acetylation and  
647 phosphorylation, respectively shown to influence p60-katanin severing activity [232] or  
648 DCC/TUBB3 interaction [233], will be of particular interest.

649

### 650 **III. Microtubules in axon pruning**

651 After reaching their terminal targets, many developing axons, including motor and sensory  
652 axons, initially form exuberant collaterals and supernumerary synapses. While a subset of  
653 them is stabilised to form the mature terminal axonal arbour, misplaced branches and  
654 synapses are pruned. Pruning comes in diverse manifestations that differ in morphology and  
655 extent as well as in their initiation signal. While in some cases, axonal pruning is initiated by  
656 growth cone exposure to conventional guidance cues, it can also be induced by the lack of  
657 sufficient trophic support or activity-dependent competition mechanisms. In some species  
658 such as in insects undergoing metamorphosis, large scale axon remodelling is induced by  
659 intrinsic genetic programs [234,235]. In all cases, obvious changes of the MT cytoskeleton  
660 architecture have been reported as a common and early event in axon pruning. Such MT  
661 remodelling has been described in various systems: the axonal branch elimination in the  
662 *Drosophila* mushroom body during metamorphosis, the selection of the single axon that  
663 innervates each muscle fibre at the mammalian neuro-muscular junction or the neurotrophin-  
664 dependent pruning of sensory axons [17,236,237]. Importantly, abnormal refinement of  
665 exuberant connectivity is likely to explain subtle connectivity defects at the origin of

666 neurodevelopmental disorders including schizophrenia, autism spectrum disorders, attention  
667 deficit hyperactivity disorders. Consistently, causal mutations in a few MT-regulatory  
668 proteins have been reported in these disorders [238]. However, although postulated as key  
669 players in axon pruning for many years, the precise contribution of MTs and their regulatory  
670 proteins to this developmental process is only starting to be deciphered.

### 671 **III.1 MT remodelling as a driving force of axon pruning**

672 During the past decades, several lines of evidence have underpinned MT breakdown as an  
673 early event required for axon pruning. In 2002, Eaton and colleagues revealed MT loss as an  
674 early cellular process in the stereotyped pruning of *Drosophila* neuromuscular junction  
675 supernumerary synapses, preceding the removal of presynaptic markers [239]. Supporting this  
676 idea, Watts et al. subsequently showed that during *Drosophila* metamorphosis, branches  
677 subjected to pruning in the mushroom body lose their tubulin content before any other  
678 reported change in their protein content including actin, presynaptic proteins or adhesion  
679 molecules. These observations establish MT remodelling as one of the earliest molecular  
680 changes occurring in axonal branches to be pruned [236]. Consistently, motor axons retreating  
681 from mammalian neuromuscular junctions are largely devoid of MTs [240]. Further  
682 supporting MT disassembly as an early and evolutionary conserved feature of axon pruning,  
683 Brill and colleagues have recently revealed that MTs are preferentially fragmented in motor  
684 axon branches undergoing retraction. Indeed, the authors show that the density of EB3-  
685 positive MT growing plus ends is reduced in branches to be pruned. Notably, this loss of MT  
686 polymerisation is correlated with a reduction in the MT mass and occurs before organelle  
687 evacuation [17]. Importantly, they further demonstrate that in this system, epothilone B-  
688 induced stabilisation of MTs delays synapse removal, demonstrating that MT loss is required  
689 for efficient axonal branch pruning [17]. This need to disassemble the MT network to drive  
690 axonal branch elimination is supported by the work of Maor-Nof and colleagues [237]. Using  
691 an *in vitro* assay based on NGF deprivation-driven DRG axon elimination, they show that the  
692 lack of NGF induces a reduction in DRG axon MT mass. The latter is correlated with a  
693 reduction in posttranscriptional modifications characteristic of stable MTs (*i.e.* detyrosination,  
694 acetylation and glutamylation). Notably, the authors demonstrate that pharmacological  
695 stabilisation of MTs with paclitaxel prevents NGF deprivation-induced changes of MT  
696 dynamics and posttranslational modifications in DRG axons, as well as their subsequent  
697 degeneration. Overall, these pharmacological studies underpin MT breakdown as a common  
698 intrinsic driving force of sensory and motor axon pruning occurring downstream of diverse  
699 initiation signals, such as neurotrophic factor deprivation or activity-based competition,  
700 respectively. However, despite this evidence, the intracellular signalling events and MT-  
701 interacting effectors that selectively drive MT disassembly in response to these pruning  
702 initiation signals remain largely unknown.

### 703 **III.2 MT-regulatory proteins required for axon pruning**

704 Several MT-regulatory proteins involved in dendritic pruning have been identified [for  
705 review, see 241]. However, the driving mechanism of axon pruning might largely differ from  
706 the one underlying dendritic branch elimination. Indeed, some MT-interactors involved in this  
707 latter process fail to influence axon remodelling. A relevant example comes from the work of  
708 Lee et al. showing that during *Drosophila* metamorphosis, the MT severer katanin-like 1 is  
709 required to drive the pruning of *da* neuron dendrites, while it is dispensable for the  
710 elimination of  $\gamma$  neuron axonal branches in the mushroom body [242]. This report highlights  
711 the difficulty to extend the mechanisms involved in dendritic pruning to axonal branch-  
712 specific elimination processes without experimental investigations.



713 Studies identifying MT-regulatory proteins required for axon pruning are scarce. However, a  
714 few modulators of MT dynamics have recently emerged as key players in this process (Table  
715 1), including the MT-severing enzyme spastin. Spastin knock-out mice exhibit a delayed  
716 elimination of polyinnervated neuromuscular junctions, highlighting the involvement of  
717 spastin in motor axon branch pruning [17,243]. An elevated MT content is found in delayed  
718 retreating motor axons of spastin-deficient mice, confirming the key contribution of the  
719 spastin severing activity to the MT cytoskeleton breakage required for efficient axon pruning  
720 [17; Figure 4A]. Interestingly, the loss of MT mass (especially polyglutamylated and stable  
721 MTs) reduces local membrane tension [244] and is therefore predicted to promote  
722 endocytosis, another driving process in neurite pruning [245,246]. Based on its molecular and  
723 functional links with the ESCRT-III (Endosomal Sorting Complex Required for Transport III  
724 proteins) proteins Ist1 and CHMP1B in the endocytic pathway [247–249], it is most likely  
725 that spastin drives axon pruning by coordinating MT disassembly with membrane  
726 remodelling.

727 Spastin is also part of a set of MT-destabilising proteins screened by Maor-Nof et al. to  
728 identify the molecular mechanisms underlying MT breakage in NGF-deprivation-induced  
729 DRG axon pruning. The screened proteins include the four members of the stathmin family,  
730 the MT severers katanin and spastin, as well as the MT-depolymerising motor Kif2A. Among  
731 these candidates, only the knock-down of Kif2A protects DRG axons from NGF deprivation-  
732 induced degeneration *in vitro*. The involvement of Kif2A in axon pruning is supported by the  
733 hyperinnervation of the skin by sensory axons in Kif2A knock-out animals [236; Figure 4B].

734 While MT regulators involved in axon pruning mentioned so far exclusively comprise MT-  
735 destabilising proteins, two recent studies have uncovered a critical role for CRMP structural  
736 MAPs in a subset of axon refinement models [146,250]. This observation is consistent with  
737 their implication in neurodevelopmental disorders associated with subtle connectivity defects  
738 [251]. Extensive anatomical analysis of neuronal connectivity in CRMP2 knockout mice has  
739 led Ziak and colleagues to uncover a critical role for CRMP2 in Sema3F-induced pruning of  
740 the infrapyramidal bundle of hippocampal mossy fibres and cortico-collicular axons. These  
741 observations are consistent with the CRMP2 requirement for Sema3F-induced axon repulsion  
742 *in vitro* [146]. By contrast, although CRMP2 also mediates Sema3A-induced axon collapse *in*  
743 *vitro* [142,143], the Sema3A-dependent pruning of hippocampo-septal axons is not affected in  
744 CRMP2 knockout mice [146], suggesting a selective role for CRMP2 as a Sema3F effector in  
745 axon pruning *in vivo* (Figure 4C). In line with this observation, Nakanishi et al. have  
746 uncovered a non-overlapping role for CRMP2 and CRMP4 in the axon pruning of mossy  
747 fibres in the mouse hippocampus and reveal a selective epistatic interaction between the  
748 Sema3F receptor, Neuropilin-2 [252] and CRMP2 in this process [250]. The counter intuitive  
749 role of these MT stabilising proteins in MT breakdown-driven axon pruning suggests that  
750 CRMP proteins might not directly operate on MTs in this process. Rather, they may be acting  
751 as key signalling hubs promoting interactions between components of the Semaphorin or  
752 Neuropilin receptor complex and downstream effectors such as MT-destabilising enzymes  
753 (*e.g.* stathmins, spastin). Addressing this issue will be of particular interest in the field.

754 Overall, a diversity of MT-regulatory proteins are involved in axon pruning, with functional  
755 specificities that might rely on the neuronal subtype considered and/or on the nature of the  
756 initiating signal (*e.g.* activity-dependent mechanisms, trophic deprivation or axon guidance  
757 cues). Deciphering the signalling events underlying the selective regulation of MT-associated  
758 proteins downstream of specific axon pruning signals, as well as their contribution to the MT  
759 breakdown that drives axon remodelling, should provide essential information on the basic  
760 mechanisms governing neuronal connectivity refinement.

761 **Conclusion**

762 Our understanding of the microtubule function as a driving force of axon guidance and  
763 pruning has considerably expanded and evolved in recent years, as highlighted by the  
764 numerous studies reported here and the complementary review from Sanchez-Huertas and  
765 Herrera that was published during the final writing steps of the present article [253]. These  
766 emerging studies reveal not only the importance of the microtubule composition and  
767 posttranslational modifications in fine tuning MT functions in neuronal circuit wiring, but also  
768 of the functional versatility of the numerous MT interactors in such process. This extensive  
769 intracellular diversity contrasts with the relatively limited number of identified extracellular  
770 cues that wire the nervous system. It therefore highlights how decisive the tubulin code and  
771 the diversity/versatility of MT-interactors are in expanding the range of axon guidance  
772 behaviours and underlying neuronal connections that can be controlled by a limited repertoire  
773 of extracellular cues. The versatility of a given MT interactor may be dependent on the  
774 neuronal subtype in which it is expressed (which defines the pool of genes expressed), the  
775 diversity of the upstream intracellular signalling events by which it is recruited to the MTs, its  
776 binding partners, but also the subpopulations of MTs (dependent on the tubulin isoforms and  
777 MT modifying enzymes expressed in this neuronal cell type) with which it interacts. Adding  
778 to this complexity, different sets of MT interactors with cooperative or antagonistic functions  
779 may be acting concomitantly downstream of a single guidance molecule to regulate different  
780 aspects of MT dynamics/organisation required for axon steering or pruning. With this in  
781 mind, dissecting the MT regulatory networks that adapt the axon responsiveness to  
782 extracellular signals, rather than focusing on individual MT-associated effectors, will  
783 represent a major challenge in the field to gain a more comprehensive and realistic  
784 understanding of the basic intracellular mechanisms underlying neuronal circuit wiring and  
785 refinement. Tackling these issues will require the combined use of global transcriptomic and  
786 proteomic approaches, as well as multigenic functional analyses (such as the one recently  
787 published by Hahn and colleagues [129]). Similarly, moving towards model organisms  
788 enabling *in vivo* live imaging of MT dynamics and genetic/optogenetic manipulation of MT  
789 regulatory proteins in navigating growth cones or retracting axons will be crucial to dissect  
790 MT functions in a physiological context, where crosstalks between several chemical but also  
791 physical and mechanical extracellular signals may influence MT functions.

792 **Legends to figures:**

793 **Figure 1: MT-regulatory proteins that drive the growth cone turn in response to**  
794 **attractive/permissive guidance cues and their associated molecular mechanism(s).**

795 **(A)** Asymmetric invasion and stabilisation of MTs within growth cone filopodia facing an  
796 attractive/permissive cue drives the growth cone turn. **B**, **C** and **D** are higher magnifications  
797 of the boxed region in **A** and depict the molecular effect(s) of Netrin-1 (**B**), Sema3E (**C**) or  
798 Sema3D/3E (**D**) on the MT cytoskeleton through different MT-regulatory proteins. While the  
799 boxed region in **A** gives an example of the growth cone region where the depicted  
800 mechanisms occur, it is important to bear in mind that these mechanisms may also be  
801 functioning in other growth cone regions, such as the transition zone or the central domain.  
802 Molecular mechanisms are schematised according to the literature cited in Table 1. Full  
803 arrows indicate the direct effect of the attractive/permissive cue on a specific receptor and its  
804 downstream effectors. Dotted arrows represent an indirect effect of the cue on a given effector  
805 through a receptor that remains unidentified. MAP: microtubule-associated protein; MTs:  
806 microtubules.

807 **Figure 2: MT-regulatory proteins that drive the growth cone turn in response to**  
808 **repulsive/non permissive guidance cues and their associated molecular mechanism(s).**

809 **A.** Asymmetric destabilisation of MTs on the growth cone side facing a repulsive/ non  
810 permissive cue drives the growth cone turn. **B-H** are higher magnifications of the boxed  
811 region in **A** and depict the different effectors engaged downstream of Sema-3A (**B**), Slit/Slit2  
812 (**C**), EphrinA2/A(**D**), NO (**E**), Draxin (**F**), Wnt5A (**G**) and Netrin-1 (**H**) to drive the MT  
813 remodelling required for growth cone repulsion. While the boxed region in **A** gives an  
814 example of the growth cone region where the depicted mechanisms occur, it is important to  
815 bear in mind that these mechanisms may also be functioning in other growth cone regions,  
816 such as the transition zone or the central domain. Molecular mechanisms are schematised  
817 according to the literature cited in Table 1. While full arrows indicate the direct effect of the  
818 repulsive/non-permissive cue on a specific receptor and its downstream effectors, dotted  
819 arrows represent an indirect effect of the cue on a given effector through a receptor that  
820 remains unidentified. MAP: Microtubule-Associated Protein; MTs: Microtubules; NO: Nitric  
821 Oxide.

822 **Figure 3: Emerging key role of the tubulin code in axon guidance.** Diagram depicting our  
823 current knowledge of the tubulin code impact on axon guidance from a single molecule scale  
824 (left hand panels) to a physiological one (right hand panels). (**A**) Selective involvement of  
825 tubulin isotypes in axon guidance pathways.  $\beta$ -II (light pink sphere) and  $\beta$ -III tubulin (dark  
826 pink sphere) isotypes control the pathfinding of specific populations of mammalian axons  
827 through their selective interaction with MT-regulatory proteins or axon guidance  
828 receptors(respectively) and their subsequent modulation of MT dynamics. (**B**)  
829 Posttranslational modifications of the C-terminal tail of  $\alpha$ -tubulin (*e.g.* tubulin tyrosination  
830 and polyglutamylation) tune the activity of specific MT-regulatory proteins, thereby  
831 influencing MT-driven growth cone steering. (**A-B**) Molecular mechanisms are schematised  
832 according to the literature cited in Table 1. (Left hand panels) The C-terminal tyrosine residue  
833 of  $\alpha$ -tubulin is indicated in blue, while glutamic acids subjected to polyglutamylation are  
834 shown in grey. Polyglutamate side chains generated by TLL6 or TLL11 are shown in  
835 yellow and red, respectively. (Middle panels) Tubulin C-terminal tails that protrude from the  
836 MT lattice are represented in grey in the absence of key PTMs. Dotted arrows indicate MT  
837 polymerisation events. Purple arrow represents Myosin IIb displacement on F-actin (red  
838 filament). Boxes are higher magnifications of the adjacent left hand dotted panel.  
839 Polymerising and depolymerising microtubules are depicted as in Figures 1 and 2. CA:  
840 callosal axons; CSA: corticospinal axons, OMA: oculomotor axons; FB: forebrain; MB:  
841 midbrain; HB: hindbrain; SC: spinal cord; FP: floor plate; DRG: dorsal root ganglia; DREZ:  
842 dorsal root entry zone; PCN: precerebellar nuclei; HM: horizontal myoseptum; MyoIIb:  
843 myosin IIb; TTL: tubulin tyrosine ligase; TLL: tubulin tyrosine ligase like; Kinesin depol:  
844 Kinesin depolymerising motor; ABP: Actin binding protein.

845 **Figure 4: MT-regulatory proteins that contribute to the MT breakage required for axon**  
846 **pruning.** (**A**) Activity-dependent retraction of supernumerary axons in polyinnervated  
847 neuromuscular junctions relies on the spastin MT-severing activity. (**B**) NGF deprivation-  
848 induced pruning of DRG axons requires the MT depolymerising motor Kif2A (**C**) Sema3F-  
849 evoked elimination of exuberant axonal branches involves the structural MAP CRMP2. Since  
850 CRMP2 mostly stabilises MTs, Sema3F might detach CRMP2 from MTs, thereby favouring  
851 their depolymerisation. Whether phospho-CRMP2 acts as a signalling hub promoting the  
852 interaction between Sema3F receptor complex components and MT-destabilising enzymes  
853 remains to be addressed. **Table 1: Summary table of the MT regulators involved in axon**  
854 **guidance and pruning.**

855

856

857 **References:**

- 858 [1] A. Chédotal, L.J. Richards, Wiring the Brain: The Biology of Neuronal Guidance, Cold Spring Harb  
859 Perspect Biol. 2 (2010) a001917. <https://doi.org/10.1101/cshperspect.a001917>.
- 860 [2] A.L. Kolodkin, M. Tessier-Lavigne, Mechanisms and Molecules of Neuronal Wiring: A Primer, Cold  
861 Spring Harb Perspect Biol. 3 (2011) a001727. <https://doi.org/10.1101/cshperspect.a001727>.
- 862 [3] E.C. Engle, Human Genetic Disorders of Axon Guidance, Cold Spring Harb Perspect Biol. 2 (2010)  
863 a001784. <https://doi.org/10.1101/cshperspect.a001784>.
- 864 [4] J.K. Chilton, S. Guthrie, Axons get ahead: Insights into axon guidance and congenital cranial  
865 dysinnervation disorders, Dev Neurobiol. 77 (2017) 861–875.  
866 <https://doi.org/10.1002/dneu.22477>.
- 867 [5] D.S. Prieur, A. Rebsam, Retinal axon guidance at the midline: Chiasmatic misrouting and  
868 consequences, Dev Neurobiol. 77 (2017) 844–860. <https://doi.org/10.1002/dneu.22473>.
- 869 [6] Q. Welniarz, I. Dusart, E. Roze, The corticospinal tract: Evolution, development, and human  
870 disorders, Dev Neurobiol. 77 (2017) 810–829. <https://doi.org/10.1002/dneu.22455>.
- 871 [7] L.A. Lowery, D. Van Vactor, The trip of the tip: understanding the growth cone machinery, Nat.  
872 Rev. Mol. Cell Biol. 10 (2009) 332–343. <https://doi.org/10.1038/nrm2679>.
- 873 [8] S. Roig-Puiggros, R.J. Vigouroux, D. Beckman, N.I. Bocai, B. Chiou, J. Davimes, G. Gomez, S. Grassi,  
874 A. Hoque, T.K. Karikari, F. Kiffer, M. Lopez, G. Lunghi, P. Mazenganya, S. Meier, M. Olguín-  
875 Albuerne, M.M. Oliveira, J. Paraíso-Luna, J. Pradhan, A. Radiske, A.B. Ramos-Hryb, M.C. Ribeiro,  
876 R. Schellino, M.C. Selles, S. Singh, P. Theotokis, A. Chédotal, Construction and reconstruction of  
877 brain circuits: normal and pathological axon guidance, J Neurochem. 153 (2020) 10–32.  
878 <https://doi.org/10.1111/jnc.14900>.
- 879 [9] Z. Wen, L. Han, J.R. Bamberg, S. Shim, G. Ming, J.Q. Zheng, BMP gradients steer nerve growth  
880 cones by a balancing act of LIM kinase and Slingshot phosphatase on ADF/cofilin, The Journal of  
881 Cell Biology. 178 (2007) 107–119. <https://doi.org/10.1083/jcb.200703055>.
- 882 [10] N.P. Boyer, S.L. Gupton, Revisiting Netrin-1: One Who Guides (Axons), Front Cell Neurosci. 12  
883 (2018) 221. <https://doi.org/10.3389/fncel.2018.00221>.
- 884 [11] D. Bentley, T.P. O'Connor, Cytoskeletal events in growth cone steering, Curr Opin Neurobiol. 4  
885 (1994) 43–48. [https://doi.org/10.1016/0959-4388\(94\)90030-2](https://doi.org/10.1016/0959-4388(94)90030-2).
- 886 [12] C.-H. Lin, C.A. Thompson, P. Forscher, Cytoskeletal reorganization underlying growth cone  
887 motility, Current Opinion in Neurobiology. 4 (1994) 640–647. [https://doi.org/10.1016/0959-4388\(94\)90004-3](https://doi.org/10.1016/0959-4388(94)90004-3).
- 889 [13] A. Hall, G. Lalli, Rho and Ras GTPases in Axon Growth, Guidance, and Branching, Cold Spring Harb  
890 Perspect Biol. 2 (2010) a001818. <https://doi.org/10.1101/cshperspect.a001818>.
- 891 [14] T.M. Gomez, P.C. Letourneau, Actin dynamics in growth cone motility and navigation, J.  
892 Neurochem. 129 (2014) 221–234. <https://doi.org/10.1111/jnc.12506>.
- 893 [15] S. Niftullayev, N. Lamarche-Vane, Regulators of Rho GTPases in the Nervous System: Molecular  
894 Implication in Axon Guidance and Neurological Disorders, Int J Mol Sci. 20 (2019) E1497.  
895 <https://doi.org/10.3390/ijms20061497>.
- 896 [16] K.B. Buck, J.Q. Zheng, Growth cone turning induced by direct local modification of microtubule  
897 dynamics, J. Neurosci. 22 (2002) 9358–9367.
- 898 [17] M.S. Brill, T. Kleele, L. Ruschkies, M. Wang, N.A. Marahori, M.S. Reuter, T.J. Hausrat, E. Weigand,  
899 M. Fisher, A. Ahles, S. Engelhardt, D.L. Bishop, M. Kneussel, T. Misgeld, Branch-Specific  
900 Microtubule Destabilization Mediates Axon Branch Loss during Neuromuscular Synapse  
901 Elimination, Neuron. 92 (2016) 845–856. <https://doi.org/10.1016/j.neuron.2016.09.049>.

- 902 [18] M.E. Rothenberg, S.L. Rogers, R.D. Vale, L.Y. Jan, Y.-N. Jan, *Drosophila* pod-1 crosslinks both actin  
903 and microtubules and controls the targeting of axons, *Neuron*. 39 (2003) 779–791.  
904 [https://doi.org/10.1016/s0896-6273\(03\)00508-7](https://doi.org/10.1016/s0896-6273(03)00508-7).
- 905 [19] H. Lee, U. Engel, J. Rusch, S. Scherrer, K. Sheard, D. Van Vactor, The microtubule plus end  
906 tracking protein Orbit/MAST/CLASP acts downstream of the tyrosine kinase Abl in mediating  
907 axon guidance, *Neuron*. 42 (2004) 913–926. <https://doi.org/10.1016/j.neuron.2004.05.020>.
- 908 [20] J.A. Del Río, C. González-Billault, J.M. Ureña, E.M. Jiménez, M.J. Barallobre, M. Pascual, L.  
909 Pujadas, S. Simó, A. La Torre, F. Wandosell, J. Avila, E. Soriano, MAP1B is required for Netrin 1  
910 signaling in neuronal migration and axonal guidance, *Curr. Biol*. 14 (2004) 840–850.  
911 <https://doi.org/10.1016/j.cub.2004.04.046>.
- 912 [21] B. Erdogan, G.M. Cammarata, E.J. Lee, B.C. Pratt, A.F. Francl, E.L. Rutherford, L.A. Lowery, The  
913 microtubule plus-end-tracking protein TACC3 promotes persistent axon outgrowth and mediates  
914 responses to axon guidance signals during development, *Neural Dev*. 12 (2017) 3.  
915 <https://doi.org/10.1186/s13064-017-0080-7>.
- 916 [22] C. Fassier, A. Fréal, L. Gasmi, C. Delphin, D. Ten Martin, S. De Gois, M. Tambalo, C. Bosc, P. Mailly,  
917 C. Revenu, L. Peris, S. Bolte, S. Schneider-Maunoury, C. Houart, F. Nothias, J.-C. Larcher, A.  
918 Andrieux, J. Hazan, Motor axon navigation relies on Fidgetin-like 1-driven microtubule plus end  
919 dynamics, *J. Cell Biol.* (2018). <https://doi.org/10.1083/jcb.201604108>.
- 920 [23] N. Jardin, F. Giudicelli, D. Ten Martín, A. Vitrac, S. De Gois, R. Allison, C. Houart, E. Reid, J. Hazan,  
921 C. Fassier, BMP- and neuropilin 1-mediated motor axon navigation relies on spastin alternative  
922 translation, *Development*. 145 (2018). <https://doi.org/10.1242/dev.162701>.
- 923 [24] S. Biswas, K. Kalil, The Microtubule-Associated Protein Tau Mediates the Organization of  
924 Microtubules and Their Dynamic Exploration of Actin-Rich Lamellipodia and Filopodia of Cortical  
925 Growth Cones, *J Neurosci*. 38 (2018) 291–307. [https://doi.org/10.1523/JNEUROSCI.2281-](https://doi.org/10.1523/JNEUROSCI.2281-17.2017)  
926 [17.2017](https://doi.org/10.1523/JNEUROSCI.2281-17.2017).
- 927 [25] P.G. Slater, G.M. Cammarata, A.G. Samuelson, A. Magee, Y. Hu, L.A. Lowery, XMAP215 promotes  
928 microtubule-F-actin interactions to regulate growth cone microtubules during axon guidance in  
929 *Xenopus laevis*, *J Cell Sci*. 132 (2019) jcs224311. <https://doi.org/10.1242/jcs.224311>.
- 930 [26] C. Sánchez-Huertas, M. Bonhomme, A. Falco, C. Fagotto-Kaufmann, J. van Haren, F. Jeanneteau,  
931 N. Galjart, A. Debant, J. Boudeau, The +TIP Navigator-1 is an actin-microtubule crosslinker that  
932 regulates axonal growth cone motility, *J Cell Biol*. 219 (2020) e201905199.  
933 <https://doi.org/10.1083/jcb.201905199>.
- 934 [27] E. Arbeille, G.J. Bashaw, Brain Tumor promotes axon growth across the midline through  
935 interactions with the microtubule stabilizing protein Apc2, *PLoS Genet*. 14 (2018) e1007314.  
936 <https://doi.org/10.1371/journal.pgen.1007314>.
- 937 [28] C. Qu, T. Dwyer, Q. Shao, T. Yang, H. Huang, G. Liu, Direct binding of TUBB3 with DCC couples  
938 netrin-1 signaling to intracellular microtubule dynamics in axon outgrowth and guidance, *J Cell*  
939 *Sci*. 126 (2013) 3070–3081. <https://doi.org/10.1242/jcs.122184>.
- 940 [29] Q. Shao, T. Yang, H. Huang, F. Alarmanazi, G. Liu, Uncoupling of UNC5C with Polymerized TUBB3  
941 in Microtubules Mediates Netrin-1 Repulsion, *J. Neurosci*. 37 (2017) 5620–5633.  
942 <https://doi.org/10.1523/JNEUROSCI.2617-16.2017>.
- 943 [30] S. Marcos, J. Moreau, S. Backer, D. Job, A. Andrieux, E. Bloch-Gallego, Tubulin Tyrosination Is  
944 Required for the Proper Organization and Pathfinding of the Growth Cone, *PLOS ONE*. 4 (2009)  
945 e5405. <https://doi.org/10.1371/journal.pone.0005405>.
- 946 [31] D.T. Martin, N. Jardin, F. Giudicelli, L. Gasmi, J. Vouigny, C. Haumaitre, X. Nicol, C. Janke, C.  
947 Fassier, J. Hazan, A key role for p60-Katanin in axon navigation is conditioned by the tubulin  
948 polyglutamylase TLL6, (2022) 2022.01.20.477127.  
949 <https://www.biorxiv.org/content/10.1101/2022.01.20.477127v1> (accessed January 21, 2022).
- 950 [32] A. Akhmanova, M.O. Steinmetz, Tracking the ends: a dynamic protein network controls the fate  
951 of microtubule tips, *Nat. Rev. Mol. Cell Biol*. 9 (2008) 309–322.  
952 <https://doi.org/10.1038/nrm2369>.

- 953 [33] T. Mitchison, M. Kirschner, Dynamic instability of microtubule growth, *Nature*. 312 (1984) 237–  
954 242. <https://doi.org/10.1038/312237a0>.
- 955 [34] L.C. Kapitein, C.C. Hoogenraad, Building the Neuronal Microtubule Cytoskeleton, *Neuron*. 87  
956 (2015) 492–506. <https://doi.org/10.1016/j.neuron.2015.05.046>.
- 957 [35] M. Breuss, D.A. Keays, Microtubules and neurodevelopmental disease: the movers and the  
958 makers, *Adv Exp Med Biol*. 800 (2014) 75–96. [https://doi.org/10.1007/978-94-007-7687-6\\_5](https://doi.org/10.1007/978-94-007-7687-6_5).
- 959 [36] M. Lasser, J. Tiber, L.A. Lowery, The Role of the Microtubule Cytoskeleton in  
960 Neurodevelopmental Disorders, *Frontiers in Cellular Neuroscience*. 12 (2018) 165.  
961 <https://doi.org/10.3389/fncel.2018.00165>.
- 962 [37] Q. Chang, H. Yang, M. Wang, H. Wei, F. Hu, Role of Microtubule-Associated Protein in Autism  
963 Spectrum Disorder, *Neurosci Bull*. 34 (2018) 1119–1126. [https://doi.org/10.1007/s12264-018-  
0246-2](https://doi.org/10.1007/s12264-018-<br/>964 0246-2).
- 965 [38] R.P. Tas, A. Chazeau, B.M.C. Cloin, M.L.A. Lambers, C.C. Hoogenraad, L.C. Kapitein,  
966 Differentiation between Oppositely Oriented Microtubules Controls Polarized Neuronal  
967 Transport, *Neuron*. 96 (2017) 1264–1271.e5. <https://doi.org/10.1016/j.neuron.2017.11.018>.
- 968 [39] C. Leterrier, P. Dubey, S. Roy, The nano-architecture of the axonal cytoskeleton, *Nat Rev*  
969 *Neurosci*. 18 (2017) 713–726. <https://doi.org/10.1038/nrn.2017.129>.
- 970 [40] Y. Guo, D. Li, S. Zhang, Y. Yang, J.-J. Liu, X. Wang, C. Liu, D.E. Milkie, R.P. Moore, U.S. Tulu, D.P.  
971 Kiehart, J. Hu, J. Lippincott-Schwartz, E. Betzig, D. Li, Visualizing Intracellular Organelle and  
972 Cytoskeletal Interactions at Nanoscale Resolution on Millisecond Timescales, *Cell*. 175 (2018)  
973 1430–1442.e17. <https://doi.org/10.1016/j.cell.2018.09.057>.
- 974 [41] D. Jurriens, V. van Batenburg, E.A. Katrukha, L.C. Kapitein, Mapping the neuronal cytoskeleton  
975 using expansion microscopy, *Methods Cell Biol*. 161 (2021) 105–124.  
976 <https://doi.org/10.1016/bs.mcb.2020.04.018>.
- 977 [42] Quantitative mapping of dense microtubule arrays in mammalian neurons | *eLife*, (n.d.).  
978 <https://elifesciences.org/articles/67925> (accessed January 9, 2022).
- 979 [43] J. Atherton, M. Stouffer, F. Francis, C.A. Moores, Visualising the cytoskeletal machinery in  
980 neuronal growth cones using cryo-electron tomography, 2021.  
981 <https://doi.org/10.1101/2021.08.06.455451>.
- 982 [44] C. Leterrier, A Pictorial History of the Neuronal Cytoskeleton, *J. Neurosci*. 41 (2021) 11–27.  
983 <https://doi.org/10.1523/JNEUROSCI.2872-20.2020>.
- 984 [45] C. Janke, M. Kneussel, Tubulin post-translational modifications: encoding functions on the  
985 neuronal microtubule cytoskeleton, *Trends Neurosci*. 33 (2010) 362–372.  
986 <https://doi.org/10.1016/j.tins.2010.05.001>.
- 987 [46] M.-J. Moutin, C. Bosc, L. Peris, A. Andrieux, Tubulin post-translational modifications control  
988 neuronal development and functions, *Developmental Neurobiology*. 81 (2021) 253–272.  
989 <https://doi.org/10.1002/dneu.22774>.
- 990 [47] C. Sánchez-Huertas, F. Freixo, R. Viais, C. Lacasa, E. Soriano, J. Lüders, Non-centrosomal  
991 nucleation mediated by augmin organizes microtubules in post-mitotic neurons and controls  
992 axonal microtubule polarity, *Nat Commun*. 7 (2016) 12187.  
993 <https://doi.org/10.1038/ncomms12187>.
- 994 [48] A. Roll-Mecak, The Tubulin Code in Microtubule Dynamics and Information Encoding, *Dev Cell*.  
995 54 (2020) 7–20. <https://doi.org/10.1016/j.devcel.2020.06.008>.
- 996 [49] C. Janke, M.M. Magiera, The tubulin code and its role in controlling microtubule properties and  
997 functions, *Nat Rev Mol Cell Biol*. 21 (2020) 307–326. [https://doi.org/10.1038/s41580-020-0214-  
3](https://doi.org/10.1038/s41580-020-0214-<br/>998 3).
- 999 [50] R.P. Tas, L.C. Kapitein, Exploring cytoskeletal diversity in neurons, *Science*. 361 (2018) 231–232.  
1000 <https://doi.org/10.1126/science.aat5992>.
- 1001 [51] P. Forscher, S.J. Smith, Actions of cytochalasins on the organization of actin filaments and  
1002 microtubules in a neuronal growth cone, *J. Cell Biol*. 107 (1988) 1505–1516.

- 1003 [52] E.W. Dent, S.L. Gupton, F.B. Gertler, The growth cone cytoskeleton in axon outgrowth and  
1004 guidance, *Cold Spring Harb Perspect Biol.* 3 (2011).  
1005 <https://doi.org/10.1101/cshperspect.a001800>.
- 1006 [53] G.M. Cammarata, E.A. Bearce, L.A. Lowery, Cytoskeletal social networking in the growth cone:  
1007 How +TIPs mediate microtubule-actin cross-linking to drive axon outgrowth and guidance,  
1008 *Cytoskeleton (Hoboken)*. 73 (2016) 461–476. <https://doi.org/10.1002/cm.21272>.
- 1009 [54] E.A. Bearce, B. Erdogan, L.A. Lowery, TIPsy tour guides: how microtubule plus-end tracking  
1010 proteins (+TIPs) facilitate axon guidance, *Front Cell Neurosci.* 9 (2015).  
1011 <https://doi.org/10.3389/fncel.2015.00241>.
- 1012 [55] R. Pinto-Costa, M.M. Sousa, Microtubules, actin and cytolinkers: how to connect cytoskeletons in  
1013 the neuronal growth cone, *Neurosci Lett.* 747 (2021) 135693.  
1014 <https://doi.org/10.1016/j.neulet.2021.135693>.
- 1015 [56] J.H. Sabry, T.P. O'Connor, L. Evans, A. Toroian-Raymond, M. Kirschner, D. Bentley, Microtubule  
1016 behavior during guidance of pioneer neuron growth cones in situ, *J Cell Biol.* 115 (1991) 381–  
1017 395. <https://doi.org/10.1083/jcb.115.2.381>.
- 1018 [57] E. Tanaka, M.W. Kirschner, The role of microtubules in growth cone turning at substrate  
1019 boundaries, *J Cell Biol.* 128 (1995) 127–137. <https://doi.org/10.1083/jcb.128.1.127>.
- 1020 [58] C.H. Lin, P. Forscher, Cytoskeletal remodeling during growth cone-target interactions, *J Cell Biol.*  
1021 121 (1993) 1369–1383. <https://doi.org/10.1083/jcb.121.6.1369>.
- 1022 [59] C.-H. Lin, P. Forscher, Growth cone advance is inversely proportional to retrograde F-actin flow,  
1023 *Neuron.* 14 (1995) 763–771. [https://doi.org/10.1016/0896-6273\(95\)90220-1](https://doi.org/10.1016/0896-6273(95)90220-1).
- 1024 [60] T. Williamson, P.R. Gordon-Weeks, M. Schachner, J. Taylor, Microtubule reorganization is  
1025 obligatory for growth cone turning, *Proc Natl Acad Sci U S A.* 93 (1996) 15221–15226.  
1026 <https://doi.org/10.1073/pnas.93.26.15221>.
- 1027 [61] J.F. Chalacombe, D.M. Snow, P.C. Letourneau, Dynamic microtubule ends are required for  
1028 growth cone turning to avoid an inhibitory guidance cue, *J. Neurosci.* 17 (1997) 3085–3095.
- 1029 [62] F.-Q. Zhou, C.M. Waterman-Storer, C.S. Cohan, Focal loss of actin bundles causes microtubule  
1030 redistribution and growth cone turning, *J Cell Biol.* 157 (2002) 839–849.  
1031 <https://doi.org/10.1083/jcb.200112014>.
- 1032 [63] T.G.A. Mack, M.P. Koester, G.E. Pollerberg, The Microtubule-Associated Protein MAP1B Is  
1033 Involved in Local Stabilization of Turning Growth Cones, *Molecular and Cellular Neuroscience.* 15  
1034 (2000) 51–65. <https://doi.org/10.1006/mcne.1999.0802>.
- 1035 [64] M.P. Koester, O. Müller, G.E. Pollerberg, Adenomatous polyposis coli is differentially distributed  
1036 in growth cones and modulates their steering, *J. Neurosci.* 27 (2007) 12590–12600.  
1037 <https://doi.org/10.1523/JNEUROSCI.2250-07.2007>.
- 1038 [65] M. Higurashi, M. Iketani, K. Takei, N. Yamashita, R. Aoki, N. Kawahara, Y. Goshima, Localized role  
1039 of CRMP1 and CRMP2 in neurite outgrowth and growth cone steering, *Dev Neurobiol.* 72 (2012)  
1040 1528–1540. <https://doi.org/10.1002/dneu.22017>.
- 1041 [66] A.W. Schaefer, N. Kabir, P. Forscher, Filopodia and actin arcs guide the assembly and transport of  
1042 two populations of microtubules with unique dynamic parameters in neuronal growth cones, *J.*  
1043 *Cell Biol.* 158 (2002) 139–152. <https://doi.org/10.1083/jcb.200203038>.
- 1044 [67] S. Geraldo, U.K. Khanzada, M. Parsons, J.K. Chilton, P.R. Gordon-Weeks, Targeting of the F-actin-  
1045 binding protein drebrin by the microtubule plus-tip protein EB3 is required for neuritogenesis,  
1046 *Nat Cell Biol.* 10 (2008) 1181–1189. <https://doi.org/10.1038/ncb1778>.
- 1047 [68] D.T. Burnette, L. Ji, A.W. Schaefer, N.A. Medeiros, G. Danuser, P. Forscher, Myosin II Activity  
1048 Facilitates Microtubule Bundling in the Neuronal Growth Cone Neck, *Developmental Cell.* 15  
1049 (2008) 163–169. <https://doi.org/10.1016/j.devcel.2008.05.016>.
- 1050 [69] F.-Q. Zhou, J. Zhou, S. Dedhar, Y.-H. Wu, W.D. Snider, NGF-induced axon growth is mediated by  
1051 localized inactivation of GSK-3 $\beta$  and functions of the microtubule plus end binding protein  
1052 APC, *Neuron.* 42 (2004) 897–912. <https://doi.org/10.1016/j.neuron.2004.05.011>.

- 1053 [70] A.W. Schaefer, V.T.G. Schoonderwoert, L. Ji, N. Mederios, G. Danuser, P. Forscher, Coordination  
1054 of actin filament and microtubule dynamics during neurite outgrowth, *Dev. Cell.* 15 (2008) 146–  
1055 162. <https://doi.org/10.1016/j.devcel.2008.05.003>.
- 1056 [71] A.C. Lee, D.M. Suter, Quantitative Analysis of Microtubule Dynamics during Adhesion-Mediated  
1057 Growth Cone Guidance, *Dev Neurobiol.* 68 (2008) 1363–1377.  
1058 <https://doi.org/10.1002/dneu.20662>.
- 1059 [72] L. Li, T. Fothergill, B.I. Hutchins, E.W. Dent, K. Kalil, Wnt5a evokes cortical axon outgrowth and  
1060 repulsive guidance by tau mediated reorganization of dynamic microtubules, *Dev Neurobiol.* 74  
1061 (2014) 797–817. <https://doi.org/10.1002/dneu.22102>.
- 1062 [73] M. Atkins, L. Gasmi, V. Bercier, C. Revenu, F. Del Bene, J. Hazan, C. Fassier, FIGNL1 associates  
1063 with KIF1B $\beta$  and BICD1 to restrict dynein transport velocity during axon navigation, *J Cell Biol.*  
1064 218 (2019) 3290–3306. <https://doi.org/10.1083/jcb.201805128>.
- 1065 [74] B. Erdogan, E.A. Bearce, L.A. Lowery, Live Imaging of Cytoskeletal Dynamics in Embryonic  
1066 *Xenopus laevis* Growth Cones and Neural Crest Cells, *Cold Spring Harb Protoc.* 2021 (2021).  
1067 <https://doi.org/10.1101/pdb.prot104463>.
- 1068 [75] Y. Mimori-Kiyosue, I. Grigoriev, G. Lansbergen, H. Sasaki, C. Matsui, F. Severin, N. Galjart, F.  
1069 Grosveld, I. Vorobjev, S. Tsukita, A. Akhmanova, CLASP1 and CLASP2 bind to EB1 and regulate  
1070 microtubule plus-end dynamics at the cell cortex, *Journal of Cell Biology.* 168 (2005) 141–153.  
1071 <https://doi.org/10.1083/jcb.200405094>.
- 1072 [76] K. Drabek, M. van Ham, T. Stepanova, K. Draegestein, R. van Horssen, C.L. Sayas, A. Akhmanova,  
1073 T. ten Hagen, R. Smits, R. Fodde, F. Grosveld, N. Galjart, Role of CLASP2 in Microtubule  
1074 Stabilization and the Regulation of Persistent Motility, *Current Biology.* 16 (2006) 2259–2264.  
1075 <https://doi.org/10.1016/j.cub.2006.09.065>.
- 1076 [77] J. Al-Bassam, H. Kim, G. Brouhard, A. van Oijen, S.C. Harrison, F. Chang, CLASP promotes  
1077 microtubule rescue by recruiting tubulin dimers to the microtubule, *Dev Cell.* 19 (2010) 245–258.  
1078 <https://doi.org/10.1016/j.devcel.2010.07.016>.
- 1079 [78] A. Aher, M. Kok, A. Sharma, A. Rai, N. Olieric, R. Rodriguez-Garcia, E.A. Katrukha, T. Weinert, V.  
1080 Olieric, L.C. Kapitein, M.O. Steinmetz, M. Dogterom, A. Akhmanova, CLASP Suppresses  
1081 Microtubule Catastrophes through a Single TOG Domain, *Dev Cell.* 46 (2018) 40–58.e8.  
1082 <https://doi.org/10.1016/j.devcel.2018.05.032>.
- 1083 [79] E.-M. Hur, Saijilafu, B.D. Lee, S.-J. Kim, W.-L. Xu, F.-Q. Zhou, GSK3 controls axon growth via  
1084 CLASP-mediated regulation of growth cone microtubules, *Genes Dev.* 25 (2011) 1968–1981.  
1085 <https://doi.org/10.1101/gad.17015911>.
- 1086 [80] U. Engel, Y. Zhan, J.B. Long, S.N. Boyle, B.A. Ballif, K. Dorey, S.P. Gygi, A.J. Koleske, D. VanVactor,  
1087 Abelson Phosphorylation of CLASP2 Modulates its Association With Microtubules and Actin,  
1088 *Cytoskeleton (Hoboken).* 71 (2014) 195–209. <https://doi.org/10.1002/cm.21164>.
- 1089 [81] J. Groden, A. Thliveris, W. Samowitz, M. Carlson, L. Gelbert, H. Albertsen, G. Joslyn, J. Stevens, L.  
1090 Spirio, M. Robertson, Identification and characterization of the familial adenomatous polyposis  
1091 coli gene, *Cell.* 66 (1991) 589–600. [https://doi.org/10.1016/0092-8674\(81\)90021-0](https://doi.org/10.1016/0092-8674(81)90021-0).
- 1092 [82] Y. Wen, C.H. Eng, J. Schmoranzner, N. Cabrera-Poch, E.J.S. Morris, M. Chen, B.J. Wallar, A.S.  
1093 Alberts, G.G. Gundersen, EB1 and APC bind to mDia to stabilize microtubules downstream of Rho  
1094 and promote cell migration, *Nat Cell Biol.* 6 (2004) 820–830. <https://doi.org/10.1038/ncb1160>.
- 1095 [83] L. Serre, V. Stoppin-Mellet, I. Arnal, Adenomatous Polyposis Coli as a Scaffold for Microtubule  
1096 End-Binding Proteins, *J Mol Biol.* 431 (2019) 1993–2005.  
1097 <https://doi.org/10.1016/j.jmb.2019.03.028>.
- 1098 [84] J. Schneikert, J. Behrens, The canonical Wnt signalling pathway and its APC partner in colon  
1099 cancer development, *Gut.* 56 (2007) 417–425. <https://doi.org/10.1136/gut.2006.093310>.
- 1100 [85] S.A. Purro, L. Ciani, M. Hoyos-Flight, E. Stamatakou, E. Siomou, P.C. Salinas, Wnt Regulates Axon  
1101 Behavior through Changes in Microtubule Growth Directionality: A New Role for Adenomatous  
1102 Polyposis Coli, *J. Neurosci.* 28 (2008) 8644–8654. <https://doi.org/10.1523/JNEUROSCI.2320-08.2008>.
- 1103



- 1104 [86] H. Nakagawa, Y. Murata, K. Koyama, A. Fujiyama, Y. Miyoshi, M. Monden, T. Akiyama, Y.  
1105 Nakamura, Identification of a Brain-specific APC Homologue, APCL, and Its Interaction with  $\beta$ -  
1106 Catenin, *Cancer Res.* 58 (1998) 5176–5181.
- 1107 [87] T. Shintani, M. Ihara, S. Tani, J. Sakuraba, H. Sakuta, M. Noda, APC2 plays an essential role in  
1108 axonal projections through the regulation of microtubule stability, *J. Neurosci.* 29 (2009) 11628–  
1109 11640. <https://doi.org/10.1523/JNEUROSCI.2394-09.2009>.
- 1110 [88] T. Watanabe, S. Wang, J. Noritake, K. Sato, M. Fukata, M. Takefuji, M. Nakagawa, N. Izumi, T.  
1111 Akiyama, K. Kaibuchi, Interaction with IQGAP1 links APC to Rac1, Cdc42, and actin filaments  
1112 during cell polarization and migration, *Dev. Cell.* 7 (2004) 871–883.  
1113 <https://doi.org/10.1016/j.devcel.2004.10.017>.
- 1114 [89] T. Shintani, Y. Takeuchi, A. Fujikawa, M. Noda, Directional neuronal migration is impaired in mice  
1115 lacking adenomatous polyposis coli 2, *J. Neurosci.* 32 (2012) 6468–6484.  
1116 <https://doi.org/10.1523/JNEUROSCI.0590-12.2012>.
- 1117 [90] M.A. Juanes, D. Isnardon, A. Badache, S. Brasselet, M. Mavrikis, B.L. Goode, The role of APC-  
1118 mediated actin assembly in microtubule capture and focal adhesion turnover, *J. Cell Biol.* 218  
1119 (2019) 3415–3435. <https://doi.org/10.1083/jcb.201904165>.
- 1120 [91] M.A. Juanes, C.P. Fees, G.J. Hoeprich, R. Jaiswal, B.L. Goode, EB1 Directly Regulates APC-  
1121 Mediated Actin Nucleation, *Curr. Biol.* 30 (2020) 4763–4772.e8.  
1122 <https://doi.org/10.1016/j.cub.2020.08.094>.
- 1123 [92] L.A. Lowery, H. Lee, C. Lu, R. Murphy, R.A. Obar, B. Zhai, M. Schedl, D. Van Vactor, Y. Zhan,  
1124 Parallel Genetic and Proteomic Screens Identify Msps as a CLASP–Abl Pathway Interactor in  
1125 *Drosophila*, *Genetics.* 185 (2010) 1311–1325. <https://doi.org/10.1534/genetics.110.115626>.
- 1126 [93] L.A. Lowery, A. Stout, A.E. Faris, L. Ding, M.A. Baird, M.W. Davidson, G. Danuser, D. Van Vactor,  
1127 Growth cone-specific functions of XMAP215 in restricting microtubule dynamics and promoting  
1128 axonal outgrowth, *Neural Development.* 8 (2013) 22. <https://doi.org/10.1186/1749-8104-8-22>.
- 1129 [94] M.J. Lee, F. Gergely, K. Jeffers, S.Y. Peak-Chew, J.W. Raff, Msps/XMAP215 interacts with the  
1130 centrosomal protein D-TACC to regulate microtubule behaviour, *Nat. Cell Biol.* 3 (2001) 643–649.  
1131 <https://doi.org/10.1038/35083033>.
- 1132 [95] K. Kinoshita, T.L. Noetzel, L. Pelletier, K. Mechtler, D.N. Drechsel, A. Schwager, M. Lee, J.W. Raff,  
1133 A.A. Hyman, Aurora A phosphorylation of TACC3/maskin is required for centrosome-dependent  
1134 microtubule assembly in mitosis, *J. Cell Biol.* 170 (2005) 1047–1055.  
1135 <https://doi.org/10.1083/jcb.200503023>.
- 1136 [96] L.L. O’Brien, A.J. Albee, L. Liu, W. Tao, P. Dobrzyn, S.B. Lizarraga, C. Wiese, The *Xenopus* TACC  
1137 Homologue, Maskin, Functions in Mitotic Spindle Assembly, *Mol. Biol. Cell.* 16 (2005) 2836–2847.  
1138 <https://doi.org/10.1091/mbc.E04-10-0926>.
- 1139 [97] I. Peset, J. Seiler, T. Sardon, L.A. Bejarano, S. Rybina, I. Vernos, Function and regulation of  
1140 Maskin, a TACC family protein, in microtubule growth during mitosis, *Journal of Cell Biology.* 170  
1141 (2005) 1057–1066. <https://doi.org/10.1083/jcb.200504037>.
- 1142 [98] B.U. Nwagbara, A.E. Faris, E.A. Bearce, B. Erdogan, P.T. Ebbert, M.F. Evans, E.L. Rutherford, T.B.  
1143 Enzenbacher, L.A. Lowery, TACC3 is a microtubule plus end-tracking protein that promotes axon  
1144 elongation and also regulates microtubule plus end dynamics in multiple embryonic cell types,  
1145 *Mol. Biol. Cell.* 25 (2014) 3350–3362. <https://doi.org/10.1091/mbc.E14-06-1121>.
- 1146 [99] B. Erdogan, R.M. St. Clair, G.M. Cammarata, T. Zaccaro, B.A. Ballif, L.A. Lowery, Investigating the  
1147 impact of the phosphorylation status of tyrosine residues within the TACC domain of TACC3 on  
1148 microtubule behavior during axon growth and guidance, *Cytoskeleton (Hoboken).* 77 (2020)  
1149 277–291. <https://doi.org/10.1002/cm.21622>.
- 1150 [100] M.J. Martínez-López, S. Alcántara, C. Mascaró, F. Pérez-Brangulí, P. Ruiz-Lozano, T. Maes, E.  
1151 Soriano, C. Buesa, Mouse neuron navigator 1, a novel microtubule-associated protein involved in  
1152 neuronal migration, *Mol. Cell. Neurosci.* 28 (2005) 599–612.  
1153 <https://doi.org/10.1016/j.mcn.2004.09.016>.
- 1154 [101] J. van Haren, K. Draegestein, N. Keijzer, J.P. Abrahams, F. Grosveld, P.J. Peeters, D. Moechars,  
1155 N. Galjart, Mammalian Navigators are microtubule plus-end tracking proteins that can reorganize

1156 the cytoskeleton to induce neurite-like extensions, *Cell Motility*. 66 (2009) 824–838.  
1157 <https://doi.org/10.1002/cm.20370>.

1158 [102] S. Hekimi, D. Kershaw, Axonal guidance defects in a *Caenorhabditis elegans* mutant reveal  
1159 cell-extrinsic determinants of neuronal morphology, *J Neurosci*. 13 (1993) 4254–4271.

1160 [103] E.M. McNeill, K.P. Roos, D. Moechars, M. Clagett-Dame, Nav2 is necessary for cranial nerve  
1161 development and blood pressure regulation, *Neural Dev*. 5 (2010) 6.  
1162 <https://doi.org/10.1186/1749-8104-5-6>.

1163 [104] P.A. Kolodziej, L.Y. Jan, Y.N. Jan, Mutations that affect the length, fasciculation, or ventral  
1164 orientation of specific sensory axons in the *Drosophila* embryo, *Neuron*. 15 (1995) 273–286.  
1165 [https://doi.org/10.1016/0896-6273\(95\)90033-0](https://doi.org/10.1016/0896-6273(95)90033-0).

1166 [105] T. Lee, L. Luo, Mosaic Analysis with a Repressible Cell Marker for Studies of Gene Function in  
1167 Neuronal Morphogenesis, *Neuron*. 22 (1999) 451–461. [https://doi.org/10.1016/S0896-6273\(00\)80701-1](https://doi.org/10.1016/S0896-6273(00)80701-1).

1168  
1169 [106] S. Lee, M. Nahm, M. Lee, M. Kwon, E. Kim, A.D. Zadeh, H. Cao, H.-J. Kim, Z.H. Lee, S.B. Oh, J.  
1170 Yim, P.A. Kolodziej, S. Lee, The F-actin-microtubule crosslinker Shot is a platform for Krasavietz-  
1171 mediated translational regulation of midline axon repulsion, *Development*. 134 (2007) 1767–  
1172 1777. <https://doi.org/10.1242/dev.02842>.

1173 [107] W.B. Dobyns, K.A. Aldinger, G.E. Ishak, G.M. Mirzaa, A.E. Timms, M.E. Grout, M.H.G.  
1174 Dremmen, R. Schot, L. Vandervore, M.A. van Slegtenhorst, M. Wilke, E. Kasteleijn, A.S. Lee, B.J.  
1175 Barry, K.R. Chao, K. Szczałuba, J. Kobori, A. Hanson-Kahn, J.A. Bernstein, L. Carr, F. D’Arco, K.  
1176 Miyana, T. Okazaki, Y. Saito, M. Sasaki, S. Das, M.M. Wheeler, M.J. Bamshad, D.A. Nickerson,  
1177 University of Washington Center for Mendelian Genomics, Center for Mendelian Genomics at  
1178 the Broad Institute of MIT and Harvard, E.C. Engle, F.W. Verheijen, D. Doherty, G.M.S. Mancini,  
1179 MACF1 Mutations Encoding Highly Conserved Zinc-Binding Residues of the GAR Domain Cause  
1180 Defects in Neuronal Migration and Axon Guidance, *Am J Hum Genet*. 103 (2018) 1009–1021.  
1181 <https://doi.org/10.1016/j.ajhg.2018.10.019>.

1182 [108] S. Lee, P.A. Kolodziej, Short Stop provides an essential link between F-actin and microtubules  
1183 during axon extension, *Development*. 129 (2002) 1195–1204.

1184 [109] N. Sanchez-Soriano, M. Travis, F. Dajas-Bailador, C. Gonçalves-Pimentel, A.J. Whitmarsh, A.  
1185 Prokop, Mouse ACF7 and drosophila short stop modulate filopodia formation and microtubule  
1186 organisation during neuronal growth, *J Cell Sci*. 122 (2009) 2534–2542.  
1187 <https://doi.org/10.1242/jcs.046268>.

1188 [110] N. Sánchez-Soriano, C. Gonçalves-Pimentel, R. Beaven, U. Haessler, L. Ofner-Ziegenfuss, C.  
1189 Ballestrem, A. Prokop, *Drosophila* growth cones: a genetically tractable platform for the analysis  
1190 of axonal growth dynamics, *Dev Neurobiol*. 70 (2010) 58–71.  
1191 <https://doi.org/10.1002/dneu.20762>.

1192 [111] J. Alves-Silva, N. Sánchez-Soriano, R. Beaven, M. Klein, J. Parkin, T.H. Millard, H.J. Bellen, K.J.T.  
1193 Venken, C. Ballestrem, R.A. Kammerer, A. Prokop, Spectraplakins Promote Microtubule-  
1194 Mediated Axonal Growth by Functioning As Structural Microtubule-Associated Proteins and EB1-  
1195 Dependent +TIPs (Tip Interacting Proteins), *J. Neurosci*. 32 (2012) 9143–9158.  
1196 <https://doi.org/10.1523/JNEUROSCI.0416-12.2012>.

1197 [112] E. Tortosa, N. Galjart, J. Avila, C.L. Sayas, MAP1B regulates microtubule dynamics by  
1198 sequestering EB1/3 in the cytosol of developing neuronal cells, *The EMBO Journal*. 32 (2013)  
1199 1293–1306. <https://doi.org/10.1038/emboj.2013.76>.

1200 [113] S. Ramirez-Rios, E. Denarier, E. Prezel, A. Vinit, V. Stoppin-Mellet, F. Devred, P. Barbier, V.  
1201 Peyrot, C.L. Sayas, J. Avila, L. Peris, A. Andrieux, L. Serre, A. Fourest-Lieuvain, I. Arnal, Tau  
1202 antagonizes end-binding protein tracking at microtubule ends through a phosphorylation-  
1203 dependent mechanism, *Mol Biol Cell*. 27 (2016) 2924–2934. <https://doi.org/10.1091/mbc.E16-01-0029>.

1204  
1205 [114] C.L. Sayas, E. Tortosa, F. Bollati, S. Ramírez-Ríos, I. Arnal, J. Avila, Tau regulates the  
1206 localization and function of End-binding proteins 1 and 3 in developing neuronal cells, *J*  
1207 *Neurochem*. 133 (2015) 653–667. <https://doi.org/10.1111/jnc.13091>.

- 1208 [115] M.D. Weingarten, A.H. Lockwood, S.Y. Hwo, M.W. Kirschner, A protein factor essential for  
1209 microtubule assembly., *Proc Natl Acad Sci U S A.* 72 (1975) 1858–1862.
- 1210 [116] R.D. Sloboda, W.L. Dentler, J.L. Rosenbaum, Microtubule-associated proteins and the  
1211 stimulation of tubulin assembly in vitro, *Biochemistry.* 15 (1976) 4497–4505.  
1212 <https://doi.org/10.1021/bi00665a026>.
- 1213 [117] A. Elie, E. Prezel, C. Guérin, E. Denarier, S. Ramirez-Rios, L. Serre, A. Andrieux, A. Fourest-  
1214 Lieuvin, L. Blanchoin, I. Arnal, Tau co-organizes dynamic microtubule and actin networks, *Sci Rep.*  
1215 5 (2015) 9964. <https://doi.org/10.1038/srep09964>.
- 1216 [118] C. Cuveillier, B. Boulan, C. Ravello, E. Denarier, J.-C. Deloulme, S. Gory-Fauré, C. Delphin, C.  
1217 Bosc, I. Arnal, A. Andrieux, Beyond Neuronal Microtubule Stabilization: MAP6 and CRMPS, Two  
1218 Converging Stories, *Frontiers in Molecular Neuroscience.* 14 (2021) 82.  
1219 <https://doi.org/10.3389/fnmol.2021.665693>.
- 1220 [119] A. Meixner, S. Haverkamp, H. Wässle, S. Führer, J. Thalhammer, N. Kropf, R.E. Bittner, H.  
1221 Lassmann, G. Wiche, F. Propst, Map1b Is Required for Axon Guidance and Is Involved in the  
1222 Development of the Central and Peripheral Nervous System, *J Cell Biol.* 151 (2000) 1169–1178.
- 1223 [120] G.B. Walters, O. Gustafsson, G. Sveinbjornsson, V.K. Eiriksdottir, A.B. Agustsdottir, G.A.  
1224 Jonsdottir, S. Steinberg, A.F. Gunnarsson, M.I. Magnusson, U. Unnsteinsdottir, A.L. Lee, A.  
1225 Jonasdottir, A. Sigurdsson, A. Jonasdottir, A. Skuladottir, L. Jonsson, M.S. Nawaz, P. Sulem, M.  
1226 Frigge, A. Ingason, A. Love, G.L. Norddhal, M. Zervas, D.F. Gudbjartsson, M.O. Ulfarsson, E.  
1227 Saemundsen, H. Stefansson, K. Stefansson, MAP1B mutations cause intellectual disability and  
1228 extensive white matter deficit, *Nat Commun.* 9 (2018) 3456. <https://doi.org/10.1038/s41467-018-05595-6>.
- 1230 [121] C. Bouquet, S. Soares, Y. von Boxberg, M. Ravaille-Veron, F. Propst, F. Nothias, Microtubule-  
1231 associated protein 1B controls directionality of growth cone migration and axonal branching in  
1232 regeneration of adult dorsal root ganglia neurons, *J Neurosci.* 24 (2004) 7204–7213.  
1233 <https://doi.org/10.1523/JNEUROSCI.2254-04.2004>.
- 1234 [122] H. Stroissnigg, A. Trancíková, L. Descovich, J. Fuhrmann, W. Kutschera, J. Kostan, A. Meixner,  
1235 F. Nothias, F. Propst, S-nitrosylation of microtubule-associated protein 1B mediates nitric-oxide-  
1236 induced axon retraction, *Nat Cell Biol.* 9 (2007) 1035–1045. <https://doi.org/10.1038/ncb1625>.
- 1237 [123] R. Meli, P. Weisová, F. Propst, Repulsive Axon Guidance by Draxin Is Mediated by Protein  
1238 Kinase B (Akt), Glycogen Synthase Kinase-3 $\beta$  (GSK-3 $\beta$ ) and Microtubule-Associated Protein 1B,  
1239 *PLoS One.* 10 (2015) e0119524. <https://doi.org/10.1371/journal.pone.0119524>.
- 1240 [124] D. Villarroel-Campos, C. Gonzalez-Billault, The MAP1B case: an old MAP that is new again,  
1241 *Dev Neurobiol.* 74 (2014) 953–971. <https://doi.org/10.1002/dneu.22178>.
- 1242 [125] S.R. Tymanskyj, T.M.E. Scales, P.R. Gordon-Weeks, MAP1B enhances microtubule assembly  
1243 rates and axon extension rates in developing neurons, *Mol Cell Neurosci.* 49 (2012) 110–119.  
1244 <https://doi.org/10.1016/j.mcn.2011.10.003>.
- 1245 [126] C. Bouquet, M. Ravaille-Veron, F. Propst, F. Nothias, MAP1B coordinates microtubule and  
1246 actin filament remodeling in adult mouse Schwann cell tips and DRG neuron growth cones, *Mol*  
1247 *Cell Neurosci.* 36 (2007) 235–247. <https://doi.org/10.1016/j.mcn.2007.07.002>.
- 1248 [127] Y. Wang, E. Mandelkow, Tau in physiology and pathology, *Nat Rev Neurosci.* 17 (2016) 5–21.  
1249 <https://doi.org/10.1038/nrn.2015.1>.
- 1250 [128] E. Prezel, A. Elie, J. Delaroche, V. Stoppin-Mellet, C. Bosc, L. Serre, A. Fourest-Lieuvin, A.  
1251 Andrieux, M. Vantard, I. Arnal, Tau can switch microtubule network organizations: from random  
1252 networks to dynamic and stable bundles, *MBoC.* 29 (2018) 154–165.  
1253 <https://doi.org/10.1091/mbc.E17-06-0429>.
- 1254 [129] I. Hahn, A. Voelzmann, J. Parkin, J.B. Fülle, P.G. Slater, L.A. Lowery, N. Sanchez-Soriano, A.  
1255 Prokop, Tau, XMAP215/Msps and Eb1 co-operate interdependently to regulate microtubule  
1256 polymerisation and bundle formation in axons, *PLoS Genet.* 17 (2021) e1009647.  
1257 <https://doi.org/10.1371/journal.pgen.1009647>.

- 1258 [130] D. Job, E.H. Fischer, R.L. Margolis, Rapid disassembly of cold-stable microtubules by  
1259 calmodulin, *Proc Natl Acad Sci U S A.* 78 (1981) 4679–4682.  
1260 <https://doi.org/10.1073/pnas.78.8.4679>.
- 1261 [131] C. Bosc, R. Frank, E. Denarier, M. Ronjat, A. Schweitzer, J. Wehland, D. Job, Identification of  
1262 novel bifunctional calmodulin-binding and microtubule-stabilizing motifs in STOP proteins, *J Biol*  
1263 *Chem.* 276 (2001) 30904–30913. <https://doi.org/10.1074/jbc.M011614200>.
- 1264 [132] J.-C. Deloulme, S. Gory-Fauré, F. Mauconduit, S. Chauvet, J. Jonckheere, B. Boulan, E. Mire, J.  
1265 Xue, M. Jany, C. Maucler, A.A. Deparis, O. Montigon, A. Daoust, E.L. Barbier, C. Bosc, N. Deglon, J.  
1266 Brocard, E. Denarier, I. Le Brun, K. Pernet-Gallay, I. Vilgrain, P.J. Robinson, H. Lahrech, F. Mann,  
1267 A. Andrieux, Microtubule-associated protein 6 mediates neuronal connectivity through  
1268 Semaphorin 3E-dependent signalling for axonal growth, *Nat Commun.* 6 (2015) 7246.  
1269 <https://doi.org/10.1038/ncomms8246>.
- 1270 [133] U. Gimenez, B. Boulan, F. Mauconduit, F. Taurel, M. Leclercq, E. Denarier, J. Brocard, S. Gory-  
1271 Fauré, A. Andrieux, H. Lahrech, J.C. Deloulme, 3D imaging of the brain morphology and  
1272 connectivity defects in a model of psychiatric disorders: MAP6-KO mice, *Sci Rep.* 7 (2017) 10308.  
1273 <https://doi.org/10.1038/s41598-017-10544-2>.
- 1274 [134] B. Boulan, C. Ravello, A. Peyrel, C. Bosc, C. Delphin, F. Appaix, E. Denarier, A. Kraut, M.  
1275 Jacquier-Sarlin, A. Fournier, A. Andrieux, S. Gory-Fauré, J.-C. Deloulme, CRMP4-mediated fornix  
1276 development involves Semaphorin-3E signaling pathway, *ELife.* 10 (2021) e70361.  
1277 <https://doi.org/10.7554/eLife.70361>.
- 1278 [135] L. Peris, M. Bisbal, J. Martinez-Hernandez, Y. Saoudi, J. Jonckheere, M. Rolland, M. Sebastien,  
1279 J. Brocard, E. Denarier, C. Bosc, C. Guerin, S. Gory-Fauré, J.C. Deloulme, F. Lanté, I. Arnal, A.  
1280 Buisson, Y. Goldberg, L. Blanchoin, C. Delphin, A. Andrieux, A key function for microtubule-  
1281 associated-protein 6 in activity-dependent stabilisation of actin filaments in dendritic spines, *Nat*  
1282 *Commun.* 9 (2018) 3775. <https://doi.org/10.1038/s41467-018-05869-z>.
- 1283 [136] C. Cuveillier, J. Delaroche, M. Seggio, S. Gory-Fauré, C. Bosc, E. Denarier, M. Bacia, G.  
1284 Schoehn, H. Mohrbach, I. Kulić, A. Andrieux, I. Arnal, C. Delphin, MAP6 is an intraluminal protein  
1285 that induces neuronal microtubules to coil, *Sci Adv.* 6 (2020) eaaz4344.  
1286 <https://doi.org/10.1126/sciadv.aaz4344>.
- 1287 [137] Y. Fukata, T.J. Itoh, T. Kimura, C. Ménager, T. Nishimura, T. Shiromizu, H. Watanabe, N.  
1288 Inagaki, A. Iwamatsu, H. Hotani, K. Kaibuchi, CRMP-2 binds to tubulin heterodimers to promote  
1289 microtubule assembly, *Nat Cell Biol.* 4 (2002) 583–591. <https://doi.org/10.1038/ncb825>.
- 1290 [138] P.-C. Lin, P.M. Chan, C. Hall, E. Manser, Collapsin response mediator proteins (CRMPs) are a  
1291 new class of microtubule-associated protein (MAP) that selectively interacts with assembled  
1292 microtubules via a taxol-sensitive binding interaction, *J Biol Chem.* 286 (2011) 41466–41478.  
1293 <https://doi.org/10.1074/jbc.M111.283580>.
- 1294 [139] F. Nakamura, T. Ohshima, Y. Goshima, Collapsin Response Mediator Proteins: Their Biological  
1295 Functions and Pathophysiology in Neuronal Development and Regeneration, *Front Cell Neurosci.*  
1296 14 (2020) 188. <https://doi.org/10.3389/fncel.2020.00188>.
- 1297 [140] Y. Goshima, F. Nakamura, P. Strittmatter, S.M. Strittmatter, Collapsin-induced growth cone  
1298 collapse mediated by an intracellular protein related to UNC-33, *Nature.* 376 (1995) 509–514.  
1299 <https://doi.org/10.1038/376509a0>.
- 1300 [141] M. Brown, T. Jacobs, B. Eickholt, G. Ferrari, M. Teo, C. Monfries, R.Z. Qi, T. Leung, L. Lim, C.  
1301 Hall, Alpha2-chimaerin, cyclin-dependent Kinase 5/p35, and its target collapsin response  
1302 mediator protein-2 are essential components in semaphorin 3A-induced growth-cone collapse, *J*  
1303 *Neurosci.* 24 (2004) 8994–9004. <https://doi.org/10.1523/JNEUROSCI.3184-04.2004>.
- 1304 [142] Y. Uchida, T. Ohshima, Y. Sasaki, H. Suzuki, S. Yanai, N. Yamashita, F. Nakamura, K. Takei, Y.  
1305 Ihara, K. Mikoshiba, P. Kolattukudy, J. Honnorat, Y. Goshima, Semaphorin3A signalling is  
1306 mediated via sequential Cdk5 and GSK3beta phosphorylation of CRMP2: implication of common  
1307 phosphorylating mechanism underlying axon guidance and Alzheimer’s disease, *Genes Cells.* 10  
1308 (2005) 165–179. <https://doi.org/10.1111/j.1365-2443.2005.00827.x>.

- 1309 [143] T. Yoshimura, Y. Kawano, N. Arimura, S. Kawabata, A. Kikuchi, K. Kaibuchi, GSK-3beta  
1310 regulates phosphorylation of CRMP-2 and neuronal polarity, *Cell*. 120 (2005) 137–149.  
1311 <https://doi.org/10.1016/j.cell.2004.11.012>.
- 1312 [144] E.M. Hedgecock, J.G. Culotti, J.N. Thomson, L.A. Perkins, Axonal guidance mutants of  
1313 *Caenorhabditis elegans* identified by filling sensory neurons with fluorescein dyes, *Dev Biol*. 111  
1314 (1985) 158–170. [https://doi.org/10.1016/0012-1606\(85\)90443-9](https://doi.org/10.1016/0012-1606(85)90443-9).
- 1315 [145] W. Li, R.K. Herman, J.E. Shaw, Analysis of the *Caenorhabditis elegans* axonal guidance and  
1316 outgrowth gene *unc-33*, *Genetics*. 132 (1992) 675–689.  
1317 <https://doi.org/10.1093/genetics/132.3.675>.
- 1318 [146] J. Ziak, R. Weissova, K. Jeřábková, M. Janikova, R. Maimon, T. Petrasek, B. Pukajova, M.  
1319 Kleisnerova, M. Wang, M.S. Brill, P. Kasperek, X. Zhou, G. Alvarez-Bolado, R. Sedlacek, T. Misgeld,  
1320 A. Stuchlik, E. Perlson, M. Balastik, CRMP2 mediates Sema3F-dependent axon pruning and  
1321 dendritic spine remodeling, *EMBO Rep*. 21 (2020) e48512.  
1322 <https://doi.org/10.15252/embr.201948512>.
- 1323 [147] Z.-Z. Liu, J. Zhu, C.-L. Wang, X. Wang, Y.-Y. Han, L.-Y. Liu, H.A. Xu, CRMP2 and CRMP4 Are  
1324 Differentially Required for Axon Guidance and Growth in Zebrafish Retinal Neurons, *Neural*  
1325 *Plasticity*. 2018 (2018) e8791304. <https://doi.org/10.1155/2018/8791304>.
- 1326 [148] M.R. Khazaei, M.-P. Girouard, R. Alchini, S. Ong Tone, T. Shimada, S. Bechstedt, M. Cowan, D.  
1327 Guillet, P.W. Wiseman, G. Brouhard, J.F. Cloutier, A.E. Fournier, Collapsin response mediator  
1328 protein 4 regulates growth cone dynamics through the actin and microtubule cytoskeleton, *J Biol*  
1329 *Chem*. 289 (2014) 30133–30143. <https://doi.org/10.1074/jbc.M114.570440>.
- 1330 [149] L. Carretero-Rodriguez, R. Guðjónsdóttir, I. Poparic, M.L. Reilly, M. Chol, I.H. Bianco, M.  
1331 Chiapello, R. Feret, M.J. Deery, S. Guthrie, The Rac-GAP alpha2-Chimaerin Signals via CRMP2 and  
1332 Stathmins in the Development of the Ocular Motor System., *The Journal of Neuroscience : The*  
1333 *Official Journal of the Society for Neuroscience*. (2021). [https://doi.org/10.1523/jneurosci.0983-](https://doi.org/10.1523/jneurosci.0983-19.2021)  
1334 [19.2021](https://doi.org/10.1523/jneurosci.0983-19.2021).
- 1335 [150] Z.-S. Ji, J.-P. Li, C.-H. Fu, J.-X. Luo, H. Yang, G.-W. Zhang, W. Wu, H.-S. Lin, Spastin interacts  
1336 with collapsin response mediator protein 3 to regulate neurite growth and branching, *Neural*  
1337 *Regen Res*. 16 (2021) 2549–2556. <https://doi.org/10.4103/1673-5374.313052>.
- 1338 [151] S. Li, J. Zhang, J. Zhang, J. Li, L. Cheng, L. Chen, C. Cha, G. Guo, Spastin Interacts with CRMP2  
1339 to Regulate Neurite Outgrowth by Controlling Microtubule Dynamics through Phosphorylation  
1340 Modifications, *CNS Neurol Disord Drug Targets*. 20 (2021) 249–265.  
1341 <https://doi.org/10.2174/1871527319666201026165855>.
- 1342 [152] Z. Ji, G. Zhang, L. Chen, J. Li, Y. Yang, C. Cha, J. Zhang, H. Lin, G. Guo, Spastin Interacts with  
1343 CRMP5 to Promote Neurite Outgrowth by Controlling the Microtubule Dynamics, *Developmental*  
1344 *Neurobiology*. 78 (2018) 1191–1205. <https://doi.org/10.1002/dneu.22640>.
- 1345 [153] N. Bahi-Buisson, I. Souville, F.J. Fourniol, A. Toussaint, C.A. Moores, A. Houdusse, J.Y.  
1346 Lemaitre, K. Poirier, R. Khalaf-Nazzal, M. Hully, P.L. Leger, C. Elie, N. Boddaert, C. Beldjord, J.  
1347 Chelly, F. Francis, SBH-LIS European Consortium, New insights into genotype-phenotype  
1348 correlations for the doublecortin-related lissencephaly spectrum, *Brain*. 136 (2013) 223–244.  
1349 <https://doi.org/10.1093/brain/aws323>.
- 1350 [154] C.A. Moores, M. Perderiset, F. Francis, J. Chelly, A. Houdusse, R.A. Milligan, Mechanism of  
1351 Microtubule Stabilization by Doublecortin, *Molecular Cell*. 14 (2004) 833–839.  
1352 <https://doi.org/10.1016/j.molcel.2004.06.009>.
- 1353 [155] S. Bodakuntla, A.S. Jijumon, C. Villablanca, C. Gonzalez-Billault, C. Janke, Microtubule-  
1354 Associated Proteins: Structuring the Cytoskeleton, *Trends Cell Biol*. 29 (2019) 804–819.  
1355 <https://doi.org/10.1016/j.tcb.2019.07.004>.
- 1356 [156] L.A. Amos, D. Schlieper, Microtubules and maps, *Adv Protein Chem*. 71 (2005) 257–298.  
1357 [https://doi.org/10.1016/S0065-3233\(04\)71007-4](https://doi.org/10.1016/S0065-3233(04)71007-4).
- 1358 [157] C.J. Bott, L.P. McMahon, J.M. Keil, C.C. Yap, K.Y. Kwan, B. Winckler, Nestin Selectively  
1359 Facilitates the Phosphorylation of the Lissencephaly-Linked Protein Doublecortin (DCX) by

1360 cdk5/p35 to Regulate Growth Cone Morphology and Sema3a Sensitivity in Developing Neurons, *J*

1361 *Neurosci.* 40 (2020) 3720–3740. <https://doi.org/10.1523/JNEUROSCI.2471-19.2020>.

1362 [158] W. Yu, L. Qiang, J.M. Solowska, A. Karabay, S. Korulu, P.W. Baas, The microtubule-severing

1363 proteins spastin and katanin participate differently in the formation of axonal branches, *Mol.*

1364 *Biol. Cell.* 19 (2008) 1485–1498. <https://doi.org/10.1091/mbc.E07-09-0878>.

1365 [159] P.I. Hanson, S.W. Whiteheart, AAA+ proteins: have engine, will work, *Nat. Rev. Mol. Cell Biol.*

1366 6 (2005) 519–529. <https://doi.org/10.1038/nrm1684>.

1367 [160] A. Roll-Mecak, R.D. Vale, Structural basis of microtubule severing by the hereditary spastic

1368 paraplegia protein spastin, *Nature.* 451 (2008) 363–367. <https://doi.org/10.1038/nature06482>.

1369 [161] F.J. McNally, A. Roll-Mecak, Microtubule-severing enzymes: From cellular functions to

1370 molecular mechanism, *J Cell Biol.* 217 (2018) 4057–4069.

1371 <https://doi.org/10.1083/jcb.201612104>.

1372 [162] Y.-W. Kuo, J. Howard, Cutting, Amplifying, and Aligning Microtubules with Severing Enzymes,

1373 *Trends Cell Biol.* 31 (2021) 50–61. <https://doi.org/10.1016/j.tcb.2020.10.004>.

1374 [163] A. Karabay, W. Yu, J.M. Solowska, D.H. Baird, P.W. Baas, Axonal growth is sensitive to the

1375 levels of katanin, a protein that severs microtubules, *J. Neurosci.* 24 (2004) 5778–5788.

1376 <https://doi.org/10.1523/JNEUROSCI.1382-04.2004>.

1377 [164] J.D. Wood, J.A. Landers, M. Bingley, C.J. McDermott, V. Thomas-McArthur, L.J. Gleadall, P.J.

1378 Shaw, V.T. Cunliffe, The microtubule-severing protein Spastin is essential for axon outgrowth in

1379 the zebrafish embryo, *Hum. Mol. Genet.* 15 (2006) 2763–2771.

1380 <https://doi.org/10.1093/hmg/ddl212>.

1381 [165] R. Butler, J.D. Wood, J.A. Landers, V.T. Cunliffe, Genetic and chemical modulation of spastin-

1382 dependent axon outgrowth in zebrafish embryos indicates a role for impaired microtubule

1383 dynamics in hereditary spastic paraplegia, *Dis Model Mech.* 3 (2010) 743–751.

1384 <https://doi.org/10.1242/dmm.004002>.

1385 [166] L. Leo, W. Yu, M. D’Rozario, E.A. Waddell, D.R. Marena, M.A. Baird, M.W. Davidson, B. Zhou,

1386 B. Wu, L. Baker, D.J. Sharp, P.W. Baas, Vertebrate Fidgetin Restrains Axonal Growth by Severing

1387 Labile Domains of Microtubules, *Cell Rep.* 12 (2015) 1723–1730.

1388 <https://doi.org/10.1016/j.celrep.2015.08.017>.

1389 [167] K.J. Evans, E.R. Gomes, S.M. Reisenweber, G.G. Gundersen, B.P. Luring, Linking axonal

1390 degeneration to microtubule remodeling by Spastin-mediated microtubule severing, *J Cell Biol.*

1391 168 (2005) 599–606. <https://doi.org/10.1083/jcb.200409058>.

1392 [168] A. Errico, A. Ballabio, E.I. Rugarli, Spastin, the protein mutated in autosomal dominant

1393 hereditary spastic paraplegia, is involved in microtubule dynamics, *Hum Mol Genet.* 11 (2002)

1394 153–163. <https://doi.org/10.1093/hmg/11.2.153>.

1395 [169] L. Qiang, W. Yu, A. Andreadis, M. Luo, P.W. Baas, Tau Protects Microtubules in the Axon from

1396 Severing by Katanin, *J Neurosci.* 26 (2006) 3120–3129. [https://doi.org/10.1523/JNEUROSCI.5392-](https://doi.org/10.1523/JNEUROSCI.5392-05.2006)

1397 05.2006.

1398 [170] D. Zhang, G.C. Rogers, D.W. Buster, D.J. Sharp, Three microtubule severing enzymes

1399 contribute to the “Pacman-flux” machinery that moves chromosomes, *J. Cell Biol.* 177 (2007)

1400 231–242. <https://doi.org/10.1083/jcb.200612011>.

1401 [171] J. Feldner, T. Becker, K. Goishi, J. Schweitzer, P. Lee, M. Schachner, M. Klagsbrun, C.G. Becker,

1402 Neuropilin-1a is involved in trunk motor axon outgrowth in embryonic zebrafish, *Dev Dyn.* 234

1403 (2005) 535–549. <https://doi.org/10.1002/dvdy.20520>.

1404 [172] R. Morimura, K. Nozawa, H. Tanaka, T. Ohshima, Phosphorylation of Dpsyl2 (CRMP2) and

1405 Dpsyl3 (CRMP4) is required for positioning of caudal primary motor neurons in the zebrafish

1406 spinal cord, *Dev Neurobiol.* 73 (2013) 911–920. <https://doi.org/10.1002/dneu.22117>.

1407 [173] A. Roll-Mecak, R.D. Vale, Making more microtubules by severing: a common theme of

1408 noncentrosomal microtubule arrays?, *J Cell Biol.* 175 (2006) 849–851.

1409 <https://doi.org/10.1083/jcb.200611149>.

1410 [174] D. Zhang, K.D. Grode, S. Stewman, D. Diaz, E. Liebling, U. Rath, T. Riera, J. Currie, D.W. Buster,

1411 A.B. Asenjo, H.J. Sosa, J. Ross, A. Ma, S.L. Rogers, D.J. Sharp, Drosophila Katanin is a microtubule

1412 depolymerase that regulates cortical-microtubule plus-end interactions and cell migration, *Nat*  
1413 *Cell Biol.* 13 (2011) 361–370. <https://doi.org/10.1038/ncb2206>.

1414 [175] J.D. Díaz-Valencia, M.M. Morelli, M. Bailey, D. Zhang, D.J. Sharp, J.L. Ross, *Drosophila katanin-*  
1415 *60* depolymerizes and severs at microtubule defects, *Biophys J.* 100 (2011) 2440–2449.  
1416 <https://doi.org/10.1016/j.bpj.2011.03.062>.

1417 [176] A. Vemu, E. Szczesna, E.A. Zehr, J.O. Spector, N. Grigorieff, A.M. Deaconescu, A. Roll-Mecak,  
1418 Severing enzymes amplify microtubule arrays through lattice GTP-tubulin incorporation, *Science*.  
1419 361 (2018) eaau1504. <https://doi.org/10.1126/science.aau1504>.

1420 [177] L.D. Belmont, T.J. Mitchison, Identification of a protein that interacts with tubulin dimers and  
1421 increases the catastrophe rate of microtubules, *Cell.* 84 (1996) 623–631.  
1422 [https://doi.org/10.1016/s0092-8674\(00\)81037-5](https://doi.org/10.1016/s0092-8674(00)81037-5).

1423 [178] S. Chauvin, A. Sobel, Neuronal stathmins: a family of phosphoproteins cooperating for  
1424 neuronal development, plasticity and regeneration, *Prog. Neurobiol.* 126 (2015) 1–18.  
1425 <https://doi.org/10.1016/j.pneurobio.2014.09.002>.

1426 [179] C.M. Drerup, S. Lusk, A. Nechiporuk, Kif1B Interacts with KBP to Promote Axon Elongation by  
1427 Localizing a Microtubule Regulator to Growth Cones, *J. Neurosci.* 36 (2016) 7014–7026.  
1428 <https://doi.org/10.1523/JNEUROSCI.0054-16.2016>.

1429 [180] R. Phillis, D. Statton, P. Caruccio, R.K. Murphey, Mutations in the 8 kDa dynein light chain  
1430 gene disrupt sensory axon projections in the *Drosophila* imaginal CNS, *Development.* 122 (1996)  
1431 2955–2963.

1432 [181] M.A. Tischfield, H.N. Baris, C. Wu, G. Rudolph, L. Van Maldergem, W. He, W.-M. Chan, C.  
1433 Andrews, J.L. Demer, R.L. Robertson, D.A. Mackey, J.B. Ruddle, T.D. Bird, I. Gottlob, C. Pieh, E.I.  
1434 Traboulsi, S.L. Pomeroy, D.G. Hunter, J.S. Soul, A. Newlin, L.J. Sabol, E.J. Doherty, C.E. de  
1435 Uzcátegui, N. de Uzcátegui, M.L.Z. Collins, E.C. Sener, B. Wabbels, H. Hellebrand, T. Meitinger, T.  
1436 de Berardinis, A. Magli, C. Schiavi, M. Pastore-Trossello, F. Koc, A.M. Wong, A.V. Levin, M.T.  
1437 Geraghty, M. Descartes, M. Flaherty, R.V. Jamieson, H.U. Møller, I. Meuthen, D.F. Callen, J.  
1438 Kerwin, S. Lindsay, A. Meindl, M.L. Gupta, D. Pellman, E.C. Engle, Human TUBB3 mutations  
1439 perturb microtubule dynamics, kinesin interactions, and axon guidance, *Cell.* 140 (2010) 74–87.  
1440 <https://doi.org/10.1016/j.cell.2009.12.011>.

1441 [182] N.D. Dwyer, D.K. Manning, J.L. Moran, R. Mudbhary, M.S. Fleming, C.B. Favero, V.M. Vock,  
1442 D.D.M. O’Leary, C.A. Walsh, D.R. Beier, A forward genetic screen with a thalamocortical axon  
1443 reporter mouse yields novel neurodevelopment mutants and a distinct *emx2* mutant phenotype,  
1444 *Neural Dev.* 6 (2011) 3. <https://doi.org/10.1186/1749-8104-6-3>.

1445 [183] K.M. Janisch, V.M. Vock, M.S. Fleming, A. Shrestha, C.M. Grimsley-Myers, B.A. Rasoul, S.A.  
1446 Neale, T.D. Cupp, J.M. Kinchen, K.F. Liem, N.D. Dwyer, The vertebrate-specific Kinesin-6, Kif20b,  
1447 is required for normal cytokinesis of polarized cortical stem cells and cerebral cortex size,  
1448 *Development.* 140 (2013) 4672–4682. <https://doi.org/10.1242/dev.093286>.

1449 [184] G.Y. Cederquist, A. Luchniak, M.A. Tischfield, M. Peeva, Y. Song, M.P. Menezes, W.-M. Chan,  
1450 C. Andrews, S. Chew, R.V. Jamieson, L. Gomes, M. Flaherty, P.E. Grant, M.L. Gupta, E.C. Engle, An  
1451 inherited TUBB2B mutation alters a kinesin-binding site and causes polymicrogyria, CFEOM and  
1452 axon dysinnervation, *Hum Mol Genet.* 21 (2012) 5484–5499.  
1453 <https://doi.org/10.1093/hmg/dds393>.

1454 [185] W. Lu, V.I. Gelfand, Moonlighting Motors: Kinesin, Dynein, and Cell Polarity, *Trends Cell Biol.*  
1455 27 (2017) 505–514. <https://doi.org/10.1016/j.tcb.2017.02.005>.

1456 [186] K.A. Myers, I. Tint, C.V. Nadar, Y. He, M.M. Black, P.W. Baas, Antagonistic Forces Generated  
1457 by Cytoplasmic Dynein and Myosin-II during Growth Cone Turning and Axonal Retraction, *Traffic.*  
1458 7 (2006) 1333–1351. <https://doi.org/10.1111/j.1600-0854.2006.00476.x>.

1459 [187] V.C. Nadar, A. Ketschek, K.A. Myers, G. Gallo, P.W. Baas, Kinesin-5 Is Essential for Growth-  
1460 Cone Turning, *Current Biology.* 18 (2008) 1972–1977. <https://doi.org/10.1016/j.cub.2008.11.021>.

1461 [188] M. Liu, V.C. Nadar, F. Kozielski, M. Kozłowska, W. Yu, P.W. Baas, Kinesin-12, a mitotic  
1462 microtubule-associated motor protein, impacts axonal growth, navigation and branching, *J*  
1463 *Neurosci.* 30 (2010) 14896–14906. <https://doi.org/10.1523/JNEUROSCI.3739-10.2010>.

- 1464 [189] V.C. Nadar, S. Lin, P.W. Baas, Microtubule Redistribution in Growth Cones Elicited by Focal  
1465 Inactivation of Kinesin-5, *J. Neurosci.* 32 (2012) 5783–5794.  
1466 <https://doi.org/10.1523/JNEUROSCI.0144-12.2012>.
- 1467 [190] O.I. Kahn, P.W. Baas, Microtubules and Growth Cones: Motors Drive the Turn, *Trends in*  
1468 *Neurosciences.* 39 (2016) 433–440. <https://doi.org/10.1016/j.tins.2016.04.009>.
- 1469 [191] C.D. Deppmann, S. Mihalas, N. Sharma, B.E. Lonze, E. Niebur, D.D. Ginty, A model for  
1470 neuronal competition during development, *Science.* 320 (2008) 369–373.  
1471 <https://doi.org/10.1126/science.1152677>.
- 1472 [192] M. Ascaño, A. Richmond, P. Borden, R. Kuruvilla, Axonal targeting of Trk receptors via  
1473 transcytosis regulates sensitivity to neurotrophin responses, *J. Neurosci.* 29 (2009) 11674–11685.  
1474 <https://doi.org/10.1523/JNEUROSCI.1542-09.2009>.
- 1475 [193] N. Yamashita, M. Yamane, F. Suto, Y. Goshima, TrkA mediates retrograde semaphorin 3A  
1476 signaling through plexin A4 to regulate dendritic branching, *J. Cell. Sci.* 129 (2016) 1802–1814.  
1477 <https://doi.org/10.1242/jcs.184580>.
- 1478 [194] H.-L. Yu, Y. Peng, Y. Zhao, Y.-S. Lan, B. Wang, L. Zhao, D. Sun, J.-X. Pan, Z.-Q. Dong, L. Mei, Y.-  
1479 Q. Ding, X.-J. Zhu, W.-C. Xiong, Myosin X Interaction with KIF13B, a Crucial Pathway for Netrin-1-  
1480 Induced Axonal Development, *J. Neurosci.* 40 (2020) 9169–9185.  
1481 <https://doi.org/10.1523/JNEUROSCI.0929-20.2020>.
- 1482 [195] T.M.M. Saez, I.F. Bessone, M.S. Rodriguez, M. Alloatti, M.G. Otero, L.E. Cromberg, V.M.P.  
1483 Devoto, G. Oubiña, L. Sosa, M.G. Buffone, D.M. Gelman, T.L. Falzone, Kinesin-1-mediated axonal  
1484 transport of CB1 receptors is required for cannabinoid-dependent axonal growth and guidance,  
1485 *Development.* 147 (2020). <https://doi.org/10.1242/dev.184069>.
- 1486 [196] T.A. Alther, E. Domanitskaya, E.T. Stoeckli, Calsyntenin 1-mediated trafficking of axon  
1487 guidance receptors regulates the switch in axonal responsiveness at a choice point,  
1488 *Development.* 143 (2016) 994–1004. <https://doi.org/10.1242/dev.127449>.
- 1489 [197] J.Y. Chua, S.J. Ng, O. Yagensky, E.E. Wanker, J.J.E. Chua, FEZ1 Forms Complexes with CRMP1  
1490 and DCC to Regulate Axon and Dendrite Development, *ENeuro.* 8 (2021) ENEURO.0193-20.2021.  
1491 <https://doi.org/10.1523/ENEURO.0193-20.2021>.
- 1492 [198] P.G. McKean, S. Vaughan, K. Gull, The extended tubulin superfamily, *J Cell Sci.* 114 (2001)  
1493 2723–2733.
- 1494 [199] S. Chakraborti, K. Natarajan, J. Curiel, C. Janke, J. Liu, The emerging role of the tubulin code:  
1495 From the tubulin molecule to neuronal function and disease, *Cytoskeleton (Hoboken).* 73 (2016)  
1496 521–550. <https://doi.org/10.1002/cm.21290>.
- 1497 [200] M.W. Breuss, I. Leca, T. Gstrein, A.H. Hansen, D.A. Keays, Tubulins and brain development -  
1498 The origins of functional specification, *Mol Cell Neurosci.* 84 (2017) 58–67.  
1499 <https://doi.org/10.1016/j.mcn.2017.03.002>.
- 1500 [201] R. Romaniello, F. Arrigoni, A.E. Fry, M.T. Bassi, M.I. Rees, R. Borgatti, D.T. Pilz, T.D. Cushion,  
1501 Tubulin genes and malformations of cortical development, *Eur J Med Genet.* 61 (2018) 744–754.  
1502 <https://doi.org/10.1016/j.ejmg.2018.07.012>.
- 1503 [202] G. Fourel, C. Boscheron, Tubulin mutations in neurodevelopmental disorders as a tool to  
1504 decipher microtubule function, *FEBS Letters.* 594 (2020) 3409–3438.  
1505 <https://doi.org/10.1002/1873-3468.13958>.
- 1506 [203] Q. Shao, T. Yang, H. Huang, T. Majumder, B.A. Khot, M.M. Khouzani, F. Alarmanazi, Y.K. Gore,  
1507 G. Liu, Disease-associated mutations in human TUBB3 disturb netrin repulsive signaling, *PLoS*  
1508 *One.* 14 (2019) e0218811. <https://doi.org/10.1371/journal.pone.0218811>.
- 1509 [204] H. Huang, T. Yang, Q. Shao, T. Majumder, K. Mell, G. Liu, Human TUBB3 Mutations Disrupt  
1510 Netrin Attractive Signaling, *Neuroscience.* 374 (2018) 155–171.  
1511 <https://doi.org/10.1016/j.neuroscience.2018.01.046>.
- 1512 [205] X.H. Jaglin, K. Poirier, Y. Saillour, E. Buhler, G. Tian, N. Bahi-Buisson, C. Fallet-Bianco, F. Phan-  
1513 Dinh-Tuy, X.P. Kong, P. Bomont, L. Castelnau-Ptakhine, S. Odent, P. Loget, M. Kossorotoff, I.  
1514 Snoeck, G. Plessis, P. Parent, C. Beldjord, C. Cardoso, A. Represa, J. Flint, D.A. Keays, N.J. Cowan,



1515 J. Chelly, Mutations in the beta-tubulin gene TUBB2B result in asymmetrical polymicrogyria, *Nat*  
1516 *Genet.* 41 (2009) 746–752. <https://doi.org/10.1038/ng.380>.

1517 [206] C. Zheng, M. Diaz-Cuadros, K.C.Q. Nguyen, D.H. Hall, M. Chalfie, Distinct effects of tubulin  
1518 isotype mutations on neurite growth in *Caenorhabditis elegans*, *Mol Biol Cell.* 28 (2017) 2786–  
1519 2801. <https://doi.org/10.1091/mbc.E17-06-0424>.

1520 [207] C. Aillaud, C. Bosc, L. Peris, A. Bosson, P. Heemeryck, J. Van Dijk, J. Le Fric, B. Boulan, F.  
1521 Vossier, L.E. Sanman, S. Syed, N. Amara, Y. Couté, L. Lafanechère, E. Denarier, C. Delphin, L.  
1522 Pelletier, S. Humbert, M. Bogyo, A. Andrieux, K. Rogowski, M.-J. Moutin, Vasohibins/SVBP are  
1523 tubulin carboxypeptidases (TCPs) that regulate neuron differentiation, *Science.* 358 (2017) 1448–  
1524 1453. <https://doi.org/10.1126/science.aaa4165>.

1525 [208] J. Nieuwenhuis, A. Adamopoulos, O.B. Bleijerveld, A. Mazouzi, E. Stickel, P. Celie, M. Altelaar,  
1526 P. Knipscheer, A. Perrakis, V.A. Blomen, T.R. Brummelkamp, Vasohibins encode tubulin  
1527 de-tyrosinating activity, *Science.* 358 (2017) 1453–1456.  
1528 <https://doi.org/10.1126/science.aaa5676>.

1529 [209] K. Ersfeld, J. Wehland, U. Plessmann, H. Dodemont, V. Gerke, K. Weber, Characterization of  
1530 the tubulin-tyrosine ligase, *J Cell Biol.* 120 (1993) 725–732.  
1531 <https://doi.org/10.1083/jcb.120.3.725>.

1532 [210] H.C. Schröder, J. Wehland, K. Weber, Purification of brain tubulin-tyrosine ligase by  
1533 biochemical and immunological methods, *J Cell Biol.* 100 (1985) 276–281.  
1534 <https://doi.org/10.1083/jcb.100.1.276>.

1535 [211] C. Erck, L. Peris, A. Andrieux, C. Meissirel, A.D. Gruber, M. Vernet, A. Schweitzer, Y. Saoudi, H.  
1536 Pointu, C. Bosc, P.A. Salin, D. Job, J. Wehland, A vital role of tubulin-tyrosine-ligase for neuronal  
1537 organization, *Proc Natl Acad Sci U S A.* 102 (2005) 7853–7858.  
1538 <https://doi.org/10.1073/pnas.0409626102>.

1539 [212] N. Wang, C. Bosc, S. Ryul Choi, B. Boulan, L. Peris, N. Olieric, H. Bao, F. Krichen, L. Chen, A.  
1540 Andrieux, V. Olieric, M.-J. Moutin, M.O. Steinmetz, H. Huang, Structural basis of tubulin  
1541 de-tyrosination by the vasohibin–SVBP enzyme complex, *Nat Struct Mol Biol.* 26 (2019) 571–582.  
1542 <https://doi.org/10.1038/s41594-019-0241-y>.

1543 [213] A.T. Pagnamenta, P. Heemeryck, H.C. Martin, C. Bosc, L. Peris, I. Uszynski, S. Gory-Fauré, S.  
1544 Couly, C. Deshpande, A. Siddiqui, A.A. Elmonairy, WGS500 Consortium, Genomics England  
1545 Research Consortium, S. Jayawant, S. Murthy, I. Walker, L. Loong, P. Bauer, F. Vossier, E.  
1546 Denarier, T. Maurice, E.L. Barbier, J.-C. Deloulme, J.C. Taylor, E.M. Blair, A. Andrieux, M.-J.  
1547 Moutin, Defective tubulin de-tyrosination causes structural brain abnormalities with cognitive  
1548 deficiency in humans and mice, *Hum Mol Genet.* 28 (2019) 3391–3405.  
1549 <https://doi.org/10.1093/hmg/ddz186>.

1550 [214] L. Lafanechere, C. Courtay-Cahen, T. Kawakami, M. Jacrot, M. Rudiger, J. Wehland, D. Job,  
1551 R.L. Margolis, Suppression of tubulin tyrosine ligase during tumor growth, *Journal of Cell Science.*  
1552 111 (1998) 171–181. <https://doi.org/10.1242/jcs.111.2.171>.

1553 [215] V. Redeker, J. Rossier, A. Frankfurter, Posttranslational modifications of the C-terminus of  
1554 alpha-tubulin in adult rat brain: alpha 4 is glutamylated at two residues, *Biochemistry.* 37 (1998)  
1555 14838–14844. <https://doi.org/10.1021/bi981335k>.

1556 [216] B.N. Smith, N. Ticozzi, C. Fallini, A.S. Gkazi, S. Topp, K.P. Kenna, E.L. Scotter, J. Kost, P. Keagle,  
1557 J.W. Miller, D. Calini, C. Vance, E.W. Danielson, C. Troakes, C. Tiloca, S. Al-Sarraj, E.A. Lewis, A.  
1558 King, C. Colombrita, V. Pensato, B. Castellotti, J. de Bellerocche, F. Baas, A.L.M.A. ten Asbroek, P.C.  
1559 Sapp, D. McKenna-Yasek, R.L. McLaughlin, M. Polak, S. Asress, J. Esteban-Pérez, J.L. Muñoz-  
1560 Blanco, M. Simpson, SLAGEN Consortium, W. van Rheenen, F.P. Diekstra, G. Lauria, S. Duga, S.  
1561 Corti, C. Cereda, L. Corrado, G. Sorarù, K.E. Morrison, K.L. Williams, G.A. Nicholson, I.P. Blair, P.A.  
1562 Dion, C.S. Leblond, G.A. Rouleau, O. Hardiman, J.H. Veldink, L.H. van den Berg, A. Al-Chalabi, H.  
1563 Pall, P.J. Shaw, M.R. Turner, K. Talbot, F. Taroni, A. García-Redondo, Z. Wu, J.D. Glass, C. Gellera,  
1564 A. Ratti, R.H. Brown, V. Silani, C.E. Shaw, J.E. Landers, Exome-wide rare variant analysis identifies  
1565 TUBA4A mutations associated with familial ALS, *Neuron.* 84 (2014) 324–331.  
1566 <https://doi.org/10.1016/j.neuron.2014.09.027>.

- 1567 [217] C. Janke, K. Rogowski, D. Wloga, C. Regnard, A.V. Kajava, J.-M. Strub, N. Temurak, J. van Dijk,  
1568 D. Boucher, A. van Dorselaer, S. Suryavanshi, J. Gaertig, B. Eddé, Tubulin polyglutamylase  
1569 enzymes are members of the TTL domain protein family, *Science*. 308 (2005) 1758–1762.  
1570 <https://doi.org/10.1126/science.1113010>.
- 1571 [218] J. van Dijk, K. Rogowski, J. Miro, B. Lacroix, B. Eddé, C. Janke, A Targeted Multienzyme  
1572 Mechanism for Selective Microtubule Polyglutamylation, *Molecular Cell*. 26 (2007) 437–448.  
1573 <https://doi.org/10.1016/j.molcel.2007.04.012>.
- 1574 [219] K. Rogowski, J. van Dijk, M.M. Magiera, C. Bosc, J.-C. Deloulme, A. Bosson, L. Peris, N.D. Gold,  
1575 B. Lacroix, M. Bosch Grau, N. Bec, C. Larroque, S. Desagher, M. Holzer, A. Andrieux, M.-J. Moutin,  
1576 C. Janke, A family of protein-deglutamylating enzymes associated with neurodegeneration, *Cell*.  
1577 143 (2010) 564–578. <https://doi.org/10.1016/j.cell.2010.10.014>.
- 1578 [220] O. Tort, S. Tanco, C. Rocha, I. Bièche, C. Seixas, C. Bosc, A. Andrieux, M.-J. Moutin, F.X. Avilés,  
1579 J. Lorenzo, C. Janke, The cytosolic carboxypeptidases CCP2 and CCP3 catalyze posttranslational  
1580 removal of acidic amino acids, *Mol Biol Cell*. 25 (2014) 3017–3027.  
1581 <https://doi.org/10.1091/mbc.E14-06-1072>.
- 1582 [221] C. Janke, M.M. Magiera, The tubulin code and its role in controlling microtubule properties  
1583 and functions, *Nat Rev Mol Cell Biol*. 21 (2020) 307–326. [https://doi.org/10.1038/s41580-020-](https://doi.org/10.1038/s41580-020-0214-3)  
1584 [0214-3](https://doi.org/10.1038/s41580-020-0214-3).
- 1585 [222] M. Sirajuddin, L.M. Rice, R.D. Vale, Regulation of microtubule motors by tubulin isotypes and  
1586 post-translational modifications, *Nat Cell Biol*. 16 (2014) 335–344.  
1587 <https://doi.org/10.1038/ncb2920>.
- 1588 [223] S. Bodakuntla, X. Yuan, M. Genova, S. Gadadhar, S. Leboucher, M.-C. Birling, D. Klein, R.  
1589 Martini, C. Janke, M.M. Magiera, Distinct roles of  $\alpha$ - and  $\beta$ -tubulin polyglutamylation in  
1590 controlling axonal transport and in neurodegeneration, *EMBO J*. 40 (2021) e108498.  
1591 <https://doi.org/10.15252/embj.2021108498>.
- 1592 [224] D.V. Lessard, O.J. Zinder, T. Hotta, K.J. Verhey, R. Ohi, C.L. Berger, Polyglutamylation of  
1593 tubulin's C-terminal tail controls pausing and motility of kinesin-3 family member KIF1A, *J Biol*  
1594 *Chem*. 294 (2019) 6353–6363. <https://doi.org/10.1074/jbc.RA118.005765>.
- 1595 [225] D. Boucher, J.-C. Larcher, F. Gros, P. Denoulet, Polyglutamylation of Tubulin as a Progressive  
1596 Regulator of in Vitro Interactions between the Microtubule-Associated Protein Tau and Tubulin,  
1597 *Biochemistry*. 33 (1994) 12471–12477. <https://doi.org/10.1021/bi00207a014>.
- 1598 [226] C. Bonnet, D. Boucher, S. Lazereg, B. Pedrotti, K. Islam, P. Denoulet, J.C. Larcher, Differential  
1599 binding regulation of microtubule-associated proteins MAP1A, MAP1B, and MAP2 by tubulin  
1600 polyglutamylation, *J Biol Chem*. 276 (2001) 12839–12848.  
1601 <https://doi.org/10.1074/jbc.M011380200>.
- 1602 [227] B. Lacroix, J. van Dijk, N.D. Gold, J. Guizetti, G. Aldrian-Herrada, K. Rogowski, D.W. Gerlich, C.  
1603 Janke, Tubulin polyglutamylation stimulates spastin-mediated microtubule severing, *J Cell Biol*.  
1604 189 (2010) 945–954. <https://doi.org/10.1083/jcb.201001024>.
- 1605 [228] S.C. Shin, S.-K. Im, E.-H. Jang, K.S. Jin, E.-M. Hur, E.E. Kim, Structural and Molecular Basis for  
1606 Katanin-Mediated Severing of Glutamylated Microtubules, *Cell Reports*. 26 (2019) 1357-1367.e5.  
1607 <https://doi.org/10.1016/j.celrep.2019.01.020>.
- 1608 [229] E.A. Zehr, A. Szyk, E. Szczesna, A. Roll-Mecak, Katanin Grips the  $\beta$ -Tubulin Tail through an  
1609 Electropositive Double Spiral to Sever Microtubules, *Developmental Cell*. 52 (2020) 118-131.e6.  
1610 <https://doi.org/10.1016/j.devcel.2019.10.010>.
- 1611 [230] M.L. Valenstein, A. Roll-Mecak, Graded Control of Microtubule Severing by Tubulin  
1612 Glutamylation, *Cell*. 164 (2016) 911–921. <https://doi.org/10.1016/j.cell.2016.01.019>.
- 1613 [231] N. Pathak, C.A. Austin, I.A. Drummond, Tubulin Tyrosine Ligase-like Genes *tll3* and *tll6*  
1614 Maintain Zebrafish Cilia Structure and Motility, *J Biol Chem*. 286 (2011) 11685–11695.  
1615 <https://doi.org/10.1074/jbc.M110.209817>.
- 1616 [232] H. Sudo, P.W. Baas, Acetylation of microtubules influences their sensitivity to severing by  
1617 katanin in neurons and fibroblasts, *J Neurosci*. 30 (2010) 7215–7226.  
1618 <https://doi.org/10.1523/JNEUROSCI.0048-10.2010>.

- 1619 [233] C. Qu, T. Dwyer, Q. Shao, T. Yang, H. Huang, G. Liu, Direct binding of TUBB3 with DCC couples  
1620 netrin-1 signaling to intracellular microtubule dynamics in axon outgrowth and guidance, *J. Cell.*  
1621 *Sci.* 126 (2013) 3070–3081. <https://doi.org/10.1242/jcs.122184>.
- 1622 [234] O. Schuldiner, A. Yaron, Mechanisms of developmental neurite pruning, *Cell Mol Life Sci.* 72  
1623 (2015) 101–119. <https://doi.org/10.1007/s00018-014-1729-6>.
- 1624 [235] C. Fassier, X. Nicol, Retinal Axon Interplay for Binocular Mapping, *Frontiers in Neural Circuits.*  
1625 15 (2021) 49. <https://doi.org/10.3389/fncir.2021.679440>.
- 1626 [236] R.J. Watts, E.D. Hoopfer, L. Luo, Axon pruning during *Drosophila* metamorphosis: evidence  
1627 for local degeneration and requirement of the ubiquitin-proteasome system, *Neuron.* 38 (2003)  
1628 871–885. [https://doi.org/10.1016/s0896-6273\(03\)00295-2](https://doi.org/10.1016/s0896-6273(03)00295-2).
- 1629 [237] M. Maor-Nof, N. Homma, C. Raanan, A. Nof, N. Hirokawa, A. Yaron, Axonal pruning is actively  
1630 regulated by the microtubule-destabilizing protein kinesin superfamily protein 2A, *Cell Rep.* 3  
1631 (2013) 971–977. <https://doi.org/10.1016/j.celrep.2013.03.005>.
- 1632 [238] F. Marchisella, E.T. Coffey, P. Hollos, Microtubule and microtubule associated protein  
1633 anomalies in psychiatric disease, *Cytoskeleton.* 73 (2016) 596–611.  
1634 <https://doi.org/10.1002/cm.21300>.
- 1635 [239] B.A. Eaton, R.D. Fetter, G.W. Davis, Dynactin is necessary for synapse stabilization, *Neuron.*  
1636 34 (2002) 729–741. [https://doi.org/10.1016/s0896-6273\(02\)00721-3](https://doi.org/10.1016/s0896-6273(02)00721-3).
- 1637 [240] D.L. Bishop, T. Misgeld, M.K. Walsh, W.-B. Gan, J.W. Lichtman, Axon branch removal at  
1638 developing synapses by axosome shedding, *Neuron.* 44 (2004) 651–661.  
1639 <https://doi.org/10.1016/j.neuron.2004.10.026>.
- 1640 [241] S. Rumpf, N. Wolterhoff, S. Herzmann, Functions of Microtubule Disassembly during Neurite  
1641 Pruning, *Trends Cell Biol.* 29 (2019) 291–297. <https://doi.org/10.1016/j.tcb.2019.01.002>.
- 1642 [242] H.-H. Lee, L.Y. Jan, Y.-N. Jan, *Drosophila* IKK-related kinase Ik2 and Katanin p60-like 1 regulate  
1643 dendrite pruning of sensory neuron during metamorphosis, *Proc Natl Acad Sci U S A.* 106 (2009)  
1644 6363–6368. <https://doi.org/10.1073/pnas.0902051106>.
- 1645 [243] M. Wang, T. Kleele, Y. Xiao, G. Plucinska, P. Avramopoulos, S. Engelhardt, M.H. Schwab, M.  
1646 Kneussel, T. Czopka, D.L. Sherman, P.J. Brophy, T. Misgeld, M.S. Brill, Completion of neuronal  
1647 remodeling prompts myelination along developing motor axon branches, *Journal of Cell Biology.*  
1648 220 (2021) e201911114. <https://doi.org/10.1083/jcb.201911114>.
- 1649 [244] S. Herzmann, R. Krumkamp, S. Rode, C. Kintrup, S. Rumpf, PAR-1 promotes microtubule  
1650 breakdown during dendrite pruning in *Drosophila*, *EMBO J.* 36 (2017) 1981–1991.  
1651 <https://doi.org/10.15252/embj.201695890>.
- 1652 [245] X.-S. Wu, S. Elias, H. Liu, J. Heureaux, P.J. Wen, A.P. Liu, M.M. Kozlov, L.-G. Wu, Membrane  
1653 Tension Inhibits Rapid and Slow Endocytosis in Secretory Cells, *Biophysical Journal.* 113 (2017)  
1654 2406–2414. <https://doi.org/10.1016/j.bpj.2017.09.035>.
- 1655 [246] T. Kanamori, J. Yoshino, K. Yasunaga, Y. Dairyo, K. Emoto, Local endocytosis triggers dendritic  
1656 thinning and pruning in *Drosophila* sensory neurons, *Nat Commun.* 6 (2015) 6515.  
1657 <https://doi.org/10.1038/ncomms7515>.
- 1658 [247] R. Allison, J.H. Lumb, C. Fassier, J.W. Connell, D. Ten Martin, M.N.J. Seaman, J. Hazan, E. Reid,  
1659 An ESCRT-spastin interaction promotes fission of recycling tubules from the endosome, *J Cell*  
1660 *Biol.* 202 (2013) 527–543. <https://doi.org/10.1083/jcb.201211045>.
- 1661 [248] R. Allison, J.R. Edgar, G. Pearson, T. Rizo, T. Newton, S. Günther, F. Berner, J. Hague, J.W.  
1662 Connell, J. Winkler, J. Lippincott-Schwartz, C. Beetz, B. Winner, E. Reid, Defects in ER-endosome  
1663 contacts impact lysosome function in hereditary spastic paraplegia, *J Cell Biol.* 216 (2017) 1337–  
1664 1355. <https://doi.org/10.1083/jcb.201609033>.
- 1665 [249] J.W. Connell, R.J. Allison, C.E. Rodger, G. Pearson, E. Zlamalova, E. Reid, ESCRT-III-associated  
1666 proteins and spastin inhibit protrudin-dependent polarised membrane traffic, *Cell. Mol. Life Sci.*  
1667 77 (2020) 2641–2658. <https://doi.org/10.1007/s00018-019-03313-z>.
- 1668 [250] Y. Nakanishi, S. Akinaga, K. Osawa, N. Suzuki, A. Sugeno, P. Kolattukudy, Y. Goshima, T.  
1669 Ohshima, Regulation of axon pruning of mossy fiber projection in hippocampus by CRMP2 and  
1670 CRMP4, *Dev Neurobiol.* (2021). <https://doi.org/10.1002/dneu.22865>.

- 1671 [251] T.T. Quach, J. Honnorat, P.E. Kolattukudy, R. Khanna, A.M. Duchemin, CRMPs: critical  
1672 molecules for neurite morphogenesis and neuropsychiatric diseases, *Mol Psychiatry*. 20 (2015)  
1673 1037–1045. <https://doi.org/10.1038/mp.2015.77>.
- 1674 [252] M.M. Riccomagno, A. Hurtado, H. Wang, J.G.J. Macopson, E.M. Griner, A. Betz, N. Brose, M.G.  
1675 Kazanietz, A.L. Kolodkin, The RacGAP  $\beta$ -Chimaerin Selectively Mediates Stereotyped  
1676 Hippocampal Axonal Pruning, *Cell*. 149 (2012) 1594–1606.  
1677 <https://doi.org/10.1016/j.cell.2012.05.018>.
- 1678 [253] C. Sánchez-Huertas, E. Herrera, With the Permission of Microtubules: An Updated Overview  
1679 on Microtubule Function During Axon Pathfinding, *Frontiers in Molecular Neuroscience*. 14  
1680 (2021). <https://www.frontiersin.org/article/10.3389/fnmol.2021.759404> (accessed January 26,  
1681 2022).  
1682

### 1683 **Conflict of Interest**

1684 The authors declare no conflict of interest.

### 1685 **Acknowledgement**

1686 We greatly thank Isabelle Arnal, Annie Andrieux, Jamile Hazan, Christine Métin, and the  
1687 Nicol lab members for fruitful discussions and suggestions.

### 1688 **Author contribution**





1689 MA, XN and CF wrote the manuscript and designed the figures. All authors approved the  
1690 submitted version.

### 1691 **Funding**









1692 The research of the Nicol lab was supported by grants from the Agence Nationale de la  
1693 Recherche to XN (ANR-18-CE16-0017) and CF (ANR-20-CE16-0019), funding from the  
1694 AFM (application n°23695), ASL-HSP France and Tom Wahlig foundations to CF and from  
1695 the FRM (Equipe FRM EQU202003010158) and Unadev/ITMO NNP to XN. MA is a  
1696 recipient of postdoctoral fellowships from the Brain and Behaviour Research Foundation and  
1697 the Jérôme Lejeune and Sisley-d’Ornano Foundations in the Metin lab.

1698 This work was performed in the framework of the LABEX LIFESENSES (ANR-10-LABX-  
1699 65) and IHU FOrReSIGHT (ANR-18-IAHU-0001) supported by French state funds managed  
1700 by the Agence Nationale de la Recherche within the Investissements d’Avenir program.

Protein	Models/Systems	Links with guidance cues	Functions in axon guidance	References	Figure
<b>Plus-end tracking proteins</b>					
CLASP	<i>Drosophila</i> nervous system	Slit/Robo	Downstream effector of Abl in Slit-mediated repulsion at the midline, regulation of MT-F-actin crosstalk	[19; 79; 80]	Fig. 2C
TACC3	<i>Xenopus</i> spinal neurons	Slit2	Regulation of MT dynamics involved in steering behaviours	[21]	Fig. 2C
APC2	Chick retinal axons	Ephrin-A2	Regulation of MT stability and retinal axon pathfinding; +TIP function?	[27; 94]	Fig. 2D
APC	Mouse DRG neurons	Wnt3a	Required for Wnt3a-induced axonal remodelling, regulation of MT plus-end growth directionality in growth cones	[92]	-
NAV1	Mouse cortical neurons	Netrin-1	Instructs growth cone steering, promotes MT perdurance in the P-domain, MT-F-actin crosslinking	[26]	Fig. 1B
XMAP215	<i>Xenopus</i> embryonic neural tube explants <i>Drosophila</i> nervous system	EphrinA5 midline repellent cues ?	Regulation of growth cone morphology, response to the EphrinA5 repellent cue, MT-F-actin crosslinking; Growth cone repulsion from the midline by antagonising CLASP and Abl activity	[25; 81] [80]	Fig. 2D -
<b>Structural MAP</b>					
MAP1B	Mouse forebrain commissural axons  DRG axons DRG axons	Netrin-1/DCC; Draxin/DCC/Gsk3 $\beta$  Non permissive substrate Nitric oxide (NO)	Netrin-1- and Draxin-evoked axon outgrowth and repulsion, respectively  Promotes tyrosinated MT dynamics, reduces EB-binding at plus ends, regulates MT-F-actin crosstalk (?) in growth cones Abnormal steering behaviour when facing a non-permissive substrate, regulation of growth cone MT dynamics (?) Required for NO-induced growth cone collapse; MT effects ?	[20; 122] [111; 124; 125] [120] [121]	Fig. 1B & 2F - Fig. 2E
MAP6/STOP	Mouse subicular neurons	Sema6E	required for Sema3E-induced attractive outgrowth; Signalling hub protein	[131]	Fig.1C
Tau	Mouse cortical neurons	Wnt5a	Required for Wnt5a-evoked axon outgrowth and repulsive turning; MT invasion, remodelling and reorientation in the P-domain; MT-F-actin crosslinking	[24; 72] [116]	Fig.2G
CRMP2	Mouse peripheral nerves and callosal axons Zebrafish oculo motor axons	Sema3A/Cdk5 ?/ $\alpha$ 2-Chimaerin	Downstream effector of Sema3A required for axon repulsion; phosphorylation-induced detachment from MTs, permissive for MT breakdown Downstream effector of the RhoGAP $\alpha$ 2-Chimaerin; concerted regulation of MT dynamics with stathmin1/2	[141; 142; 145] [148]	Fig.2B
CRMP4	Zebrafish retinal axons	Neuropilin-1?	Required for midline crossing of retinal axons; MT effects (?)	[133; 146]	Fig.1D
DCX	Mouse cortical neurons	Sema3A/Nestin/Cdk5	Regulation of growth cone morphology and sensitivity to Sema3A; MT breakdown through phosphorylation-induced detachment from MTs (?)	[156]	Fig.2B
<b>MT severing/depolymerizing enzymes</b>					
M1 spastin	Zebrafish spinal motor neuron axons	BMPs	Controls rostrally-projecting axon pathfinding through BMP inhibition; MT dynamics (?), BMP receptor trafficking (?)	[23]	-
M61 spastin	Zebrafish spinal motor neuron axons	Neuropilin-1	Downstream effector of Neuropilin-1, controls steering behaviours at guidance choice points, Promotes growth cone MT dynamics (?)	[23; 164]	Fig. 2B
p60-Katanin	Zebrafish spinal motor neuron axons	?	Required for proper dorsal motor nerve pathfinding/targeting, Promotes MT dynamics in growth cones?	[31; 164]	Fig.3B
Fidgetin-like 1	Zebrafish spinal motor neuron axons	?	Controls growth cone morphology and steering behaviours at choice points, Regulates MT-plus-end composition and dynamics; Regulates MT-F-actin crosstalk (?)	[22]	-
Stathmin 1/2	Zebrafish oculo motor nerves	Sema3A/? $\alpha$ 2-Chimaerin	Required for $\alpha$ 2-Chimaerin-dependent ocular motor axon pathfinding/targeting; Concerted regulation of MT dynamics with CRMP2?	[148]	Fig.2B
<b>Molecular motors/regulatory proteins</b>					
kinesin-5	Rat primary neurons; <i>Xenopus</i> neural tube explants	NGF ?	Antagonises dynein-mediated forces to prevent MT invasion in the P-domain on the opposite of the growth cone turning side	[186; 188]	-
kinesin-12	Rat primary neurons	?	Restricts MT bidirectional transport in the axon, as well as MT invasion of the growth cone P-domain; Regulates growth cone morphology via its interaction with the actin cytoskeleton (?)	[187]	-
Figl1/KIF1bB (kinesin-3)/bicd1 complex	Zebrafish spinal motor neuron axons	?	Controls rostral motor nerve pathfinding; Dynein speed limiter; regulates bidirectional axonal transport of Rab3 vesicles	[73]	-
Kif13B (kinesin-3)	Mouse cortical neurons	Netrin-1	Controls Myo X anterograde transport (later required for DCC transport), Required to promote axon initiation and branching/targeting in a Netrin-1 dependent manner	[193]	Fig.1B
kinesin-1	Mouse cortical and hippocampal neurons	?	Regulates mouse growth cone remodelling and corticofugal axon pathfinding by mediating CB1R anterograde axonal transport	[194]	-
<b>Tubulin isoforms</b>					
TUBB2	Mouse/ human cortical neuron axons, ocular nerves	?	Required for proper axon pathfinding; regulates MT dynamics and MT-depolymerase targeting at MT plus ends	[183]	Fig.3A
TUBB3	Mouse/chicken commissural neurons DRG axons	Netrin-1/DCC Netrin-1/UNC-5	Controls Netrin-1/DCC-induced axon attraction of commissural neurons, promotes MT dynamics in growth cones Required for Netrin-1/UNC5-evoked axon repulsion and DRG axon pathfinding; promotes MT dynamics in growth cones	[28] [29; 202]	Fig.3A
<b>Tubulin modifying enzymes</b>					
TTL	Mouse cortical/hippocampal axons and olivary fibers	?	Required for proper navigational behaviour of olivary fibres at the floor plate and cortico-thalamic loop formation; Regulates growth cone morphology; recruitment of MT-F-actin crosstalk mediators in growth cones	[30; 210]	Fig.3B
TTL6	Zebrafish spinal motor neuron axons	?	Required for proper dorsal nerve pathfinding/targeting; regulator of p60-Katanin activity	[31]	Fig.3B
TLL11	Zebrafish spinal motor neuron axons	?	Required for proper rostral nerve pathfinding/targeting, concerted regulation of spastin activity with TTL6?	[31]	Fig.3B
<b>Table 1</b>					
Protein	Models/systems	Axon pruning mechanism	Functions in axon pruning	References	Figure
<b>MT severing/depolymerizing enzymes</b>					
spastin KIF2A	Mouse neuromuscular junction Mouse KO/ primary cultures	activity-dependent competition NGF deprivation	Elimination of supernumerary synaptic connections; MT severing Control of skin innervation; MT depolymerisation	[17] [236]	Fig.4A Fig.4B
<b>Structural MAP</b>					
CRMP2	Mouse hippocampus and visual cortex	Sema3F	Branch-specific elimination; MT dynamics/ disassembly	[145]	Fig.4C




**Figure 1**



**Figure 2**

## A Tubulin isotypes



## B Tubulin posttranslational modifications




Figure 3



