



# Hydrological classification by clustering approach of time-integrated samples at the outlet of the Rhône River: application to $\Delta^{14}\text{C}$ -POC

Nathan Bodereau, Adrien Delaval, Hugo Lepage, Frederique Eyrolle, Patrick Raimbault, Yoann Copard

## ► To cite this version:

Nathan Bodereau, Adrien Delaval, Hugo Lepage, Frederique Eyrolle, Patrick Raimbault, et al.. Hydrological classification by clustering approach of time-integrated samples at the outlet of the Rhône River: application to  $\Delta^{14}\text{C}$ -POC. Water Research, 2022, 220, pp.118652. 10.1016/j.watres.2022.118652 . hal-03799428

**HAL Id: hal-03799428**

**<https://hal.science/hal-03799428>**

Submitted on 5 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Hydrological classification by clustering approach of time-integrated samples at the outlet of the Rhône River: application to $\Delta^{14}\text{C}$ -POC

Nathan Bodereau<sup>1\*</sup>, Adrien Delaval<sup>1,2</sup>, Hugo Lepage<sup>1</sup>, Frederique Eyrolle<sup>1</sup>, Patrick Raimbault<sup>3</sup>, Yoann Copard<sup>4</sup>

<sup>1</sup> Institute for Radioprotection and Nuclear Safety (IRSN), PSE-ENV/LRTA, PSE-ENV/LMRE, BP 3, 13 115, Saint-Paul-lez-Durance, France

<sup>2</sup> Adict Solutions, Campus INP ENSAT, Avenue de l'Agrobiopole, BP 32 0607, 31 326 Castanet-Tolosan, France

<sup>3</sup> Aix Marseille Université, CNRS/INSU, Université de Toulon, IRD, Mediterranean Institute of Oceanography (MIO) UM 110, 13288, Marseille, France

<sup>4</sup> Normandie Univ, UNIROUEN, UNICAEN, CNRS, M2C, France, 76000 Rouen, France

\* Corresponding author.

E-mail address: bodereau.nathan@gmail.com

## Highlights

- The classification of the Rhône River hydrology was obtained by fuzzy C-mean logic
- A 5-Cluster configuration is efficient to characterize Rhône River hydrology
- $\Delta^{14}\text{C}$ -POC distribution among each cluster confirms the main assessments in literature
- This approach allows to identify the contribution of each hydrology to samples

## 21 Abstract

22 Within the framework of the Rhône Sediment Observatory, monthly time-integrated  
23 samples have been collected by Particle Traps in the last decade to monitor particulate  
24 contaminants in the Rhône River and its main tributaries. In this watershed with a contrasted  
25 hydrology, a clustering approach is used to classify the samples according to the main  
26 hydrological events. This approach has been applied to riverine particulate organic  
27 radiocarbon signatures ( $\Delta^{14}\text{C}$ -POC) that are strongly affected by the origin of the material  
28 and the occurrence of nuclear power plant releases. Suspended Particulate Matter (SPM)  
29 samples were collected near the outlet of the Rhône River and analysed for  $^{14}\text{C}$  along with  
30 particulate organic carbon (POC), chlorophyll a and tritium contents to confirm  $\Delta^{14}\text{C}$ -POC  
31 origins. Cluster Analysis, coupled to Principal Component Analysis, were performed based on  
32 monthly average water discharges of the Upper Rhône River and the five main tributaries.  
33 The classification obtained by fuzzy C-mean logic of the Rhône River hydrology into 5 clusters  
34 is similar to that already observed in the literature with Mediterranean/Cevenol flood,  
35 oceanic pluvial flood, nival flood, low-water level and baseflow clusters. The contribution of  
36 each cluster among the  $\Delta^{14}\text{C}$ -POC values demonstrate the complexity of hydrological  
37 classification of time-integrated samples. First, the samples with a unique and significantly  
38 dominant cluster are easily explained with negative  $\Delta^{14}\text{C}$ -POC values observed in the flood  
39 clusters due to input of  $^{14}\text{C}$ -depleted material from soil or rock weathering, and positive  
40 values observed in the low-water level and baseflow clusters due to anthropogenic input by  
41 nuclear industry. Second, samples that present a homogeneous mixture between several  
42 clusters demonstrate the occurrence of different hydrological events during the sampling  
43 periods. This tool appears as a solution to estimate the contribution of each hydrological  
44 event classification in time-integrated samples.

**Keywords** : Hydrological distribution, Cluster Analysis, Radiocarbon, Particulate Organic Carbon.

## 1. Introduction

Rivers are unquestionably links between the Earth system reservoirs as they connect 87 % of the Earthland surface to the oceans (Meybeck, 1982). They are also the receptacle of pollutants such as organic and inorganic micropollutants due to agricultural, industrial, domestical and mining activities (Meybeck, 2003) but also artificial radionuclides that are released by nuclear industries (Eyrolle et al., 2020).

Suspended Particulate Matter (SPM) is one of the aquatic fractions and belongs to sedimentary systems with processes of particles burying and remobilisation within river systems entering in the long-term cycle (Leithold et al., 2016). Photosynthesis, microbial oxidation, floodplain sediment remobilisation, sediment introduction are so many processes that can affect pollutant composition across the river (Meybeck et al., 2003; Walling et al., 2003). Thus, monitoring riverine SPM dynamic is of a great challenge in the perspective of health risk assessment, ecological restauration or river management (Walling et al., 2003). The development of network monitoring of SPM and pollutant fluxes has been performed over these last decades using SPM discrete sampling (Horowitz et al., 2001) but their frequencies of acquisition are still be constrained to event with a fixed periodicity, such as flood, and variations of SPM concentration ( $C_{SPM}$ ) that can occur between samples are often omitted (Misset et al., 2019) .

The deployment of Particle Traps (PT) has been described as a serious alternative from direct sample as they are time-integrative, low cost and easily handled (Phillips et al., 2000; Schulze et al., 2007). However, they can cover periods that might integrate several hydrological

events and affect our knowledge on contaminant dynamic (Poulier et al., 2019). Thus, constraining hydrosedimentary behaviours of river systems using time-integrated sample is challenging. Multivariate analysis and clustering to characterize catchment seasonal hydrology could be linked with time-integrated sample. The Principal Component Analysis (PCA) coupled to Clustering Analysis (CA) has been widely used in Earth and Environmental sciences (e.g., Delaval et al., 2021, Raux et al., 2011). It is also an usual method for data mining when it comes to identify groups and classify individuals. However, many algorithms exist and present different results and convergence speeds (Jain et al., 1999). Fuzzy c-mean algorithms are usually considered as reliable algorithms in environmental sciences (Delaval et al., 2021, Kim et al., 2011). Their success is due to their simple geometric criteria and their robustness without any prior information on the cluster structure such as possible overlap or sphericity (Kamel and Selim, 1994). In addition, the fuzzy approach provides cluster membership degrees (probability for an observation to belong to a cluster) instead of sharp cluster boundaries. As a result, the ambiguity of the data can be preserved and these probabilities can be used later for post treatments or deeper interpretations (Kim et al., 2011).

The Rhône River is one of the major sources of freshwater and SPM to the Mediterranean Sea (Milliman and Farnsworth, 2011). It is also a strongly anthropized catchment where inorganic and organic pollutants are monitored for several decades (e.g., Delile et al., 2020; Eyrolle et al., 2018; Launay et al., 2019; Ollivier et al., 2011; Poulier et al., 2019; Radakovitch et al., 2008). Within the Rhône Sediment Observatory (OSR) framework, SPM are collected every month on the Rhône River and its main tributaries using PT (Lepage et al., 2022). While the results demonstrate a strong link between the contaminant concentrations and the hydroclimatic heterogeneities of this catchment including different tributary regimes (e.g.,

[Delile et al., 2020; Poulhier et al., 2019](#)), the occurrence of different events in a same time-integrated sample may lead to a bias in interpretation.

In this study, we propose to classify time-integrated samples according to hydrological events using a clustering approach and to investigate the distribution of the clusters among the concentrations of  $\Delta^{14}\text{C}$  (Carbon 14 reported to 1950 standard) in Particulate Organic Carbon (POC). Here, the challenge is that while the hydrology of the Rhône River was already classified based on hourly or daily water discharge values by previous studies ([e.g., Mourier et al., 2014; Ollivier et al., 2011; Poulhier et al., 2019; Radakovitch et al., 2008; Zebracki et al., 2015](#)), time-integrated samples, that can cover periods from two weeks to one month, might integrate several hydrological events ([Delile et al., 2020; Poulhier et al., 2019](#)). The hydrology of the Rhône River main tributaries during time-integrated sampling (164 samples) has been classified with a fuzzy-C mean algorithm. Carbon-14 ( $^{14}\text{C}$ ), expressed as  $\Delta^{14}\text{C}$ , in riverine POC, is a radioisotope occurring naturally in the upper parts of the atmosphere and from anthropogenic sources such as atmospheric nuclear weapon tests (1945-1990) or releases of nuclear industries ([Eyrolle et al., 2018](#)). The  $^{14}\text{C}$ -POC is selected here because its dynamic in the Rhône River is well documented:  $^{14}\text{C}$ -labelled POC by nuclear industry releases during low-water level is opposite to depleting  $^{14}\text{C}$ -POC by introduction of  $^{14}\text{C}$ -dead Carbon, during floods, coming from rock or soil weathering ([Copard et al., 2018 ; Eyrolle et al., 2018; in review](#)). Geochemical indicator distributions have been then discussed such as POC content in SPM (%POC) and Chlorophyll a content in POC (%Chl.a) to study the distribution of the clusters among the  $\Delta^{14}\text{C}$ -POC. To identify potential artificial  $^{14}\text{C}$  releases, tritium concentrations collected in water masses were also investigated.


## 114 2. Material and methods

### 115 2.1. Study area

116 The Rhône River basin covers a 98 500 km<sup>2</sup> area (**Figure 1**), with a river stream length of 816  
117 km. The watershed boundaries are delimited with the presence of the Alps, Jura, Cevennes  
118 and Vosges mountains leading to a strong topographic gradient and hydroclimatic  
119 heterogeneities ([Olivier et al., 2009](#)). It encompasses tributaries with contrasted hydrologies  
120 :

121 i) oceanic pluvial regime observed at the North of the watershed in winter (Saône and Ain  
122 rivers);

123



124

125 Figure 1 – Map of the Rhône watershed with the delimitations of main tributary basins discussed in this study.

126 The locations of SORA and Jons stations, the tributary stations and the nuclear industries are also added.



127

128 ii) flash Mediterranean rainfalls, coming from Mediterranean Sea, and generally observed in  
129 autumn, that occur in the Southeastern part of the watershed (Durance, Drôme and Ouvèze  
130 rivers);

131 iii) flash Cevenol rainfalls that conduct to the same sharp flow variation than Mediterranean  
132 floods but occur early in fall season and are more located around the Cevennes mountains  
133 (Ardèche, Cèze and Gardon rivers);

134 iv) alpine regimes linked to the snow melt and generally occurring in Alps tributaries (Arve,  
135 Fier, Isère and Durance rivers) during spring;

136 v) generalized floods that encompass all flood events ([Delile et al., 2020](#), [Pont et al., 2002](#),  
137 [Sadaoui et al., 2016](#)).

138 In addition, fourteen nuclear reactors and various nuclear industries are distributed along  
139 the Rhône River making it as one of the most nuclearized river in the world ([Eyrolle et al.,](#)  
140 [2020](#)).

## 141 2.2. Sampling methodology

142 Samples were collected at the SORA monitoring station that is located at the outlet of the  
143 Rhône River, 47 km upstream from the mouth, and has been designed to monitor river  
144 inputs into the Mediterranean Sea ([Lepage et al., 2022](#)). It is located on the “Grand Rhône”  
145 branches where 90% of the water and sediment fluxes of the Rhône River are transiting,  
146 while the remaining 10% transit through the “Petit Rhône” ([Pont et al., 2002](#)). Water intake  
147 is performed on a floatable structure at 7 m from the bank and 0.5 m under the surface

regardless the water discharge and continuously supplies the sampling devices (Lepage et al., 2022).

The SPM are collected over a two-week to one-month period in PT, designed with the same characteristics as described by Schulze et al. (2007), deep-frozen (-18°C), freeze-dried, and finally homogenized by grinding in an agate mortar. SPM samples are stored in the dark and at ambient temperature before analysis (Lepage et al., 2022). A total of 164 samples were collected at SORA from 2007 to 2020 and  $\Delta^{14}\text{C}$ -POC was measured among 54 of them.

POC and Chlorophyll a are separately measured in SPM filtered in daily composite water samples. Sampling is achieved using a cooled automatic water sampler that fills a daily bottle with 150 mL every 90 minutes to constitute a composite sample each day (Eyrolle et al., 2020). The sampling frequency is reduced to every 30 minutes to constitute a composite sample every 4 hours when water discharge is greater than  $3000 \text{ m}^3 \text{ s}^{-1}$ . Samples are poisoned with  $\text{HgCl}_2$  and kept at 5°C until they are filtered on a 0.7 Whatman pre-conditioned glass fiber filters GF/F (dried at 500°C for 4 hours). The samples are then frozen, respectively, until laboratory analysis (Lepage et al., 2022). For tritium analysis, a few millimeters of water is collected per 80 minutes to obtain a monthly composite sample. Water is then filtered through  $0.22 \mu\text{m}$  GF/F before analysis (Eyrolle et al., 2020).

## 2.3. Principal Component Analysis (PCA) and Cluster Analysis (CA)

### 2.3.1. Input data

With the perspective to trigger all hydrologic regimes occurring on the Rhône according to the literature, the water discharges (Q) of the five main tributaries (Saône, Isère, Ardèche, Durance and Gardon rivers) as well as the measures conducted at the Jons monitoring

station, considered as the reference station to evaluate water discharge from the Upper Rhône River, were used to define the clusters (**Figure 1**). For those stations, daily averaged water discharge data are continuously measured ([Thollet et al., 2021](#)) at the exception of the Upper Rhône station (Jons) where  $Q$  is estimated from a 1D hydrodynamic model with water discharges of upstream main tributaries as input data ([Launay et al., 2019](#)).

Mean water discharge values ( $Q_{\text{mean}}$ ) for each tributary and Jons monitoring stations were estimated from the start to the end of each 164 PT samples and submitted in PCA. We assumed that this is enough to characterized tributary hydrograph shapes for each period of acquisition ([Hannah et al., 2000](#), [Ouyang et al., 2010](#)).

In order to better interpret the results of the PCA and the clustering, hydrological data ( $Q$  and  $C_{\text{SPM}}$ ) at the SORA station was also used so as the  $C_{\text{SPM}}$  of the tributary stations if available. At SORA, daily averaged water discharge data are continuously measured ([Thollet et al., 2021](#)) while the SPM concentration ( $C_{\text{SPM}}$ ) at SORA station is measured by time-integrating weighing of daily automatic samples ([AFNOR, 2005](#)). At tributaries, averaged daily  $C_{\text{SPM}}$  is estimated from turbidity and  $C_{\text{SPM}}$  rating curves with 10-minute frequency of acquisition and monthly automatic water samples ([Lepage et al., 2022](#)). A mean value ( $C_{\text{SPM-mean}}$ ) was calculated for each PT sample.

### 2.3.2. PCA and CA algorithms

The PCA has been performed using R programming language including the R packages “FactoMineR” for PCA ([Husson et al., 2010](#)), “e1071” ([Meyer et al., 2021](#)) and “fclust” for fuzzy c-mean algorithms ([Ferraro et al., 2019](#)). The goal of PCA is to describe a data set using a smaller number of uncorrelated variables, while retaining as much information (variance) as possible. The reduction is achieved by transforming the data into a new set of continuous

variables named the principal components. The reduction of dimensionality provides a framework to visualize the Rhône hydrology. The essential of the information being on the first components ([Husson et al., 2010](#)), PCA is usually used as a pre-processing tool before clustering analysis ([Delaval et al., 2021](#)).

Then, clustering algorithms such as c-mean algorithms (crisp or fuzzy) classify individuals according to their variable similarities and allows to identify scenarios and trends in river hydrology and hydrochemistry. This is done by randomly defining  $i$  centroids in the same coordinate systems as the individuals. Each individual  $x$  (total of  $k$ ) is assigned to the closest centroid center  $C_i$ . The barycenter of each subgroup is then calculated and becomes the new centroid. This procedure converges when the centroids are no longer moving. In addition, the fuzzy alternative introduces two new parameters. The first one is the membership coefficient. This membership represents how closely the  $k_{th}$  data object ( $x_k$ ) is located from the  $i_{th}$  Cluster center. Membership score indicates the degree (in %) to which a sample belongs to each cluster. Each sample is assigned to the cluster with the highest membership among the others.

The second parameter  $m$  is the fuzzifier. It is greater than 1 and usually dependent on the dataset structure because it represents the degree of overlap of the clusters ([Klawonn and Höppner, 2003](#)). When  $m$  is close to one, the solution of the fuzzy c-means algorithm is similar to the one of c-means. Elements are simply assigned to only the nearest cluster and membership to other clusters is nearly existent. When  $m$  is large, fuzziness is also large and clusters are blurred which can be a more accurate description of the system. More details about c-mean algorithms are provided in [Bezdek et al., \(1988\)](#).

216 The best combination of  $\{C, m\}$  parameters is not determined by the algorithm. One  
217 approach is to run different simulations with different  $\{C, m\}$  pairs and to check the efficiency  
218 of clustering with a quality criteria (Ramze Rezaee et al., 1998). River hydrology related to  
219 clustering studies have a number of clusters falling in the interval  $[2, 8]$  (Delaval et al., 2021;  
220 Javed et al., 2021; Ouyang et al., 2010). Our number of  $\Delta^{14}\text{C}$ -POC observations being limited  
221 ( $n=54$ ), we are expecting 8 clusters at most (Chaimontree et al., 2010) and we will set the  
222 test interval for the number of clusters to  $[2, 8]$ .

223 The fuzzifier  $m$  can be in theory any real number between 1 and  $\infty$ .  $m$  is usually set to 2 by  
224 default but can be any real number between 1 and  $\infty$  usually decreasing with dataset size  
225 (Klawonn and Höppner, 2003). In our tests, we will take 1 as the lower limit for  $m$ . The upper  
226 limit is  $m=3$  according to the empirical threshold equation based on the length and  
227 dimensions of the dataset proposed by Schwämmle and Jensen (2010). As a result, we will  
228 perform clustering on the interval  $[2, 8]$  for  $C$ , the number of clusters and on the interval  $]1, 3]$   
229 for  $m$  the fuzzifier.

## 230 2.4. Analyses

### 231 2.4.1. Carbon-14 measurement

232 The  $^{14}\text{C}$  measurement in SPM are performed using a 3 MV NEC PELLOTRON Accelerator Mass  
233 Spectrometry (Artemis AMS facility, Dumoulin et al., 2017). Briefly, SPM samples are firstly  
234 decarbonated with 0.5 M HCl, placed in Pyrex sealed tubes and calcined ( $850^\circ\text{C}$  for 5 hours).  
235  $\text{CO}_2$  is collected into a glass line under vacuum ( $< 10^{-5}$  mbar) and cryogenically trapped via  
236 liquid nitrogen. The sample  $\text{CO}_2$  is reduced with hydrogen and iron powder as a catalyser  
237 following the protocol of Voegel et al., (1984) and pressed into an aluminium cathode (1 mm  
238 hole diameter) to form a graphite target.

In the AMS, the target is submitted to high bombing with Caesium sputter ion sources and sequentially sent, through a 90° magnet, to an accelerator tube. Here, Carbon isotopes are stripped in an argon gas and distinguished through a 45° analyser according to their weight. <sup>14</sup>C ions are counted in an ionization chambers (Cottureau et al., 2007).

After AMS counting, the <sup>14</sup>C/<sup>12</sup>C ratio R is normalized (R<sub>n</sub>) to a δ<sup>13</sup>C of -25‰ in order to take into account carbon isotopic fractionation (Stuiver and Polach, 1977) . A 1950 standard R<sub>s</sub> (i.e., oxalic acid II, a crop of sugar beet) is also incorporated in the sample batch and normalized as R<sub>sn</sub>. Following Stuiver and Polach (1977) formulation, Δ<sup>14</sup>C is privileged in order to take into account radiodecay (**Equation 1**) with y, the year of measurement and the Godwin's mean life (8267) estimated from soil and water samples (Godwin et al., 1962).

$$\Delta^{14}\text{C} = \left( \frac{R_n}{R_{sn}} \times e^{\left(\frac{y-1950}{8267}\right)} - 1 \right) \times 1\,000 \quad \text{Equation 1}$$

Carbon, after the 1950 standard, has a Δ<sup>14</sup>C up to 0 ‰ and could integrate the contribution of anthropogenic sources such as the effects of thermonuclear weapon tests (1945-1990) or releases of nuclear industries. Values reaching -1000‰ are linked to material old of up to 50 000 years BP (Trumbore et al., 2016).

#### 2.4.2. Particulate nutrient and tritium analyses

The POC concentration measurements were carried out at the Mediterranean Institute of Oceanography (M.I.O.) and were performed on daily decarbonated SPM by using high combustion procedure (950°C) on a Carbon Nitrogen (CN) Integra mass spectrometer (serCon Ltd, Crewe, UK) as described in Raimbault et al. (2008). Chlorophyll a measurement is conducted through fluorimetry methodology after its extraction with methanol according

to Raimbault et al. (2004). Tritium analyses ( $\text{Bq l}^{-1}$ ) were carried out at the LMRE/IRSN by liquid scintillation counting (AFNOR, 2015).

### 3. Results

#### 3.1. Data description

The distributions of environmental data treated by CA are available in *Dataset S1*. The  $Q_{\text{mean}}$  and  $C_{\text{SPM-mean}}$  histories are respectively presented in **Figures 2.a.** and **2.b.** and  $\Delta^{14}\text{C-POC}$  values in **Figure 2.c.**




Figure 2 - Distribution of a.)  $Q_{\text{mean}}$  ( $n = 164$ ), b.)  $C_{\text{SPM-mean}}$  ( $n = 163$ ) and c.)  $\Delta^{14}\text{C-POC}$  values ( $n = 54$ ) over the 2007-2020 period.

274

275 During the whole 2007-2020 period, for all 164 samples,  $Q_{\text{mean}}$ , at Arles varied between 558  
276 and  $3\,668\text{ m}^3\text{ s}^{-1}$  (mean=  $1\,565\text{ m}^3\text{ s}^{-1}$ , s.d. =  $687\text{ m}^3\text{ s}^{-1}$ ) while  $C_{\text{SPM-mean}}$  varied between 2.9 mg  
277  $\text{l}^{-1}$  and  $879.7\text{ mg l}^{-1}$  (mean=  $68.3\text{ mg l}^{-1}$ , s.d. =  $117.4\text{ mg l}^{-1}$ ).  $\Delta^{14}\text{C-POC}$  signatures are ranged  
278 from -742 to 699 ‰ (mean = -84.5 ‰, s.d. = 346 ‰). The signatures are unequally  
279 distributed over the period (**Figure 2.c.**) because  $\Delta^{14}\text{C-POC}$  analyses in SPM have been  
280 continuously performed since 2014 despite some exceptional measurements performed  
281 around 2009 and 2012 (Eyrolle et al., 2018).

282 The average POC content in SPM (%POC) is estimated between 1.03% and 10.37% (mean=  
283 3.95 %, s.d. = 1.73 %). The average chlorophyll a content in POC (%Chl.a) is varying between  
284 0.01% and 2.15 % (mean = 0.34 %, s.d. = 0.42 %), while tritium concentrations in water  
285 between 0.4 and  $20.7\text{ Bq l}^{-1}$  (mean =  $6.1\text{ Bq l}^{-1}$ ).

### 286 3.2. Principal Component Analysis

287 The PCA transformed the variables (mean Q for the five tributaries and Jons) into six vectors  
288 (**Figures 3a and b**).

289






Figure 3 – Principal component analyses 1 and 2 of the dataset. The PCA were conducted on the mean of the water discharge of the five main tributaries of the Rhône River (Saône, Isère, Ardèche, Gardon and Durance rivers) and the station of Jons that monitors the Upper Rhône River. Clusters are highlighted by colours.

The F1 axis explains 43.2 % of the variance of the dataset and all tributary vectors are oriented toward the right side of the PCA. The F2 axis, explaining 31.4% of the variance, distinguishes, from the top to the bottom, southern tributaries (Ardèche, Gardon and Durance rivers) and northern ones (Saône, Isère and Upper Rhône rivers) (**Figure 3.a**). Through the F1 × F2 axis configuration (PCA1), the observations are concentrated around the 0 axis on the left side of F2 while they are scattered on the right side. Through the F2 × F3 axis (14.2% of the variance) configuration (PCA2), the Isère and Durance vectors are located on the top side of the PCA whereas other vectors are located on the bottom side. As observed for PCA1, Ardèche and Gardon vectors are grouped. The F4 axis explains 6.6 % of the variability and separates Durance and Saône rivers from other tributaries (*Supplementary information, Figure S1*). Through F3 × F4 axis configuration (PCA3), samples are centred and staggered across the F4 axis.

308

### 309 3.3. Cluster number assessment

310 The fuzzy clustering was performed with the first four components of the PCA in order to  
311 cumulate an explained variance of up to 95 % (*Figure S2*). The performances were estimated  
312 through Xie and Beni index (XB) (to minimize) according to different {C,m} simulations (*Table*  
313 *S1*). The five-cluster configuration showed the lowest score ( $m = 13$ ,  $XB = 0.296$ ) and will be  
314 retained for the remainder of this work.

### 315 3.4. Hydrodynamic distribution for each cluster

316 Respectively, 12.1 %, 20.1 %, 11.6 %, 31.2 % and 25.0 % of  $Q_{mean}$  are distributed among  
317 clusters 1, 2, 3, 4 and 5. The  $Q_{mean}$  and  $C_{SPM-mean}$  distributions at Arles, Jons and tributaries  
318 are respectively available in *Tables S2 and S3*. In both PCA1 and PCA2, Cluster 1 includes  
319 Ardèche and Gardon vectors (**Figures 3a and 3b**). For clusters 2 and 3, the results are  
320 contrasted. In PCA1, Cluster 2 integrates the three northern stations while in PCA2 the Isère  
321 vector is excluded. For Cluster 3, only the Isère vector is included in PCA2 while the Saône  
322 vector is included in PCA1. Finally, in both configurations, the Durance vector is located  
323 between clusters 1 and 3.

324 Clusters 1, 2 and 3 are characterized by a maximum  $Q_{mean}$  above the daily flood threshold at  
325 Arles ( $3000 \text{ m}^3 \text{ s}^{-1}$ ) with high standard deviations (up to  $400 \text{ m}^3 \text{ s}^{-1}$ ). For clusters 1 and 3,  
326 similar results were observed for the  $C_{SPM-mean}$  with average concentrations close to  $200 \text{ mg l}^{-1}$   
327 at Arles and high values in the Ardèche (mean  $C_{SPM-mean} = 27 \text{ mg l}^{-1}$ ), Durance (mean  $C_{SPM-}$   
328  $mean = 661 \text{ mg l}^{-1}$ ) and Gardon (mean  $C_{SPM-mean} = 58 \text{ mg l}^{-1}$ ) rivers for Cluster 1 and Isère (mean  
329  $C_{SPM-mean} = 333 \text{ mg l}^{-1}$ ) and Durance (mean  $C_{SPM-mean} = 498 \text{ mg l}^{-1}$ ) rivers for Cluster 3. For

330 Cluster 2, while the  $C_{SPM-mean}$  observed at Arles (mean  $C_{SPM-mean} = 77 \text{ mg l}^{-1}$ ) are lower than  
331 clusters 1 and 3 (mean  $C_{SPM-mean} > 150 \text{ mg l}^{-1}$ ).

332 Located on the left side of the PCA1, Cluster 4 presents lowest values of  $Q_{mean}$  with a mean  
333 at Arles of  $849 \text{ m}^3 \cdot \text{s}^{-1}$ . Cluster 4 is also characterized by the lowest  $C_{SPM-mean}$  at Arles and for  
334 all tributaries. Cluster 5 is centred between F1 and F2 with a predominance of the Isère and  
335 Durance rivers (**Figure 3b**) and also of the Saône River but their discharges remain lower  
336 than those observed in clusters 2 and 3. Cluster 5 presents moderate  $C_{SPM-mean}$  at Arles  
337 (mean=  $33 \text{ mg l}^{-1}$ ) with low variability and moderate increases of the Durance and Isère  
338 rivers  $C_{SPM-mean}$  (77 and  $54 \text{ mg l}^{-1}$  respectively).

339 Percentages of occurrence for calendar months are presented in **Figure 4**. The samples for  
340 Cluster 1 are generally centred in from October to December (maximum of 47%), February  
341 (25%) and March (23%). Cluster 2 samples are mainly centred from December to April  
342 (maximum of 67%). Cluster 3 is mainly observed from April to June (up to 46%). Cluster 4  
343 samples are generally observed from July to November (up to 85%). Finally, Cluster 5 mainly  
344 occurs from March to August with a peak in July (58%).




Figure 4 – Inter-annual occurrences of each cluster among the 164 samples. Flood delimitations were estimated from Delile et al., 2020.

### 3.5. $\Delta^{14}\text{C}$ -POC distribution among the clusters

Distribution of  $\Delta^{14}\text{C}$ -POC among the clusters are presented in **Figure 5** where each value was associated to the cluster with the highest contribution (**Figure 6**). While  $\Delta^{14}\text{C}$ -POC values cover all hydrological events, the number of  $\Delta^{14}\text{C}$ -POC data is unequally distributed among clusters (7, 7, 8, 12 and 20 respectively).





Figure 5 –  $\Delta^{14}\text{C-POC}$  (n=54) distribution among the 5 clusters over the 2007-2020 period. Each value was associated to the cluster with the highest membership score. Dotted horizontal line represents the year 1950 for which the value is 0 ‰ (cf. equation 1).

Cluster 1, Cluster 2 and Cluster 3 include the lowest values with a median of -223, -367 and -487‰. Cluster 2 includes the largest distribution of  $\Delta^{14}\text{C-POC}$  signatures from -613 to 655 ‰ while clusters 1 and 3 are concentrated between -614 and 142 ‰ and -742 and -40‰. Most of the values of the Cluster 4 are above 0‰ with a median of 174 ‰, excepted for three negative values (-420, -354 and -185 ‰). The Cluster 5 is equally centred around 0‰ with a median of -24 ‰.

The dominant or attributed cluster in each sample is characterized by its membership degree ranging from 25.8% to 84.6% (**Figure 6**). A few samples show a homogeneous cluster membership score distribution highlighting that they could belong to another cluster than the attributed one. For example, the sample with the highest  $\Delta^{14}\text{C-POC}$  in Cluster 3 (-40‰) presents membership degrees ranging from 13.3 to 26.7 %. Respectively 29%, 57%, 25%,

371 75% and 35% of the samples have an appurtenance degree to their attributed cluster higher  
 372 than 50% in Cluster 1 to 5.

373



374

375 Figure 6 – Distribution of cluster memberships for  $\Delta^{14}\text{C-POC}$  values ( $n = 54$ ) among the classification. Each  
 376 membership score is expressed in %.

### 3.6. Geochemistry distribution

The  $\Delta^{14}\text{C}$ -POC relationships with  $C_{\text{SPM-mean}}$ , average %POC, %Chl.a and tritium concentrations, among the 5 clusters, are presented in **Figure 7**. Overall,  $\Delta^{14}\text{C}$ -POC shows positive relationships with %POC and a negative relationship with  $C_{\text{SPM-mean}}$ . High  $\Delta^{14}\text{C}$ -POC are correlated with high %Chl.a (up to 0.6 %) and high tritium concentrations (up to 5 Bq l<sup>-1</sup>). Cluster 1, 2 and 3 gather low %POC while clusters 4 and 5 present higher and more variable records. Highest %POC, %Chl.a. and tritium records are observed for Cluster 4 and Cluster 5 and some exceptions in Cluster 2.




Figure 7 –  $\Delta^{14}\text{C}$ -POC relationships with a.)  $C_{\text{SPM-mean}}$ , b.) average %POC, c.) average %Chl.a. and d.) tritium concentrations according to the cluster.

## 4. Discussion

### 4.1. Characterisation of the clusters

#### 4.1.1. PCA interpretation

The PCA1 made with time-integrated samples is similar to the one given by [Sadaoui et al., \(2016\)](#) with daily water discharge (**Figure 3a**). The F1 axis contains the information of water discharge variation for all tributaries while F2 axis separates the northern tributaries from the southern tributaries that have contrasted hydrologies in accordance with [Pont et al., \(2002\)](#). In PCA 2, F3 axis explains the temporal variability (spring vs autumn) and opposes alpine tributary hydrology to other hydrologies (**Figure 3b**).

#### 4.1.2. Cluster hydrodynamic characterisation

Despite the use of integrated samples (and associated hydrological data) in this study that may aggregate several hydrological events, the five clusters defined are similar to the hydrological classification of the Rhône River made from daily water discharge ([Pont et al., 2002](#), [Sadaoui et al., 2016](#)).

Cluster 1 is linked to southern tributary vectors (**Figure 3a**) and gathers southern flash-flood events ([Pont et al., 2012](#), [Eyrolle et al., 2012](#), [Delile et al., 2020](#)). No distinction between Cevenol and Mediterranean floods are proposed by the algorithm. The high standard deviations of average  $Q_{\text{mean}}$  and  $C_{\text{SPM-mean}}$  highlight sharp variabilities. In the literature, southern events contribute the most on large export of SPM at the Rhône River scale. [Eyrolle](#)



et al., (2012) or Copard et al., (2018) demonstrated that, the Durance River could contribute to sediment load from < 1% to 80% during Mediterranean flood events. However, numerous studies have also estimated sediment budgets using geochemical methods and agree on the exceptional but non-negligible contribution of the Cevenol floods, that could occasionally reach more than 40% (Ollivier et al., 2011; Radakovitch et al., 2008; Zebracki et al., 2015).

Cluster 2 gathers oceanic regime with  $Q_{mean}$  distribution at Arles occurring during winter and the activation of the Saône and Upper Rhône rivers. The contrast between average  $C_{SPM-mean}$  and  $Q_{mean}$  is explained by the main roles of these two tributaries as water suppliers. At the interannual scale, their contribution on water fluxes reaches more than 60% while their contribution in term of SPM (11 and 6 % respectively) are lower (Delile et al., 2020; Poulhier et al., 2019; Sadaoui et al., 2016).

Cluster 3 observations occur during the end of spring season (**Figure 4**) and gather nival flood regime with net predominance of alpine rivers (Isère and Durance rivers) (**Figure 3b**). Their contribution can exceed 80 % in water fluxes and over 100% in SPM fluxes (Delile et al., 2020, Poulhier et al., 2019).

Cluster 4 observations gather low-water level, during summer (**Figure 4**), with lowest  $Q_{mean}$  and  $C_{SPM-mean}$ . During such period, the production of autochthonous organic matter is increased as observed with the increase of %POC and %Chl.a (**Figure 4**).

Cluster 5, located at the center of the PCA1 (**Figure 3.a.**), gathers baseflow regime with medium  $Q_{mean}$  and  $C_{SPM-mean}$  values. It occurs over spring and summer seasons (**Figure 4**) and the slight increases of  $Q_{mean}$  and  $C_{SPM-mean}$  in the Durance and Isère rivers indicate nival episodes that might happen during these periods. As for Cluster 4, high values of %POC and %Chl.a indicate a production of autochthonous POC.

## 4.2. Classification of $\Delta^{14}\text{C}$ -POC

The  $\Delta^{14}\text{C}$ -POC depletion observed for clusters 1 and 3 confirms the main knowledge presented in the literature about  $^{14}\text{C}$  dynamic in the Rhône (Copard et al., 2018, Eyrolle et al., 2018; Eyrolle et al., in review). Indeed, the increase of  $\text{C}_{\text{SPM-mean}}$  and the decrease of %POC (**Figure 7**) confirm the organic matter dilution with particulate mineral fractions by the weathering of carbonate rocks (Higuera et al., 2014). Then, rocks such as i) Black Marls, composed bare Mesozoic Badlands located among the French alps arc (Graz et al., 2011) or ii) Coal mines observed in slag heaps of Cevennes or mining operation sites in Briançonnais area of Alps, containing high contents of organic matter old of up 50 000 years BP. Because they are characterized by low vegetation density and violent hydroclimatic conditions, they could be considered as main sediment hotspots and could release consequent amount of organic matter (Eyrolle et al., in review). However, their contribution in sediment budget at the Rhône scale still poorly documented, excepted for the Durance River (Copard et al., 2018). Another source of such  $^{14}\text{C}$ -depleted material might be the remobilisation of aging sediment POC trapped in flood plain across the main network (e.g., Leithold et al., 2016).

The case of Cluster 2 is different as the oceanic floods (Saône and Upper Rhône rivers) are characterized by a low level of  $\text{C}_{\text{SPM-mean}}$ . The range of  $\Delta^{14}\text{C}$ -POC values and their disparity around 0 (**Figure 5**) can be explained by the fact that the SPM sampled for the measurement of  $\Delta^{14}\text{C}$ -POC do not specifically come from these two tributaries. Since the samples are integrated over a period of about one month, they are very sensitive to floods that generate a lot of material or to anthropogenic inputs (e.g., dam flushing).

Cluster 4 and Cluster 5 present more enriched  $\Delta^{14}\text{C}$ -POC (**Figure 5**), displaying a privileged autochthonous origin, related to high %Chl.a, for positive signatures (after 1950) showing

aquatic plants development preferentially higher when water flow and turbidity are low (Figure 7) (Higuera et al., 2014). High tritium concentrations, combined to high  $\Delta^{14}\text{C}$ -POC values in clusters 4 and 5 might be related to the releases from the nuclear facilities (Figure 7). At the outlet of the Rhône River, the possibility of nuclear release increasing  $\Delta^{14}\text{C}$ -POC is significant when %Chl.a is high suggesting a contamination of POC when primary productivity is important. Several studies have highlighted that  $^{14}\text{C}$  is released as dissolved inorganic forms (DIC) because of oxidation processes applied in effluents before releases (e.g., Kang et al., 2019) and it could integrate POC through photosynthesis. However additional investigations might be conducted as the sampling methods were not the same for these two parameters and could lead to bias in the interpretation.

#### 4.3. Relevance of Cluster analyses for classification of $\Delta^{14}\text{C}$ -POC signatures.

Figure 6 displays the complexity of hydrological classification of time-integrated sampling as a few samples are characterized by a mixture of clusters including several hydrological events and  $\Delta^{14}\text{C}$ -POC signatures including different origins. Moreover, for the outliers (Figure 5), the relationship between the cluster classification and the  $\Delta^{14}\text{C}$ -POC values is complex to explain and requires looking at the hydrograph during the sampling periods. For example, southern quick flood events have been registered in samples of low-water level related clusters (4 and 5) with depleted values of  $\Delta^{14}\text{C}$ -POC. For Cluster 1, the positive value of  $\Delta^{14}\text{C}$ -POC is observed in a sample with a southern flash flood occurring at the beginning of the sampling period but followed by a low-water level period favourable of  $\Delta^{14}\text{C}$ -artificial POC. Cluster 1 samples also have consequent appurtenances to clusters 2 and 3. The sum of the membership degrees for the flood-related clusters (1, 2 and 3) is always higher than 58% and highlights that i) northern floods could also simultaneously occur than southern floods in autumn and ii) some southern floods could be observed during spring season as the same

time than nival flood events (**Figure 4**). The complex relationship of Cluster 2 with  $\Delta^{14}\text{C}$ -POC values is due to the low  $C_{\text{SPM}}$  load of the Saône and Upper Rhône river floods and the consequent sensitivity to other events such as the Durance river floods or industrial discharges. The main solution to reduce the disparity of the values would have been to use the flux of SPM instead of the water discharge as the input data to determine the clusters. Therefore, this would improve the classification of  $\Delta^{14}\text{C}$ -POC values measured on SPM. Unfortunately, such data was not available from all the tributaries and for the entire period of study (starting since 2011), while clustering using water discharge was already commonly used (Delaval et al., 2021). Finally, samples in the Cluster 5 seem to be a mix between the Cluster 4 (low-water level cluster) and the flood-related clusters (1, 2 and 3). For this cluster, centred in the PCA (**Figure 3**), the absence of relationship with the  $\Delta^{14}\text{C}$ -POC values is directly related to the complexity to categorize samples collected during baseflow.

## 5. Conclusion

This study presents a first classification of time-integrated samples by clustering approach according to the Rhône hydrology. Principal Component Analysis (PCA) followed by fuzzy-C mean Cluster Analysis have been performed using time average of Q among the five major water tributaries and the Upper Rhône River monitored at Jons. We used  $\Delta^{14}\text{C}$ -POC data collected in Particle Traps (PT) over a two-week to one-month sampling period as a candidate to validate the classification. The classification exhibited 5 groups of hydrology similar to that described in the literature: southern floods with Mediterranean and Cevenol hydrologies confounded (Cluster 1), northern oceanic pluvial floods (Cluster 2), nival floods (Cluster 3), but also baseflows (Cluster 5) and low-water levels (Cluster 4). Overall,  $\Delta^{14}\text{C}$ -POC distribution in time-integrated sample confirms the clustering classification according to

flood-related, baseflow and low-water level clusters. High  $\Delta^{14}\text{C}$ -POC signatures, correlated to high tritium concentrations, are observed in clusters 4 and 5, incorporating nuclear releases, and are opposed against depleted signatures occurring during flood events observed in clusters 1 and 3 because of the  $\Delta^{14}\text{C}$ -depleted material mobilisation from tributaries (e.g., black marls). The  $\Delta^{14}\text{C}$ -POC outliers observed at each cluster show the sensitivity of PT samples to short events remobilising high sedimentary material or nuclear releases during a long sampling period especially for clusters 2 and 4. Furthermore, the overlaps of different events making difficult the Cluster Analysis to attribute the origins of  $\Delta^{14}\text{C}$ -POC especially for flood clusters. Indeed, homogenous distribution of membership scores at some samples in clusters 1 to 3 shows a mixture of contrasted flood events over the sampling period. Because Cluster 5 is a mix of flood and low-water level events, some  $\Delta^{14}\text{C}$ -POC values are incorrectly attributed here. If this method of classification with  $Q_{\text{mean}}$  showed pertinent results for  $\Delta^{14}\text{C}$ -POC, applying SPM fluxes would improve the CA abilities to classify  $\Delta^{14}\text{C}$ -POC according to the Rhône River hydrology. For this purpose, other signatures are expected for the coming years.

## 6. Acknowledgements

This study was supported by l'Observatoire des Sédiments du Rhône (OSR), a multi-partner research program funded by European Development Fund (Feder) and the Plan Rhône.  $^{14}\text{C}$  and  $\delta^{13}\text{C}$  analyses were carried out by le Laboratoire des Sciences du Climat et de l'Environnement (LSCE), le Laboratoire de Mesure du Carbone 14 (LMC14) (Saclay, France), and the MOOSE programm (<https://www.moose-network.fr/fr/>). We strongly thank the INRAE colleagues for providing the data. We are grateful to our IRSN colleagues, Franck Giner and David Mourier who managed the SPM sampling at Arles and southern OSR

stations. We also thank LSCE/LMC14 lab. colleagues, Jean-Pascal Dumoulin, Ingrid Caffy, Marc Sieudat, Emmanuelle Delque-Kolic and Christophe Moreau for the sample storage, preparation, graphitization and  $^{14}\text{C}$  measurement.

## References

AFNOR, 2005. NF EN 872 : Water Quality - Determination of suspended solids - Method by filtration through glass fibre filters. 14 (In French). pp.10.

Bezdek, J.C., Ehrlich, R., Full, W., 1984. FCM: The fuzzy c-means clustering algorithm. Comput. Geosci. 10 (2-3), 191–203. [https://doi.org/10.1016/0098-3004\(84\)90020-7](https://doi.org/10.1016/0098-3004(84)90020-7)

Chaimontree, S., Atkinson, K., Coenen, F., 2010. Best Clustering Configuration Metrics: Towards Multiagent Based Clustering, in Advanced data mining and applications Part 1, Springer (2010), pp. 48-60.

Copard, Y., Eyrolle, F., Radakovitch, O., Poirel, A., Raimbault, P., Gairoard, S., Di-Giovanni, C., 2018. Badlands as a hot spot of petrogenic contribution to riverine particulate organic carbon to the Gulf of Lion (NW Mediterranean Sea). Earth Surf. Process. Landforms 43, 2495–2509. <https://doi.org/10.1002/esp.4409>

Cottureau, E., Arnold, M., Moreau, C., Baqué, D., Bavay, D., Caffy, I., Comby, C., Dumoulin, J.P., Hain, S., Perron, M., Salomon, J., Setti, V., 2007. Artemis, the new  $^{14}\text{C}$  AMS at LMC14 in Saclay, France. Radiocarbon 49, 291–299. <https://doi.org/10.1017/S0033822200042211>

Delaval, A., Duffa, C., Pairaud, I., Radakovitch, O., 2021. A fuzzy classification of the hydrodynamic forcings of the Rhone River plume: An application in case of accidental

552 release of radionuclides. Environ. Model. Softw. 140, 105005.  
 553 <https://doi.org/10.1016/j.envsoft.2021.105005>

554 Delile, H., Masson, M., Miège, C., Le Coz, J., Poulier, G., Le Bescond, C., Radakovitch, O.,  
 555 Coquery, M., 2020. Hydro-climatic drivers of land-based organic and inorganic  
 556 particulate micropollutant fluxes: The regime of the largest river water inflow of the  
 557 Mediterranean Sea. Water Res. 185, 116067.  
 558 <https://doi.org/10.1016/j.watres.2020.116067>

559 Ducros, L., 2018. Origine et devenir du tritium au sein des hydrosystèmes continentaux  
 560 méditerranéens français. (Doctoral dissertation). Aix-Marseille University, Marseille,  
 561 France.

562 Dumoulin, J.P., Comby-Zerbino, C., Delqué-Količ, E., Moreau, C., Caffy, I., Hain, S., Perron, M.,  
 563 Thellier, B., Setti, V., Berthier, B., Beck, L., 2017. Status Report on Sample Preparation  
 564 Protocols Developed at the LMC14 Laboratory, Saclay, France: From Sample Collection  
 565 to  $^{14}\text{C}$  AMS Measurement. Radiocarbon 59, 713–726.  
 566 <https://doi.org/10.1017/RDC.2016.116>

567 Eyrolle, F., Lepage, H., Antonelli, C., Morereau, A., Cossonnet, C., Boyer, P., Gurriaran, R.,  
 568 2020. Radionuclides in waters and suspended sediments in the Rhone River (France) -  
 569 Current contents, anthropic pressures and trajectories. Sci. Total Environ. 723.  
 570 <https://doi.org/10.1016/j.scitotenv.2020.137873>

571 Eyrolle, F., Lepage, H., Copard, Y., Ducros, L., Claval, D., Saey, L., Cossonnet, C., Giner, F.,  
 572 Mourier, D., 2018. A brief history of origins and contents of Organically Bound Tritium  
 573 (OBT) and  $^{14}\text{C}$  in the sediments of the Rhône watershed. Sci. Total Environ. 643, 40–51.  
 574 <https://doi.org/10.1016/j.scitotenv.2018.06.074>

Eyrolle, F., Radakovitch, O., Copard, Y., Lepage, H., Bodereau, N., Raimbault, P., Dabrin, A.,  
 Lagadec, V., Le Corre, C., n.d. Tritium and <sup>14</sup>C dependencies upon Particulate Organic  
 Matter within the nuclearized Rhone River (France). *Journal of Soil and Sediment*, in  
 review.

Ferraro, M.B., Giordani, P., Serafini, A., 2019. Fclust: An R package for fuzzy clustering. *R J.*  
 11, 1–18. <https://doi.org/10.32614/rj-2019-017>

Godwin, H., Willis, E.H., Hill, S., 1962. Natural Radiocarbon Measurements V. *Radiocarbon* 4,  
 57–70.

Graz, Y., Di-Giovanni, C., Copard, Y., Elie, M., Faure, P., Laggoun Defarge, F., Lévêque, J.,  
 Michels, R., Olivier, J.E., 2011. Occurrence of fossil organic matter in modern  
 environments: Optical, geochemical and isotopic evidence. *Appl. Geochemistry* 26,  
 1302–1314. <https://doi.org/10.1016/j.apgeochem.2011.05.004>

Hannah, D.M., Smith, B.P.G., Gurnell, A.M., McGregor, G.R., 2000. An approach to  
 hydrograph classification. *Hydrol. Process.* 14, 317–338.  
[https://doi.org/10.1002/\(SICI\)1099-1085\(20000215\)14:2<317::AID-HYP929>3.0.CO;2-T](https://doi.org/10.1002/(SICI)1099-1085(20000215)14:2<317::AID-HYP929>3.0.CO;2-T)

Higuera, M., Kerhervé, P., Sanchez-Vidal, A., Calafat, A., Ludwig, W., Verdoit-Jarraya, M.,  
 Heussner, S., Canals, M., 2014. Biogeochemical characterization of the riverine  
 particulate organic matter transferred to the NW Mediterranean Sea. *Biogeosciences*  
 11, 157–172. <https://doi.org/10.5194/bg-11-157-2014>

Horowitz, A.J., Elrick, K.A., Smith, J.J., 2001. Estimating suspended sediment and trace  
 element fluxes in large river basins: Methodological considerations as applied to the  
 NASQAN programme. *Hydrol. Process.* 15, 1107–1132. <https://doi.org/10.1002/hyp.206>



597 Husson, F., Josse, J., Pagès, J., 2010. L'analyse de données avec R - Complémentarité des  
598 Méthodes d'analyse factorielle et de classification. 42èmes Journées de Statistique,  
599 2010, Marseille, France, France. inria-00494779

600 Jain, A.K., Murty, M.N., Flynn, P.J., 1999. Data clustering: A review. *ACM Comput. Surv.* 31,  
601 264–323. <https://doi.org/10.1145/331499.331504>

602 Javed, A., Hamshaw, S.D., Lee, B.S., Rizzo, D.M., 2021. Multivariate event time series analysis  
603 using hydrological and suspended sediment data. *J. Hydrol.* 593, 125802.  
604 <https://doi.org/10.1016/j.jhydrol.2020.125802>

605 Jean-Baptiste, P., Fontugne, M., Fourré, E., Marang, L., Antonelli, C., Charmasson, S., Siclet,  
606 F., 2018. Tritium and radiocarbon levels in the Rhône river delta and along the French  
607 Mediterranean coastline. *J. Environ. Radioact.* 187, 53–64.  
608 <https://doi.org/10.1016/j.jenvrad.2018.01.031>

609 Kamel, M., Selim, S., 1994. New Algorithms for Solving the Fuzzy Clustering Problem. *Pattern*  
610 *Recognit.* 27, 421–428.

611 Kang, D., Lee, J., Kim, S., Noh, S., & Kang, S. (2019). Evaluation on  $^{14}\text{C}$  emission patterns of  
612 various PWR types in South Korea and effectiveness of inventory reduction driving  
613 (IRD). *Journal of Radioanalytical and Nuclear Chemistry*, 322(3), 1409–1415.  
614 <https://doi.org/10.1007/s10967-019-06805-1>

615 Kim, H.S., Kim, J.H., Ho, C.H., Chu, P.S., 2011. Pattern classification of typhoon tracks using  
616 the fuzzy c-means clustering method. *J. Clim.* 24, 488–508.  
617 <https://doi.org/10.1175/2010JCLI3751.1>

618 Klawonn, F., Höppner, F., 2003. What Is Fuzzy about Fuzzy Clustering? Understanding and

619 Improving the Concept of the Fuzzifier, in: *Advances in Intelligent Data Analysis V.* pp.  
620 254–264.

621 Launay, M., Dugué, V., Faure, J.B., Coquery, M., Camenen, B., Le Coz, J., 2019. Numerical  
622 modelling of the suspended particulate matter dynamics in a regulated river network.  
623 *Sci. Total Environ.* 665, 591–605. <https://doi.org/10.1016/j.scitotenv.2019.02.015>

624

625 Lepage, H., Gruat, A., Thollet, F., Le Coz, J., Coquery, M., Masson, M., Dabrin, A.,  
626 Radakovitch, O., Eyrolle, F., Labille, J., Ambrosi, J.-P., Delanghe, D., Raimbault, P., 2022.  
627 Concentrations and fluxes of suspended particulate matters and associated  
628 contaminants in the Rhône River from Lake Geneva to the Mediterranean Sea. *Earth*  
629 *Syst. Sci. Data* 7, 1–22. <https://doi.org/10.5194/essd-2021-350>

630

631 Leithold, E.L., Blair, N.E., Wegmann, K.W., 2016. Source-to-sink sedimentary systems and  
632 global carbon burial: A river runs through it. *Earth-Science Rev.* 153, 30–42.  
633 <https://doi.org/10.1016/j.earscirev.2015.10.011>

634 Masson, M., Angot, H., Le Bescond, C., Launay, M., Dabrin, A., Miège, C., Le Coz, J., Coquery,  
635 M., 2018. Sampling of suspended particulate matter using particle traps in the Rhône  
636 River: Relevance and representativeness for the monitoring of contaminants. *Sci. Total*  
637 *Environ.* 637–638, 538–549. <https://doi.org/10.1016/j.scitotenv.2018.04.343>

638 Meybeck, M., 2003. Global analysis of river systems: From Earth system controls to  
639 Anthropocene syndromes. *Philos. Trans. R. Soc. B Biol. Sci.* 358, 1935–1955.  
640 <https://doi.org/10.1098/rstb.2003.1379>

641 Meybeck, M., 1982. Carbon, nitrogen, and phosphorus transport by world rivers. *Am. J. Sci.*  
642 <https://doi.org/10.2475/ajs.282.4.401>

643 Meyer, D., Dimitriadou, E., Hornik, K., Weingessel, A., Leisch, F., Chang, C.-C., Lin, C.-C., 2021.  
644 Package 'e1071' : Misc Functions of the Department of Statistics, Probability Theory  
645 Group (Formerly: E1071), TU Wien Imports.

646 Milliman, J.D., Farnsworth, K.L., 2011. River discharge to the coastal ocean: A global  
647 synthesis, *River Discharge to the Coastal Ocean: A Global Synthesis*.  
648 <https://doi.org/10.1017/CBO9780511781247>

649 Misset, C., Recking, A., Legout, C., Valsangkar, N., Bodereau, N., Zanker, S., Poirel, A.,  
650 Borgniet, L., 2019. The Dynamics of Suspended Sediment in a Typical Alpine Alluvial  
651 River Reach: Insight From a Seasonal Survey. *Water Resour. Res.* 55, 10918–10934.  
652 <https://doi.org/10.1029/2019WR025222>

653 Mourier, B., Desmet, M., Van Metre, P.C., Mahler, B.J., Perrodin, Y., Roux, G., Bedell, J.P.,  
654 Lefèvre, I., Babut, M., 2014. Historical records, sources, and spatial trends of PCBs along  
655 the Rhône River (France). *Sci. Total Environ.* 476–477, 568–576.  
656 <https://doi.org/10.1016/j.scitotenv.2014.01.026>

657 Olivier, J.M., Carrel, G., Lamouroux, N., Dole-Olivier, M.J., Malard, F., Bravard, J.P., Amoros,  
658 C., 2009. The Rhône River Basin, *Rivers of Europe*. [https://doi.org/10.1016/B978-0-12-](https://doi.org/10.1016/B978-0-12-369449-2.00007-2)  
659 [369449-2.00007-2](https://doi.org/10.1016/B978-0-12-369449-2.00007-2)

660 Ollivier, P., Radakovitch, O., Hamelin, B., 2011. Major and trace element partition and fluxes  
661 in the Rhône River. *Chem. Geol.* 285, 15–31.  
662 <https://doi.org/10.1016/j.chemgeo.2011.02.011>

663 Ouyang, R., Ren, L., Cheng, W., Zhou, C., 2010. Similarity search and pattern discovery in  
 664 hydrological time series data mining. *Hydrol. Process.* 24, 1198–1210.  
 665 <https://doi.org/10.1002/hyp.7583>

666 Phillips, J.M., Russell, M.A., Walling, D.E., 2000. Time-integrated sampling of fluvial  
 667 suspended sediment: A simple methodology for small catchments. *Hydrol. Process.* 14,  
 668 2589–2602. [https://doi.org/10.1002/1099-1085\(20001015\)14:14<2589::AID-](https://doi.org/10.1002/1099-1085(20001015)14:14<2589::AID-HYP94>3.0.CO;2-D)  
 669 [HYP94>3.0.CO;2-D](https://doi.org/10.1002/1099-1085(20001015)14:14<2589::AID-HYP94>3.0.CO;2-D)

670 Pont, D., Simonnet, J.-P., Walter, A.V., 2002. Medium-term Changes in Suspended Sediment  
 671 Delivery to the Ocean: Consequences of Catchment Heterogeneity and River  
 672 Management (Rhône River, France). *Estuar. Coast. Shelf Sci.* 54, 1–18.  
 673 <https://doi.org/10.1006/ecss.2001.0829>

674 Poulier, G., Launay, M., Le Bescond, C., Thollet, F., Coquery, M., Le Coz, J., 2019. Combining  
 675 flux monitoring and data reconstruction to establish annual budgets of suspended  
 676 particulate matter, mercury and PCB in the Rhône River from Lake Geneva to the  
 677 Mediterranean Sea. *Sci. Total Environ.* 658, 457–473.  
 678 <https://doi.org/10.1016/j.scitotenv.2018.12.075>

679 Radakovitch, O., Roussiez, V., Ollivier, P., Ludwig, W., Grenz, C., Probst, J.L., 2008. Input of  
 680 particulate heavy metals from rivers and associated sedimentary deposits on the Gulf of  
 681 Lion continental shelf. *Estuar. Coast. Shelf Sci.* 77, 285–295.  
 682 <https://doi.org/10.1016/j.ecss.2007.09.028>

683 Raimbault P., Neveux J., Lantoiné F., 2004. Dosage rapide de la chlorophylle a et des  
 684 phaeopigments a par fluorimétrie après extraction au méthanol. Comparaison avec la  
 685 méthode classique d'extraction à l'acétone. *Oceanis*, 30(2) : 189-205.

686 Raimbault, P., Garcia, N., Cerutti, F., 2008. Distribution of inorganic and organic nutrients in  
687 the South Pacific Ocean – evidence for long-term accumulation of organic matter in  
688 nitrogen-depleted waters. *Biogeosciences* 5, 281–298. [https://doi.org/10.5194/bg-5-](https://doi.org/10.5194/bg-5-281-2008)  
689 281-2008

690 Ramze Rezaee, M., Lelieveldt, B.P.F., Reiber, J.H.C., 1998. A new cluster validity index for the  
691 fuzzy c-mean. *Pattern Recognit. Lett.* 19, 237–246. [https://doi.org/10.1016/S0167-](https://doi.org/10.1016/S0167-8655(97)00168-2)  
692 [8655\(97\)00168-2](https://doi.org/10.1016/S0167-8655(97)00168-2)

693 Raux, J., Copard, Y., Laignel, B., Fournier, M., Masseï, N., 2011. Classification of worldwide  
694 drainage basins through the multivariate analysis of variables controlling their  
695 hydrosedimentary response. *Glob. Planet. Change* 76, 117–127.  
696 <https://doi.org/10.1016/j.gloplacha.2010.12.005>

697 Raymond, P.A., Bauer, J.E., 2001. Use of <sup>14</sup>C and <sup>13</sup>C natural abundances for evaluating  
698 riverine, estuarine, and coastal DOC and POC sources and cycling: A review and  
699 synthesis. *Org. Geochem.* 32, 469–485. [https://doi.org/10.1016/S0146-6380\(00\)00190-](https://doi.org/10.1016/S0146-6380(00)00190-X)  
700 X

701 Sadaoui, M., Ludwig, W., Bourrin, F., Raimbault, P., 2016. Controls, budgets and variability  
702 of riverine sediment fluxes to the Gulf of Lions (NW Mediterranean Sea). *J. Hydrol.* 540,  
703 1002–1015. <https://doi.org/10.1016/j.jhydrol.2016.07.012>

704 Schulze, T., Ricking, M., Schröter-Kermani, C., Körner, A., Denner, H.-D., Weinfurtner, K.,  
705 Winkler, A., Pekdeger, A., 2007. The German Environmental Specimen Bank - Sampling,  
706 processing, and archiving sediment and suspended particulate matter. *J. Soils*  
707 *Sediments* 7, 361–367. <https://doi.org/10.1065/jss2007.08.248>

708 Schwämmle, V., Jensen, O.N., 2010. A simple and fast method to determine the parameters  
 709 for fuzzy c-means cluster analysis. *Bioinformatics* 26, 2841–2848.  
 710 <https://doi.org/10.1093/bioinformatics/btq534>

711 Stuiver, M., Polach, H.-A., 1977. Reporting of <sup>14</sup>C data. *Radiocarbon* 19(3), 355-363.  
 712 <https://doi.org/10.1017/S0033822200003672>

713 Thollet, F., Le Bescond, C., Lagouy, M., Gruat A., Grisot, G., Le Coz, J., Coquery, M., Lepage,  
 714 H., Gairoard, S., Gattacceca, J.C., Ambrosi, J.-P., Radakovitch, O., Dur, G., Richard, L.,  
 715 Giner, F., Eyrolle, F., Angot, H., Mourier, D., Bonnefoy, A., Dugué, V., Launay, M.,  
 716 Troudet, L., Labille, J., Kieffer, L. (2021): Développement de l'application informatique  
 717 BDOH pour l'Observatoire des Sédiments du Rhône [Research report] INRAE. 2021.  
 718 <https://dx.doi.org/10.17180/OBS.OSR>

719

720 Trumbore, S.E., Sierra, C.A., Hick Pries, C.E., 2016. Radiocarbon Nomenclature, Theory,  
 721 Models, and Interpretation : Measuring Age,determining cycling rates, and Tracing  
 722 Source Roles, in: Schuur, E.A.G., Druffel, E.R.M., Trumbore, S.E. (Eds.), Radiocarbon and  
 723 Climate Change, Mechanisms, Applications and Laboratory Techniques. Springer  
 724 International Publishing, pp. 45–82.

725 Voegel, J.S., Southon, J.R., Nelson, D.E., Brown, T.A., 1984. Performance of catalytically  
 726 condensed carbon for use in Accelerator Mass Spectrometry. *Nucl. Instruments*  
 727 *Methods Phys. Res.* 5, 289–293.

728 Walling, D.E., Owens, P.N., Carter, J., Leeks, G.J.L., Lewis, S., Meharg, A.A., Wright, J., 2003.  
 729 Storage of sediment-associated nutrients and contaminants in river channel and

floodplain systems. *Appl. Geochemistry* 18, 195–220. [https://doi.org/10.1016/S0883-2927\(02\)00121-X](https://doi.org/10.1016/S0883-2927(02)00121-X)

Zebracki, M., Eyrolle-Boyer, F., Evrard, O., Claval, D., Mourier, D., Gairoard, S., Cagnat, X., Antonelli, C., 2015. Tracing the origin of suspended sediment in a large Mediterranean river by combining continuous river monitoring and measurement of artificial and natural radionuclides. *Sci. Total Environ.* 502, 122–213.

