Electronic Supplementary Material

Article: Ectophagous folivores do not profit from rich resources on phylogenetically isolated trees

Journal: Oecologia

Authors: Soumen Mallick, Freerk Molleman, Benjamin Yguel, Richard Bailey, Jörg Müller, Frédéric Jean, and Andreas Prinzing

Corresponding author: Soumen Mallick, Centre National de la Recherche Scientifique, Université de Rennes 1, Research Unit UMR 6553, Ecosystèmes Biodiversité Evolution (ECOBIO), Campus de Beaulieu, 35042 Rennes, France, soumen.mallick@univ-rennes1.fr Online Resource 1. Phylogenetic distances and calculation of phylogenetic isolation

Online Resource 2. Effects of tree pairs on resource use

- **Online Resource 3.** Influence of crown position and budburst phenology on the effect of phylogenetic isolation on proportional use of resources
- **Online Resource 4.** Distribution of phylogenetic isolation, distance to the nearest oak and oak circumference
- **Online Resource 5.** Community composition and species traits of ectophages
- **Online Resource 6.** Role of different tree traits and leaf traits to shape species composition in ectophage communities
- **Online Resource 7.** Effects of distance to the nearest oak, oak circumference and percentage of pines in the neighbourhood on proportional use of resources
- Online Resource 8. Logistic regression of presence/absence of species with wingless females

Online Resource 1. Phylogenetic distances and calculation of phylogenetic isolation

Phylogenetic distance between two tree species is the estimated time (in MYBP) since the evolutionary establishment of the clades of a given neighbouring tree species and of oaks. These phylogenetic distances were taken from Vialatte et al. (2010) and Yguel et al. (2011), based on Magallon et al., 1999, Manos et al., 1999, Wikström et al., 2001, Poinar et al., 2007. Note that this is not the most recent common ancestor, as this would give gymnosperms such as pines an extreme weight (Savard et al., 1994), and would essentially render our parameter a simple percentage of pines in the surroundings of the oaks. Rather, this is the age when both sister clades had established their particular characteristics as hosts for insects (i.e. phylogenetic crown-age of the younger of the two lineages and not stem age). Thus, phylogenetic distance corresponds to the smaller of the two crown ages of the two lineages involved (i.e. of oak and of the other tree species) at the corresponding phylogenetic rank (ranks within Angiosperms inferred from APG 2003, 2009, and checked against 2016).

Table S1: Phylogenetic d	istance between or	aks and otl	her trees in	our data set	based on	crown	ages
(taken from Vialatte et al.	. 2010 and Yguel e	et al. 2011)).				

Species		Phyloge	netic rank	of separation v	with oak		Distance (MY)
Chamaecyparis sp.	Spermatophytes	-	-	-	-	-	140
Pinus sylvestris	Spermatophytes	-	-	-	-	-	140
Abies sp.	Spermatophytes	-	-	-	-	-	140
<i>Ilex</i> sp.	Angiosperms	Asterids	-	-	-	-	128
<i>Tilia</i> sp.	Angiosperms	Rosids	Malvids	-	-	-	89.5
Salix caprea	Angiosperms	Rosids	Fabids	Malpighiales	-	-	68
Populus tremula	Angiosperms	Rosids	Fabids	Malpighiales	-	-	68
Rhamnus sp.	Angiosperms	Rosids	Fabids	Rosales	-	-	58.5
Prunus sp.	Angiosperms	Rosids	Fabids	Rosales	-	-	58.5
Sorbus sp	Angiosperms	Rosids	Fabids	Rosales	-	-	58.5
Pyrus sp.	Angiosperms	Rosids	Fabids	Rosales	-	-	58.5
Malus sp.	Angiosperms	Rosids	Fabids	Rosales	-	-	58.5
Ulmus minor	Angiosperms	Rosids	Fabids	Rosales	-	-	58.5
Alnus glutinosa	Angiosperms	Rosids	Fabids	Fagales	Betulaceae	-	54
Corylus avellana	Angiosperms	Rosids	Fabids	Fagales	Betulaceae	-	54
<i>Betula</i> sp.	Angiosperms	Rosids	Fabids	Fagales	Betulaceae	-	54
Carpinus betulus	Angiosperms	Rosids	Fabids	Fagales	Betulaceae	-	54
Fagus sylvatica	Angiosperms	Rosids	Fabids	Fagales	Fagaceae	Fagus	40
Castanea sativa	Angiosperms	Rosids	Fabids	Fagales	Fagaceae	Castanea	40

Degree of phylogenetic isolation of a tree was calculated as mean phylogenetic distances to the trees with which its crown was in contact (or the projections of crowns of trees on the ground were in contact). Hence, the formula to calculate the phylogenetic isolation of an oak is Phylogenetic isolation = $\frac{\sum_{l=1}^{n} Phylogenetic distance}{\sum_{l=1}^{n} Phylogenetic distance}$; where *n* is the number of trees in crown contact of п that oak. As an example, the case given in the figure below is a Quercus petraea surrounded by 3 Pinus sylvestris and 3 Fagus sylvatica. So, the total number of trees in crown contact with the oak is 6. Phylogenetic distance between a Quercus petraea and a Pinus sylvestris is 140 MY, and between a Quercus petraea and a Fagus sylvatica is 40 MY. Hence, the phylogenetic isolation of this oak will be $\frac{140+140+140+40+40+40}{140+140+40+40} = 90 \text{ MY}$. 6



References:

Angiosperm Phylogeny Group (2003) An update of the Angiosperm phylogeny group classification for the orders and families of flowering plants: APG II. Bot J Linn Soc 141: 399–436.

Angiosperm Phylogeny Group (2009) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III", Bot J Linn Soc 161: 105–121

Angiosperm Phylogeny Group (2016) An update of the Angiosperm phylogeny group classification for the orders and families of flowering plants: APG IV. Bot J Linn Soc 181: 1–20.

Magallon S, Crabe PR, Herendeen PS (1999) Phylogenetic pattern, diversity, and diversification of eudicots. Ann Missouri Bot Garden 86: 1407–1419.

Manos PS, Doyle JJ, Nixon KC (1999) Phylogeny, biogeography, and processes of molecular differentiation in *Quercus* subgenus *Quercus* (Fagaceae). Mol Phyl Evol 12: 333–349.

Poinar G, Chambers KL, Buckley R (2007) *Eoepigynia burmensis* gen. and sp. nov., an early Cretaceous Eudicot flower (Angiospermae) in Burmese Amber. J Bot Res Inst Texas 1: 91–96.

Savard L, Li P, Strauss SH, Chase MW, Michaud M, Bousquet J (1994). Chloroplast and nuckear genessequences indicate late Pennsylvanian time for the last common ancestor of extant seed plants. Proc Natl Acad Sci USA 91: 5163-5167.

Vialatte A, Bailey RI, Vasseur C, Matocq A, Gossner MM, Everhart D, Vitrac X, Belhadj A, Ernoult A & Prinzing A (2010) Phylogenetic isolation of host trees affects assembly of local Heteroptera communities. Proc Biol Sci 277: 2227-2236.

Wikström, N, Savolainen, V & Chase, M W (2001) Evolution of the angiosperms: calibrating the family tree. Proc R Soc. Lond. B 268, 2211–2220.

Yguel B, Bailey R, Tosh N, et al (2011) Phytophagy on phylogenetically isolated trees: Why hosts should escape their relatives. Ecol Lett 14:1117–24. https://doi.org/10.1111/j.1461-0248.2011.01680.x

Online Resource 2. Effects of tree pairs on resource use

Species composition and abundance of insects may vary spatially in a forest, so that more proximate trees have more similar insect communities. Therefore, we selected trees that differ in phylogenetic isolation in pairs of spatially proximate trees, rather than selecting the trees randomly (Legendre et al. 2004). To test whether the tree pair defines the various level of use of resources by ectophages, leaf gallers, leaf miners and parasitoids of ectophagous caterpillars, we performed an ANOVA. We found that pair did not significantly affect folivore abundance or folivory (Table S2). So, we do not account for "pair" in further analyses.

Group	Dependent variable	Year	DF	F	р
Ectophages	Per-leaf folivory	2006	8	1.574	0.256
		2010	10	0.378	0.932
		2011	10	0.381	0.930
	Per-individual folivory	2010	10	0.873	0.581
		2011	10	1.390	0.298
	Per-leaf abundance	2010	10	0.214	0.989
	v	2011	10	0.986	0.505
Leaf gallers	Per-leaf abundance	2006	8	1.239	0.376
e	0	2010	10	1.472	0.267
		2011	10	1.584	0.231
Leaf miners	Per-leaf abundance	2010	10	1.033	0.475
	5	2011	10	0.719	0.695
Parasitoids	Parasitised caterpillar	2010	10	0.699	0.710
	Ĩ	2011	10	0.756	0.667

Table S2. Results of ANOVA testing for the effect of tree pair on various level of use of resources by ectophages, leaf gallers, leaf miners and parasitoids of ectophagous caterpillars.

References:

Legendre, P., Dale, M.R.T., Fortin, M.J., Casgrain, P. & Gurevitch, J. (2004). Effects of spatial structures on the results of field experiments. Ecology, 85, 3202–3214.

Online Resource 3. Influence of crown position and budburst phenology on the effect of phylogenetic isolation on proportional use of resources

Crown position and budburst phenology did not change the overall effect of phylogenetic isolation on proportional use of resources. After accounting for crown position and budburst phenology, phylogenetic isolation still statistically reduced the tracking of leaf size and not leaf quality by ectophages. Specifically, for ectophages, tracking of leaf size was still reduced at *Perleaf folivory* level in all three years, and at *Per-individual folivory* level in 2011 but not in 2010 - only the signal at the level of *Per-leaf abundance* was lost. In addition, for ectophages, phylogenetic isolation still consistently did not affect the proportional use of leaf quality. For leaf gallers and leaf miners, phylogenetic isolation still did not consistently affect the proportional use of resources. Moreover, for parasitoids, phylogenetic isolation still affected the proportional use of caterpillars in 2010 but not in 2011. Overall, the models accounting for the effects of crown position and budburst phenology (Table S3) support the results of the models that do not account for these co-variables (Table 2).

Table S3. Statistics of the multiple linear regressions that accounted for the effects of crown position (in 2006) and budburst phenology (in 2010 and 2011). Note that the statistics given in the table is of only the interaction term in a model, but the models also account for the main effects of the variables in the interaction term and the variable given in the column 'Accounted for'. LA = leaf area, LDMC = leaf dry-matter content, C:N = carbon-nitrogen ratio, LPC = leaf phenolics content, TC = total caterpillars, PI = phylogenetic isolation, CP = crown position, BB = budburst.

Group	Use of	Dependent variable	Effects of	Year	Accounted for	t	р	DF
Ectophages	Leaf size	Per-leaf folivory	$LA \times PI$	2006	СР	-2.495	0.028	12
			$LA \times PI$	2010	BB	-3.315	0.004	17
			$LA \times PI$	2011	BB	-2.193	0.043	16
		Per-individual	$LA \times PI$	2010	BB	-0.956	0.352	17
		folivory	$LA \times PI$	2011	BB	-2.518	0.023	16
		Per-leaf	$LA \times PI$	2010	BB	-1.166	0.260	17
		abundance	$LA \times PI$	2011	BB	1.490	0.154	17
	Leaf quality	Per-leaf folivory	$LDMC \times PI$	2006	CP	0.435	0.672	11
			$LDMC \times PI$	2010	BB	0.739	0.470	17
			C:N imes PI	2006	CP	-1.408	0.187	11
			C:N imes PI	2010	BB	1.011	0.326	17
			$LPC \times PI$	2006	CP	0.165	0.872	13
			$LPC \times PI$	2010	BB	1.352	0.194	17
		Per-individual	$LDMC \times PI$	2010	BB	-0.721	0.481	17
		folivory	$C:N \times PI$	2010	BB	-0.460	0.651	17
			$LPC \times PI$	2010	BB	-0.300	0.768	17
		Per-leaf	$LDMC \times PI$	2010	BB	1.171	0.259	16
		abundance	$C:N \times PI$	2010	BB	-0.120	0.906	16
			$LPC \times PI$	2010	BB	1.278	0.219	16
Leaf	Leaf size	Per-leaf	$LA \times PI$	2006	CP	-0.512	0.617	13
gallers		abundance	$LA \times PI$	2010	BB	-0.379	0.709	16
			$LA \times PI$	2011	BB	-1.579	0.133	17
	Leaf quality	Per-leaf	$LDMC \times PI$	2006	CP	0.763	0.459	13
	1 4	abundance	$LDMC \times PI$	2010	BB	1.610	0.127	16
			$C:N \times PI$	2006	CP	0.459	0.654	13
			C:N imes PI	2010	BB	5.419	<0.001	16
			$LPC \times PI$	2006	CP	1.824	0.093	12
			$LPC \times PI$	2010	BB	0.980	0.342	16
Leaf	Leaf size	Per-leaf	$\mathrm{LA} \times \mathrm{PI}$	2010	BB	-0.694	0.498	16
miners		abundance	$LA \times PI$	2011	BB	-0.983	0.339	17
	Leaf quality	Per-leaf	$LDMC \times PI$	2010	BB	0.671	0.512	16
		abundance	$C:N \times PI$	2010	BB	1.238	0.233	17
			$LPC \times PI$	2010	BB	2.096	0.052	17
Parasitoids	Caterpillars	Parasitised	$TC \times PI$	2010	BB	-2.173	0.044	17
	-	caterpillars	$\mathrm{TC} \times \mathrm{PI}$	2011	BB	-0.003	0.998	17

Online Resource 4. Distribution of phylogenetic isolation, distance to the nearest oak and oak circumference

In our data, the phylogenetic isolation of individual trees, their distance to the nearest oak and oak circumference had continuous distributions (Fig. S4), and were treated as such in our analyses. Phylogenetic isolation has a binary representation in the illustrations in the main body of the article (Fig. 1, 2, 3 and 4) only to permit an easier interpretation.



Fig. S4: The distribution of (a) phylogenetic isolation, (b) distance to the nearest oak, and (c) oak circumference for First Set (studied in 2006) and Second Set of trees (studied in 2010 and 2011). The phylogenetic isolation ranged from 5.71 to 106.67 million years in First Set, and from 10 to 125.67 million years in Second Set of trees. The spatial distance from the study trees to the nearest adult oak ranged from 2.5 to 18.9 m in First Set and from 0.70 to 9.40 m in Second Set of trees. The circumference of the study trees was measured at breast height, and ranged from 57.75 to 133.1 cm in First Set, and from 37.8 to 91.4 cm in Second Set of trees.

Online Resource 5. Community composition and species traits of ectophages

Year	Tree	Acrobasis consociella	Acrobasis repandana	Aglia tau	Agriopis aurantaria	Aleimma loeflingiana	Archips Podana	Archips sp	Biston strataria	Carcina quercana	Catocala promissa	Catocala sponsa	Conistra sp	Cyclophora linearia	Ennomos quercinaria	Epirrita autumnata	Erannis defoliaria	Eupithecia abbreviata	Eupithecia sp	Eupsilia transversa	Hedya nubifera	Lymantria dispar	Lymantria monacha	Operophtera brumata	Orgyia antiqua	Orthosia cerasi	Orthosia incerta	Orthosia munda	Pammene argyrana	Phycita roborella	Poecilocampa populi	Polyploca ridens	Spilonota ocellana	Tortrix viridana	Undetermined	Ypsolopha parenthesella	Ypsolopha ustella	Zeiraphera isertana
2010	2.1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	0	0	0	0	0	0	4	5	1	0	2
	2.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0
	4.1	0	0	0	1	0	0	4	0	2	0	0	0	0	0	0	1	0	0	1	1	0	0	0	0	2	0	0	0	0	0	0	0	1	1	0	0	1
	4.2	0	0	0	0	0	1	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	1	0	1
	5.1	0	0	0	0	0	0	4	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	2	0	0	0	0	0	0	0	2	3	1	1	1
	5.2	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	1	0	0	2	0	0	0	0	0	0	0	0	0	0	1	0	3	1	1	0	0
	6.1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	3	0	0	0	0	0	0	0	0	3	1	0	0
	6.2	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0
	7.1	0	2	0	0	1	0	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	2	0	1	0	0	0	0	0	0	0	0	2	0	0	0
	7.2	0	0	0	0	0	0	2	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	1	1
	8.1	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	1	1	0	0	0	0	1	0	0	0	0	0	0	0	2	2	0	0	0
	8.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0
	9.1	0	0	0	1	0	0	6	1	0	0	0	3	0	0	0	0	0	0	0	1	0	1	0	0	1	0	0	0	0	0	1	0	0	2	0	0	0
	9.2	0	0	0	0	0	0	1	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0
	10	0	0	0	1	0	1	8	0	0	0	0	1	0	0	0	0	0	0	0	0	0	1	1	0	1	0	0	0	0	0	0	0	1	0	1	0	0
	10	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	2	0	0	0	0	2	0	0	0	1	0	0	1	1	1	0	0	1
	13	0	1	0	2	0	0	1	0	0	0	0	0	0	1	0	0	0	0	0	1	0	1	0	0	2	0	0	0	0	0	0	0	4	5	0	0	0
	14	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	3	1	0	0
	14	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
	X 1	0	1	1	0	0	0	ő	ő	0	ő	0	ő	0	0	ő	0	0	0	õ	4	0	0	ő	1	0	0	ő	0	ő	0	0	0	6	2	0	0	1
	X.2	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	õ	0	0	0	0	0	õ	0	0	0	0	0	0	0	0
2011	2.1	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0
	2.2	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1
	4.1	0	0	0	0	0	0	0	0	3	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	2	3	0	0	0
	4.2	0	0	0	1	0	0	0	0	2	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0
	5.1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	1
	5.2	2	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	0	1	0	0	0	0	1	0	0	1	0	0	0	0	1	5	2	0	1
	6.1	2	0	0	0	0	0	2	0	2	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0
	6.2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0
	7.1	0	0	0	0	0	0	2	1	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	5	0	0	0
	7.2	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	0	0	1	0	0	0	0	0	0	0	0	2	0	0	0
	8.1	0	0	1	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	2	0	0	1	0	0	0	0	0	0	1	0	0
	8.2	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	1	1	0	0	0	1	0	0	0	0	0	0	0	0	2	1	0	0
	9.1	0	0	0	0	0	0	3	0	3	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	1	5	0	0	2
	9.2	1	0	0	1	0	0	1	0	2	0	0	0	0	0	0	0	0	1	0	1	1	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0
	10	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	4	0	0	1	0	0	0	1	0	1	2	0	0
	10	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	1	3	0	0	1
	13	0	0	0	0	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	2	7	0	0	0
	13	1	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	3	2	0	0	0
	14	0	0	0	0	0	0	3	0	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	3	1	0	0	0	0	0	0	0	9	0	0	0
	14 V 1	1	0	0	0	0	0	2	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	3	0	0	0
	л.1 х 2	1	0	0	0	0	0	1	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	2	5	0	0	0
	X .2	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0	0

Table S5A: Community composition of ectophages

Table S5B: Traits of ectophage species that are present in the community. Information are taken from the websites Catalogue of the Lepidoptera of Belgium (De Prins and Steeman 2021), and Guide to the moths of Great Britain and Ireland (Kimber 2021).

Species	Average	Host-plant	Fly between	Wingless
species	wingspan (mm)	specialisation	June-October	females
Acrobasis consociella	20.5	Specialist	Yes	No
Acrobasis repandana	22.5	Specialist	Yes	No
Aglia tau	72.0	Generalist	No	No
Agriopis aurantaria	31.0	Generalist	No	Yes
Aleimma loeflingiana	16.5	Generalist	Yes	No
Archips Podana	22.0	Specialist	Yes	No
Archips sp	20.8	Specialist	Yes	No
Bena bicolorana	42.5	Specialist	Yes	No
Carcina quercana	19.0	Generalist	Yes	No
Catocala promissa	62.5	Specialist	Yes	No
Catocala sponsa	65.0	Specialist	Yes	No
Conistra sp	33.9	Generalist	No	No
Cyclophora linearia	29.5	Specialist	Yes	No
Ennomos quercinaria	46.0	Generalist	Yes	No
Epirrita autumnata	42.0	Generalist	Yes	No
Erannis defoliaria	35.0	Generalist	No	Yes
Eupithecia abbreviata	20.5	Generalist	No	No
Eupithecia sp	17.5	Generalist	No	No
Eupsilia transversa	37.0	Generalist	No	No
Hedya nubifera	18.0	Generalist	Yes	No
Lymantria dispar	43.5	Generalist	Yes	No
Lymantria monacha	40.0	Generalist	Yes	No
Operophtera brumata	25.0	Generalist	No	Yes
Orgyia antiqua	27.5	Generalist	Yes	Yes
Orthosia cerasi	37.0	Generalist	No	No
Orthosia incerta	37.5	Generalist	No	No
Orthosia munda	41.0	Generalist	No	No
Pammene argyrana	11.5	Specialist	No	No
Phycita roborella	26.5	Specialist	Yes	No
Poecilocampa populi	37.5	Generalist	No	No
Polyploca ridens	32.5	Specialist	No	No
Spilonota ocellana	14.5	Generalist	Yes	No
Tortrix viridana	20.5	Specialist	Yes	No
Ypsolopha parenthesel	18.0	Generalist	Yes	No
Ypsolopha ustella	17.5	Specialist	No	No
Zeiraphera isertana	17.0	Specialist	Yes	No

References:

De Prins W, Steeman C (2021) Catalogue of the Lepidoptera of Belgium.

https://projects.biodiversity.be/lepidoptera/. Accessed 24 Feb 2022

Kimber I (2021) UKMoths | Guide to the moths of Great Britain and Ireland. https://ukmoths.org.uk/. Accessed 24 Feb 2022 **Online Resource 6.** Role of different tree traits and leaf traits to shape species composition in ectophage communities

In both the years 2010 and 2011, phylogenetic isolation has contributed the most to shape the species composition in ectophage communities on individual trees, albeit only significantly in 2010.

Table S6: Contributions of different tree traits and leaf traits to shape the species composition of ectophagous communities on individual trees. Tested by a Permutational Multivariate Analysis of Variance (PERMANOVA) using the R package *vegan*. PI = phylogenetic isolation, DIS = Distance to nearest oak, TC = Tree circumference, BB = budburst, LA = leaf area, LDMC = leaf dry-matter content, C:N = carbon-nitrogen ratio, LPC = leaf phenolics content

Effects of	Year:			201	0			2011								
Effects of		Df	SumsOfSqs	MeanSqs	F.Model	R2	Pr(>F)	Df	SumsOfSqs	MeanSqs	F.Model	R2	Pr(>F)			
PI		1	0.537	0.537	1.934	0.089	0.040	1	0.308	0.308	1.363	0.064	0.213			
DIS		1	0.444	0.444	1.600	0.073	0.110	1	0.273	0.273	1.208	0.057	0.260			
TC		1	0.281	0.281	1.011	0.046	0.422	1	0.179	0.179	0.793	0.037	0.616			
BB		1	0.193	0.193	0.695	0.032	0.753	1	0.203	0.203	0.898	0.042	0.549			
LA		1	0.134	0.134	0.481	0.022	0.893	1	0.250	0.250	1.108	0.052	0.374			
LDMC		1	0.283	0.283	1.018	0.047	0.408	NA	NA	NA	NA	NA	NA			
C:N		1	0.177	0.177	0.638	0.029	0.781	NA	NA	NA	NA	NA	NA			
LPC		1	0.397	0.397	1.429	0.066	0.158	NA	NA	NA	NA	NA	NA			
Residuals		13	3.611	0.278	0.596			16	3.616	0.226	0.749					
Total		21	6.058	1.000				21	4.830	1.000						

The figure represents the ordination plots of Nonmetric Multidimensional Scaling (NMDS), where datapoints are individual trees, and the arrows represent a tree trait or a leaf trait.



Online Resource 7. Effects of distance to the nearest oak, oak circumference and percentage pines in the neighbourhood on proportional use of resources

(A) Distance to the nearest oak and oak circumference: We compared effects of phylogenetic isolation of individual host trees on the proportional use of resources to that of two more commonly studied variables, isolation in space and availability of the resource in time. Isolation in space was measured as the spatial distance of the study oaks to its nearest adult oak. We used circumference of the tree at breast height as a proxy of the age of a tree. Neither the distance to the nearest oak (Table S7A.a), nor the circumference of the trees (Table S7A.b) showed consistent significant effect on proportional use of resources. Note that the statistics given in the table is of only the interaction term in a model, but the models also accounted for the main effects of the variables in the interaction term. LA = leaf area, LDMC = leaf dry-matter content, C:N = carbon-nitrogen ratio, LPC = leaf phenolics content, TC = total caterpillars, DIS = distance to the nearest oak, OC = oak circumference.

Group	Use of	Dependent variable	Effects of	Year	t	р	DF
Ectophages	Leaf size	Per-leaf folivory	$LA \times DIS$	2006	1.532	0.148	14
			$LA \times DIS$	2010	-1.010	0.326	18
			$LA \times DIS$	2011	-1.039	0.312	18
		Per-individual folivory	$LA \times DIS$	2010	0.487	0.633	17
			$LA \times DIS$	2011	-0.887	0.387	18
		Per-leaf abundance	$LA \times DIS$	2010	-0.778	0.447	18
		·	$LA \times DIS$	2011	0.713	0.485	18
	Leaf	Per-leaf folivory	$LDMC \times DIS$	2006	0.569	0.579	13
	quality		$LDMC \times DIS$	2010	0.143	0.888	17
			$C:N \times DIS$	2006	-0.588	0.567	13
			$C:N \times DIS$	2010	1.049	0.310	16
			$LPC \times DIS$	2006	0.166	0.871	14
			$LPC \times DIS$	2010	-0.921	0.369	18
		Per-individual folivory	$LDMC \times DIS$	2010	-2.672	0.016	18
			$C:N \times DIS$	2010	1.893	0.075	18
			$LPC \times DIS$	2010	Inva	lid mode	1
		Per-leaf abundance	$LDMC \times DIS$	2010	1.371	0.188	18
			$C:N \times DIS$	2010	-1.227	0.236	18
			$LPC \times DIS$	2010	0.882	0.389	18
Leaf gallers	Leaf size	Per-leaf abundance	$LA \times DIS$	2006	-2.783	0.016	13
			$LA \times DIS$	2010	0.541	0.595	17
			$LA \times DIS$	2011	Inva	lid mode	1
	Leaf	Per-leaf abundance	$LDMC \times DIS$	2006	0.653	0.525	14
	quality		$LDMC \times DIS$	2010	-1.757	0.097	17
			$C:N \times DIS$	2006	0.943	0.362	14
			$C:N \times DIS$	2010	2.927	0.009	17
			$LPC \times DIS$	2006	0.792	0.442	14
			$LPC \times DIS$	2010	-1.997	0.063	16
Leaf miners	Leaf size	Per-leaf abundance	$LA \times DIS$	2010	0.541	0.595	17
			$LA \times DIS$	2011	-1.536	0.143	17
	Leaf	Per-leaf abundance	$LDMC \times DIS$	2010	-1.757	0.097	17
	quality		$C:N \times DIS$	2010	2.927	0.009	17
			$LPC \times DIS$	2010	-1.997	0.063	16
Parasitoids	Caterpillars	Parasitised caterpillars	$TC \times DIS$	2010	-0.673	0.509	18
	-	-	$\text{TC} \times \text{DIS}$	2011	-1.075	0.299	16

Table S7A.a: Effects of distance to the nearest oak on proportional use of resources.

Group	Use of	Dependent variable	Effects of	Year	t	р	DF
Ectophages	Leaf size	Per-leaf folivory	$LA \times OC$	2006	1.444	0.171	14
			$LA \times OC$	2010	-0.266	0.793	18
			$LA \times OC$	2011	1.907	0.074	17
		Per-individual folivory	$LA \times OC$	2010	1.489	0.156	16
			$LA \times OC$	2011	0.716	0.483	18
		Per-leaf abundance	$LA \times OC$	2010	-0.988	0.336	18
		·	$LA \times OC$	2011	-0.380	0.708	18
	Leaf	Per-leaf folivory	$LDMC \times OC$	2006	-0.805	0.434	14
	quality	55 5	$LDMC \times OC$	2010	0.935	0.363	17
	1 5		$C:N \times OC$	2006	0.213	0.834	14
			$C:N \times OC$	2010	-0.402	0.692	17
			$LPC \times OC$	2006	-1.910	0.077	14
			$LPC \times OC$	2010	0.290	0.775	18
		Per-individual folivory	$LDMC \times OC$	2010	Inva	lid mode	1
		0 0	$C:N \times OC$	2010	0.800	0.434	18
			$LPC \times OC$	2010	Inva	lid mode	1
		Per-leaf abundance	$LDMC \times OC$	2010	0.808	0.430	18
		·	$C:N \times OC$	2010	-0.770	0.451	18
			$LPC \times OC$	2010	0.036	0.972	18
Leaf gallers	Leaf size	Per-leaf abundance	$LA \times OC$	2006	-0.334	0.743	14
			$LA \times OC$	2010	-1.489	0.155	17
			$LA \times OC$	2011	Inva	lid mode	1
	Leaf	Per-leaf abundance	$LDMC \times OC$	2010	Inva	lid mode	1
	quality	-	$LDMC \times OC$	2006	-1.743	0.105	13
			$C:N \times OC$	2006	0.913	0.377	14
			$C:N \times OC$	2010	1.211	0.243	17
			$LPC \times OC$	2010	Inva	lid mode	1
			$LPC \times OC$	2006	-0.419	0.682	14
Leaf miners	Leaf size	Per-leaf abundance	$LA \times OC$	2010	-1.489	0.155	17
			$LA \times OC$	2011	-1.225	0.237	17
	Leaf	Per-leaf abundance	$LDMC \times OC$	2010	Inva	lid mode	1
	quality		$C:N \times OC$	2010	1.211	0.243	17
			$LPC \times OC$	2010	Inva	lid mode	1
Parasitoids	Caterpillars	Parasitised caterpillars	$TC \times OC$	2010	-1.909	0.073	17
		*	$TC \times OC$	2011	-1.060	0.303	18

Table S7A.b: Effects of	of oak	circumference on	proportional	use of resources.
-------------------------	--------	------------------	--------------	-------------------

(B) Percentage pines in the neighbourhood: Phylogenetic isolation is the mean phylogenetic distance to the neighbours and the value can be determined mainly by the neighbours having large phylogenetic distance to the focal tree. Our focal oak has the maximum phylogenetic distance to pines (among the neighbouring trees). Hence, we compared effects of phylogenetic isolation of individual host trees on the proportional use of resources to that of percentage of pines in the neighbourhood. We conducted these analyses for all groups of insects (extophages, leaf gallers, leaf miners and parasitoids), but not for the subgroups within ectophages. We found that percentage pine has a weaker effect than phylogenetic isolation. Percentage pines has a significant interaction effect in 9 out of those 35 cases (Table S7B), whereas phylohenetic isolation has a significant interaction and percentage pines have a significant interaction effect, there are 4 cases where phylogenetic isolation has a smaller p-value than percentage pines (compare between Table 2 and Table S7B). Hence, Phylogenetic isolation explains proportional use of resources more

consistently and more strongly than percentage pines in the neighbourhood. Note that the statistics given in the table is of only the interaction term in a model, but the models also accounted for the main effects of the variables in the interaction term. LA = leaf area, LDMC = leaf dry-matter content, C:N = carbon-nitrogen ratio, LPC = leaf phenolics content, TC = total caterpillars, % Pines= percentage pines.

Group	Use of	Dependent variable	Effects of	Year	t	р	DF
Ectophages	Leaf size	Per-leaf folivory	$LA \times \%$ Pines	2006	-0.031	0.975	14
			$LA \times \%$ Pines	2010	-3.159	0.005	18
			$LA \times \%$ Pines	2011	-2.538	0.021	17
		Per-individual folivory	$LA \times \%$ Pines	2010	0.549	0.590	18
			$LA \times \%$ Pines	2011	-2.197	0.042	17
		Per-leaf abundance	$LA \times \%$ Pines	2010	-2.268	0.037	17
			$LA \times \%$ Pines	2011	1.390	0.181	18
	Leaf	Per-leaf folivory	LDMC \times % Pines	2006	0.555	0.588	14
	quality		$LDMC \times \%$ Pines	2010	0.389	0.702	18
			$C:N \times \%$ Pines	2006	-0.452	0.658	14
			$C:N \times \%$ Pines	2010	1.158	0.262	18
			LPC \times % Pines	2006	0.338	0.740	14
			LPC \times % Pines	2010	0.744	0.467	18
		Per-individual folivory	$LDMC \times \%$ Pines	2010	-1.342	0.197	17
			$C:N \times \%$ Pines	2010	-0.545	0.592	18
			LPC \times % Pines	2010	-0.403	0.692	18
		Per-leaf abundance	$LDMC \times \%$ Pines	2010	1.417	0.174	17
			$C:N \times \%$ Pines	2010	0.209	0.837	17
			LPC \times % Pines	2010	0.615	0.615	18
Leaf gallers	Leaf	Per-leaf abundance	$LA \times \%$ Pines	2006	-0.686	0.504	14
	quality		$LA \times \%$ Pines	2010	0.222	0.827	17
			$LA \times \%$ Pines	2011	0.222	0.827	17
	Quality	Per-leaf abundance	LDMC \times % Pines	2006	0.867	0.400	14
			$LDMC \times \%$ Pines	2010	Inva	lid model	
			$C:N \times \%$ Pines	2006	0.987	0.340	14
			$C:N \times \%$ Pines	2010	5.473	<0.001	17
			$LPC \times \%$ Pines	2006	1.018	0.326	14
			$LPC \times \%$ Pines	2010	Inva	lid model	
Leaf miners	Leaf size	Per-leaf abundance	$LA \times \%$ Pines	2010	-1.155	0.263	18
			$LA \times \%$ Pines	2011	-1.179	0.255	17
	Leaf	Per-leaf abundance	LDMC \times % Pines	2010	0.4	0.694	18
	quality		$C:N \times \%$ Pines	2010	1.505	0.150	18
			LPC \times % Pines	2010	1.351	0.194	17
Parasitoids	Caterpillars	Parasitised caterpillars	$TC \times \%$ Pines	2010	-2.436	0.026	17
	-		$TC \times \%$ Pines	2011	-0.904	0.378	18

Table S7B: Effects of percentage of pines on proportional use of resources.

Online Resource 8. Logistic regression of presence/absence of species with wingless females

We conducted a logistic regression to test for an effect of phylogenetic isolation on the presence of species with wingless females. Phylogenetic isolation had a negative effect in 2010 (z = -2.079, p = 0.038, df = 20), but no effect in 2011 (z = 0.834, p = 0.404, df = 20).

