

#### From footprint morphometrics to the stature of fossil hominins: experimental regressions and uncertainties Jérémy Duveau

#### ▶ To cite this version:

Jérémy Duveau. From footprint morphometrics to the stature of fossil hominins: experimental regressions and uncertainties. 12th European Society for the study of Human Evolution meeting, Sep 2022, Tübingen, Germany. 2022. hal-03802103

#### HAL Id: hal-03802103 https://hal.science/hal-03802103

Submitted on 6 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.







# From footprint morphometrics to the stature of fossil hominins: experimental regressions and uncertainties

## Jérémy Duveau<sup>1,2</sup>

<sup>1</sup>DFG Center for Advanced Studies "Words, Bones, Genes, Tools: Tracking Linguistic, Cultural and Biological Trajectories of the Human Past", Eberhard Karls University of Tübingen, Tübingen, Germany. <sup>2</sup>UMR 7194 Histoire Naturelle de l'Homme Préhistorique, CNRS, Muséum National d'Histoire Naturelle, Université Perpignan Via Domitia, Paris, France

## Introduction

-

IV

Stature is one of the most used characteristics in paleoanthropology. Commonly inferred from long bones, it can also be estimated from footprints. It is usually estimated from footprint length by using the well-known foot length to stature ratio (14-16%). However, the morphology of the footprints is not only impacted by the dimensions of the foot but also by the biomechanical features of the individuals, the nature of the substrate in which they were left and taphonomic agents. Therefore, it is necessary to know the relationships between individual stature and footprint morphometry and not just foot morphometry. This poster presents a morphometric study performed on experimental footprints left in a dune environment, a common context in the fossil record.

> Issues & objectives

What are the relationships between footprint morphometrics and stature? > Use the most appropriate variable to best estimate the stature of an individual

For each variable, what is the intraindividual variation? > Quantify uncertainties to better estimate stature from isolated footprints.



## Material

- This study is based on the morphometric study of **175 experimental footprints** made by **20 individuals**.
- The age (11-36 years old) of each participant was recorded and their stature (146-182 cm) was measured.





- Each individual left footprints by walking on an experimental area composed of wet sand (fine to medium granulometry) from a dune environment.
- Each footprint was photographed, described and measured in situ.
- 49 footprints were digitized in 3D by photogrammetry in order to control the in situ measurements and to measure surfaces and volumes.

#### Methodological approach

The morphometric study is based on 24 variables measured on the 175 experimental footprints. The use of variables other than length is uncommon, but provides guidance for researchers wishing to estimate stature from incomplete footprints for which length is no longer measurable.



- $\succ$  12 lengths
- $\succ$  4 widths

## 5 areas:

- $\succ$  4 areas determined from linear dimensions
- > Area 5: accurate area determined on 3D models
- **3 volumes** determined from the 3D models:
  - Volume 1: heel maximum depth x Area 5
  - Volume 2: forefoot maximum depth x Area 5
  - > Volume 3: accurate volume determined on 3D models



Linear dimensions





Area 4

## From footprint morphometrics to stature

- Methods: The relationship between stature and each mean morphometric variable was investigated by determining the Pearson correlation coefficient and the linear regression equation giving the best prediction.
- **Results:** Total and tarsometatarsal lengths, especially those associated with the first three toes, are the variables giving the best predictions. Widths, surfaces, but especially volumes have a much lower correlation.



Morphometric	Pearson correlation	Linear equation for
variables (x)	coefficient (r)	estimating stature (y)
L <sub>tot1</sub> (cm)	0.69	y = 4.1x + 66.9
L <sub>tot2</sub> (cm)	0.69	y = 4.1x + 67.6
L <sub>tot3</sub> (cm)	0.69	y = 4.5x + 62.5
L <sub>tot4</sub> (cm)	0.68	y = 4.6x + 66.5
L <sub>tot5</sub> (cm)	0.65	y = 4.8x + 70.1
L <sub>tmt1</sub> (cm)	0.70	y = 4.8x + 70.6
L <sub>tmt2</sub> (cm)	0.69	y = 4.5x + 72.5
L <sub>tmt3</sub> (cm)	0.70	y = 4.5x + 75.8
L <sub>tmt4</sub> (cm)	0.66	y = 4.3x + 83.1
L <sub>tmt5</sub> (cm)	0.64	y = 4.5x + 85.6
L <sub>med</sub> (cm)	0.62	y = 5.6x + 67.2
L <sub>lat</sub> (cm)	0.65	y = 5.9x + 68.2
W <sub>prox</sub> (cm)	0.53	y = 10.3x + 103.7
W <sub>mid</sub> (cm)	0.39	y = 3.2x +153.6
W <sub>dist</sub> (cm)	0.28	y = 4.9x + 121.4
W <sub>toes</sub> (cm)	0.64	y = 9.6x + 78.9
Area 1 (cm²)	0.39	y = 0.1x + 142.2
Area 2 (cm <sup>2</sup> )	0.56	y = 0.2x + 133.7
Area 3 (cm²)	0.48	y = 0.2x + 135.8
Area 4 (cm <sup>2</sup> )	0.52	y = 0.2x + 138.5
Area 5 (cm²)	0.59	y = 0.2x +137.9
Volume 1 (cm <sup>3</sup> )	0.30	y = 0.01x + 165.6
Volume 2 (cm <sup>3</sup> )	0.35	y = 0.04x + 159.8
Volume 3 (cm <sup>3</sup> )	0 17	$y = 0.02y \pm 164.4$

#### Intraindividual variation V

- **Methods:** The intraindividual variation was studied for each individual and each variable by calculating the average coefficient of variation and the maximum deviation from the individual mean. They have not been calculated for Area 5 and the 3 volumes, the 3D models from which they have been determined are too few.
- **Results:** Lengths are the variables with the least intraindividual variation. They are therefore the variables for which there will be the least uncertainties when estimating stature from an isolated footprint. The widths but especially the areas have much larger intra-individual variations.



#### VI **Discussion & Conclusion**

This study shows that footprint lengths, particularly those between the base of the heel and the tips of the first three toes, are the morphometric variables of footprints that best predict stature. However, even these lengths are subject to significant uncertainties, not only residual errors in the regressions but also uncertainties related to intraindividual variation. This intraindividual variation is particularly important when estimating stature on isolated footprints, where the differences with the real stature of the individuals can be significant (~15% for the estimates based on footprint lengths).

In addition to these residual errors in the regressions and to the intraindividual variation, there is a third uncertainty linked to variations in body proportions between the different hominin species. Like other body proportions (intermembral index, crural index...), the foot length to stature ratio may have varied during human evolution which would impact the estimation of stature from footprints. However, the fossil record is too fragmentary to directly and accurately quantify this ratio for any species other than Homo sapiens, including the well-known Neandertals.





### More information about this research?

If you want to learn more about this experimental study, it is detailed in an article that will be published in L'Anthropologie in October 2022.

Acknowledgements: I am particularly grateful to the volunteers who participated in the experiments. This research would not have been possible without the support and technical help of D. Cliquet during the preparation of the experimentation and the data collection. My thanks also go to J. Guériel for his indispensable help during the construction of the experimental area. I would like to thank A. Gicqueau, C. Biets, B. Albouy and M. Tudal for their help in the recording of the experimental footprints. I would also like to thank G. Berillon and C. Verna for their helpful comments in improving this experimental study and its interpretations. The experimental study was validated by prefectural decree #28-2017-339 (05/17/2017, Normandy, France). This work is part of a research project on the biological and biomechanical characteristics of hominins funded by the FYSSEN foundation. The experiments were funded by the CNRS - Institut Ecologie et Environment International Research Network IRN-GDRI0870. The metric analyses were conducted under the ANR-18-CE27-0010-01 HoBiS programme.

