



Interferometric millimeter observations of water vapor on Mars

Thierry Fouchet, Raphaël Moreno, Emmanuel Lellouch, Franck Montmessin,
Marco Giuranna, Vittorio Formisano

► To cite this version:

Thierry Fouchet, Raphaël Moreno, Emmanuel Lellouch, Franck Montmessin, Marco Giuranna, et al..
Interferometric millimeter observations of water vapor on Mars. EPSC-DPS Joint Meeting 2011, Oct
2011, Nantes, France. pp.EPSC-DPS2011-787. hal-03802544

HAL Id: hal-03802544

<https://hal.science/hal-03802544>

Submitted on 30 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Interferometric millimeter observations of water vapor on Mars

T. Fouchet (1,4,5), R. Moreno (1), E. Lellouch (1), F. Montmessin (2), M. Giuranna (3) and V. Formisano (3)
 (1) Observatoire de Paris, LESIA, Meudon, France (thierry.fouchet@obspm.fr), (2) LATMOS, IPSL, Guyancourt, France, (3) IFSI-INAF, Roma, Italy, (4) Université Pierre et Marie Curie, UMR8109, Paris, France, (5) Institut Universitaire de France, Paris, France

Abstract

We present interferometric mapping of the 225.9-GHz HDO and 203.4-GHz H_2^{18}O lines on Mars obtained with the IRAM Plateau de Bure facility (PdBI). The observations were performed during martian year 28 (MY28), at $L_s = 320.3^\circ$ for the HDO line, and at $L_s = 324.3^\circ$ for the H_2^{18}O line. The HDO line is detected at the eastern (morning) and western (evening) limbs of the northern hemisphere, corresponding to a water column density in the range 3–6 pr.- μm (Fig. 1). The H_2^{18}O line is not detected, which is compatible with the column densities derived from the HDO line (Fig. 2). Quasi-simultaneous far infrared measurements obtained by the Planetary Fourier Spectrometer (PFS) onboard the Mars Express spacecraft confirm our PdBI results, yielding a 5 ± 1 pr.- μm meridionally constant water column abundance (Fig 3).

Such a low water abundance during the southern mid-autumn of MY28 does not correspond to the standard martian climatology as observed during the previous years. It was however already retrieved from near-infrared observations performed by the Compact Reconnaissance Imaging Spectrometer for Mars (CRISM) onboard the Mars Reconnaissance Orbiter spacecraft [Smith, M. D., Wolff, M. J., Clancy, R. T., Murchie, S. L. 2009. CRISM observations of water vapor and carbon monoxide. *J. Geophys. Res.* **114**, doi:10.1029/2008JE003288]. Our observations thus confirm that the planet-encircling dust storm that occurred during MY28 significantly affected the martian water cycle. Our observations also demonstrate the usefulness of interferometric submillimeter observations to survey the martian water cycle from ground-based facilities.




Figure 1: Spectral map in the HDO line at 225.897 GHz (black histogram lines), compared with synthetic models. At beam positions where the HDO line is detected the best-fit model is displayed in red along with the best-fit water column density. At beam positions where the HDO line is not detected a synthetic spectrum calculated for 5 pr.- μm is displayed in green. RA, Dec offsets (in ") are indicated for each beam position. The surface, equator, central meridian and terminator of Mars are indicated along with the sub-Earth and subsolar points, and the synthesized beam.




Figure 2: Spectral map in the H_2^{18}O line at 203.407 GHz (black histogram lines), compared with synthetic models calculated for 5 μm (green lines). RA, Dec offsets (in ") are indicated for each beam position. The surface, equator, central meridian and terminator of Mars are indicated along with the sub-Earth and subsolar points, and the synthesized beam.




Figure 3: Left panel: Surface temperature (black diamond) and atmospheric temperature at 100 Pa (red triangles) retrieved from PFS spectra obtained during orbit #4791. Right panel: Water column abundance retrieved from the same dataset.