



## The 1.10- and 1.18-micron nightside windows of Venus observed by SPICAV-IR aboard Venus Express

Bruno Bézard, Anna Fedorova, Jean-Loup Bertaux, Alexander Rodin, Oleg Korablev

### ► To cite this version:

Bruno Bézard, Anna Fedorova, Jean-Loup Bertaux, Alexander Rodin, Oleg Korablev. The 1.10- and 1.18-micron nightside windows of Venus observed by SPICAV-IR aboard Venus Express. EPSC-DPS Joint Meeting 2011, Oct 2011, Nantes, France. pp.EPSC-DPS2011-544. hal-03803165

**HAL Id: hal-03803165**

**<https://hal.science/hal-03803165>**

Submitted on 30 Nov 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The 1.10- and 1.18- $\mu\text{m}$ nightside windows of Venus observed by SPICAV-IR aboard Venus Express

B. Bézard (1), A. Fedorova (2), J.-L. Bertaux (3), A. Rodin (2) and O. Korabiev (2)

(1) LESIA, Observatoire de Paris, Meudon, France ([bruno.bezard@obspm.fr](mailto:bruno.bezard@obspm.fr) / Fax: +33-1-45072806), (2) Space Research Institute IKI, Moscow, Russia, (3) LATMOS, Guyancourt, France

## 1. Introduction

Thermal emission from the deep atmosphere of Venus observed on the night side in the 1.10- and 1.18- $\mu\text{m}$  windows provides information on the water vapor abundance in the altitude range 0-25 km. The  $\text{H}_2\text{O}$  mole fraction near the surface is still controversial and the various analyses of in situ measurements and observations of the nightside windows do not yield consistent results [4]. We present here an analysis of the 1.10- and 1.18- $\mu\text{m}$  nightside windows observed by the infrared channel of the SPICAV (Spectroscopy for the investigation of the characteristics of the atmosphere of Venus) instrument aboard Venus Express [3]. We investigated the influence of the various sources of opacity ( $\text{CO}_2$  and  $\text{H}_2\text{O}$  line lists,  $\text{CO}_2$  far wing and “continuum” opacity) on the spectrum and of the associated uncertainties or unknowns. The results were used to propose a best representation of the  $\text{CO}_2$  and  $\text{H}_2\text{O}$  opacity and to derive a best fit  $\text{H}_2\text{O}$  mixing ratio.

## 2. Observations

We used nadir observations by the LW range of the visible-infrared channel (Vis-IR) of SPICAV, a single pixel spectrometer based on AOTF technology [3]. The spectral resolution is about  $5.2 \text{ cm}^{-1}$  (0.7 nm at 1.18  $\mu\text{m}$ ). We made a large average of LW spectra recorded during Orbits 800 and 825, yielding a S/N ratio of 500 at the peak of the 1.18- $\mu\text{m}$  window (Fig. 1).

## 3. Spectral modelling

Synthetic spectra were generated using a line-by-line radiative transfer model with scattering [4] and compared with the SPICAV selection. We allowed the use of a constant scaling factor as a free parameter to account for the unknown cloud optical depth.

We first investigated the  $\text{CO}_2$  far wing lineshape, which is constrained by the high-wavelength wing of the 1.18- $\mu\text{m}$  window. We found that the line profile model of Afanasenko and Rodin [1] underestimates the  $\text{CO}_2$  absorption and we derived instead an empirical sub-lorentzian profile that reproduces this wing. It provides more absorption than those previously derived for  $\text{CO}_2$  bands near 2.3  $\mu\text{m}$  at room temperature.

The variation of the 1.10- and 1.185- $\mu\text{m}$  radiance with surface elevation, as observed by the Venus Express / VIRTIS-M instrument, requires an additional “continuum” opacity, besides that provided by the  $\text{CO}_2$  and  $\text{H}_2\text{O}$  allowed bands. We have re-analyzed the dataset of Bézard et al. [4] to constrain this additional absorption at 1.10 and 1.185  $\mu\text{m}$  and derived a constant absorption coefficient, equal to  $0.7 \pm 0.2 \times 10^{-9} \text{ cm}^{-1} \text{ amagat}^{-2}$ .

We have modeled the SPICAV spectra using either the high-T [5] or CDSD [6]  $\text{CO}_2$  line lists. The latter provides less absorption than “High-T” in the interval 1110-1190 nm and reproduces worse the observed spectrum, due to the absence of the  $5\nu_1 + \nu_3$  series of  $\text{CO}_2$  bands. On the other hand, CDSD agrees better with the observations shortward of 1100 nm although the data clearly show evidence for a missing absorption between 1085 and 1100 nm. In this work, we have used a composite  $\text{CO}_2$  line list combining “High-T” and CDSD.

Absorption by HDO had not been previously added in the modeling of the 1.10- and 1.18- $\mu\text{m}$  windows. Using a recently published line list (VTT) [7], we found that the influence of HDO is significant in the range 1140-1190 nm, inducing a 7% drop in radiance at 1185 nm and slightly improving the fit of the SPICAV spectrum. We also confirm that the  $\text{H}_2^{16}\text{O}$  BT2 database [6] yields more opacity than the GEISA or HITRAN databases, being more accurate and more complete in high-energy lines [2].

Using the sources of atmospheric opacity described above, we determined a best-fit value for the H<sub>2</sub>O mole fraction of  $30^{+10}_{-5}$  ppm (assumed constant below the clouds) (Fig. 1). The emission in the H<sub>2</sub>O band is most sensitive to the region 5-25 km.




Figure 1: The SPICAV spectrum is compared to synthetic spectra calculated with H<sub>2</sub>O mole fractions of 20, 30 and 45 ppm using our best estimate for the representation of the H<sub>2</sub>O/HDO and CO<sub>2</sub> opacity. *Upper panel*: raw calculations; *Lower panel*: spectra renormalized to the observed radiance at 1185 nm.

## 4. Discussion and conclusion

We have used observations of the night side of Venus by the Venus Express / SPICAV-IR instrument in the 1.10- and 1.18- $\mu$ m windows to better characterize the various sources of molecular opacity at these wavelengths and determine the average mole fraction of water vapor in the 5-25 km altitude range.

The 30-ppm mole fraction we derived is smaller than that derived from VIRTIS-M spectra ( $44 \pm 9$  ppm) [4], even though the error bars overlap. This value is fully consistent with the mole fraction derived at 30-45 km from the 2.3- $\mu$ m window and at 15-25 km from the 1.74- $\mu$ m window. This result gives strong evidence that the water mole fraction is constant

from approximately 5 to 40 km, as predicted by chemical models.

Our best fit model still shows some discrepancies with the SPICAV spectrum, e.g. between 1085 and 1100 nm. They likely result from an inaccurate representation of the CO<sub>2</sub> opacity. Laboratory measurements would help in this regard.

## Acknowledgements

We are grateful to V. Perevalov and A. Campargue for extremely useful discussions on the CDSD databank and for providing positions and relative intensities of the  $5\nu_1 + \nu_3$  CO<sub>2</sub> bands. We thank J. Bailey for providing useful information on the water vapor databases. B.B and J.-L.B. acknowledge support from CNES and CNRS. A.F., A.R., and O.K. acknowledge RFBR grants 10-02-93116 and 08-02-01383.

## References

- [1] Afanasenko, T.S. and Rodin, A.V.: Interference of spectral lines in thermal radiation from the lower atmosphere of Venus. *Astron. Lett.*, Vol. 33, pp. 203-210, 2007.
- [2] Bailey, J.: A comparison of water vapor line parameters for modeling the Venus deep atmosphere. *Icarus*, Vol. 201, pp. 44-453, 2007.
- [3] Bertaux, J.L., and 35 colleagues: SPICAV on Venus Express: Three spectrometers to study the global structure and composition of the Venus atmosphere, *Planet. Space Sci.*, Vol. 55, pp. 1673-1700, 2007.
- [4] Bézard, B., Tsang, C.C.C., Carlson, R.W., Piccioni, G., Marcq, E., and Drossart, P.: Water vapor abundance near the surface of Venus from Venus Express/VIRTIS observations. *J. Geophys. Res.*, Vol. 114, E00B39, doi: 10.1029/2008JE003251, 2009.
- [5] Pollack, J.B., and 11 colleagues: Near-infrared light from Venus' nightside: A spectroscopic analysis. *Icarus*, Vol. 103, pp. 1-42, 1993.
- [6] Rothman, L.S., and 8 colleagues: HITRAN, the high-temperature molecular spectroscopic database. *J. Quant. Spectrosc. Trans.*, Vol. 111, pp. 2139-2150, 2010.
- [7] Voronin, B.A., Tennyson, J., Tolchenov, R.N., Lugovskoy, A.A., and Yurchenko, S.N.: A high accuracy computed line list for the HDO molecule. *Mon. Not. R. Astron. Soc.*, Vol. 402, pp. 492-496, 2010.