



Ultrasound processing to enhance the functionality of plant-based beverages and proteins

Meliza Lindsay Rojas, Mirian T K Kubo, Alberto Claudio Miano, Pedro E D Augusto

► To cite this version:

Meliza Lindsay Rojas, Mirian T K Kubo, Alberto Claudio Miano, Pedro E D Augusto. Ultrasound processing to enhance the functionality of plant-based beverages and proteins. *Current Opinion in Food Science*, In press, 10.1016/j.cofs.2022.100939 . hal-03807784

HAL Id: hal-03807784

<https://hal.science/hal-03807784>

Submitted on 10 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rojas, M. L., Kubo, M. T., Miano, A. C., & Augusto, P. E. D. (2022). Ultrasound processing to enhance the functionality of plant-based beverages and proteins. *Current Opinion in Food Science*, 100939. DOI: <http://dx.doi.org/10.1016/j.cofs.2022.100939>

Ultrasound processing to enhance the functionality of plant-based beverages and proteins

Meliza Lindsay Rojas¹, Mirian T. K. Kubo², Alberto Claudio Miano¹, Pedro E. D. Augusto^{3*}

¹ Dirección de Investigación, Innovación y Responsabilidad Social, Universidad Privada del Norte (UPN), Trujillo, Peru.

² Université de Technologie de Compiègne, UPJV, CNRS, Enzyme and Cell Engineering, Centre de recherche Royallieu - CS 60319 - 60 203 Compiègne Cedex, France.

³ Université Paris-Saclay, CentraleSupélec, Laboratoire de Génie des Procédés et Matériaux, SFR Condorcet FR CNRS 3417, Centre Européen de Biotechnologie et de Bioéconomie (CEBB), 3 rue des Rouges Terres 51110 Pomacle, France.

* pedro.augusto@centralesupelec.fr

Highlights

- Ultrasound processing of plant-based beverages are discussed, including juices, milk analogues and proteins.
- Ultrasound promotes structural modifications, impacting the beverage properties.
- Enhanced nutritional and bioactive properties can be obtained.
- Ultrasound processing can substitute the addition of some ingredients focusing on clean labels products.
- The new functionalities can result in new products.


Abstract

This opinion paper discusses how ultrasound can be used to modify the structure of plant-based beverages (juices and “milk”) and proteins, achieving new functionalities. Both positive (such as increasing the nutrient and bioactive compounds bioaccessibility, improving protein solubility and modifying its digestibility) and negative (such as degradation of nutrient and bioactive compounds by exposing them to the environment, or the limited microbial inactivation) aspects are discussed. It is clear that ultrasound technology can be used as a valuable tool to improve plant-based beverages properties, helping to achieve clean label products and positively impacting wellbeing. Its scale-up

to industry, however, is still a concern that needs both scientific studies and technological development.

Keywords: ultrasound, ultrasonication, juices, vegetable milk, proteins, plant-based, beverages, non-thermal technologies, emerging technologies

Graphical Abstract



1. Introduction and contextualization

Fruits and vegetables are important parts of the human diet as sources of nutrients and bioactive compounds, such as carotenoids, phenolics (or polyphenols), flavonoids, vitamins and minerals, being juices a practical and relevant way to ingest those plant components.

Furthermore, there is a rising demand for plant-based products (such as milk analogues) and ingredients (such as proteins). The expansion of the plant-based market is a worldwide trend and is related to several reasons, such as the growth of the vegan, vegetarian and 'flexi-vegans' public, the concern with the environmental impact of food production, sustainability and animal welfare, in addition to the health aspect (e.g. allergenicity of milk proteins).

Therefore, this paper describes the current research on using ultrasound processing to enhance the functionality of plant-based beverages, considering the three most representative examples in the context: (i) fruit and vegetable juices, produced from top-down (i.e., by disassembling the plant organism towards a smaller scale); (ii) plant-based "milk", produced from top-down or bottom-up (i.e., by assembling a new product from ingredients, then using (iii) vegetable proteins). In those products, the main objectives of using ultrasound are preserving nutrients and bioactive compounds, increasing their digestibility and accessibility to the human body, improving physical stability, rheology and interaction with water, thus achieving desirable sensorial impact, better healthy aspects and promoting wellbeing.

More information about using ultrasound technology in food products, ingredients and processing can be obtained in the recent review articles of [1-6].

2. Ultrasound processing of fruit and vegetable juices

Fruit and vegetable juices are particulate solid-liquid systems, constituted by a suspension of cells and their fragments, including fibres and cell walls, in the serum (water+soluble material). Ultrasound processing modify this structure in different stages, as described by Rojas, Leite, Cristianini, Alvim and Augusto [7], impacting their physical, chemical, sensorial, and nutritional modifications – as recently detailed by Rojas, Kubo, Caetano-Silva and Augusto [4••]. Table 1 presents recent studies in ultrasound processed juices, where the main results are discussed.

Table 1. Ultrasound processing and main reported impact on fruit and vegetable juices, regarding physical modifications, compound content and quality: **Increase or Improvement**, **Decrease or Negative effect**, and **No variation**, compared to untreated or fresh juices.

Juice source	Processing conditions	Main results of ultrasound processing	
Spinach [8•]	TS [30 kHz; 60 °C; 20 min; 200-600 W; 50% dc] Vs [100 mL]	Physical stability, rheological properties, bioactive compounds	Particle size
Apple [9]	TS [30-70 °C; 5-12 min; 525-1125 W] Vs [200 mL]	Colour, cloudiness, Physical stability, rheological properties	Particle size
Kiwifruit [10]	US [20 kHz; ~ 4 °C; 4-16 min; 400 W; 50% dc] Vs [100 mL]	Colour, cloudiness, rheological properties	Pectin
Carrot [11]	US [20 kHz; 30 °C; 10 min; 2s on-2s off; 221 - 321 W] Vs [25 mL]	Apparent viscosity	Colour, carotenoids
Plum [12]	TS [40 kHz; 40 - 60 °C; 5 - 30 min; 2s on-2s off; 0.348 W/cm ³]	Colour, cloudiness, flavonoids, phenolics, carotenoids, ascorbic acid	Compounds degradation at T > 60°C
Tomato [13•]	TS [20 kHz; 37 - 52 °C; 2 - 10 min; 28 - 40 W/cm ²] Vs [150 mL]	Lycopene	Ascorbic acid, Lycopene
Sweet potato [14]	TS [26 kHz; 8 min; 0.66 W/cm ²] Vs [150 mL]	Bioaccessibility of β -carotene	Colour
Guava [15]	US [20 kHz; 25 °C; 0 - 9 min; 15 W/cm ²] Vs [200 mL]	Physical stability, bioaccessibility of lycopene	Lycopene
Strawberry [16]	TS [40 kHz; 25-50 °C; 5 -15 min; 110 W] Vs [50 mL]	Ascorbic acid, anthocyanin	pH, °Brix, titrable acidity
Orange [17]	US [20 kHz; 10 min; ~ 0 °C; 300-1200 W] Vs [10 mL]	Ascorbic acid	Xylooligosaccharides
Mango [18]	US [20 kHz; ~ 0 °C; 0-40 min; 5 s on - 5s off; 0-600 W]	Consistency and flow behaviour index	Carotenoid, phenolics

	Vs [100 mL]			
Mandarin [19]	US [19 kHz; 50 °C; 36 min; 750 W]	Colour	Sugar, acid components	
Orange [20].	US [19 kHz; 37 °C; 15-150 s; 15-86 J/mL] Vs [30 mL]	Cloudiness	Ascorbic acid	
Watermelon [21]	US [26 kHz; 4-16 min; 80 W; 50%A] Vs [100 mL]	Lycopene, phenolics, flavonoids, antioxidant capacity	Ascorbic acid	pH, °Brix, titrable acidity
Açaí (<i>Euterpe precatoria</i>), buriti (<i>Mauritia flexuosa</i>) [22]	US [20 kHz; 70 °C; 10 min; 0.9-3.6 kJ/cm ³] Vs [100 mL]	Fatty acids, Bioaccessibility and concentration of anthocyanins, carotenoids	Fatty acids, Colour	pH, °Brix, titrable acidity
Açaí [23•]	US [19 kHz; 25 °C; 2-10 min; 75.34-272.93 W/cm ²] Vs [150 mL]	Sugar, betaine, bioaccessibility of ascorbic acid.		
Orange [24]	TS [28 kHz; 10-50 °C; 5-25 min; 180-900 W/L]	Aroma, glycosidically bound volatiles		
<i>Berberis amurensis</i> Rupr. [25]	US [20 kHz; 10 min; 140 W; 70%A]	Colour, anthocyanins		
Red pitaya [26]	TS [56 °C; 20 min; 475 W] US [10 °C; 20 min; 475 W] Vs [60 mL]	Colour, betacyanins		
Strawberry [27]	TS [24 kHz; 23.5-56.5 °C; 3 min; 0.29 W/mL]	Anthocyanins, soluble solids, phenolics		
Sugar cane [28]	TS [20 kHz; 80 °C; 23 min; 750 W]	Phenolics, colour		

TS = termosonication processing conditions; US = ultrasound processing conditions. Regarding processing conditions, it is presented: the frequency, temperature, time, power or acoustic density or acoustic intensity, duty cycle (dc) and amplitude (A). Vs= volume of sample.

The cell disruption and reductions in particle size promote new particle-particle and particle-serum interactions that allows increasing of cloudiness, suspension stability and variation on rheological properties [8-11, 18]. The rheological properties can be changed temporally or permanently, depending on the ultrasound energy applied, processing time, pulp content and temperature [7, 29]. Therefore, a different impact on the rheological properties is expected, and the processing conditions must be evaluated depending on the defined target.

On the other hand, colour variations are reported, which are related to the decrease in particle size, extraction and/or degradation of main pigments. In fact, once the cell wall is impacted, the intracellular content is exposed and some compounds become more assessable (to the human body, as a positive aspect, or to oxygen and the environment, as a drawback). The variation in the content of the compound (Table 1) is a balance of improved extraction (causing an apparent increase in the quantified concentration) and exposition to deteriorating conditions (decreasing their concentration). In fact, ultrasound can modify the juice structure with potential health benefits, which can be particularly relevant by considering food products for special needs (Rojas et al., 2021).

Even so, some ambiguous results are reported, such as for lycopene and ascorbic acid, once they can be easily degraded due to exposure to heat, light, and oxygen. The decrease in such compounds is due to long times, high temperatures and/or powers of process, while at inverse conditions, increase or no variation occur. In either case, despite the decrease in some compounds, ultrasonic processing is more effective than conventional thermal processing regarding compound preservation [12, 16, 20, 23•] – although this comparison is always difficult, being necessary an evaluation of process target and conditions.

Moreover, ultrasound alone is rarely effective for the required microbial inactivation in juices, even considering microbial resistance to ultrasound is small at low pH [30]. Therefore, ultrasound has been combined with other technologies to improve its microbial inactivation, such as natural antimicrobials [11, 31, 32], pulsed light ([33], ozone [34], heat under pressure [35] and, especially, the combination with a thermal process (thermo-sonication) at mild temperatures [9, 27]. In fact, ultrasound can turn the microorganisms more sensitivity to thermal inactivation, such as during the thermo-sonication to inactivate *Neosartorya fischeri* ascospores [36] and *Saccharomyces cerevisiae* [37] in apple juice, or *Escherichia coli* in apple cider [38]. Therefore, from a microbiological perspective, the use of ultrasound in combination with mild heating or other technologies for industrial use is promissory, although the impact on juice compounds must be evaluated since the combination of unsuitable conditions can be detrimental to sensitive compounds.

Summarising, ultrasound can improve the functionality of fruit and vegetable juices by improving the stability, physical properties, and bioaccessibility of some bioactive compounds. In addition, it preserves the thermolabile compounds or those added to the juices to increase functionality [17]. However, some aspects should be improved in future studies such as the calculation of the actual ultrasound power density in the products, the control of the process temperature, evaluation of the stability throughout the storage, and studies of bioaccessibility, bioavailability and bioactivity.

3. Ultrasound processing of vegetable alternatives to milk

There is a rising trend in producing vegetable beverages to be similar to animal milk, concerning the main sensorial and/or nutritional properties. Those beverages, often called plant-based “milks” (and here referenced as that, to avoid confusions with other plant-based beverages, such as juices, nectars, soft drinks, etc), are suspensions and emulsions from sources such as soybean, rice, almond, peanut, oat, among others – thus being physical stability an important challenge.

Ultrasound has demonstrated being useful for improving the physical stability of almond “milk” using 5 min – 300 W [39], coconut “milk” using 13 min – 55W/cm² [40•] and peanut “milk” using 3 min – 400W [41•], which can be related to changes on particle size (proteins and fats) and rheology.

These beverages are reported to present bioactive compounds such as isoflavones, phytosterol and great antioxidant activity [42]. However, plant-based “milk” can present antinutritional factors such as phytates, saponins and protease inhibitors [43]. In fact, ultrasound can positively impact the nutritional and bioactive properties of plant-based “milk”. For instance, it increased the protein digestibility by 7.4%, reducing the trypsin inhibitor by 52% in soybean milk - 400 W, 25 kHz, 16 min [44]. However, no significant improvement was found for almond milk using ultrasound (400 W, 20 kHz, 16 min) [45]. Moreover, ultrasound (400 W, 24 kHz) treatment for 20 min applied to hydrated soybeans produced “milk” with 63–84% higher isoflavone content than untreated beans [46].

Different works report that ultrasound alone is rarely effective for the required microbial inactivation in plant-based “milk”. For instance, a reduction of 0.9 log (total plate count) was obtained using a probe of 400 W for 3 min on peanut “milk” [41•], while a reduction up to 1.3 log was found for *Escherichia coli* and *Listeria monocytogenes* when ultrasound (130 W, 20 kHz) with 80% of power and 8 min treatment was applied to almond “milk” [47]. On the other hand, some works report microbial inactivation similar to pasteurization processes, which can be an alternative for this product commercialization under refrigeration. For example, Campaniello, Bevilacqua, Speranza, Sinigaglia and Corbo [48•] reported a reduction on *Salmonella enterica* load

up to 8 log in rice “milk” - although samples experienced up to 48°C/10 min during processing, being thus relevant the contribution of thermal energy. However, by being low acid products (pH close to neutrality), the main target of vegetable “milk” are bacteria spores. Therefore, further studies are needed considering the safety and stability aspects of microbiology, with a focus on specific microbial targets and a combination of technologies.

Summarizing, ultrasound technology has been demonstrated to be beneficial to process plant-based milk obtaining beverages with physical stability during storage and with some nutritional enhancements. The most used device was the ultrasonic probe of different nominal power (130–400W) and frequency (20-25 kHz), reporting different parameter effects as power percentage application (20–100%), time (1–60 min), amplitude (50–100%) and temperature (25–60°C). In contrast, ultrasonic bath has not been used in the last 5 years for plant-based milk processing. This probably occurred due to ultrasonic probe is more powerful than the ultrasonic bath leading to used shorter processing time. Similarly to other products, more studies of processing equipment and conditions, focusing on scale-up, are needed. Particularly, the potential of ultrasound to improve stability, physical, sensorial and nutritional properties of vegetable “milk” is just starting to be revealed, and further studies are needed involving different sources, processing approaches and targets, the effect in other antinutritional factors as saponins and phytates, and how ultrasound treatment affects further processes for elaborating analogues of cheese, yoghurt, or ice cream.

4. Ultrasound processing of plant proteins

Proteins are complex biopolymers made up of amino acids and are of great nutritional, sensory and functional importance. They present different functional properties depending on factors such as their source, amino acid composition and sequence, molecular weight, structure, conformation, and surface hydrophobicity [49]. Among the various techno-functional properties of proteins, one can highlight the properties related to solubility, thickening, water/oil holding capacity, emulsification, foaming, and gelation. The variety of properties makes proteins important components in the formulation of food products, either as constituents from raw materials (e.g. soy proteins in soy “milk”) or as ingredients (protein concentrates or isolates). However, vegetable proteins might present some limitations due to the unsuitability of their properties for specific applications and formulations in the food industry. To overcome these limitations, chemical, biological and physical methods have been employed for protein modification, including enzymatic hydrolysis, fermentation, heat treatment, high-pressure processing, ultrasound [50-53]. Comprehensive reviews on the different

approaches for the modification of plant-based proteins can be found in the literature [54, 55••].

Recently, the use of ultrasound has attracted attention to modify, improve and modulate the functional properties of protein concentrates and isolates from different plant sources. Several studies have reported that the application of high-intensity ultrasound to dispersed/suspended proteins resulted in structural changes and thus, modifications of the functional properties of these proteins. Some recent studies are presented in Table 2. These changes are attributed to cavitation, shear stress, turbulence effects and localized temperature rise, which can promote modification even at a molecular level [6, 56-58].

For example, Flores-Jiménez, Ulloa, Urías-Silvas, Ramírez-Ramírez, Bautista-Rosales and Gutiérrez-Leyva [59•] found that ultrasound altered the secondary and tertiary structures of guamuchil seed protein isolates, which increased their surface hydrophobicity and molecular flexibility. These structural changes led to a significant improvement of solubility, oil absorption capacity, emulsification, foaming and gelation properties. Similarly, Hussain, Qayum, Zhang, Hao, Liu, Wang, Hussain and Li [60] investigated the effect of high-intensity ultrasound on the bioactive, functional and structural properties of potato protein and reported that emulsifying ability index, emulsifying stability, foaming ability, solubility and digestibility were significantly enhanced.

However, it is worth mentioning that depending on the process parameters used, different results can be obtained, and may result in an increase or decrease in protein functionality. In general, if conditions are too intense, especially long sonication times and very high intensities, the (over-)processing can result in an excessive molecular aggregation leading to the obtention of proteins with poor functional properties [61]. For example, Karabulut and Yemiş [61] observed an increase in solubility of hemp seed protein isolates after ultrasound processing at the range of 45–65% amplitude (6–11 W/cm², 27–32 W/cm²) for 5–10 min, which was attributed to the partial unfolding of protein molecules and greater protein-water interaction. However, at higher amplitudes and longer times, the protein solubility was reduced, probably due to the formation of aggregates with higher molecular weights. Similarly, Tang, Du and Fu [62] observed that solubility, foaming properties and emulsifying properties of *Moringa oleifera* seed protein increased or decreased depending on the ultrasonic power. Therefore, to achieve the desired level of functionality, the operating parameters must be chosen/optimized.

In summary, ultrasound processing is an interesting and promising approach for protein modification, which may allow the utilization of a wider variety of vegetable proteins and the expansion of their applications in the food industry. Nevertheless, it

should be noted that the focus has been on using sonication for modification of isolated proteins (in suspension), but the modification of proteins in complex food matrices and the potential interaction of proteins with other components should not be neglected.

Future developments are expected to demonstrate the increase of functionality of plant proteins of different sources, for different purposes (including as ingredients for other food products and improvement of nutritional properties of emerging sources).

Table 2. Examples of recent studies on the modification of plant proteins by ultrasound processing and the respective improvements in techno-functional properties: increasing in S: solubility; E: emulsifying; G: gelling; OA: oil absorption capacity; WH: water holding capacity; F: foaming.

Type of protein	Improved properties						References
	S	E	G	OA	WH	F	
black gram protein							[63]
guamuchil seed protein isolate							[59]
hemp seed protein isolate							[61]
lupin protein isolate							[64]
<i>Moringa oleifera</i> seed protein							[62]
pea protein concentrate							[65]
perilla seed protein isolate							[58]
potato proteins							[60]
potato protein isolates							[66]
pumpkin-seed protein isolates							[57]

5. Conclusions and perspectives

Ultrasound processing induces a sequence of structural modifications in plant-based products, impacting their properties. The modification involved different steps from the intact tissue to cell and macromolecules spatial changes to disruption. Consequently, different properties are altered, positively or negatively, such as improving the health benefits or sensorial perception, but also degrading compounds of interest due to exposition to detrimental conditions. Therefore, more studies are needed to describe how processing conditions can affect the plant beverage functionalities – considering different vegetable sources and food products for different needs.

In particular, a persistent challenge is scaling up the ultrasound processing at industrial levels – in special considering that most of the studies apply the probe reactor.

Studies of reactor engineering are needed to provide equipment able to process the volumes of the food industry, as well as studies in material science, are needed to develop more resistant probe tips (which wear out by erosion).

We consider the ultrasound technology can be used as a valuable tool to improve the properties of plant-based beverages, helping to achieve clean label products and positively impacting wellbeing. Its industrial application, though, is close to be effective, although depending on the next scientific studies and technological developments.

Acknowledgements

This study was carried out in the Centre Européen de Biotechnologie et de Bioéconomie (CEBB), supported by the Région Grand Est, Département de la Marne, Greater Reims (France) and the European Union. In particular, the authors would like to thank the Département de la Marne, Greater Reims, Région Grand Est and the European Union along with the European Regional Development Fund (ERDF Champagne Ardenne 2014-2020) for their financial support of the Chair of Biotechnology of CentraleSupélec.

Author contributions

Meliza Lindsay Rojas: Conceptualization; Data curation; Formal analysis; Investigation; Project administration; Visualization; Writing - original draft; Writing - review & editing.

Mirian T. K. Kubo: Conceptualization; Data curation; Formal analysis; Investigation; Project administration; Visualization; Writing - original draft; Writing - review & editing.

Alberto Claudio Miano: Conceptualization; Data curation; Formal analysis; Investigation; Project administration; Visualization; Writing - original draft; Writing - review & editing.

Pedro E. D. Augusto: Conceptualization; Data curation; Formal analysis; Investigation; Project administration; Visualization; Writing - original draft; Writing - review & editing.

References

Papers of particular interest, published within the period of review, have been highlighted as: of special interest (•), or of outstanding interest (••)

- 1 Bhargava N, Mor RS, Kumar K, Sharanagat VS: **Advances in application of ultrasound in food processing: A review.** *Ultrasonics Sonochemistry* 2021; **70**:105293.
- 2 Cui R, Zhu F: **Ultrasound modified polysaccharides: A review of structure, physicochemical properties, biological activities and food applications.** *Trends in Food Science & Technology* 2021; **107**:491-508.
- 3 Gallo M, Ferrara L, Naviglio D: **Application of Ultrasound in Food Science and Technology: A Perspective.** *Foods* 2018; **7**:164.

- 4•• Rojas ML, Kubo MTK, Caetano-Silva ME, Augusto PED: **Ultrasound processing of fruits and vegetables, structural modification and impact on nutrient and bioactive compounds: a review.** *International Journal of Food Science & Technology* 2021; **56**:4376-4395.
- This recent review discuss and correlate how the ultrasound processing impacts the fruits and vegetables structure, nutrients and bioactive compounds.
- 5 Singla M, Sit N: **Application of ultrasound in combination with other technologies in food processing: A review.** *Ultrasonics Sonochemistry* 2021; **73**:105506.
- 6 Téllez-Morales JA, Hernández-Santo B, Rodríguez-Miranda J: **Effect of ultrasound on the techno-functional properties of food components/ingredients: A review.** *Ultrasonics sonochemistry* 2020; **61**:104787.
- 7 Rojas ML, Leite TS, Cristianini M *et al.*: **Peach juice processed by the ultrasound technology: Changes in its microstructure improve its physical properties and stability.** *Food Research International* 2016; **82**:22-33.
- 8• Manzoor MF, Xu B, Khan S *et al.*: **Impact of high-intensity thermosonication treatment on spinach juice: Bioactive compounds, rheological, microbial, and enzymatic activities.** *Ultrasonics Sonochemistry* 2021; **78**:105740.
- This work shows the effect of thermosonication (200-600 W, 30 kHz, 60 °C, 20 min) on physical stability, rheological properties, bioactive compounds, microorganisms, and peroxidase and polyphenoloxidase enzymes of spinach juice.
- 9 Shen Y, Zhu D, Xi P *et al.*: **Effects of temperature-controlled ultrasound treatment on sensory properties, physical characteristics and antioxidant activity of cloudy apple juice.** *LWT* 2021; **142**:111030.
- 10 Wang J, Wang J, Vanga SK, Raghavan V: **High-intensity ultrasound processing of kiwifruit juice: Effects on the microstructure, pectin, carbohydrates and rheological properties.** *Food Chemistry* 2020; **313**:126121.
- 11 Bi X, Zhou Z, Wang X *et al.*: **Changes in the Microbial Content and Quality Attributes of Carrot Juice Treated by a Combination of Ultrasound and Nisin During Storage.** *Food and Bioprocess Technology* 2020; **13**:1556-1565.
- 12 Oladunjoye AO, Adeboyejo FO, Okekunbi TA, Aderibigbe OR: **Effect of thermosonication on quality attributes of hog plum (*Spondias mombin* L.) juice.** *Ultrasonics Sonochemistry* 2021; **70**:105316.
- 13••Starek A, Kobus Z, Sagan A *et al.*: **Influence of ultrasound on selected microorganisms, chemical and structural changes in fresh tomato juice.** *Scientific Reports* 2021; **11**:3488.

- This work evaluated and described the ultrasound effects on microstructure of tomato juice, also effects on microorganisms and compounds such as lycopene and ascorbic acid.
- 14 Rios-Romero EA, Ochoa-Martínez LA, Bello-Pérez LA *et al.*: **Effect of ultrasound and steam treatments on bioaccessibility of β -carotene and physicochemical parameters in orange-fleshed sweet potato juice.** *Heliyon* 2021; 7:e06632.
- 15 Campoli SS, Rojas ML, do Amaral JEPG *et al.*: **Ultrasound processing of guava juice: Effect on structure, physical properties and lycopene in vitro accessibility.** *Food Chemistry* 2018; **268**:594-601.
- 16 Menelli GS, Fracalossi KL, Lepaus BM, De São José JFB: **Effects of high-intensity ultrasonic bath on the quality of strawberry juice.** *CyTA - Journal of Food* 2021; **19**:501-510.
- 17 Silva EK, Arruda HS, Pastore GM *et al.*: **Xylooligosaccharides chemical stability after high-intensity ultrasound processing of prebiotic orange juice.** *Ultrasonics Sonochemistry* 2020; **63**:104942.
- 18 Wang J, Liu Q, Xie B, Sun Z: **Effect of ultrasound combined with ultraviolet treatment on microbial inactivation and quality properties of mango juice.** *Ultrasonics Sonochemistry* 2020; **64**:105000.
- 19 Cheng C-x, Jia M, Gui Y, Ma Y: **Comparison of the effects of novel processing technologies and conventional thermal pasteurisation on the nutritional quality and aroma of Mandarin (Citrus unshiu) juice.** *Innovative Food Science & Emerging Technologies* 2020; **64**:102425.
- 20 Gomes A, Costa ALR, Rodrigues PD *et al.*: **Sonoprocessing of freshly squeezed orange juice: Ascorbic acid content, pectin methylesterase activity, rheological properties and cloud stability.** *Food Control* 2022; **131**:108391.
- 21 Yıkmış S: **Sensory, physicochemical, microbiological and bioactive properties of red watermelon juice and yellow watermelon juice after ultrasound treatment.** *Journal of Food Measurement and Characterization* 2020; **14**:1417-1426.
- 22 de Souza Carvalho, L.M., Lemos, M.C.M., Sanches, E.A. *et al.*: **Improvement of the bioaccessibility of bioactive compounds from Amazon fruits treated using high energy ultrasound.** *Ultrasonics Sonochemistry* 2020, **67**:105148.
- 23• Linhares MdFD, Alves Filho EG, Silva LMA *et al.*: **Thermal and non-thermal processing effect on açai juice composition.** *Food Research International* 2020; **136**:109506.
- In this work, the bioaccessibility of biocompounds is evaluated by applying conventional thermal treatment or emerging technologies, including ultrasound.

- 24 Sun Y, Peng W, Zeng L *et al.*: **Using power ultrasound to release glycosidically bound volatiles from orange juice: A new method.** *Food Chemistry* 2021; **344**:128580.
- 25 Radziejewska-Kubzdela E, Szwengiel A, Ratajkiewicz H, Nowak K: **Effect of ultrasound, heating and enzymatic pre-treatment on bioactive compounds in juice from *Berberis amurensis* Rupr.** *Ultrasonics Sonochemistry* 2020; **63**:104971.
- 26 Zhu W, Ai Y, Fang F, Liao H: **Application of Thermosonication in Red Pitaya Juice Processing: Impacts on Native Microbiota and Quality Properties during Storage.** *Foods* 2021; **10**.
- 27 Yildiz S, Pokhrel PR, Unluturk S, Barbosa-Cánovas GV: **Shelf life extension of strawberry juice by equivalent ultrasound, high pressure, and pulsed electric fields processes.** *Food Research International* 2021; **140**:110040.
- 28 Rodrigues NP, Brochier B, de Medeiros JK *et al.*: **Phenolic profile of sugarcane juice: Effects of harvest season and processing by ohmic heating and ultrasound.** *Food Chemistry* 2021; **347**:129058.
- 29 Bi X, Hemar Y, Balaban MO, Liao X: **The effect of ultrasound on particle size, color, viscosity and polyphenol oxidase activity of diluted avocado puree.** *Ultrasonics Sonochemistry* 2015; **27**:567-575.
- 30 Salleh-Mack SZ, Roberts JS: **Ultrasound pasteurization: The effects of temperature, soluble solids, organic acids and pH on the inactivation of *Escherichia coli* ATCC 25922.** *Ultrasonics Sonochemistry* 2007; **14**:323-329.
- 31 Ferrante S, Guerrero S, Alzamora SM: **Combined Use of Ultrasound and Natural Antimicrobials To Inactivate *Listeria monocytogenes* in Orange Juice.** *Journal of Food Protection* 2007; **70**:1850-1856.
- 32 Hashemi SMB, Jafarpour D: **Ultrasound and malic acid treatment of sweet lemon juice: Microbial inactivation and quality changes.** *Journal of Food Processing and Preservation* 2020; **44**:e14866.
- 33 Ferrario M, Alzamora SM, Guerrero S: **Study of the inactivation of spoilage microorganisms in apple juice by pulsed light and ultrasound.** *Food Microbiology* 2015; **46**:635-642.
- 34 Fonteles TV, Barroso MKdA, Alves Filho EdG *et al.*: **Ultrasound and Ozone Processing of Cashew Apple Juice: Effects of Single and Combined Processing on the Juice Quality and Microbial Stability.** *Processes* 2021; **9**:2243.

- 35 Arroyo C, Cebrián G, Pagán R, Condón S: **Synergistic combination of heat and ultrasonic waves under pressure for *Cronobacter sakazakii* inactivation in apple juice.** *Food Control* 2012; **25**:342-348.
- 36 Evelyn, Kim HJ, Silva FVM: **Modeling the inactivation of *Neosartorya fischeri* ascospores in apple juice by high pressure, power ultrasound and thermal processing.** *Food Control* 2016; **59**:530-537.
- 37 Marx G, Moody A, Bermúdez-Aguirre D: **A comparative study on the structure of *Saccharomyces cerevisiae* under nonthermal technologies: High hydrostatic pressure, pulsed electric fields and thermo-sonication.** *International Journal of Food Microbiology* 2011; **151**:327-337.
- 38 Ugarte-Romero E, Feng H, Martin SE *et al.*: **Inactivation of *Escherichia coli* with Power Ultrasound in Apple Cider.** *Journal of Food Science* 2006; **71**:E102-E108.
- 39 Maghsoudlou Y, Alami M, Mashkour M, Shahraki MH: **Optimization of ultrasound-assisted stabilization and formulation of almond milk.** *Journal of food processing and preservation* 2016; **40**:828-839.
- 40• Lu X, Chen J, Zheng M *et al.*: **Effect of high-intensity ultrasound irradiation on the stability and structural features of coconut-grain milk composite systems utilizing maize kernels and starch with different amylose contents.** *Ultrasonics Sonochemistry* 2019; **55**:135-148.
- Besides studying ultrasound effect on coconut milk, this work adds high-amylose maize starch and evaluate the effect on the mix properties.
- 41• Salve AR, Pegu K, Arya SS: **Comparative assessment of high-intensity ultrasound and hydrodynamic cavitation processing on physico-chemical properties and microbial inactivation of peanut milk.** *Ultrasonics Sonochemistry* 2019; **59**:104728.
- This work evidenced by optical microscopy the globules size reduction by ultrasound. In addition, showed photos of sedimentation index analysis demonstrating ultrasound improvement.
- 42 Aydar EF, Tutuncu S, Ozcelik B: **Plant-based milk substitutes: Bioactive compounds, conventional and novel processes, bioavailability studies, and health effects.** *Journal of Functional Foods* 2020; **70**:103975.
- 43 Rasika DMD, Vidanarachchi JK, Rocha RS *et al.*: **Plant-based milk substitutes as emerging probiotic carriers.** *Current Opinion in Food Science* 2021; **38**:8-20.
- 44 Vanga SK, Wang J, Raghavan V: **Effect of ultrasound and microwave processing on the structure, in-vitro digestibility and trypsin inhibitor activity of soymilk proteins.** *LWT* 2020; **131**:109708.

- 45 Vanga SK, Wang J, Orsat V, Raghavan V: **Effect of pulsed ultrasound, a green food processing technique, on the secondary structure and in-vitro digestibility of almond milk protein.** *Food Research International* 2020; **137**:109523.
- 46 Morales-de la Peña M, Martín-Belloso O, Welti-Chanes J: **High-power ultrasound as pre-treatment in different stages of soymilk manufacturing process to increase the isoflavone content.** *Ultrasonics Sonochemistry* 2018; **49**:154-160.
- 47 Iorio MC, Bevilacqua A, Corbo MR *et al.*: **A case study on the use of ultrasound for the inhibition of Escherichia coli O157:H7 and Listeria monocytogenes in almond milk.** *Ultrasonics Sonochemistry* 2019; **52**:477-483.
- 48• Campaniello D, Bevilacqua A, Speranza B *et al.*: **Inactivation of Salmonella enterica in a Rice Beverage by Ultrasound: Study of the Parameters Affecting the Antibacterial Effect.** *Food and Bioprocess Technology* 2018; **11**:1139-1148.
- In contrast to other works that only study the microbiological inactivation on the beverage by using ultrasound, this work studied microbiological stability of rice milk during storage.
- 49 Kinsella JE, Melachouris N: **Functional properties of proteins in foods: A survey.** *C R C Critical Reviews in Food Science and Nutrition* 1976; **7**:219-280.
- 50 Bakwo Bassogog CB, Nyobe CE, Ngui SP *et al.*: **Effect of heat treatment on the structure, functional properties and composition of Moringa oleifera seed proteins.** *Food Chemistry* 2022; **384**:132546.
- 51 Saricaoglu FT: **Application of high-pressure homogenization (HPH) to modify functional, structural and rheological properties of lentil (Lens culinaris) proteins.** *International Journal of Biological Macromolecules* 2020; **144**:760-769.
- 52 Shen Y, Hong S, Singh G *et al.*: **Improving functional properties of pea protein through “green” modifications using enzymes and polysaccharides.** *Food Chemistry* 2022; **385**:132687.
- 53 Xu X, Qiao Y, Shi B, Dia VP: **Alcalase and bromelain hydrolysis affected physicochemical and functional properties and biological activities of legume proteins.** *Food Structure* 2021; **27**:100178.
- 54 Avelar Z, Vicente AA, Saraiva JA, Rodrigues RM: **The role of emergent processing technologies in tailoring plant protein functionality: New insights.** *Trends in Food Science & Technology* 2021; **113**:219-231.
- 55•• Nikbakht Nasrabadi M, Sedaghat Doost A, Mezzenga R: **Modification approaches of plant-based proteins to improve their techno-functionality and use in food products.** *Food Hydrocolloids* 2021; **118**:106789.

- This interesting review presents and discusses the different modification methods for plant proteins, including their advantages and disadvantages.

56 Arzeni C, Martinez K, Zema P *et al.*: **Comparative study of high intensity ultrasound effects on food proteins functionality.** *Journal of Food Engineering* 2012; **108**:463-472.

57 Du H, Zhang J, Wang S *et al.*: **Effect of high-intensity ultrasonic treatment on the physicochemical, structural, rheological, behavioral, and foaming properties of pumpkin (*Cucurbita moschata* Duch.)-seed protein isolates.** *LWT* 2022; **155**:112952.

58 Zhao Q, Xie T, Hong X *et al.*: **Modification of functional properties of perilla protein isolate by high-intensity ultrasonic treatment and the stability of o/w emulsion.** *Food Chemistry* 2022; **368**:130848.

59• Flores-Jiménez NT, Ulloa JA, Urías-Silvas JE *et al.*: **Influence of high-intensity ultrasound on physicochemical and functional properties of a guamuchil *Pithecellobium dulce* (Roxb.) seed protein isolate.** *Ultrasonics Sonochemistry* 2022; **84**:105976.

- This paper provides a good example of using ultrasound processing to modify proteins, evaluating and correlating the structural and physicochemical changes with functional properties.

60 Hussain M, Qayum A, Zhang X *et al.*: **Improvement in bioactive, functional, structural and digestibility of potato protein and its fraction patatin via ultrasonication.** *LWT* 2021; **148**:111747.

61 Karabulut G, Yemiş O: **Modification of hemp seed protein isolate (*Cannabis sativa* L.) by high-intensity ultrasound treatment. Part 1: Functional properties.** *Food Chemistry* 2022; **375**:131843.

62 Tang S-Q, Du Q-H, Fu Z: **Ultrasonic treatment on physicochemical properties of water-soluble protein from *Moringa oleifera* seed.** *Ultrasonics Sonochemistry* 2021; **71**:105357.

63 Kamani MH, Semwal J, Meera MS: **Functional modification of protein extracted from black gram by-product: Effect of ultrasonication and micronization techniques.** *LWT* 2021; **144**:111193.

64 Lo B, Kasapis S, Farahnaky A: **Effect of low frequency ultrasound on the functional characteristics of isolated lupin protein.** *Food Hydrocolloids* 2022; **124**:107345.

65 Omura MH, de Oliveira APH, Soares LdS *et al.*: **Effects of protein concentration during ultrasonic processing on physicochemical properties and techno-functionality of plant food proteins.** *Food Hydrocolloids* 2021; **113**:106457.

- 66 Zhao R, Liu X, Liu W *et al.*: **Effect of high-intensity ultrasound on the structural, rheological, emulsifying and gelling properties of insoluble potato protein isolates.** *Ultrasonics Sonochemistry* 2022:105969.