



HAL
open science

Direct numerical simulations of turbulent jets: vortex-interface-surfactant interactions

C.R. Constante-Amores, T. Abadie, L. Kahouadji, S. Shin, J. Chergui, D.
Juric, A.A. Castrejon-Pita, O.K. Matar

► **To cite this version:**

C.R. Constante-Amores, T. Abadie, L. Kahouadji, S. Shin, J. Chergui, et al.. Direct numerical simulations of turbulent jets: vortex-interface-surfactant interactions. *Journal of Fluid Mechanics*, 2023, 955, pp.A42. 10.1017/jfm.2022.1056 . hal-03808070v2

HAL Id: hal-03808070

<https://hal.science/hal-03808070v2>

Submitted on 24 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Direct numerical simulations of turbulent jets: vortex-interface-surfactant interactions

C. R. Constante-Amores^{1,2}, T. Abadie¹, L. Kahouadji¹, S. Shin³, J. Chergui⁴, D. Juric^{4,5}, A. A. Castrejon-Pita² and O. K. Matar¹†

¹Department of Chemical Engineering, Imperial College London, London SW7 2AZ, UK

²Department of Engineering Science, University of Oxford, Oxford OX1 3PJ, UK

³Department of Mechanical and System Design Engineering, Hongik University, KR

⁴ Université Paris Saclay, Centre National de la Recherche Scientifique (CNRS), Laboratoire Interdisciplinaire des Sciences du Numérique (LISN), 91400 Orsay, France

⁵Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge CB3 0WA, UK

(Received xx; revised xx; accepted xx)

We study the effect of insoluble surfactants on the spatio-temporal evolution of turbulent jets. We use three-dimensional numerical simulations and employ an interface-tracking/level-set method that accounts for surfactant-induced Marangoni stresses. The present study builds on our previous work (Constante-Amores *et al.*, 2021, *J. Fluid Mech.*, **922**, A6) in which we examined in detail the vortex-surface interaction in the absence of surfactants. Numerical solutions are obtained for a wide range of Weber and elasticity numbers in which vorticity production is generated by surface deformation and surfactant-induced Marangoni stresses. The present work demonstrates, for the first time, the crucial role of Marangoni stresses, brought about by surfactant concentration gradients, in the formation of coherent, hairpin-like vortex structures. These structures have a profound influence on the development of the three-dimensional interfacial dynamics. We also present theoretical expressions for the mechanisms that influence the rate of production of circulation in the presence of surfactants for a general, three-dimensional, two-phase flow and highlight the dominant contribution surfactant-induced Marangoni stresses.

1. Introduction

The atomisation of a liquid jet has driven interest in the fluid mechanics community because of its occurrence in both natural and industrial applications (e.g., propellant combustion, pharmaceutical sprays, etc.). The process results in a ‘cascade mechanism’ for fluid fragmentation (Plateau 1873; Eggers 1997; Marmottant & Villermaux 2004; Constante-Amores *et al.* 2020a): from the growth of linear modes through a Kelvin-Helmholtz instability to the development of nonlinearities leading to capillary breakup events via long filament pinch-off that can be modulated by a Rayleigh-Plateau instability or controlled by an ‘end-pinching’ mechanism. The understanding of the interfacial dynamics relies on the characterisation of the vortex-interface interactions. For instance, Jarrahbashi *et al.* (2016), Zandian *et al.* (2018, 2019) and Constante-Amores *et al.* (2021a) reported that their interplay determines the interfacial dynamics for turbulent jets; Hoepffner & Paré (2013) showed that vorticity production results in a change in the capillary retraction of a liquid thread. Theoretically, Longuet-Higgins (1992), Wu (1995), Lundgren & Koumoutsakos (1999) demonstrated that vorticity production depends on the

† Email address for correspondence: o.matar@imperial.ac.uk

velocity field and the interfacial curvature for the condition of zero shear stress at a free surface. Additionally, Brøns *et al.* (2014) and Terrington *et al.* (2020, 2021) extended the previous results to show that interfacial curvature effects, viscosity and density difference across the interface are the only mechanisms driving vorticity production. Recently, Fuster & Rossi (2021) also demonstrated the role of interfacial curvature and density differences across the interface with identical dynamical viscosity via two-dimensional, non-axisymmetric numerical studies.

We note that the studies mentioned in the foregoing involve a constant surface tension and therefore do not support the formation of Marangoni gradients. Liquid streams, however, are invariably contaminated with surface-active-agents (surfactants), deliberately-placed or naturally-occurring, which give rise to surface tension gradients, and subsequently Marangoni-induced flow (Manikantan & Squires 2020). While the atomisation of uncontaminated liquid jets has received significant attention in the literature (Herrmann 2010; Desjardins & Pitsch 2010; Jarrahbashi & Sirignano 2014; Jarrahbashi *et al.* 2016; Zandian *et al.* 2018, 2019; Constante-Amores *et al.* 2020*b*, 2021*a*), the effect of surfactant on their dynamics remains far less studied. The multi-scale nature of the flow, and the complex coupling between the surfactant concentration fields and interfacial topology complicate its experimental scrutiny. This can be alleviated via the use of high-fidelity simulations which can unravel the delicate interplay among the different physical mechanisms across the relevant scales.

Through the use of state-of-the-art imaging techniques, Kooij *et al.* (2018), Sijs & Bonn (2020), and Sijs *et al.* (2021) showed that the presence of surfactants influences the interfacial fragmentation during atomisation and decreases the mean-droplet size in agreement with Ellis *et al.* (2001) and Ariyapadi *et al.* (2004). All the previous studies, however, have not reported the role of Marangoni stresses which the present paper will address for the case of an insoluble surfactant. Although the presence of surfactants can also induce both shear and dilatational surface rheological effects (discussed below), these effects will not be considered in this study. Nonetheless, we will use transient numerical simulations to demonstrate that the Marangoni stresses influence the production of vorticity near the interface, and modify the interface-vortex interactions and the three-dimensional destabilisation of the jet. In order to focus on the role of Marangoni stresses in the jet dynamics, we will study the case of a jet of one fluid issuing into another characterised by equal densities and viscosities.

There has been significant scientific interest in studying the role of surfactants in the destabilization and fragmentation of non-turbulent liquid jets of pure Newtonian fluids (see for example Eggers (1993); Lister & Stone (1998); Craster *et al.* (2002); Liao *et al.* (2004); Craster *et al.* (2009)). Those authors have shown the existence of multiple intermediate or transient scaling regimes which are not altered by the presence of surfactants as they are convected away from the pinch-off region. However, McGough & Basaran (2006) and Kamat *et al.* (2018) showed the formation of micro threads, which connect drops during the surfactant-induced thinning. Additionally, the presence of surfactants not only give rise to gradients in surface tension and hence tangential interfacial stresses, but also induce both shear and dilatational surface rheological effects. Recently, work by Wee *et al.* (2021) and Martínez-Calvo & Sevilla (2020) have analysed theoretically the influence of surface viscosities on the pinch-off dynamics of a jet of an incompressible Newtonian liquid that is surrounded by a passive gas.

The rest of this paper is structured as follows: in Section 2, the problem formulation, governing dimensionless parameters, and numerical method are introduced. Section 3 provides a discussion of the results, and concluding remarks are given in Section 4.




Figure 1: (a) Initial interfacial shape, highlighting the computational domain of size $(5D)^3$ in a three-dimensional Cartesian space $\mathbf{x} = (x, y, z)$; (b) schematic representation of the problem in the $x - y$ ($z = 2.5D$) plane showing the initial ($t = 0$) streamwise velocity profile, u_x , and a representation of a monolayer of an insoluble surfactant.

2. Problem formulation and numerical method

Since the aim here is to shed light on the different mechanisms that influence the production of vorticity near the interface in the presence of surfactants, we present a general theoretical description of vorticity and circulation in a three-dimensional control volume enclosing an interface using Lighthill's and Lyman's flux definitions Terrington *et al.* (2021). We also provide a brief description of the numerical technique which is used to carry out the computations. Finally, we provide motivation for the choice of physical and physico-chemical parameters made in the present work.

2.1. Problem formulation and numerical method

Figure 1 shows a representation of the flow configuration considered in this study in a three-dimensional Cartesian domain $\mathbf{x} = (x, y, z)$: a liquid segment is initialised as a cylinder of diameter D , with a finite length, i.e. $5D$, in the positive x –(streamwise) direction. Such an approach has been used by Desjardins & Pitsch (2010), Jarrahbashi *et al.* (2016), and Zandian *et al.* (2018) for planar and cylindrical jets. Appendix A shows the effect of varying the domain size. We will focus on the case of insoluble surfactants, which enables us to isolate the surfactant-induced Marangoni dynamics during the atomisation of the jet. We acknowledge, however, that experimental studies feature soluble surfactants which are dissolved in the liquid that issues from a nozzle to form the jet and that the sorption kinetics control the surfactant interfacial concentration adding extra layers of complexity.

The dimensional governing equations, which can be found in the work of Shin *et al.* (2018), are rendered dimensionless using the following scalings:

$$\tilde{\mathbf{x}} = \frac{\mathbf{x}}{D}, \quad \tilde{t} = \frac{t}{t_r}, \quad \tilde{\mathbf{u}} = \frac{\mathbf{u}}{U}, \quad \tilde{p} = \frac{p}{\rho U^2}, \quad \tilde{\sigma} = \frac{\sigma}{\sigma_s}, \quad \tilde{\Gamma} = \frac{\Gamma}{\Gamma_\infty}, \quad (2.1)$$

where, t , \mathbf{u} , and p stand for time, velocity, and pressure, respectively; here, the dimensionless variables are designated using tildes. The physical parameters correspond to the liquid density ρ , viscosity, μ , surface tension, σ , surfactant-free surface tension, σ_s , initial jet diameter, D , and injection velocity, U . Hence, the characteristic time scale based on

the injection velocity is $t_r = D/U$. The interfacial surfactant concentration, Γ , is scaled with the saturation interfacial concentration, Γ_∞ .

Using the relations in Eq. (2.1), the dimensionless form of the continuity and momentum equations is respectively expressed as:

$$\nabla \cdot \tilde{\mathbf{u}} = 0, \quad (2.2)$$

$$\begin{aligned} \tilde{\rho} \left(\frac{\partial \tilde{\mathbf{u}}}{\partial \tilde{t}} + \tilde{\mathbf{u}} \cdot \nabla \tilde{\mathbf{u}} \right) &= -\nabla \tilde{p} + \frac{1}{Re} \nabla \cdot [\tilde{\mu}(\nabla \tilde{\mathbf{u}} + \nabla \tilde{\mathbf{u}}^T)] \\ &+ \frac{1}{We} \int_{\tilde{A}(\tilde{t})} (\tilde{\sigma} \tilde{\kappa} \hat{\mathbf{s}} + \nabla_s \tilde{\sigma}) \delta(\tilde{\mathbf{x}} - \tilde{\mathbf{x}}_f) d\tilde{A}, \end{aligned} \quad (2.3)$$

where $\tilde{\kappa}$ represents the interface curvature, ∇_s the surface gradient operator, and $\hat{\mathbf{s}}$ the outward-pointing unit normal to the interface. Here, $\tilde{\mathbf{x}}_f$ is the parametrization of the time-dependent interface area $\tilde{A}(\tilde{t})$, where $\delta(\tilde{\mathbf{x}} - \tilde{\mathbf{x}}_f)$ is the three-dimensional Dirac delta function. The density, $\tilde{\rho}$, and viscosity, $\tilde{\mu}$, are given by the following expressions

$$\tilde{\rho}(\tilde{\mathbf{x}}, \tilde{t}) = \frac{\rho_g}{\rho_l} + \left(1 - \frac{\rho_g}{\rho_l}\right) H(\tilde{\mathbf{x}}, \tilde{t}), \quad \tilde{\mu}(\tilde{\mathbf{x}}, \tilde{t}) = \frac{\mu_g}{\mu_l} + \left(1 - \frac{\mu_g}{\mu_l}\right) H(\tilde{\mathbf{x}}, \tilde{t}), \quad (2.4)$$

where $H(\tilde{\mathbf{x}}, \tilde{t})$ represents a smoothed Heaviside function; this is zero in the gas phase and unity in the liquid phase, while the subscripts l and g designate the individual liquid and gas phases, respectively.

The dimensionless surfactant transport is given by:

$$\frac{\partial \tilde{\Gamma}}{\partial \tilde{t}} + \nabla_s \cdot (\tilde{\Gamma} \tilde{\mathbf{u}}_t) = \frac{1}{Pe_s} \nabla_s^2 \tilde{\Gamma}, \quad (2.5)$$

where $\tilde{\mathbf{u}}_t = (\tilde{\mathbf{u}}_s \cdot \mathbf{t})\mathbf{t}$ is the tangential velocity vector in which $\tilde{\mathbf{u}}_s$ is the surface velocity and \mathbf{t} is the unit tangent to the interface.

The scaling results in the following dimensionless groups:

$$Re = \frac{\rho U D}{\mu}, \quad We = \frac{\rho U^2 D}{\sigma_s}, \quad Pe_s = \frac{U D}{\mathcal{D}_s}, \quad \beta_s = \frac{\Re T \Gamma_\infty}{\sigma_s}, \quad (2.6)$$

where Re , We , and Pe_s denote the Reynolds, Weber, and (interfacial) Peclet numbers, respectively, while β_s is a surfactant elasticity number which represents a measure of the sensitivity of σ to Γ ; here, \Re is the ideal gas constant value $8.314 \text{ J K}^{-1} \text{ mol}^{-1}$, T denotes temperature and \mathcal{D}_s refers to the diffusion coefficient.

To describe the relation between $\tilde{\sigma}$ and $\tilde{\Gamma}$, we use the non-linear Langmuir equation:

$$\tilde{\sigma} = 1 + \beta_s \ln(1 - \tilde{\Gamma}). \quad (2.7)$$

Surface tension gradients are expressed as a function of $\tilde{\Gamma}$ as

$$\nabla_s \tilde{\sigma} / We = -Ma / (1 - \tilde{\Gamma}) \nabla_s \tilde{\Gamma}, \quad (2.8)$$

where $Ma = \beta_s / We = \Re T \Gamma_\infty / \rho U^2 D$ is a Marangoni parameter.

The three-dimensional numerical simulations were performed by solving the two-phase Navier-Stokes equations in the Cartesian domain $\mathbf{x} = (x, y, z)$. A hybrid front-tracking/level-set method was used to treat the interface where surfactant transport was resolved in the plane of the interface (Shin *et al.* 2018). The simulations are initialised with a turbulent velocity profile in the liquid jet segment (i.e., $u(r) = 15/14 U(1 - (r/(D/2))^{28})$ (Constante-Amores *et al.* 2021a). Solutions are sought subject to Neumann boundary

conditions on all variables at the lateral boundaries, and periodic boundary conditions in the x –(streamwise) direction. The computational domain is a cube with dimensions $(5D)^3$ globally resolved by a uniform grid of $(786)^3$ cells; see Appendix of Constante-Amores *et al.* (2021a) for details of mesh-refinement studies and validation of the numerical method. This method has also been widely tested for surfactant-laden flows (Shin *et al.* 2018; Constante-Amores *et al.* 2020a, 2021b, 2022; Batchvarov *et al.* 2021) and the numerical simulations in this study conserve fluid volume and surfactant mass with a relative error of less than $10^{-3}\%$.

Next, we motivate the values of material properties by looking into the sources for vorticity production at an interface in a three-dimensional framework. These sources are due to differences in density (i.e., baroclinic effect) and viscosity, surface tension forces (due to gradients of curvature along the interface), and Marangoni stresses. Thus, to unravel the importance of the surfactant-induced Marangoni stresses on the vortex-surface-surfactant interactions, we focus on situations in which surface tension forces and Marangoni stresses are the only physical mechanisms responsible for vorticity production at the interface, i.e., the jump in material properties across the interface is zero (Fuster & Rossi 2021). This is a realistic assumption for immiscible liquid-liquid systems exemplified by the silicone oil-water pairing used by Ibarra (2017) and Ibarra *et al.* (2020) in their two-phase, stratified pipe flow experiments.

The values of the dimensionless quantities are consistent with experimentally-realizable systems and are chosen to ensure a full coupling between surfactant-induced Marangoni stresses and interfacial diffusion, and inertia. We set $Re = 5000$ to ensure a rich dynamics (Constante-Amores *et al.* 2021a) and focus on the range $50 < We < 1000$ to account for realistic values of σ_s , i.e. $\mathcal{O}(10^{-3}) < \sigma_s < \mathcal{O}(10^{-1})$ N m⁻¹. The parameter β_s is related to Γ_∞ and therefore the critical micelle concentration (CMC), i.e. $\Gamma_\infty \sim \mathcal{O}(10^{-6})$ mol m⁻² for NBD-PC (1-palmitoyl-2-12-[(7-nitro-2-1,3-benzoxadiazol-4-yl)amino]dodecanoyl-sn-glycero-3-phosphocholine) (Strickland *et al.* 2015); thus, we have explored the range of $0.1 < \beta_s < 0.9$ which corresponds to CMC in the range $\mathcal{O}(10^{-7}) < \text{CMC} < \mathcal{O}(10^{-6})$ mol m⁻², for typical values of σ_s . We have set $Pe_s = 10^2$ following Batchvarov *et al.* (2020) and Constante-Amores *et al.* (2020a) who showed that the interfacial dynamics are weakly-dependent on Pe_s beyond this value.

2.2. Vorticity and circulation

This section aims to present a general description of vorticity generation in a three-dimensional framework. We present a theoretical formulation which builds upon the inviscid theory presented by Morton (1984) for near-interface vorticity generation in three dimensions. For inviscid fluids, the rate of generation of vorticity is a result of the relative tangential acceleration of fluid on each side of the interface, which is caused by tangential pressure gradients or body forces. The present theoretical formulation is expressed as a conservation law for circulation in a control volume that includes a general surface. The total circulation is expressed as the vorticity from the fluids from both sides of the interface as well as circulation contained in the interface.

It is well known that curvature induces the generation of vorticity as the normal viscous stress at an interface is balanced by the capillary pressure. However, the presence of surfactant leads to a reduction in surface tension, which influences this mechanism. Furthermore, surfactant interfacial concentration variations induce surface tension gradients, and, as we will show, lead to a new route for vorticity generation near the interface. Once we have presented our theoretical expressions for a general three-dimensional surface, we will simplify them for the limiting case in which the jump in the tangential and normal components of the velocity across the interface vanish; this is the




Figure 2: Schematic showing a volume V with a surface ∂V which encloses two fluids separated by an interface surface I . Here, the two smaller control volumes V_1 and V_2 refer to the control volume of each fluid. Local unit vectors to the interface are $\hat{\mathbf{b}}$, $\hat{\mathbf{s}}$ and $\hat{\mathbf{t}}$; $\hat{\mathbf{n}}$ corresponds to the unit normal vector to the control volume ∂V , $\hat{\mathbf{b}}$ is a vector tangent to I , but orthogonal to ∂I , and $\hat{\mathbf{t}}$ is the unit tangent vector to the boundary curve ∂I .

case for identical material properties such as density and viscosity. This assumption will help to shed some light on the crucial role of the Marangoni-induced vorticity generation mentioned above. Future studies should extend our work to situations featuring density and viscosity contrasts.

In order to examine the effect of the surfactant on the vorticity near the interface, we consider a fixed, three-dimensional (3D) control volume V bounded by a closed surface of area ∂V with an outward-pointing unit normal $\hat{\mathbf{n}}$ (see figure 2). This volume encloses regions of the incompressible fluids 1 and 2, of volumes, V_1 and V_2 , separated by an interfacial surface I whose intersection with V defines the curve ∂I . The vector $\hat{\mathbf{s}}$ is the outward-pointing unit normal to the surface I while $\hat{\mathbf{t}}$ and $\hat{\mathbf{b}}$ are two orthogonal unit tangent vectors to the interface. We proceed below using dimensional variables and then apply the scalings in equation 2.1 to render the final equations dimensionless.

For fluid ‘i’, it is possible to write down expressions for $\omega_{b,i}$ and $\omega_{t,i}$, which represent the components of the vorticity $\boldsymbol{\omega}_i$ in the $\hat{\mathbf{b}}$ and $\hat{\mathbf{t}}$ directions, respectively:

$$\omega_{b,i} = (\hat{\mathbf{s}} \times \hat{\mathbf{t}}) \cdot (\nabla \times \mathbf{u}_i), \quad (2.9)$$

$$\omega_{t,i} = (\hat{\mathbf{s}} \times \hat{\mathbf{b}}) \cdot (\nabla \times \mathbf{u}_i), \quad (2.10)$$

where \mathbf{u}_i denotes the velocity fields. These expressions may be recast as follows †

$$\omega_{b,i} = \hat{\mathbf{s}} \cdot \nabla \mathbf{u}_i \cdot \hat{\mathbf{t}} - \hat{\mathbf{t}} \cdot \nabla \mathbf{u}_i \cdot \hat{\mathbf{s}}, \quad (2.11)$$

$$\omega_{t,i} = \hat{\mathbf{s}} \cdot \nabla \mathbf{u}_i \cdot \hat{\mathbf{b}} - \hat{\mathbf{b}} \cdot \nabla \mathbf{u}_i \cdot \hat{\mathbf{s}}. \quad (2.12)$$

In the presence of interfacial stresses arising from gradients of surface tension σ due to

† Using $(\mathbf{a} \times \mathbf{b}) \cdot (\mathbf{c} \times \mathbf{d}) = (\mathbf{a} \cdot \mathbf{c})(\mathbf{b} \cdot \mathbf{d}) - (\mathbf{a} \cdot \mathbf{d})(\mathbf{b} \cdot \mathbf{c})$, valid for any vector \mathbf{a} , \mathbf{b} , \mathbf{c} , and \mathbf{d} .

surfactant concentration gradients, the interfacial shear stress conditions are given by

$$[[\hat{\mathbf{t}} \cdot \mathbf{T} \cdot \hat{\mathbf{s}}]] = -\hat{\mathbf{t}} \cdot \nabla \sigma, \quad (2.13)$$

$$[[\hat{\mathbf{b}} \cdot \mathbf{T} \cdot \hat{\mathbf{s}}]] = -\hat{\mathbf{b}} \cdot \nabla \sigma, \quad (2.14)$$

$[[q]] = q_2 - q_1$ represents the jump across the interface of a quantity q , $\mathbf{T}_i = -p_i + \mu_i \mathbf{D}_i$ is the total stress in fluid 'i' in which p_i is the pressure, $\mathbf{D}_i = (\nabla \mathbf{u}_i + \nabla \mathbf{u}_i^T)/2$ is the rate of deformation tensor, and μ_i denote the viscosities, whence

$$[[\mu (\hat{\mathbf{t}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}} + \hat{\mathbf{s}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{t}})]] = -2\hat{\mathbf{t}} \cdot \nabla \sigma, \quad (2.15)$$

$$[[\mu (\hat{\mathbf{b}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}} + \hat{\mathbf{s}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{b}})]] = -2\hat{\mathbf{b}} \cdot \nabla \sigma. \quad (2.16)$$

Substitution of these results into Eqs. (2.11) and (2.12) yields

$$[[\mu (\omega_b + 2\hat{\mathbf{t}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}})]] = -2\hat{\mathbf{t}} \cdot \nabla \sigma, \quad (2.17)$$

$$[[\mu (\omega_t + 2\hat{\mathbf{b}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}})]] = -2\hat{\mathbf{b}} \cdot \nabla \sigma. \quad (2.18)$$

For the case $[[\mu]] = 0$, which is the focus of this paper, we obtain

$$[[w_b]] = -\frac{2}{\mu} \nabla \sigma \cdot \hat{\mathbf{t}} - 2 [[\hat{\mathbf{t}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}}]], \quad (2.19)$$

$$[[w_t]] = -\frac{2}{\mu} \nabla \sigma \cdot \hat{\mathbf{b}} - 2 [[\hat{\mathbf{b}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}}]], \quad (2.20)$$

where $\mu_2 = \mu_1 = \mu$. Noting that $\hat{\mathbf{t}} \cdot \nabla = \partial/\partial s$ and $\hat{\mathbf{b}} \cdot \nabla = \partial/\partial b$, it can be shown that

$$[[w_b]] = -\frac{2}{\mu} \frac{\partial \sigma}{\partial s} - 2 \left[\frac{\partial}{\partial s} [[\mathbf{u} \cdot \hat{\mathbf{s}}]] - \kappa_1 [[\mathbf{u} \cdot \hat{\mathbf{t}}]] \right], \quad (2.21)$$

$$[[w_t]] = -\frac{2}{\mu} \frac{\partial \sigma}{\partial b} - 2 \left[\frac{\partial}{\partial b} [[\mathbf{u} \cdot \hat{\mathbf{s}}]] - \kappa_2 [[\mathbf{u} \cdot \hat{\mathbf{b}}]] \right], \quad (2.22)$$

where the curvatures κ_1 and κ_2 are defined as follows

$$\kappa_1 = \hat{\mathbf{t}} \cdot \frac{\partial \hat{\mathbf{s}}}{\partial s}, \quad \kappa_2 = \hat{\mathbf{b}} \cdot \frac{\partial \hat{\mathbf{s}}}{\partial b}. \quad (2.23)$$

From continuity of the normal and tangential components of the velocity at the interface, i.e., $[[\mathbf{u} \cdot \hat{\mathbf{s}}]] = 0$, and $[[\mathbf{u} \cdot \hat{\mathbf{t}}]] = [[\mathbf{u} \cdot \hat{\mathbf{b}}]] = 0$, respectively, it is seen that the interfacial jumps in the vorticity components are directly related to the Marangoni stresses:

$$[[w_b]] = -\frac{2}{\mu} \frac{\partial \sigma}{\partial s}, \quad (2.24)$$

$$[[w_t]] = -\frac{2}{\mu} \frac{\partial \sigma}{\partial b}. \quad (2.25)$$

We now consider the circulation vector $\boldsymbol{\Omega}$ for 3D flows given by

$$\boldsymbol{\Omega} = \int_V \boldsymbol{\omega} dV, \quad (2.26)$$

for the fixed 3D control volume V shown in figure 2. The 3D vorticity equation is given by

$$\frac{\partial \boldsymbol{\omega}}{\partial t} + \nabla \cdot (\mathbf{u}\boldsymbol{\omega}) = \nabla \cdot (\boldsymbol{\omega}\mathbf{u}) + \nu \nabla^2 \boldsymbol{\omega}, \quad (2.27)$$

and the total rate of change of $\boldsymbol{\Omega}$ is then expressed by

$$\begin{aligned} \frac{D\boldsymbol{\Omega}}{Dt} &= \int_V \frac{D\boldsymbol{\omega}}{Dt} dV = \frac{D}{Dt} \int_V \boldsymbol{\omega} dV = \int_V \nabla \cdot (\boldsymbol{\omega}\mathbf{u} + \nu\nabla\boldsymbol{\omega}) dV \\ &= \int_{\partial V} \hat{\mathbf{n}} \cdot (\boldsymbol{\omega}\mathbf{u}) dS + \int_{\partial V} \hat{\mathbf{n}} \cdot (\nu\nabla\boldsymbol{\omega}) dS. \end{aligned} \quad (2.28)$$

The first term on the RHS of Eq. (2.28) corresponds to vortex stretching/tilting and is present only in 3D. We now write

$$\begin{aligned} \frac{D}{Dt} \int_{V_1UV_2} \boldsymbol{\omega} dV &= \oint_{\partial V_1} \hat{\mathbf{n}} \cdot (\boldsymbol{\omega}\mathbf{u} + \nu\nabla\boldsymbol{\omega}) dS + \oint_{\partial V_2} \hat{\mathbf{n}} \cdot (\boldsymbol{\omega}\mathbf{u} + \nu\nabla\boldsymbol{\omega}) dS \\ &\quad + \oint_{\partial V'_1} \hat{\mathbf{n}} \cdot (\boldsymbol{\omega}_1\mathbf{u}_1 + \nu_1\nabla\boldsymbol{\omega}_1) dS + \oint_{\partial V'_2} \hat{\mathbf{n}} \cdot (\boldsymbol{\omega}_2\mathbf{u}_2 + \nu_2\nabla\boldsymbol{\omega}_2) dS, \end{aligned} \quad (2.29)$$

and let $V_1UV_2 \rightarrow V$, $\hat{\mathbf{n}} \rightarrow \hat{\mathbf{s}}$ from fluid 1, $\hat{\mathbf{n}} \rightarrow -\hat{\mathbf{s}}$ from fluid 2, and $(\partial V_1, \partial V_2) \rightarrow I$, then it follows that

$$\frac{D}{Dt} \int_V \boldsymbol{\omega} dV = \oint_{\partial V} \hat{\mathbf{n}} \cdot (\boldsymbol{\omega}\mathbf{u} + \nu\nabla\boldsymbol{\omega}) dS - \left[\oint_I [[\hat{\mathbf{s}} \cdot (\boldsymbol{\omega}\mathbf{u})]] dS + \oint_I [[\nu\hat{\mathbf{s}} \cdot \nabla\boldsymbol{\omega}]] dS \right]. \quad (2.30)$$

It is important to establish a connection between $\oint_I [[\nu\hat{\mathbf{s}} \cdot \nabla\boldsymbol{\omega}]] dS$, which represents the jump across the plane of the interface of the vorticity flux, and the momentum conservation equation given by

$$\frac{D\mathbf{u}}{Dt} = -\frac{\nabla p}{\rho} - \nu\nabla \times \boldsymbol{\omega}. \quad (2.31)$$

In order to relate this term to the $\nu\nabla \times \boldsymbol{\omega}$ term in Eq. (2.31), we first write down the following general result †

$$\begin{aligned} -\oint_{\partial V} \hat{\mathbf{s}} \cdot \nabla\boldsymbol{\omega} dS &= -\int_V \nabla^2\boldsymbol{\omega} dV = -\int_V (\nabla(\nabla \cdot \boldsymbol{\omega}) - \nabla \times \nabla \times \boldsymbol{\omega}) dV = \int_V \nabla \times \nabla \times \boldsymbol{\omega} dV \\ &= -\oint_{\partial V} (\nabla \times \boldsymbol{\omega}) \times \hat{\mathbf{s}} dS = \oint_{\partial V} \hat{\mathbf{s}} \times \nabla \times \boldsymbol{\omega} dS. \end{aligned} \quad (2.32)$$

Note that this relation links *Lighthill's* vorticity flux to *Lyman's* flux, the latter being another form of the former (see Terrington *et al.* (2021) and references therein).

Inspired by the form of Lyman's flux, the natural way to proceed is to take the cross product of $\hat{\mathbf{s}} = \hat{\mathbf{t}} \times \hat{\mathbf{b}}$ with the LHS of Eq. (2.31) and its pressure gradient term ‡ and a cross product of $\hat{\mathbf{s}}$ with its $\nu\nabla \times \boldsymbol{\omega}$ term to arrive at

$$\begin{aligned} -\nu\hat{\mathbf{s}} \times \nabla \times \boldsymbol{\omega} &= \hat{\mathbf{b}}\hat{\mathbf{t}} \cdot \frac{D\mathbf{u}}{Dt} - \hat{\mathbf{b}}\hat{\mathbf{t}} \cdot \nabla \left(\frac{p}{\rho} \right) \\ &= \hat{\mathbf{b}} \left[\left(\frac{D}{Dt} (\mathbf{u} \cdot \hat{\mathbf{t}}) - \mathbf{u} \cdot \frac{D\hat{\mathbf{t}}}{Dt} \right) + \hat{\mathbf{t}} \cdot \nabla \left(\frac{p}{\rho} \right) \right] \\ &= \nu\hat{\mathbf{s}} \cdot \nabla\boldsymbol{\omega}; \end{aligned}$$

† We have used the vector identity $\int_V \nabla \times \mathbf{A} dV = -\oint_{\partial V} \mathbf{A} \times d\mathbf{S} = -\oint_{\partial V} \mathbf{A} \times \mathbf{n} dS = \oint_{\partial V} \mathbf{n} \times \mathbf{A} dS$, for any vector \mathbf{A} , and volume V enclosed by a surface ∂V with a unit normal \mathbf{n} .

‡ We have exploited the fact that $\hat{\mathbf{t}} \times \hat{\mathbf{b}} \times \mathbf{c} = \hat{\mathbf{b}}(\hat{\mathbf{t}} \cdot \mathbf{c}) - \mathbf{c}(\hat{\mathbf{t}} \cdot \hat{\mathbf{b}}) = \hat{\mathbf{b}}(\hat{\mathbf{t}} \cdot \mathbf{c})$ since $\hat{\mathbf{t}} \cdot \hat{\mathbf{b}} = 0$.

here, we note that the sources of vorticity are due to acceleration in the plane of the interface, which we can think of as a vortex sheet, and interfacial pressure gradients. Making use of this relation in Eq. (2.30), we arrive at

$$\begin{aligned} \frac{D}{Dt} \left[\int_V \boldsymbol{\omega} dV + \hat{\mathbf{b}} \oint_I [[\mathbf{u} \cdot \hat{\mathbf{t}}]] dS \right] &= \oint_{\partial V} \hat{\mathbf{n}} \cdot (\boldsymbol{\omega} \mathbf{u} + \nu \nabla \boldsymbol{\omega}) dS \\ &\quad - \oint_I [[\hat{\mathbf{s}} \cdot (\boldsymbol{\omega} \mathbf{u})]] dS + \oint_I \hat{\mathbf{b}} [[\mathbf{u} \cdot \frac{D\hat{\mathbf{t}}}{Dt}]] dS - \oint_I \hat{\mathbf{b}} \frac{\partial}{\partial s} \left[\frac{p}{\rho} \right] dS, \end{aligned} \quad (2.33)$$

where we have set $\hat{\mathbf{t}} \cdot \nabla(p/\rho) = \partial(p/\rho)/\partial s$. An expression for $\mathbf{u} \cdot (D\hat{\mathbf{t}}/Dt)$ can be developed given by (the details are in Appendix B)

$$\mathbf{u} \cdot \frac{D\hat{\mathbf{t}}}{Dt} = \frac{1}{2} \frac{\partial}{\partial s} \left[(\mathbf{u} \cdot \hat{\mathbf{s}})^2 + (\mathbf{u} \cdot \hat{\mathbf{b}})^2 \right] + \frac{1}{2} \frac{\partial}{\partial b} \left[(\mathbf{u} \cdot \hat{\mathbf{s}})^2 + (\mathbf{u} \cdot \hat{\mathbf{b}})^2 \right] - \kappa_1 (\mathbf{u} \cdot \hat{\mathbf{t}}) (\mathbf{u} \cdot \hat{\mathbf{s}}). \quad (2.34)$$

Furthermore, for $[[p]] = 0$, the remaining term required to close equation 2.33 is one for $[[p]]$ (the details are in Appendix C):

$$[[p]] = -\sigma(\kappa_1 + \kappa_2) - 2[[\mu] \left(\frac{\partial}{\partial s} [(\mathbf{u} \cdot \hat{\mathbf{t}}) + (\mathbf{u} \cdot \hat{\mathbf{b}})] + (\kappa_1 + \kappa_2)(\mathbf{u} \cdot \hat{\mathbf{s}}) \right)]. \quad (2.35)$$

To collapse these equations to their two-dimensional (2D) equivalents, we first note that $\hat{\mathbf{s}} \cdot \boldsymbol{\omega} = \hat{\mathbf{n}} \cdot \boldsymbol{\omega} = \mathbf{u} \cdot \hat{\mathbf{b}} = 0$ in 2D, and set $\partial/\partial b = 0$; the latter leads to $\kappa_2 = 0$. We then take a dot product of Eq. (2.33) with $\hat{\mathbf{b}}$ (and convert the volume and area integrals to area and line integrals, respectively) to arrive at a 2D analogue involving the vorticity scalar ω . Moreover, in the case studied here, characterised by $[[\mu]] = 0$, $[[\mathbf{u} \cdot \hat{\mathbf{s}}]] = 0$, $[[\mathbf{u} \cdot \hat{\mathbf{t}}]] = 0$, and $[[\mathbf{u} \cdot \hat{\mathbf{b}}]] = 0$, equation (2.33) reduces to

$$\frac{D}{Dt} \left[\int_V \boldsymbol{\omega} dV \right] = \oint_{\partial V} \hat{\mathbf{n}} \cdot (\boldsymbol{\omega} \mathbf{u} + \nu \nabla \boldsymbol{\omega}) dS - \oint_I [[\hat{\mathbf{s}} \cdot (\boldsymbol{\omega} \mathbf{u})]] dS + \frac{1}{\rho} \oint_I \hat{\mathbf{b}} \frac{\partial}{\partial s} (\sigma [\kappa_1 + \kappa_2]) dS. \quad (2.36)$$

We note that the term involving $[[\hat{\mathbf{s}} \cdot (\boldsymbol{\omega} \mathbf{u})]]$ on the right-hand-side of this equation is zero. To see this, we first note that $[[\hat{\mathbf{s}} \cdot \boldsymbol{\omega} \mathbf{u}]]$ can be re-expressed as

$$\begin{aligned} [[\hat{\mathbf{s}} \cdot \boldsymbol{\omega} \mathbf{u}]] &= (\hat{\mathbf{s}} \cdot \boldsymbol{\omega}_2) \mathbf{u}_2 - (\hat{\mathbf{s}} \cdot \boldsymbol{\omega}_1) \mathbf{u}_1 \\ &= (\hat{\mathbf{s}} \cdot \boldsymbol{\omega}_2 - \hat{\mathbf{s}} \cdot \boldsymbol{\omega}_1) \mathbf{u}_1 \\ &= (\hat{\mathbf{s}} \cdot \boldsymbol{\omega}_2 - \hat{\mathbf{s}} \cdot \boldsymbol{\omega}_1) \mathbf{u}_2 \\ &= [[\hat{\mathbf{s}} \cdot \boldsymbol{\omega}]] \mathbf{u}_1 = [[\hat{\mathbf{s}} \cdot \boldsymbol{\omega}]] \mathbf{u}_2, \end{aligned} \quad (2.37)$$

since $[[\mathbf{u}]] = 0$. We also note that $\hat{\mathbf{s}} \cdot \boldsymbol{\omega} = (\hat{\mathbf{b}} \times \hat{\mathbf{t}}) \cdot (\nabla \times \mathbf{u})$, which can be re-written as

$$\begin{aligned} \hat{\mathbf{s}} \cdot \boldsymbol{\omega} &= \hat{\mathbf{b}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{t}} - \hat{\mathbf{t}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{b}} \\ &= \hat{\mathbf{b}} \cdot \frac{\partial \mathbf{u}}{\partial s} - \hat{\mathbf{t}} \cdot \frac{\partial \mathbf{u}}{\partial b} \\ &= \frac{\partial}{\partial s} (\hat{\mathbf{b}} \cdot \mathbf{u}) - \frac{\partial}{\partial b} (\hat{\mathbf{t}} \cdot \mathbf{u}), \end{aligned} \quad (2.38)$$

since $\hat{\mathbf{b}} \neq \hat{\mathbf{b}}(s)$ and $\hat{\mathbf{t}} \neq \hat{\mathbf{t}}(b)$. Thus, we can write

$$\begin{aligned} [[\hat{\mathbf{s}} \cdot \boldsymbol{\omega}]] &= [[\frac{\partial}{\partial s}(\hat{\mathbf{b}} \cdot \mathbf{u})]] - [[\frac{\partial}{\partial b}(\hat{\mathbf{t}} \cdot \mathbf{u})]] \\ &= \frac{\partial}{\partial s}[[\hat{\mathbf{b}} \cdot \mathbf{u}]] - \frac{\partial}{\partial b}[[\hat{\mathbf{t}} \cdot \mathbf{u}]] = 0, \end{aligned} \quad (2.39)$$

since $[[\hat{\mathbf{b}} \cdot \mathbf{u}]] = 0$ and $[[\hat{\mathbf{t}} \cdot \mathbf{u}]] = 0$, whence $[[\hat{\mathbf{s}} \cdot \boldsymbol{\omega}]] = 0$. Inspection of the terms remaining in equation 2.36 suggests that circulation is influenced by vorticity diffusion, vortex tilting/stretching, and gradients of curvature and interfacial tension.

The dimensionless versions of equations (2.25) and (2.24) are then expressed by

$$[[\tilde{\omega}_t]] = -2ReMa \frac{1}{(1 - \tilde{\Gamma})} \frac{\partial \tilde{\Gamma}}{\partial b}, \quad (2.40)$$

$$[[\tilde{\omega}_b]] = -2ReMa \frac{1}{(1 - \tilde{\Gamma})} \frac{\partial \tilde{\Gamma}}{\partial s}, \quad (2.41)$$

and the dimensionless equation (2.36) reads

$$\frac{D}{Dt} \left[\int_{\tilde{V}} \tilde{\omega} d\tilde{V} \right] = \oint_{\partial \tilde{V}} \hat{\mathbf{n}} \cdot \left(\tilde{\omega} \tilde{\mathbf{u}} + \frac{1}{Re} \nabla \tilde{\omega} \right) d\tilde{S} + \frac{1}{We} \oint_I \hat{\mathbf{b}} \frac{\partial}{\partial s} (\tilde{\sigma} [\tilde{\kappa}_1 + \tilde{\kappa}_2]) d\tilde{S}, \quad (2.42)$$

and the tildes are dropped henceforth.

Note that in the case of non-isothermal systems, $\tilde{\sigma}$ has a linear dependence on the local temperature \mathcal{T} , and a linear equation of state describes $\tilde{\sigma}(\tilde{\mathcal{T}})$ (see, for example, Williams *et al.* (2021)). It is possible to relate the present, surfactant-laden case to that involving thermal gradients by linearising our equation of state, $\sigma = 1 + \beta_s \ln(1 - \Gamma)$, for $\Gamma \ll 1$ such that it reads $\sigma = 1 - \beta_s \Gamma$. Although this analogy is useful, it is, however, incomplete since the non-isothermal case does not involve a surface species whose concentration evolves spatio-temporally for which a transport equation must be solved.

3. Results

Figure 3 shows a flow regime map for $Re = 5000$ that depicts the interfacial morphology associated with various regions of the $\beta_s - We$ parameter space generated by over 100 transient simulations performed in the ranges $100 < We < 1000$ and $0.1 < \beta_s < 0.9$. We have divided the map into two distinct regions depending on the morphology: for small We , capillary forces control the interfacial dynamics preventing the development of lobes which could result in the formation of large droplets; for large We , inertial forces dominate the dynamics triggering the formation of interfacial lobes whose thinning eventually results in the generation of holes and eventually droplets. The resulting non-uniform surfactant distribution generates gradients in surface tension affecting the local dynamics. Surfactant accumulation takes places in high-curvature regions giving rise to Marangoni stresses that drive surfactant redistribution from high- to low-concentration regions. Marangoni stresses, therefore, oppose the shear stresses produced by the flow field, the former exerting a restoring effect and the latter a perturbing effect in the local surfactant concentration field. The dimensionless Marangoni velocities induced by surface tension differences $\Delta\sigma$ are of $O(ReWe^{-1}(\Delta\sigma/\sigma_s))$. Similarly, the dimensionless Marangoni stresses, τ , are of $O(We^{-1}\tilde{\nabla}\tilde{\sigma})$, or, equivalently, $O(\beta_s We^{-1}\tilde{\nabla}\tilde{\Gamma})$, viz. equation (2.8), while capillary forces and shear stresses are of $O(We^{-1})$ and $O(Re^{-1})$, respectively.




Figure 3: Regime map of the interfacial morphology in the $\beta_s - We$ space for $Re = 5000$, $Pe_s = 100$, and $\Gamma_o = \Gamma_\infty/2$. The capillary-dominated and inertia-dominated regimes, and their boundaries are clearly demarcated; the symbols represent simulations carried out at the transition lines separating these regimes. Three-dimensional representations of the interface for both regimes are also shown.

Furthermore, from equations (2.40) and (2.41), it is clear that the Marangoni-induced vorticity jumps across the interface are of $O(Re \beta_s We^{-1})$. Inspection of figure 3, which was generated for a fixed Re value, reveals that the presence of Marangoni stresses counteracts the transition from the low- to high- We regimes as the critical We increases with β_s with a quasi-linear dependence. The latter is consistent with the scaling highlighted above, $\tau \sim \beta_s We^{-1}$, which demonstrates that increasing β_s and decreasing We serve to enhance the restoring influence of the Marangoni stresses. The boundary demarcated in figure 3 was generated by examining the temporal evolution of the interfacial area normalised by its initial value over a range of We and with β_s varying parametrically and $Re = 5000$; this shows that the normalised area is maximised for an intermediate value of We , for fixed β_s (and Re) which heralds the transition towards an inertia-dominated regime.

To assess the effect of Marangoni-induced flow, we have analysed the flow physics of the surfactant-free and surfactant-laden flows characterised by $Re = 5000$ and $We = 500$. We start with the surfactant-free case depicted in Figure 4 which shows the spatio-temporal interfacial dynamics for the surfactant-free case through the Q -criterion (e.g., a measure of the dominance of vorticity $\boldsymbol{\omega}$ over strain \mathbf{s} , i.e., $Q = (||\boldsymbol{\omega}||^2 - ||\mathbf{s}||^2)/2$ (Hunt *et al.* 1988)). At early times, we observe the formation of a periodic array of quasi-symmetric Kelvin-Helmholtz (KH)-driven vortex rings as a result of the difference in velocity in the shear layer located under the interface (see figure 4a). With increasing time, the three-dimensional instability starts with the deformation of the vortex-rings leading to a mutual-induction between two consecutive vortex rings resulting in their ‘knitting’ (see figure 4b); similar vortex-pairing has been reported by Broze & Hussain (1996) and da Silva & Métais (2002). With increasing time, we observe the formation of inner and outer hairpin vortices whose pairing brings about a region where both overlap. The cascade mechanism resulting in the formation of hairpin-vortices from KH-rings is triggered by the magnitude of the streamwise vorticity, ω_x , which becomes comparable to its azimuthal counterpart, ω_y , in agreement with Jarrahbashi *et al.* (2016) and Constante-Amores *et al.* (2021a), as shown in figure 4b.




Figure 4: Spatio-temporal representation of the interfacial dynamics and the coherent vortical structures for $Re = 5000$ and $We = 500$ at $t = (23.28, 28.12, 31.25, 31.50)$ shown in (a)-(d), respectively. For (c) and (d), the first row represents the 3D coherent structures with the location of the interface, the middle row represents only the interface location, and the bottom row shows two transversal cuts of the interface coloured by the magnitude of ω_x (the white lines represent the interface location). The three-dimensional coherent structures are visualised by the Q -criterion with values of $Q = (3, 220, 320, 320)$, where the colour represents the streamwise vorticity field, ω_x . In each panel, we also show ω_x in the $y-z$ plane for each sampling location.

To provide more conclusive evidence of the existence of inner/outer hairpin vortices in the jet dynamics, a careful study of the distribution of vortex signs shows the assembling into counter-rotating vortex pairs (see ω_x in the $y-z$ plane for each sampled location of the panels in figure 4). By analysing the distribution of streamwise vorticity between the ring and braid regions of the jet core (see figure 4a), we observe that their distribution is π -out-of-phase. The arrangement of the vorticity comes from vortex induction arguments, similar to those explained by Jarrahbashi *et al.* (2016), Zandian *et al.* (2018) and Constante-Amores *et al.* (2021a), i.e., the upstream hairpin vortex from the ring overtakes the upstream hairpin vortex from the braid as the mutual induction takes place. Finally, the vortex-surface interaction triggers the formation of the interfacial structure as the interface adopts the shape of the vortex which is in its vicinity (see figure 4b-d, ‘HV’ stands

for hairpin vortices). The mutual induction between outer and inner hairpin vortices eventually leads to the thinning of the lobes to ultimately form inertia-induced holes whose capillary-driven expansion gives rise to the formation of droplets (Jarrahbashi *et al.* 2016; Zandian *et al.* 2018; Constante-Amores *et al.* 2021a).

Next, we turn our attention to the effect of surfactants on the flow dynamics. Figure 5 shows the early interfacial surfactant concentration together with the three-dimensional coherent vortical structures via the Q -criterion. Similarly to the surfactant-free case, we observe the formation of a periodic array of quasi-axisymmetric KH-vortex rings. These rings induce the formation of interfacial waves that are characterised by regions of radially converging and diverging motion that lead to higher and lower interfacial areas, and subsequently to lower and higher surfactant concentration regions, respectively; accumulation of Γ is observed in the vicinity of the KH rings (see figure 5a). Figure 5c presents the interfacial concentration Γ , and Marangoni stresses τ along an arc length, s , corresponding to $t = 32.03$. We observe that the non-uniform distribution of Γ gives rise to Marangoni-induced flow, which drives fluid motion from ring-1, ‘VR1’, ($\tau > 0$) to ring-2, ‘VR2’, and vice versa (i.e, flow from VR2 to VR1, $\tau < 0$). This flow is therefore accompanied by the retardation of the development of the interfacial waves and a subsequent delay of the onset of the three-dimensional instability of the jet observed in the surfactant-free case in figure 4.

Additionally, these Marangoni stresses promote jumps in the vorticity across the interface which we can calculate using equations 2.24 and 2.25 in the location which coincides with the formation of vortex $SV1$ and $SV2$ from figure 6 at $t = 32.81$. Figure 5d shows a three-dimensional representation of the interface together with an x - z plane at $y = 2.875$ colored by the magnitude of vorticity, $|\omega|$. Figure 5e,f show respectively the variation of the interface location and the Γ profiles, and of the distribution of $[[\omega_b]]$ and $[[\omega_t]]$, along the arc length, s (not to be confused with \hat{s} the unit vector in figure 2), in the plane cutting the interface shown in figure 5d. From figure 5e, it is seen that the surfactant accumulates in the down-sloping region immediately downstream of an interfacial wave peak; here, the gradients in Γ , and therefore in σ , are smallest corresponding to the weakest vorticity jumps, while the largest such jumps are in the wave peak and trough regions where the Γ (and σ) gradients are highest, as shown in figure 5f. Inspection of figure 5f also shows that $[[\omega_b]] \gg [[\omega_t]]$, that is, near-interface vorticity production in the azimuthal direction is dominant. This acts to disrupt the dynamics of vortex-pairing relative to the surfactant-free case as the ‘knitting process’ is promoted by streamwise rather than azimuthal vorticity production and the vortex-ring deformation is replaced by vortex-reconnection and merging in the azimuthal direction in the surfactant-laden case.

For increasing time, figure 6 shows the formation of surfactant-induced inner hairpin-like vortical structures. The shear stress, which is generated to balance the gradients in σ gives rise to counter-rotating streamwise vortices of similar magnitude to the KH rings (labelled ‘ $SV1$ ’ and ‘ $SV2$ ’ in figure 6b). These structures grow in the x -direction into a combination of streamwise vortices close to the interface, i.e. legs, and a hairpin-like head close to the center-plane of the jet (see figure 6d). The hairpin-legs extend from the regions of high-to-low values of Γ on the surface, while the hairpin-head points down in the positive x -direction (labelled ‘ $HV1$ ’ and ‘ $HV2$ ’ in figure 6e). To complete the presentation of these hairpin-like vortical structures, figure 6f,g show the direction of flow rotation of the legs and head for $HV1$. For comparison, we have added arrows to show velocity direction and to prove that this coherent vortical structure exhibits the same qualitative behaviour as the HV proposed by Theodorsen (1952) for near-wall turbulence. To the best of our knowledge, the formation of hairpin-like vortical structures induced by surfactant effects has not been reported yet. We have also observed surfactant-driven outer




Figure 5: Effect of surfactants on the early interfacial dynamics for $Re = 5000$, $We = 500$, $\beta_s = 0.5$, $Pe_s = 100$ and $\Gamma_o = \Gamma_\infty/2$ at $t = 32.03$, (a), and 32.81 , (b). The top and bottom panels represent the interface coloured by Γ and the coherent vortical structures visualised via Q -criterion with $Q = 10$. Panel (c) shows a 2D representation of Γ , and τ , with respect to the arc length s (see inset) at $t = 32.03$. Panel (d) shows a 2D representation of the magnitude of vorticity $|\omega|$ in the x - z plane ($y = 2.875$) at $t = 32.81$; interface location and Γ , and $[[\omega_b]]$ and $[[\omega_t]]$ vs the arc length s (e.g., s corresponds to the x - z plane ($y = 2.875$) intersecting the interface) shown in (e) and (f), respectively. The center of the jet core corresponds to $z = 2.5$




Figure 6: Surfactant-driven hairpin-vortical structures: temporal development of the $HV1$ and $HV2$ hairpin-like vortical structures via Q -criterion with $Q = 200$ together with the interfacial location coloured by Γ at time $t = (32.81, 33.59, 34.37, 36.71, 39.06)$, corresponding to panels (a)-(e), respectively. In panel (e), we show the two transversal slices displayed in panels (f) and (g) which depict the streamwise vorticity ω_x through the legs and head of $HV1$, respectively; arrows of in-plane velocity vectors have been added; the white lines represent the interface location. The parameter values are the same as in figure 5.

hairpin-like vortical structures (not shown) whose heads are in the negative x -direction (in the frame of reference of the legs).

At later times, figure 7a-d shows the variation with arc length of the interfacial location, Γ , and $[[\omega_t]]$ and $[[\omega_b]]$ at $t = 36.51$ and $t = 44.68$; corresponding three-dimensional representations of the interface are also shown in figure 7e,f for $t = 44.68$ coloured by the magnitude of Γ and the Q -criterion, respectively. The flow is accompanied by radially-converging and diverging motion due to vortex-surface-interaction; interfacial convection drives surfactant towards the inner lobes (interfacial contraction), and away from the outer lobes (interfacial expansion). Vorticity jumps are highest in the interfacial regions with the largest gradients in Γ . As time evolves, the ratio of these Marangoni-driven $[[\omega_t]]$ to $[[\omega_b]]$ reduces and this results in large coherent structures which merge to form counter-rotating streamwise vortical rings that eventually ‘knit’ with the adjacent vortex ring located in the x -direction (labelled ‘ $VR1$ - $VR4$ ’ in figure 7f); this pairing is similar to the surfactant-free case (in agreement with Urbin & Métais (1997) and da Silva & Métais (2002)).




Figure 7: Panels (a-b) and (c-d) show the location of the interface together with the surfactant concentration and the jumps of the vorticity across the interface for $t = 36.81$ and $t = 44.68$, respectively. Panels (e-f) show a three dimensional representation of the interface location coloured by Γ at $t = 44.68$, and vortex knitting visualised via Q -criterion with value of $Q = 1600$ where the colour represents ω_x , respectively. The arc length s corresponds to the x - z plane ($y = 2.5$) intersecting the interface. The center of the jet core corresponds to $z = 2.5$. The parameter values are the same as in figure 5.

We now examine the dynamics of the circulation Ω by considering equation 2.42 which we express as follows:

$$\frac{D\Omega}{Dt} = I_{\text{tilt}} + I_{\text{diff}} + I_{\text{curv}}, \quad (3.1)$$

where I_{tilt} , I_{diff} , and I_{curv} are defined as

$$I_{\text{tilt}} \equiv \oint_{\partial\tilde{V}} \hat{\mathbf{n}} \cdot \boldsymbol{\omega} \mathbf{u} dS, \quad I_{\text{diff}} \equiv \frac{1}{Re} \oint_{\partial\tilde{V}} \hat{\mathbf{n}} \cdot \nabla \boldsymbol{\omega} dS, \quad I_{\text{curv}} \equiv \frac{1}{We} \oint_I \hat{\mathbf{b}} \cdot \frac{\partial}{\partial s} (\sigma [\kappa_1 + \kappa_2]) dS, \quad (3.2)$$

which correspond to vortex tilting/stretching, diffusion of vorticity, and circulation variation due to gradients in curvature and interfacial tension (in the case of surfactant-laden systems). Figure 8 shows the temporal evolution of $D\Omega/Dt$, I_{tilt} , I_{diff} , and I_{curv}




Figure 8: Total rate of change of circulation, Ω , according to equation 2.42: $D\Omega/Dt$, vortex diffusion (I_{diff}), vortex tilting (I_{tilt}), and surface tension (I_{curv}) are shown in rows one to four, respectively; see equation (3.2) for the definitions of I_{tilt} , I_{diff} , and I_{curv} . Surfactant-laden and surfactant-free cases correspond to left and right panels, respectively. For panel (g), we represent the contributions that arise from the gradients of curvature (solid lines) and the gradients of surface tension (dashed lines) to underscore the relative importance of the Marangoni stresses. Red, blue, and black colored lines represent component x , y and z of $D\Omega/Dt$, I_{tilt} , I_{diff} , and I_{curv} . The parameters are $Re = 5000$, $We = 500$ (and for the surfactant-laden case) $\beta_s = 0.5$, $Pe_s = 100$ and $\Gamma_o = \Gamma_\infty/2$.




Figure 9: Spatio-temporal evolution of the control volume V with a surface ∂V which encloses an interface plane I used to calculate the rate of change of the circulation in figure 8. Panels (a)-(d) and (e)-(h) correspond to the surfactant-laden and surfactant-free cases, respectively, at $t = (25.20, 35.50, 40.37, 43.75)$ and $t = (27.12, 30.43, 34.37, 40.55)$, and the same parameters as in figure (8).

which allows us to identify the dominant physical mechanisms that contribute to the creation and dissipation of circulation. In figure 9 we also show snapshots of the three-dimensional representation of the interface corresponding to the volume used to carry out the computations necessary to calculate $D\Omega/Dt$ and its constituent terms for the surfactant-laden and surfactant-free cases; this allows one to pinpoint the mechanisms primarily responsible for the interfacial structures observed. It is clearly seen from figure 8 that during the early stages of the flow, Ω remains approximately constant. Inspection of panels (c)-(h) of figure 8 shows clearly that the rate of change of circulation is dominated by the mechanisms related to vortex diffusion I_{diff} and curvature I_{curv} , with vortex tilting/shielding playing a relatively minor role. It is also clear that in the surfactant-laden jet case, the Marangoni contribution to I_{curv} dominates that associated with curvature derivatives. This observation further bolsters the claim that Marangoni stresses drive vorticity generation in the jet dynamics.

The snapshots depicted in figure 9 for the surfactant-laden (panels (a)-(d)) and surfactant-free (panels (e)-(h)) cases have been chosen carefully so as to link the various stages of jet destabilisation to the prominent changes in the temporal variation of I_{diff} , I_{tilt} , I_{curv} , and $D\Omega/Dt$. Given the dominance of I_{curv} over the time range considered ($0 \leq t \leq 40$), we focus on the variations in this quantity and its signature effects on the interfacial shape. Inspection of figures 8(g) and 9(a) reveals that the relatively gentle interfacial undulations are linked to variations of the Marangoni contribution to I_{curv} in the $x - y$ plane. The development of the more complex interfacial shapes, on the other hand, is accompanied by a concomitant rise in three-dimensionality of I_{curv} (in addition to significant contributions from the x -component of I_{curv}). In the surfactant-free case,




Figure 10: Temporal evolution of (a) surface area, A , (b) kinetic energy, $E_k = \rho \int_V \mathbf{u}^2/2dV$, and (c) enstrophy, $\varepsilon = \int_V |\boldsymbol{\omega}^2|dV$, scaled by the initial interfacial area, A_0 , kinetic energy, E_{k0} , and enstrophy, ε_0 , respectively. The parameter values are the same as in figure 5.

inspection of figures (8)(d) and (h), and (9)(e)-(h) shows that the interfacial jet evolution is accompanied by large variations in the x -component of I_{curv} and vorticity diffusion characterised by I_{diff} .

Lastly, we plot in figure 10 the effect of surfactants on the interfacial area, kinetic energy, defined as $E_k = \rho \int_V \mathbf{u}^2/2dV$, and the enstrophy, $\varepsilon = \int_V |\boldsymbol{\omega}^2|dV$, normalised by their initial values, A_0 , E_{k0} , ε_0 , respectively. After the onset of destabilization (defined when the interfacial surface has reached $A = 1.025$), we observe that the surfactant-induced effects discussed above, which include the interfacial vorticity jumps brought about by Marangoni stresses, and their effect on the production of circulation, and jet destabilisation mechanisms associated with vortex formation and spanwise reconnection, promote the delay in increase and subsequent reduction in interfacial area; these effects also lead to a delay in the decay of the jet kinetic energy as well as its enstrophy.

4. Concluding remarks

Three-dimensional numerical simulations of jet destabilisation and atomisation in the presence of a monolayer of insoluble surfactants have been carried out for the first time. A phase diagram in the space of dimensionless surfactant elasticity and Weber number in the inertia-dominated region is presented in the limiting case where there is no vorticity production associated with jumps in material properties such as fluid density and viscosity; in the present work, surface tension forces and Marangoni stress

give rise to variations in vorticity and circulation in addition to the vortex tilting/shielding and diffusion mechanisms. We have also derived formulae for the vorticity jumps across the interface due to Marangoni stresses, and equations that provide a breakdown of the rate of production of circulation within the jet into constituent terms which we associate with vortex tilting/shielding, diffusion, and gradients in interfacial curvature and surface tension. The present theoretical formulation is expressed as a conservation law for circulation. We have focused on the limiting case where there is no vorticity production associated with jumps in material properties. Future studies should examine situations characterised by fluids with different material properties.

Then, we have analysed in details the vortex-interface-surfactant interactions in the flow dynamics. At early times, the presence of surfactants induces spanwise vortex reconnections brought about Marangoni-induced flow resulting in the delay of the onset of destabilisation to the three-dimensional interfacial instabilities. We also show that surfactant-induced Marangoni-stresses trigger the formation of hairpin-like structures whose head and legs extend in the streamwise direction. Lastly, we have attempted to link the changes in interfacial topology to the mechanisms that influence the production of vorticity and circulation demonstrating a balance between curvature gradients and diffusion for surfactant-free jets, and the dominance of Marangoni stresses in the surfactant-laden cases.

The present results have been obtained for insoluble surfactants, and we acknowledge that experimental and numerical studies feature soluble surfactants which are dissolved in the liquid that issues from a nozzle to form the jet (Sijts *et al.* 2021; Constante-Amores 2021). It is well known that the addition of surfactant-solubility will lead to additional richness and complexity. Although they do not affect the governing equations that describe the bulk fluid, they will change the boundary conditions that constrain them, resulting in a change in the flow dynamics. We can anticipate that a change of flow in the vicinity of the interface will have a detrimental effect on the coherent structures that emerge, subsequently affecting the close interplay between interface-vorticity-surfactant. These challenges will be the subject of future work.

Declaration of Interests. The authors report no conflict of interest.

This work is supported by the Engineering and Physical Sciences Research Council, United Kingdom, through the EPSRC MEMPHIS (EP/K003976/1) and PREMIERE (EP/T000414/1) Programme Grants. O.K.M. acknowledges funding from PETRONAS and the Royal Academy of Engineering for a Research Chair in Multiphase Fluid Dynamics. We acknowledge HPC facilities provided by the Research Computing Service (RCS) of Imperial College London for the computing time. AAC-P acknowledge the support from the Royal Society through a University Research Fellowship (URF/R/180016), an Enhancement Grant (RGF/EA/181002) and two NSF/CBET-EPSRC grants (Grant Nos. EP/S029966/1 and EP/W016036/1). D.J. and J.C. acknowledge support through HPC/AI computing time at the Institut du Développement et des Ressources en Informatique Scientifique (IDRIS) of the Centre National de la Recherche Scientifique (CNRS), coordinated by GENCI (Grand Équipement National de Calcul Intensif) Grant 2022 A0122B06721.

Appendix A. Effect of domain size

To provide conclusive evidence that the lateral size of the domain is sufficiently large to avoid finite-size effects, we have performed additional simulations accounting for different domain sizes, and measured the wavelength between the emergent crests. The selected

wavelength for panels for domain sizes of $(4D)^3$, $(5D)^3$ and $(6D)^3$ correspond to $\lambda \sim 1.52D$, $\lambda \sim 1.55D$ and $\lambda \sim 1.60D$, respectively. Thus, the wavelength values are very weakly dependent on the domain size indicating the absence of finite-size effects. We also note that the size of the computational domain is in agreement with previous studies (see for example Jarrahbashi & Sirignano (2014); Jarrahbashi *et al.* (2016), Zandian *et al.* (2016) and Desjardins & Pitsch (2010)), which have also used periodic boundary conditions for all three components of velocity in the streamwise direction.

Appendix B. Kinematics

We first develop an expression for $D\hat{\mathbf{t}}/Dt$. We consider the motion of an infinitesimal fluid parcel in the plane of the interface, which is treated as a material surface. The position vector is $\mathbf{x} = \mathbf{x}(s, b, t)$ where s and b represent arc length distances along the $\hat{\mathbf{t}}$ and $\hat{\mathbf{b}}$ directions, respectively. At time, $t + \delta t$, to leading order in δt , we can write the following expression for the tangent to the interface at the fluid parcel which at time t was located at $\mathbf{x}(0, 0, t)$

$$\tilde{\mathbf{t}}(t + \delta t) = \frac{\partial \mathbf{x}}{\partial s}(0, 0, t) + \nabla \mathbf{u} \cdot \left(\frac{\partial \mathbf{x}}{\partial s} + \frac{\partial \mathbf{x}}{\partial b} \right) \delta t. \quad (\text{B1})$$

Noting that $\hat{\mathbf{t}} = \partial \mathbf{x} / \partial s$, $\hat{\mathbf{b}} = \partial \mathbf{x} / \partial b$, $\hat{\mathbf{t}} \cdot \nabla \mathbf{u} = \partial \mathbf{u} / \partial s$, and $\hat{\mathbf{b}} \cdot \nabla \mathbf{u} = \partial \mathbf{u} / \partial b$, this equation can be re-expressed as

$$\tilde{\mathbf{t}}(t + \delta t) = \hat{\mathbf{t}} + \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \delta t. \quad (\text{B2})$$

The magnitude of $\tilde{\mathbf{t}}(t + \delta t)$ is given by

$$|\tilde{\mathbf{t}}| = 1 + \hat{\mathbf{t}} \cdot \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \delta t + O(\delta t)^2, \quad (\text{B3})$$

and normalisation of $\tilde{\mathbf{t}}(t + \delta t)$ by this magnitude gives the following tangent unit vector $\hat{\mathbf{t}}(t + \delta t)$

$$\begin{aligned} \hat{\mathbf{t}}(t + \delta t) &= \frac{\tilde{\mathbf{t}} + \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \delta t}{|\tilde{\mathbf{t}}(t + \delta t)|} = \frac{\hat{\mathbf{t}} + \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \delta t}{1 + \hat{\mathbf{t}} \cdot \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \delta t} \\ &= \left(\hat{\mathbf{t}} + \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \delta t \right) \left(1 - \hat{\mathbf{t}} \cdot \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \delta t + O(\delta t)^2 \right) \\ &= \hat{\mathbf{t}} + \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} - \hat{\mathbf{t}} \left(\hat{\mathbf{t}} \cdot \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \right) \right) \delta t + O(\delta t)^2. \end{aligned} \quad (\text{B4})$$

From this expression, we can arrive at an approximate formula for $D\hat{\mathbf{t}}/Dt$:

$$\frac{D\hat{\mathbf{t}}}{Dt} \sim \frac{\hat{\mathbf{t}}(t + \delta t) - \hat{\mathbf{t}}(t)}{\delta t} = \frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} - \hat{\mathbf{t}} \left(\hat{\mathbf{t}} \cdot \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right) \right) + O(\delta t). \quad (\text{B5})$$

We now insert the following expression for \mathbf{u} into $\mathbf{u} \cdot D\hat{\mathbf{t}}/Dt$

$$\mathbf{u} = (\mathbf{u} \cdot \hat{\mathbf{s}})\hat{\mathbf{s}} + (\mathbf{u} \cdot \hat{\mathbf{t}})\hat{\mathbf{t}} + (\mathbf{u} \cdot \hat{\mathbf{b}})\hat{\mathbf{b}}, \quad (\text{B6})$$

which yields

$$\mathbf{u} \cdot \frac{D\hat{\mathbf{t}}}{Dt} = (\mathbf{u} \cdot \hat{\mathbf{s}})\hat{\mathbf{s}} \cdot \frac{D\hat{\mathbf{t}}}{Dt} + (\mathbf{u} \cdot \hat{\mathbf{b}})\hat{\mathbf{b}} \cdot \frac{D\hat{\mathbf{t}}}{Dt}. \quad (\text{B7})$$

Substitution of Eq. (B 5) into $\hat{\mathbf{s}} \cdot D\hat{\mathbf{t}}/Dt$ and $\hat{\mathbf{b}} \cdot D\hat{\mathbf{t}}/Dt$ gives

$$\hat{\mathbf{s}} \cdot \frac{D\hat{\mathbf{t}}}{Dt} = \hat{\mathbf{s}} \cdot \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right), \quad (\text{B 8})$$

$$\hat{\mathbf{b}} \cdot \frac{D\hat{\mathbf{t}}}{Dt} = \hat{\mathbf{b}} \cdot \left(\frac{\partial \mathbf{u}}{\partial s} + \frac{\partial \mathbf{u}}{\partial b} \right), \quad (\text{B 9})$$

where we have made use of $\hat{\mathbf{s}} \cdot \hat{\mathbf{t}} = 0$ and $\hat{\mathbf{b}} \cdot \hat{\mathbf{t}} = 0$. We can re-express the RHS of Eqs. (B 8) and (B 9) as follows

$$\hat{\mathbf{s}} \cdot \frac{\partial \mathbf{u}}{\partial s} = \frac{\partial}{\partial s} (\hat{\mathbf{s}} \cdot \mathbf{u}) - \mathbf{u} \cdot \frac{\partial \hat{\mathbf{s}}}{\partial s}, \quad (\text{B 10})$$

$$\hat{\mathbf{s}} \cdot \frac{\partial \mathbf{u}}{\partial b} = \frac{\partial}{\partial b} (\hat{\mathbf{s}} \cdot \mathbf{u}) - \mathbf{u} \cdot \frac{\partial \hat{\mathbf{s}}}{\partial b}, \quad (\text{B 11})$$

$$\hat{\mathbf{b}} \cdot \frac{\partial \mathbf{u}}{\partial s} = \frac{\partial}{\partial s} (\hat{\mathbf{b}} \cdot \mathbf{u}) - \mathbf{u} \cdot \frac{\partial \hat{\mathbf{b}}}{\partial s}, \quad (\text{B 12})$$

$$\hat{\mathbf{b}} \cdot \frac{\partial \mathbf{u}}{\partial b} = \frac{\partial}{\partial b} (\hat{\mathbf{b}} \cdot \mathbf{u}) - \mathbf{u} \cdot \frac{\partial \hat{\mathbf{b}}}{\partial b}. \quad (\text{B 13})$$

Inserting Eq. (B 6) into the second term on the RHS of Eqs. (B 10)-(B 13), we obtain

$$\mathbf{u} \cdot \frac{\partial \hat{\mathbf{s}}}{\partial s} = \kappa_1 (\mathbf{u} \cdot \hat{\mathbf{t}}), \quad (\text{B 14})$$

$$\mathbf{u} \cdot \frac{\partial \hat{\mathbf{s}}}{\partial b} = \kappa_2 (\mathbf{u} \cdot \hat{\mathbf{b}}), \quad (\text{B 15})$$

$$\mathbf{u} \cdot \frac{\partial \hat{\mathbf{b}}}{\partial s} = 0, \quad (\text{B 16})$$

$$\mathbf{u} \cdot \frac{\partial \hat{\mathbf{b}}}{\partial b} = -\kappa_2 (\mathbf{u} \cdot \hat{\mathbf{s}}), \quad (\text{B 17})$$

where the curvatures κ_1 and κ_2 are defined as follows

$$\kappa_1 = \hat{\mathbf{t}} \cdot \frac{\partial \hat{\mathbf{s}}}{\partial s}, \quad \kappa_2 = \hat{\mathbf{b}} \cdot \frac{\partial \hat{\mathbf{s}}}{\partial b}. \quad (\text{B 18})$$

In deriving Eqs. (B 14)-(B 17), we have noted that $\hat{\mathbf{t}} \neq \hat{\mathbf{t}}(b)$ and $\hat{\mathbf{b}} \neq \hat{\mathbf{b}}(s)$. Substitution of Eqs. (B 14)-(B 17) into Eqs. (B 10)-(B 13) and the resultant relations into Eqs. (B 8) and (B 9) respectively yields the following expressions for $\hat{\mathbf{s}} \cdot (D\hat{\mathbf{t}}/Dt)$ and $\hat{\mathbf{b}} \cdot (D\hat{\mathbf{t}}/Dt)$

$$\hat{\mathbf{s}} \cdot \frac{D\hat{\mathbf{t}}}{Dt} = \frac{\partial}{\partial s} (\mathbf{u} \cdot \hat{\mathbf{s}}) + \frac{\partial}{\partial b} (\mathbf{u} \cdot \hat{\mathbf{s}}) - \kappa_1 (\mathbf{u} \cdot \hat{\mathbf{t}}) - \kappa_2 (\mathbf{u} \cdot \hat{\mathbf{b}}), \quad (\text{B 19})$$

$$\hat{\mathbf{b}} \cdot \frac{D\hat{\mathbf{t}}}{Dt} = \frac{\partial}{\partial s} (\mathbf{u} \cdot \hat{\mathbf{b}}) + \frac{\partial}{\partial b} (\mathbf{u} \cdot \hat{\mathbf{b}}) + \kappa_2 (\mathbf{u} \cdot \hat{\mathbf{s}}). \quad (\text{B 20})$$

Substitution of Eqs. (B 19) and (B 20) into Eq. (B 7) and re-arranging yields

$$\mathbf{u} \cdot \frac{D\hat{\mathbf{t}}}{Dt} = \frac{1}{2} \frac{\partial}{\partial s} \left[(\mathbf{u} \cdot \hat{\mathbf{s}})^2 + (\mathbf{u} \cdot \hat{\mathbf{b}})^2 \right] + \frac{1}{2} \frac{\partial}{\partial b} \left[(\mathbf{u} \cdot \hat{\mathbf{s}})^2 + (\mathbf{u} \cdot \hat{\mathbf{b}})^2 \right] - \kappa_1 (\mathbf{u} \cdot \hat{\mathbf{t}}) (\mathbf{u} \cdot \hat{\mathbf{s}}). \quad (\text{B 21})$$

Appendix C. Near-interface normal stress jump

In order to generate a 3D version of the pressure gradient term in Eq. (2.33), we first consider the jump in the normal stress across the plane of the interface:

$$p_2 - p_1 = -\sigma(\kappa_1 + \kappa_2) + [[\mu \hat{\mathbf{s}} \cdot \mathbf{D} \cdot \hat{\mathbf{s}}], \quad (\text{C 1})$$

where κ_1 and κ_2 are given by Eqs. (B 18). Substitution of Eq. (B 6) into $\nabla \cdot \mathbf{u} = 0$ yields

$$\begin{aligned} \hat{\mathbf{s}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}} &= -\hat{\mathbf{t}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{t}} - \hat{\mathbf{b}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{b}} \\ &= -\hat{\mathbf{t}} \cdot \frac{\partial \mathbf{u}}{\partial s} - \hat{\mathbf{b}} \cdot \frac{\partial \mathbf{u}}{\partial b}, \end{aligned} \quad (\text{C } 2)$$

where we have set $\hat{\mathbf{t}} \cdot \nabla \mathbf{u} = \partial \mathbf{u} / \partial s$ and $\hat{\mathbf{b}} \cdot \nabla \mathbf{u} = \partial \mathbf{u} / \partial b$. We can re-express $\hat{\mathbf{t}} \cdot (\partial \mathbf{u} / \partial s)$ and $\hat{\mathbf{b}} \cdot (\partial \mathbf{u} / \partial b)$ as follows

$$\hat{\mathbf{t}} \cdot \frac{\partial \mathbf{u}}{\partial s} = \frac{\partial}{\partial s} (\mathbf{u} \cdot \hat{\mathbf{t}}) - \mathbf{u} \cdot \frac{\partial \hat{\mathbf{t}}}{\partial s}, \quad (\text{C } 3)$$

$$\hat{\mathbf{b}} \cdot \frac{\partial \mathbf{u}}{\partial b} = \frac{\partial}{\partial b} (\mathbf{u} \cdot \hat{\mathbf{b}}) - \mathbf{u} \cdot \frac{\partial \hat{\mathbf{b}}}{\partial b}. \quad (\text{C } 4)$$

Substitution of Eq. (B 6) into $\mathbf{u} \cdot (\partial \hat{\mathbf{t}} / \partial s)$ and $\mathbf{u} \cdot (\partial \hat{\mathbf{b}} / \partial s)$ leads to

$$\mathbf{u} \cdot \frac{\partial \hat{\mathbf{t}}}{\partial s} = -\kappa_1 (\mathbf{u} \cdot \hat{\mathbf{s}}), \quad (\text{C } 5)$$

$$\mathbf{u} \cdot \frac{\partial \hat{\mathbf{b}}}{\partial b} = -\kappa_2 (\mathbf{u} \cdot \hat{\mathbf{s}}), \quad (\text{C } 6)$$

where, again, we have made use of the fact that $\hat{\mathbf{t}} \neq \hat{\mathbf{t}}(b)$ and $\hat{\mathbf{b}} \neq \hat{\mathbf{b}}(s)$. Substitution of Eqs. (C 5) and (C 6) into Eqs. (C 3) and (C 4) and the resultant relations into Eq. (C 2) gives

$$\hat{\mathbf{s}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}} = -\frac{\partial}{\partial s} \left[(\mathbf{u} \cdot \hat{\mathbf{t}}) + (\mathbf{u} \cdot \hat{\mathbf{b}}) \right] - (\kappa_1 + \kappa_2) (\mathbf{u} \cdot \hat{\mathbf{s}}). \quad (\text{C } 7)$$

Since $\hat{\mathbf{s}} \cdot \mathbf{D} \cdot \hat{\mathbf{s}} = 2\hat{\mathbf{s}} \cdot \nabla \mathbf{u} \cdot \hat{\mathbf{s}}$, it follows that

$$\hat{\mathbf{s}} \cdot \mathbf{D} \cdot \hat{\mathbf{s}} = -2 \frac{\partial}{\partial s} \left[(\mathbf{u} \cdot \hat{\mathbf{t}}) + (\mathbf{u} \cdot \hat{\mathbf{b}}) \right] - 2(\kappa_1 + \kappa_2) (\mathbf{u} \cdot \hat{\mathbf{s}}). \quad (\text{C } 8)$$

Substitution of this equation into Eq. (C 1) yields the following expression for the pressure jump

$$[[p]] = -\sigma(\kappa_1 + \kappa_2) - 2 \left[\left[\mu \left(\frac{\partial}{\partial s} \left[(\mathbf{u} \cdot \hat{\mathbf{t}}) + (\mathbf{u} \cdot \hat{\mathbf{b}}) \right] + (\kappa_1 + \kappa_2) (\mathbf{u} \cdot \hat{\mathbf{s}}) \right) \right] \right]. \quad (\text{C } 9)$$

REFERENCES

- ARIYAPADI, S., BALACHANDAR, R. & BERRUTI, F. 2004 Effect of surfactant on the characteristics of a droplet-laden jet. *Chem Eng Process* **43** (4), 547 – 553.
- BATCHVAROV, A., KAHOUADJI, L., CONSTANTE-AMORES, C. R., NORÕES GONÇALVES, G. F., SHIN, S., CHERGUI, J., JURIC, D., CRASTER, R. V. & MATAR, O. K. 2021 Three-dimensional dynamics of falling films in the presence of insoluble surfactants. *J. Fluid Mech.* **906**, A16.
- BATCHVAROV, A., KAHOUADJI, L., MAGNINI, M., CONSTANTE-AMORES, C. R., CRASTER, R. V., SHIN, S., CHERGUI, J., JURIC, D. & MATAR, O. K. 2020 Effect of surfactant on elongated bubbles in capillary tubes at high reynolds number. *Phys. Rev. Fluids* **5**, 093605.
- BROZE, G. & HUSSAIN, F. 1996 Transitions to chaos in a forced jet: intermittency, tangent bifurcations and hysteresis. *J. Fluid Mech.* **311**, 37–71.
- BRØNS, M., THOMPSON, M. C., LEWEKE, T. & HOURIGAN, K. 2014 Vorticity generation and conservation for two-dimensional interfaces and boundaries. *J. Fluid Mech.* **758**, 63–93.
- CONSTANTE-AMORES, C.R., CHERGUI, J., SHIN, S., JURIC, D., CASTREJÓN-PITA, J.R. & CASTREJÓN-PITA, A.A. 2022 Role of surfactant-induced marangoni stresses in retracting liquid sheets. *J. Fluid Mech.* **949**, A32.

- CONSTANTE-AMORES, C.R., KAHOUADJI, L., BATCHVAROV, A., SHIN, S., CHERGUI, J., JURIC, D. & MATAR, O.K. 2021a Direct numerical simulations of transient turbulent jets: vortex-interface interactions. *J. Fluid Mech.* **922**, A6.
- CONSTANTE-AMORES, C. R. 2021 Three-dimensional computational fluid dynamics simulations of complex multiphase flows with surfactants. *Imperial College London PhD Thesis*.
- CONSTANTE-AMORES, C. R., KAHOUADJI, L., BATCHVAROV, A., SHIN, S., CHERGUI, J., JURIC, D. & MATAR, O. K. 2020a Dynamics of retracting surfactant-laden ligaments at intermediate ohnesorge number. *Phys. Rev. Fluids* **5**, 084007.
- CONSTANTE-AMORES, C. R., KAHOUADJI, L., BATCHVAROV, A., SHIN, S., CHERGUI, J., JURIC, D. & MATAR, O. K. 2020b Rico and the jets: Direct numerical simulations of turbulent liquid jets. *Phys. Rev. Fluids* **5**, 110501.
- CONSTANTE-AMORES, C. R., KAHOUADJI, L., BATCHVAROV, A., SHIN, S., CHERGUI, J., JURIC, D. & MATAR, O. K. 2021b Dynamics of a surfactant-laden bubble bursting through an interface. *J. Fluid Mech.* **911**, A57.
- CRASTER, R. V., MATAR, O. K. & PAPAGEORGIU, D. T. 2002 Pinchoff and satellite formation in surfactant covered viscous threads. *Phys. Fluids* **14** (4), 1364–1376.
- CRASTER, R. V., MATAR, O. K. & PAPAGEORIOU, D. T. 2009 Breakup of surfactant-laden jets above the critical micelle concentration. *J. Fluid Mech.* **629**, 195–219.
- DESJARDINS, O. & PITTSCH, H. 2010 Detailed numerical investigation of turbulent atomization of liquid jets. *Atomiz. Sprays* **20** (4), 311–336.
- EGGERS, J. 1993 Universal pinching of 3d axisymmetric free-surface flow. *Phys. Rev. Lett.* **71**, 3458–3460.
- EGGERS, J. 1997 Nonlinear dynamics and breakup of free-surface flows. *Rev. Mod. Phys.* **69** (3), 865–929.
- ELLIS, M.C.B., TUCK, C.R. & MILLER, P.C.H. 2001 How surface tension of surfactant solutions influences the characteristics of sprays produced by hydraulic nozzles used for pesticide application. *Colloids Surf* **180** (3), 267 – 276.
- FUSTER, D. & ROSSI, M. 2021 Vortex-interface interactions in two-dimensional flows. *Int. J. Multiph* **143**, 103757.
- HERRMANN, M. 2010 A parallel eulerian interface tracking/lagrangian point particle multi-scale coupling procedure. *J. Comput. Phys* **229** (3), 745 – 759.
- HOEPFFNER, J. & PARÉ, G. 2013 Recoil of a liquid filament: escape from pinch-off through creation of a vortex ring. *J. Fluid Mech.* **734** (183–197).
- HUNT, J., WRAY, A. & MOIN, P. 1988 Eddies, streams, and convergence zones in turbulent flows. *Studying Turbulence Using Numerical Simulation Databases* **1**, 193–208.
- IBARRA, E., SHAFFER, F. & SAVAŞ, O. 2020 On the near-field interfaces of homogeneous and immiscible round turbulent jets. *J. Fluid Mech.* **889**, A4.
- IBARRA, R. 2017 Horizontal and low-inclination oil-water flow investigations using laser-based diagnostic techniques. *Imperial College London PhD Thesis*.
- JARRAHBASHI, D. & SIRIGNANO, W. A. 2014 Vorticity dynamics for transient high-pressure liquid injection. *Phys. Fluids* **26** (10), 101304.
- JARRAHBASHI, D., SIRIGNANO, W. A., POPOV, P. P. & HUSSAIN, F. 2016 Early spray development at high gas density: hole, ligament and bridge formations. *J. Fluid Mech.* **792**, 186–231.
- KAMAT, P. M., WAGONER, B. W., THETE, S. S. & BASARAN, O. A. 2018 Role of marangoni stress during breakup of surfactant-covered liquid threads: Reduced rates of thinning and microthread cascades. *Phys. Rev. Fluids* **3**, 043602.
- KOOIJ, S., SIJS, R., DENN, M. M., VILLERMAUX, E. & BONN, D. 2018 What determines the drop size in sprays? *Phys. Rev. X* **8**, 031019.
- LIAO, Y. C., SUBRAMANI, H. J., FRANSES, E. I. & BASARAN, O. A. 2004 Effects of soluble surfactants on the deformation and breakup of stretching liquid bridges. *Langmuir* **20** (23), 9926–9930.
- LISTER, J. R. & STONE, H. A. 1998 Capillary breakup of a viscous thread surrounded by another viscous fluid. *Phys. Fluids* **10** (11), 2758–2764.
- LONGUET-HIGGINS, MICHAEL S. 1992 Capillary rollers and bores. *J. Fluid Mech.* **240**, 659–679.
- LUNDGREN, T. & KOUMOUTSAKOS, P. 1999 On the generation of vorticity at a free surface. *J. Fluid Mech.* **382**, 351–366.

- MANIKANTAN, H. & SQUIRES, T. M. 2020 Surfactant dynamics: hidden variables controlling fluid flows. *J. Fluid Mech.* **892**, P1.
- MARMOTTANT, P. & VILLERMAUX, E. 2004 On spray formation. *J. Fluid Mech.* **498**, 73 – 111.
- MARTÍNEZ-CALVO, A. & SEVILLA, A. 2020 Universal thinning of liquid filaments under dominant surface forces. *Phys. Rev. Lett.* **125**, 114502.
- MCGOUGH, P. T. & BASARAN, O. A. 2006 Repeated formation of fluid threads in breakup of a surfactant-covered jet. *Phys. Rev. Lett.* **96**, 054502.
- MORTON, B. R. 1984 The generation and decay of vorticity. *Geophysical & Astrophysical Fluid Dynamics* **28** (3-4), 277–308.
- PLATEAU, J. 1873 *Experimental and Theoretical Statics of Liquids Subject to Molecular Forces Only*, , vol. 1. Gand et Leipzig: F. Clemm.
- SHIN, S., CHERGUI, J., JURIC, D., KAHOUADJI, L., K. MATAR, O. & CRASTER, R. V. 2018 A hybrid interface tracking - level set technique for multiphase flow with soluble surfactant. *J. of Comp. Phys.* **359**, 409–435.
- SIJS, R. & BONN, D. 2020 The effect of adjuvants on spray droplet size from hydraulic nozzles. *Pest Management Science* **76** (10), 3487–3494.
- SIJS, R., KOUIJ, S., HOLTERMAN, H. J., VAN DE ZANDE, J. & BONN, D. 2021 Drop size measurement techniques for sprays: Comparison of image analysis, phase doppler particle analysis, and laser diffraction. *AIP Advances* **11** (1), 015315.
- DA SILVA, C. B. & MÉTAIS, O. 2002 Vortex control of bifurcating jets: A numerical study. *Physics of Fluids* **14** (11), 3798–3819.
- STRICKLAND, S. L., SHEARER, M. & DANIELS, K. E. 2015 Spatio-temporal measurement of surfactant distribution on gravity–capillary waves. *J. Fluid Mech.* **777**, 523–543.
- TERRINGTON, S. J., HOURIGAN, K. & THOMPSON, M. C. 2020 The generation and conservation of vorticity: deforming interfaces and boundaries in two-dimensional flows. *J. Fluid Mech.* **890**, A5.
- TERRINGTON, S. J., HOURIGAN, K. & THOMPSON, M. C. 2021 The generation and diffusion of vorticity in three-dimensional flows: Lyman’s flux. *J. Fluid Mech.* **915**, A106–6–42.
- THEODORSEN, T. 1952 Mechanism of turbulence. *Proceedings of the Midwestern Conference Fluid Mechanics, 1-19*. .
- URBIN, G & MÉTAIS 1997 Large-eddy simulations of three-dimensional spatially-developing round jets. *ERCOTAC Series. Springer* **5**.
- WEE, HANSOL, WAGONER, BRAYDEN W., GARG, VISHRUT, KAMAT, PRITISH M. & BASARAN, OSMAN A. 2021 Pinch-off of a surfactant-covered jet. *J. Fluid Mech.* **908**, A38.
- WILLIAMS, A. G. L., KARAPETSAS, G., MAMALIS, D., SEFIANE, K., MATAR, O. K. & VALLURI, P. 2021 Spreading and retraction dynamics of sessile evaporating droplets comprising volatile binary mixtures. *J. Fluid Mech.* **907**, A22.
- WU, J. Z. 1995 A theory of three-dimensional interfacial vorticity dynamics. *Phys. Fluids* **7** (10), 2375–2395.
- ZANDIAN, A., SIRIGNANO, W. A. & HUSSAIN, F. 2016 Three-dimensional liquid sheet breakup: vorticity dynamics. *54th AIAA Aerospace Sciences Meeting*. .
- ZANDIAN, A., SIRIGNANO, W. A. & HUSSAIN, F. 2018 Understanding liquid-jet atomization cascades via vortex dynamics. *J. Fluid Mech.* **843**, 293–354.
- ZANDIAN, A., SIRIGNANO, W. A. & HUSSAIN, F. 2019 Vorticity dynamics in a spatially developing liquid jet inside a co-flowing gas. *J. Fluid Mech.* **877**, 429–470.