



HAL
open science

Developmental programming of hypothalamic melanocortin circuits

Sebastien Bouret

► **To cite this version:**

Sebastien Bouret. Developmental programming of hypothalamic melanocortin circuits. *Experimental & Molecular Medicine*, 2022, 54 (4), pp.403-413. 10.1038/s12276-021-00625-8 . hal-03808093

HAL Id: hal-03808093

<https://hal.science/hal-03808093>

Submitted on 7 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Developmental Programming of Hypothalamic Melanocortin Circuits

Sebastien G. Bouret ^{1,2*}

¹ *Inserm, Laboratory of Development and Plasticity of the Neuroendocrine Brain, Lille Neuroscience & Cognition Research Center, UMR-S 1172, Lille, 59000, France*

² *University of Lille, FHU 1,000 Days for Health, Lille, 59000, France*

Running title: Development of melanocortin circuits

Total number of words: 5,199 words

***Corresponding author:** Sebastien G. Bouret, E-mail: sebastien.bouret@inserm.fr

Abstract

The melanocortin system plays a critical role in the central regulation of food intake and energy balance. It consists of neurons producing pro-opiomelanocortin (POMC), melanocortin receptors (MC4R), and the endogenous antagonist agouti-related peptide (AgRP). *Pomc* and *Mc4r* deficiency in rodents and humans causes early onset of obesity, whereas the loss of *Agrp* function is associated with leanness. There is accumulating evidence that many chronic diseases, including obesity, might originate during early life. The melanocortin system develops during a relatively long period beginning during embryonic life with the birth of POMC and AgRP neurons and continuing postnatally with the assembly of their neuronal circuitry. The development of the melanocortin system requires the tight temporal regulation of molecular factors, such as transcription factors and axon guidance molecules, and cellular mechanisms, such as autophagy. It also involves a complex interplay of endocrine and nutritional factors. Disruption of one or more of these developmental factors can lead to abnormal maturation and function of the melanocortin system and have profound metabolic consequences in later life.

Introduction

The melanocortin system is a critical component of brain pathways that regulate feeding behavior and energy homeostasis. The brain's melanocortin system consists of pro-opiomelanocortin (POMC) neurons and the neurons that produce agouti-related peptide (AgRP), an endogenous antagonist of melanocortins. There are two anatomically distinct POMC neuronal populations in the brain. The largest population of POMC neurons resides in the arcuate nucleus of the hypothalamus (ARH). A discrete population of POMC neurons is also found in the nucleus of the solitary tract (NTS) in the brainstem. In contrast, AgRP neurons are exclusively located in the ARH and co-express neuropeptide Y (NPY). Pharmacological approaches revealed more than three decades ago that melanocortins have a potent and long-lasting inhibitory effect on feeding ¹. More recent, neuron- and gene-specific genetic studies showed the importance of POMC neurons in mediating the physiological actions of metabolic hormones, such as leptin and insulin ^{2, 3}. Opto- and chemogenetic experiments further indicated that although arcuate POMC neurons appear to be more involved in integrating long-term adiposity signals, the contribution of hindbrain POMC neurons seem to be more specific to the integration of short-term satiety signals ^{4, 5}. In contrast, optogenetic stimulation of AgRP neurons rapidly evokes feeding ⁴, whereas genetic ablation of AgRP neurons in adult animals causes starvation ^{6, 7}.

The POMC-derived peptide alpha-melanocyte-stimulating hormone (α MSH) and AgRP modulate the activity of the melanocortin 4 receptor (MC4R). α MSH activates MC4R while AgRP acts as an endogenous inverse agonist of MC4R, meaning that it suppresses constitutive MC4R activity and simultaneously antagonizes the effects of α MSH. The highest expression of MC4Rs is found in the hypothalamus and the brainstem ⁸⁻¹⁰. Target neurons expressing MC4Rs include neurons producing oxytocin, thyrotropin-releasing hormone, and corticotrophin-releasing hormone synthesizing neurons in the

paraventricular nucleus of the hypothalamus (PVH) ^{10, 11}. Remarkably, mutations in the MC4R gene are the most common monogenic disorders that cause obesity in humans ¹².

In this review, we will first describe the major steps of the melanocortin system development. We will also discuss the hormonal, molecular, cellular, and nutritional factors influencing the development of POMC and AgRP neurons.

Ontogenesis of the hypothalamic melanocortin system

POMC and AgRP neurons are derived from precursor cells located in the proliferative zone located in the inner and the lower portion of the third ventricle, also known as the neuroepithelium of the third ventricle ¹³ (**Figure 1**). Birth dating approaches using the thymidine analog bromodeoxyuridine 5-bromo-2'-deoxyuridine (BrdU) indicated that the majority of POMC and AgRP/NPY neurons in the mouse ARH are born primarily on E11-E12 ^{14, 15} (**Figure 1**). However, some POMC neurons, which are located more laterally in the ARH, are generated as late as E13. Gene expression studies showed that neurons in the presumptive ARH begin to express *Pomc* mRNA on E10–E12, whereas *Npy* mRNA expression is not observed until E14 ¹⁵. These observations are consistent with the early determination of *Pomc* cell fate. Intriguingly, genetic cell lineage tracing studies revealed that only a portion of embryonic *Pomc*-expressing precursors adopts a POMC fate in adult mice. About half of the *Pomc*-expressing precursors acquire a non-POMC fate in adult mice, and nearly one-quarter of the mature NPY neurons in the ARH share a common progenitor with POMC cells ¹⁵. These data show the unique property of *Pomc*-expressing progenitors with respect to giving rise to antagonistic neuronal populations. Differentiated neurons then send out axonal processes to other target nuclei to convey neuronal information and control behavior. The majority of arcuate POMC and AgRP neurons have relatively short axons and connect primarily to neurons within the hypothalamus ¹⁶. The development of axonal projections from the arcuate nucleus develop postnatally and

provide inputs to the PVH between postnatal day (P) 8 and P10 in mice ¹⁶ (**Figure 1**). Using immunohistochemical techniques, Grove and colleagues confirmed that projections immunopositive for AgRP are immature at birth and develop mainly during the second week of postnatal life in rats ¹⁷. The same temporal pattern was observed for the development of POMC projections in mice ¹⁸.

Mc4r mRNA is first expressed at E12 in the proliferative zone surrounding the lower portion of the third ventricle (also known as the neuroepithelium), and its expression peaks at E16 ¹⁹. These findings are particularly interesting because, as described above, it is known that neurons that compose various hypothalamic nuclei in adults are primarily derived from precursors that originate from this proliferative zone, raising the possibility that MC4R could be involved in hypothalamic neurogenesis. However, further studies are needed to determine the developmental stage during which MC4R becomes functional. Nevertheless, the observation that peripheral injection of the MC4R agonist melanotan II (MTII) reduces milk intake and body weight as early as during the first two weeks of postnatal life, suggests that MC4R receptors are present and functional in the hypothalamus at least soon after birth ²⁰. Consistent with this idea, *in situ* hybridization analysis showed that *Mc4r* mRNA is abundant in the hypothalamus, especially in the PVH at P10. That peripheral injection of MTII induces strong induction of cFos immunoreactivity (a marker of neuronal activation) in the PVH at P5-P15, further supports the functionality of MC4R in the PVH during early postnatal life ²⁰. Moreover, Melnick and colleagues showed an age-dependent increase in the electrophysiological response of specific sets of PVH neurons to melanocortins, with a maximal response observed at P28-P35 ²¹, suggesting that synapses between POMC and AgRP axons and PVH MC4R⁺ target neurons continue to build until puberty (**Figure 1**).

Consideration of species differences in melanocortin circuits development

The marked differences in the normal ontogeny of hypothalamic development between rodents and human and non-human primates warrant attention. First, the regional development of the rodent hypothalamus proceeds on a timeline of days in rodents *versus* weeks to months in human and non-human primates. Second, although rodents exhibit considerable postnatal hypothalamic development, human and non-human primates undergo considerably more prenatal maturation of hypothalamic circuits. For example, although the hypothalamus is not mature until after weaning in rodents, hypothalamic neurogenesis and axon growth occur primarily during intrauterine life in non-human primates, including humans. Hypothalamic development has been studied in detail by Kevin Grove and collaborators in Japanese macaques. They reported that *Pomc* and *Npy* mRNA-containing neurons are found in the ARH of NHP at gestational day (G) 100. Although only a few AgRP/NPY fibers and no POMC fibers are detected in the PVH at this age, the density of AgRP/NPY fibers innervating the PVH markedly increased at G130 and G170, and POMC fibers are found in the PVH at G170²². In human fetuses, NPY-immunoreactive fibers are detected in the ARH and the PVH as early as 21 weeks of gestation²³.

Hormonal factors that influence melanocortin neurons development

Leptin

The discovery of leptin led to a paradigm shift in understanding how food intake and body weight can be powerfully and dynamically regulated by hormonal signals²⁴⁻²⁶. In 1994, Friedman and colleagues used positional cloning and found that the *ob* gene encodes the hormone leptin secreted by the adipose tissue in proportion to its mass²⁴. Subsequently, other groups reported that leptin administration reduces body weight and food intake in leptin-deficient mice and humans²⁷⁻²⁹. A few years later, Ahima and colleagues reported

that circulating leptin levels exhibit a distinct surge between P8 and P12 in mice ³⁰, yet exogenous leptin does modulate food intake, growth, or energy expenditure at this developmental stage ³¹⁻³⁴. Instead of regulating food intake and body weight acutely, neonatal leptin appears to be an important neurodevelopmental factor that influences hypothalamic circuits development (**Figure 2**). Labeling of ARH axons combined with immunohistochemical analyses showed that POMC and AgRP neuronal projections are disrupted in leptin-deficient (*ob/ob*) mice ³⁵. The site of action for the developmental effects of leptin includes at least a direct action on ARH neurons because leptin induces neurite extension from isolated organotypic explants of the ARH *ex vivo* ³⁵. Remarkably, leptin appears to exert its developmental action on POMC neural projections during a discrete developmental critical period: exogenous leptin treatment up to P28 rescues AgRP projections in *ob/ob* mice ^{35, 36}. In contrast, leptin treatment of *ob/ob* mice after P28 is relatively ineffective because it did not increase the density of either POMC or AgRP fibers in the PVH to levels that are characteristic of wild-type mice ^{35, 36}. Together, these observations suggest the existence of a critical period for the neurotrophic effect of leptin on POMC and AgRP circuits that closes around puberty. More in-depth studies have examined leptin receptor signaling pathways that mediate the axonotrophic effect of leptin. The leptin receptor exists in several alternatively spliced isoforms, of which only the long-form (LepRb) associates with Janus kinase 2 to mediate intracellular signaling. LepRb initiates multiple intracellular signal transduction pathways upon leptin binding that result in the activation of STAT family transcription factors, extracellular signal-regulated kinases (ERK), and phosphoinositol-3 kinase/Akt. Developing POMC and AgRP neurons express LepRb ³⁷, and leptin administration in mouse neonates results in the activation of major LepRb signaling pathways, including STAT3, ERK, and Akt ^{37, 38}. Disruption of POMC and AgRP axonal projections is observed in mice or rats that lack functional LepRb signaling (*db/db* mice and *fa/fa* rats, respectively) ^{38, 39}. Moreover, lack of functional LepRb→STAT3

signaling *in vivo* (s/s mice) or *in vitro* results in a reduced density of POMC fibers without altering the development of AgRP projections, showing the importance of this signaling pathway, specifically, in the development of POMC neural projections³⁸. However, not all LepRb signaling pathways play a role in the formation of ARH projections. For example, mice that lack LepRb→ERK signaling (*l/l* mice) display comparable densities of POMC- and AgRP- axons in the PVH compared to wild-type mice³⁸.

Ghrelin and growth hormone

Ghrelin is a 28 amino acid hormone peptide that is mainly produced by the stomach and is an endogenous ligand for the growth hormone secretagogue receptor (GHSR). It is one of the most potent orexigenic signals that exerts its action on food intake by stimulating AgRP/NPY neurons that in turn inhibit POMC neurons^{40, 41}. Although the marked orexigenic effect of ghrelin is not yet present before weaning in mice or rats^{42, 43}, ghrelin in early postnatal life does have a lasting developmental effect on the hypothalamic circuits involved in energy homeostasis⁴³ (**Figure 2**). Mice injected with an anti-ghrelin compound during neonatal life display increased densities of POMC- and AgRP-containing axons innervating the PVH. These structural alterations are accompanied by long-term metabolic defects, including elevated body weight, increased adiposity, and hyperglycemia⁴³. However, if adult mice are treated with the anti-ghrelin compound, it does not alter POMC and AgRP circuits⁴³. These findings suggest that, similar to leptin, the developmental action of ghrelin on arcuate projections is restricted to a neonatal critical window. The site of action for the developmental effects of ghrelin likely includes direct action on arcuate neurons because direct exposure of isolated ARH explants to ghrelin inhibits axonal outgrowth⁴³. It also interacts with LepRb→STAT3 signaling to block the neurotrophic effect of leptin⁴³.

Ghrelin is a potent stimulator of growth hormone secretion ⁴⁴. Based on the documented finding that ghrelin influences hypothalamic development and can interact with leptin receptor signaling ⁴³, it is not surprising that deletion of the growth hormone receptor in *Lep^{rb}*-expressing cells also alters the development of POMC and AgRP neuronal circuits ⁴⁵. In addition, selective loss of the growth hormone receptor in AgRP neurons affects AgRP axonal projections without affecting POMC circuits ⁴⁵, demonstrating a cell-autonomous effect of growth hormone on AgRP neuronal development.

GLP1

The incretin hormone glucagon-like peptide 1 (GLP1) is secreted postprandially by intestinal enteroendocrine cells to promote satiety and glucose-induced insulin release ⁴⁶. Administration of the GLP1-R agonist exendin-4 during the first week of postnatal life decreases the density of NPY fibers innervating the PVH and has a protective effect against both age-related and diet-induced obesity ⁴⁷. Moreover, genetic deletion of *Glp1r* in *Sim1* neurons of the PVH reduces AgRP/NPY projections while it increases POMC projections to the PVH ⁴⁷.

Amylin

Amylin is a hormone produced by pancreatic β -cells and is co-released with insulin in response to caloric intake. The amylin receptor comprises the core calcitonin receptor (CTR), which heterodimerizes with one or several receptors activity-modifying proteins (RAMP-1, -2, and -3). The primary role of amylin in adults is to reduce food intake by promoting meal-ending satiation and maintaining glucose homeostasis. During development, amylin is detected in the blood circulation of embryos, where it appears to act through RMAP1-3 to influence neurogenesis of POMC neurons ⁴⁸ (**Figure 2**). During

postnatal life, amylin continues to be secreted in the blood circulation⁴⁹, and loss of *amylin* or *Ramp1/3* disrupts the development of POMC and AgRP projections to the PVH⁵⁰.

Molecular programs of melanocortin neurons development

Transcription factors

Homeobox genes belong to a class of transcription factors that play important roles in regionalization, patterning, and cell differentiation during embryogenesis and organ development. The homeobox genes *orthopedia (Otp)*, *Nkx2.1*, and *Bsx* are highly expressed in the ventral hypothalamus during embryonic development. Loss of function studies indicated that while *Nkx2.1* and *Otp* are essential for the normal morphological development of the hypothalamus, including the ARH, *Bsx* is not required⁵¹⁻⁵³ (**Figure 2**). The homeobox gene *Nkx2-1*, also known as thyroid transcription factor 1 (*Ttf-1*), plays a particularly important role in ARH specification. Ablation of *Nkx2-1* impairs the formation of the ventral hypothalamic primordium resulting in the absence of ARH formation^{51, 54}. However, the VMH, DMH, and LHA are present. Expression of *Nkx2-1* in postmitotic cells suggests that it further plays a role in differentiation and maintenance of ARH neurons in the ventral portion of the hypothalamus^{55, 56}. Deficiency in *Nkx2.1* prior to the onset of *Pomc* expression markedly reduces *Pomc* cell numbers⁵⁷. However, the number of NPY neurons was not affected in *Nkx2.1* knockout mice, and POMC neuronal cell number was not affected if *Nkx2.1* deletion occurs in postmitotic *Pomc* neurons⁵⁷. The LIM-homeodomain transcription factor *Islet 1 (Isl1)* is up-regulated in hypothalamic *Nkx2.1* progenitor cells at E10, *i.e.*, just before the onset of neuropeptide expression^{58, 59}. Consistent with the role of *Isl1* in the phenotypic determination of ARH neurons, loss of function studies revealed that *Isl1* promotes the terminal differentiation of *Pomc*, *Agrp*, and *Npy* expression^{58, 59}. In contrast, deletion of the transcription factors *Dlx1/2* in *Nkx2.1*-expressing progenitors increases *Agrp* expression without affecting *Pomc* expression⁶⁰.

More in-depth molecular studies revealed that *Dlx1/2* controls *Agrp* expression by binding to and repressing the expression of the homeodomain transcription factor orthopedia (*Otp*) that is also known to influence ARH morphogenesis^{52, 53, 60}. Thus, the loss of *Otp* in *Agrp* neurons results in a dramatic reduction in the number of *Agrp* mRNA-expressing cells⁶¹. Notably, *Otp* expression is absent in the hypothalamus of *Isl1* knockout embryos⁵⁸ suggesting that, in addition to *Dlx1/2*, *Isl1* is also required for the expression of *Otp* in the future ARH. Sonic hedgehog (*Shh*), SIX Homeobox 3 (*Six3*), and the retinal anterior neural fold homeobox (*Rax*) were also identified as critical regulators of ventral hypothalamus development and POMC development. *Shh* signaling increases *Nkx2-1* expression⁶² and deletion of *Shh* in *Nkx2-1* progenitors affects the development of POMC neurons⁶³. *Six3* is a regulator of forebrain development, including the hypothalamus⁶⁴, and is required for *Shh* expression⁶⁵. Finally, *Rax* is also important for the formation of the ventral neural tube. Mice lacking *Rax* in *Six3*-expressing cells do not show ventral hypothalamic *Nkx2.1* expression and never express POMC⁶⁶.

The Oligodendrocyte transcription factor family (*Olig1* and *Olig2*) are basic helix-loop-helix (bHLH) transcription factors highly expressed in the periventricular regions of the brain, such as the hypothalamus^{67, 68}. Lineage tracing experiments indicate that a number of POMC and NPY cells derive from *Olig1* progenitors⁶⁹. The majority of *Olig1* progenitors also express Bone morphogenetic protein receptor 1A (*Bmpr1A*)⁶⁹ and when *Bmpr1A* is deleted in *Olig1*-expressing cells, it decreased and increased the number of POMC and AgRP neurons, respectively⁶⁹. Neurogenin 3 (*Ngn3*) is another bHLH transcription factor expressed in hypothalamic progenitors, and it plays an opposite role in the specification of *Pomc* and *Npy* neurons: while it promotes the embryonic development of *Pomc* neurons, it inhibits *Npy* neuronal development^{70, 71}. However, not all POMC neurons are derived from *Ngn3* progenitors. Using a mouse model of *Mash1* deficiency, McNay and colleagues further reported that this bHLH transcription factor has

a pro-neural function and acts upstream of *Ngn3* to regulate neurogenesis in the ventral hypothalamus ⁷². Loss of *Mash1* results in a disappearance of *Ngn3* expression in ARH progenitors ⁷². Loss of *Mash1* blunts *Ngn3* expression in ARH progenitors and is associated with a dramatic reduction in the number of POMC and NPY neurons ⁷². The Notch signaling pathway also appears to mediate its developmental effects on POMC and NPY neuronal development through *Mash1*. Mice lacking Notch signaling in *Nkx2.1*-expressing cells display an increased number in POMC and NPY neurons associated with an induction in *Mash1* expression ^{72,73}. In addition, mice with a constitutively active *Notch1* intracellular domain show a complete loss in POMC and NPY neurons ⁷³, mirroring the effects of *Mash1* deficient mice ⁷².

As described above, *Pomc*-expressing progenitors in the ARH have the unique property to differentiate into functional mature NPY neurons ¹⁵. Recent data from our laboratory investigated the molecular mechanisms involved in this developmental switch and identified miR-103/107 as candidates involved in *Pomc* progenitors differentiation (**Figure 2**). Loss of the microRNA(miRNA)-processing enzyme *Dicer* increases the proportion of *Pomc* progenitors acquiring an NPY phenotype ⁷⁴. Moreover, silencing of miR-103/107, specifically, decreases the number of *Pomc*-expressing cells and increases the proportion of *Pomc* progenitors differentiating into NPY neurons ⁷⁴. Postnatal maintenance of *Pomc* and *Npy* peptidergic identity also depends on the expression of the transcription factor T-box 3 (*Tbx3*) ⁷⁵. Because the majority of miRNAs exert their effects on gene expression by targeting transcription factors, it would be interesting to study whether there is a link between miRNAs, *Tbx3*, and *Pomc* and *Npy* gene expression.

Axon guidance molecules

Axons grow by sending out a highly plastic and sensitive structure called a “growth cone,” which travels toward the target and trails behind it the elongating neurite. As described

above, metabolic hormones, including leptin, are critical factors influencing initial POMC and AgRP/NPY axon outgrowth. Growing POMC axons must then choose a path to follow and decide the direction to go on this path to innervate the proper nucleus (e.g., the PVH). The pathways are defined by cell-cell interactions and diffusible chemorepulsive and chemoattractive cues ⁷⁶. The diffusible axon guidance cues Semaphorins are highly expressed in the PVH during development, and POMC neurons express the semaphorin receptors neuropilin 1 and 2 ⁷⁷. Supporting a role for neuropilins/semaphorins in POMC axon guidance, a loss of neuropilin 2 receptors in POMC neurons disrupts the development of POMC axonal projections to the PVH, specifically ⁷⁷. These structural alterations are accompanied by metabolic dysregulation, including increased body weight and glucose intolerance ⁷⁷. Notably, exome sequencing experiments identified variants in the semaphorin and neuropilin families associated with severe obesity in humans ⁷⁷, demonstrating the translational importance of these findings. The formation of POMC neural neuronal connectivity also involves cell cell-contacting proteins. Supporting this idea, POMC neurons are enriched in cell-to-cell contact molecules *Efnb1* (EphrinB1) and *Efnb2* (EphrinB2) during postnatal development, and loss of *Efnb1* or *Efnb2* in *Pomc*-expressing progenitors decreases the amount of excitatory glutamatergic inputs ⁷⁸. In addition, mice that are deficient in *contactin*, a cell adhesion molecule involved in the formation of axonal projections, have a reduced density of POMC fibers in the PVH during postnatal development ⁷⁹.

Cellular factors underlying melanocortin neurons development

Autophagy

Axonal growth involves a dynamic remodeling of cytosolic structures and requires protein degradation and turnover to replace damaged organelles and proteins. The maintenance of cell function and growth is achieved with autophagy, which is an important cellular

degradation system that engulfs parts of the cytoplasm and organelles within double-membrane vesicles, known as autophagosomes, to turnover and recycle these cellular constituents⁸⁰. This cellular process is also critical in the supply of nutrients for survival during starvation⁸¹. Constitutive autophagy is detected in the hypothalamus, including in arcuate POMC neurons, during critical periods of axon growth and development⁸². Loss of the autophagy-related protein (*Atg*) gene *Atg7* disrupts the maturation of POMC axonal projections and causes lifelong metabolic perturbations⁸² (**Figure 2**). As described above, leptin is a critical neurotrophic factor for POMC circuits, and direct crosstalk has been described between leptin and hypothalamic autophagy during perinatal life⁸³. Supporting a role for autophagy in mediating the trophic effects of leptin, the loss of autophagy in POMC neurons exacerbates the metabolic and neurodevelopmental deficits observed in leptin-deficient *ob/ob* mice⁸³.

Primary cilia

Another cellular signaling system that plays an important role in brain development and function is the primary cilium, which as described above is an organelle found in the cell surface of most mammalian cells, including hypothalamic neurons⁸⁴. For example, during embryonic development, primary cilia are important mediators of sonic hedgehog signaling, which is a critical regulator of the ventral patterning of the hypothalamus⁸⁵. Moreover, a strong interaction between primary cilia and autophagy has been reported, including in the developing hypothalamus⁸⁶. Cilia begin to be observed in hypothalamic neurons on E12, and the number and length of primary cilia gradually increase thereafter to reach an adult-like pattern at P14⁸⁶. Disruption of cilia formation in developing POMC neurons, but not in adult POMC neurons, increases body weight, fat mass, food intake, reduces energy expenditure, and alters glucose homeostasis during adult life⁸⁶. Neuroanatomically, a reduction in the number of POMC neurons was observed in juvenile

mice lacking primary cilia on POMC neurons, but POMC cell numbers were normal in adult mutant mice. This reduction in POMC cell numbers during the pre-weaning periods is attributed to a decrease in neurogenesis during embryonic development (in opposition to an effect on apoptosis) and the adult normalization in POMC cell number to a compensatory increase in neurogenesis after weaning⁸⁶. Loss of primary cilia also alters axonal and dendritic growth resulting in a reduced density of POMC fibers innervating the PVH, DMH, and LHA⁸⁶ (**Figure 2**). It also blocks the ability of leptin to promote the development of POMC projections in *ob/ob* mice⁸⁶. Because a sub-population of *Pomc* progenitor cells also give birth to NPY neurons (see above and¹⁵), it is not surprising that mice lacking primary cilia from embryonic POMC neurons also display a reduction in NPY cells numbers and NPY fibers innervating the PVH⁸⁶.

Effect of perinatal nutrition on the development of melanocortin system

Obesity has reached alarming rates worldwide and is associated with several life-threatening conditions, such as hypertension and type 2 diabetes. Obesity among pregnant women is also at an all-time high, and it represents a significant risk factor for the development of metabolic diseases in the offspring. Obesity is determined by genetics and obesogenic environments, such as diets rich in fat and/or sugar. Maternal high-fat diet (HFD) feeding during pregnancy and/or lactation in rodents is a useful experimental approach for studying the mechanisms underlying maternal obesity. Similar to what is observed in humans, offspring born to obese females fed an HFD (45–60% of calories from fat) during gestation and/or lactation becomes progressively overweight, hyperphagic, and glucose intolerant, and they display an increase in adiposity^{87,88}. These metabolic alterations are associated with a disrupted development of POMC and AgRP/NPY projections to the PVH in both mice and rats⁸⁸⁻⁹¹ (**Figure 2**). Notably, maternal consumption of HFD during lactation alone (but not during pregnancy alone) appears

sufficient to cause obesity and diabetes and alter the development of POMC and AgRP projections⁸⁹, showing the importance of postnatal nutrition, specifically, in hypothalamic programming. A similar reduction in AgRP axonal projections has been reported in non-human primates following maternal HFD exposure⁹². The rat model of diet-induced obesity (DIO) developed by Barry Levin also provides a valuable tool for studying obesity, in part because Levin's DIO rats share several features with human obesity, including polygenic inheritance⁹³. Therefore, this animal model is particularly well suited for studying the relative contribution of genetic *versus* environmental factors in metabolic programming. Animals born to genetically obese-prone DIO dams display a reduction in the number of POMC and AgRP axons innervating the PVH⁹⁴. In addition, a significant remodeling of synapses onto POMC neurons has been observed in DIO rats, particularly in response to nutritional challenges⁹⁵. DIO rats fed a chow diet display increased inhibitory inputs to POMC neurons compared to obesity-resistant rats. In addition, DIO rats fed an HFD display a loss of synapses onto POMC neurons, whereas high-fat feeding in control obesity-resistant rats causes an increase in POMC synaptic coverage⁹⁵.

The precise mechanisms that underlie the maternal obesity-induced alterations in melanocortin system development only begin to be elucidated. Several studies have indicated that abnormal leptin and insulin signaling during postnatal development may represent a likely cause for the HFD- and DIO-induced alterations in hypothalamic development. For example, DIO rats and animals born to obese dams display an abnormal organization of projections pathways derived from the ARH that appear to be the result of the diminished responsiveness of ARH neurons to the trophic actions of leptin during critical periods of postnatal development⁹⁴. Moreover, animals born to obese dams are hyperleptinemic associated with hypothalamic leptin resistance and improving leptin sensitivity with the endoplasmic reticulum stress-relieving drug tauroursodeoxycholic acid normalized metabolic and neurodevelopmental deficits in these animals⁹⁰. At the cellular

level, a reduction in cilia length and frequency has been reported in the ARH of pups born to obese dams, suggesting that alteration of this cellular system critical for hypothalamic development and leptin signaling could contribute to obesity-induced perturbations of hypothalamic development⁸⁶. Changes in insulin signaling could also mediate the neurodevelopmental effects of maternal obesity. Mothers fed an HFD and their offspring are hyperinsulinemic, and deleting the insulin receptor in POMC neurons prevents the diet-induced disruption of POMC projections⁸⁹. More recently, Dearden and colleagues reported that offspring of obese dams display a reduction in POMC cell numbers that likely results from a diminished neurogenic action of insulin during embryonic development⁹⁶. Similarly, amylin, which is co-released with insulin by pancreatic β -cells, appears to be involved in the nutritional programming of melanocortin circuits. Offspring of obese dams are hyperamylinemic from embryonic age throughout adulthood. Amylin fails to activate hypothalamic amylin receptor signaling if animals are born to obese mothers, and it was associated with an inability of amylin to promote POMC neurogenesis⁴⁸. Similarly, knocking down CTR in the ventromedial part of the hypothalamus of DR rats (a region that encompasses the ARH+VMH) alters the development of POMC circuits⁹⁷. In contrast, neonatal amylin treatment in DIO rats enhances STAT3 signaling in the ARH accompanied with a restoration of AgRP and POMC fibers⁹⁷.

In part because of the importance of postnatal brain development in rodents, including that of POMC and AgRP neurons, animal models of postnatal metabolic programming have also been extensively studied. The reduction of litter size in rodents has proven to be very useful model to manipulate postnatal diet, specifically. This model was first characterized by pioneer works of Kennedy, Widdowson, and McCance that reported more than 60 years ago that pups raised in small litters (SL) display an almost four-fold difference in pre-weaning growth compared to pups raised in large litters^{98, 99}. This weight difference persisted throughout life and even increased as the rats were

weaned onto a regular chow diet. Similar to animals born to obese dams, pups raised in SL display early life hyperleptinemia associated with hypothalamic leptin resistance^{100, 101}. Postnatal overnutrition also results in reduced responsiveness of hypothalamic neurons to ghrelin during neonatal life that appears to be the consequence of an altered transport of the hormone across the blood-brain barrier¹⁰². These endocrine changes during critical periods of postnatal development are associated with dysregulation of the melanocortin system (**Figure 2**). A reduced arcuate *Pomc* expression has been reported in rats raised in SL^{103 87}. It is known that epigenetic mechanisms of gene regulation, such as DNA methylation and histone modifications, such as methylation, regulate gene expression in response to environmental stimuli. Andreas Plagemann and colleagues therefore studied the methylation status of CpG dinucleotides of the *Pomc* promoter in rats raised in SL. They found that postnatal overfeeding causes alterations in the methylation of the *Pomc* promoter, particularly on the CpG dinucleotides within the two *Sp1*-related binding sequences (*Sp1*, NF- κ B) that are critical for leptin and insulin's effects on *Pomc* gene expression¹⁰⁴. Similarly, human studies showed that childhood obesity is associated with *Pomc* hypermethylation¹⁰⁵. In addition to alter *Pomc* gene expression in the ARH, postnatal overfeeding also disrupts the neurophysiological responses of PVH neurons to α MSH and AgRP¹⁰⁶.

Conclusion

It is now clear from a variety of epidemiological and experimental studies that many chronic diseases, including obesity and type 2 diabetes, may have their roots during neonatal development. A perturbed environment in early life is thought to elicit a range of cellular adaptive responses in key systems, the melanocortin system being one of them. In rodents, the development of the melanocortin system is initiated during mid-gestation and continues during the postnatal period. These developmental windows represent

important periods of vulnerability during which perturbations in the perinatal environment may lead to abnormal POMC and AgRP neuron development, causing lifelong metabolic diseases. Therefore, it will be critical to have a comprehensive knowledge of factors that are detrimental or beneficial for the development of the melanocortin system if we want to design intervention studies or treatments. The marked species difference in terms of hypothalamic developmental trajectories is important to consider as rodents and humans may exhibit different periods of vulnerability to developmental insults or different responses to therapeutic interventions based on the temporal and regional maturation patterns of the hypothalamus. Nevertheless, the mature hypothalamus whether it is from rodents or primates can still exhibit neuroplastic responses, although the degree and the nature of hypothalamic remodeling may differ between adults and neonates.

Acknowledgments

Work from the author's laboratory is supported by the French National Institute of Health (Inserm, grant 1172) and the Foundation for Prader-Willi Research. Figures were created with BioRender.com

Conflict of interest

The author declared that no competing interests exist

References

1. Poggioli, R., Vergoni, A.V. & Bertolini, A. ACTH-(1-24) and alpha-MSH antagonize feeding behavior stimulated by kappa opiate agonists. *Peptides* **7**, 843-848 (1986).
2. Belgardt, B.F. & Brüning, J.C. CNS leptin and insulin action in the control of energy homeostasis. *Annals of the New York Academy of Sciences* **1212**, 97-113 (2010).
3. Williams, K.W. & Elmquist, J.K. From neuroanatomy to behavior: central integration of peripheral signals regulating feeding behavior. *Nature Neuroscience* **15**, 1350-1355 (2012).
4. Aponte, Y., Atasoy, D. & Sternson, S.M. AGRP neurons are sufficient to orchestrate feeding behavior rapidly and without training. *Nat Neurosci* **14**, 351-355 (2011).
5. Zhan, C. *et al.* Acute and Long-Term Suppression of Feeding Behavior by POMC Neurons in the Brainstem and Hypothalamus, Respectively. *The Journal of Neuroscience* **33**, 3624-3632 (2013).
6. Luquet, S., Perez, F.A., Hnasko, T.S. & Palmiter, R.D. NPY/AgRP Neurons Are Essential for Feeding in Adult Mice but Can Be Ablated in Neonates. *Science* **310**, 683-685 (2005).
7. Groppe, E. *et al.* Agouti-related peptide-expressing neurons are mandatory for feeding. *Nat Neurosci* **8**, 1289-1291 (2005).
8. Mountjoy, K.G., Mortrud, M.T., Low, M.J., Simerly, R.B. & Cone, R.D. Localization of the melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the brain. *Molecular Endocrinology* **8**, 1298-1308 (1994).
9. Kishi, T. *et al.* Expression of melanocortin 4 receptor mRNA in the central nervous system of the rat. *The Journal of Comparative Neurology* **457**, 213-235 (2003).
10. Liu, H. *et al.* Transgenic mice expressing green fluorescent protein under the control of the melanocortin-4 receptor promoter. *J Neurosci* **23**, 7143-7154 (2003).
11. Lu, X.Y., Barsh, G.S., Akil, H. & Watson, S.J. Interaction between alpha-melanocyte-stimulating hormone and corticotropin-releasing hormone in the regulation of feeding and hypothalamo-pituitary-adrenal responses. *J Neurosci* **23**, 7863-7872 (2003).
12. Coll, A.P., Farooqi, I.S. & O'Rahilly, S. The Hormonal Control of Food Intake. *Cell* **129**, 251-262 (2007).
13. Sauer, F.C. Mitosis in the neural tube. *The Journal of Comparative Neurology* **62**, 377-405 (1935).
14. Khachaturian, H. *et al.* Development of hypothalamic opioid neurons: A combined immunocytochemical and [3H]thymidine autoradiographic study. *Neuropeptides* **5**, 477-480 (1985).
15. Padilla, S.L., Carmody, J.S. & Zeltser, L.M. Pomc-expressing progenitors give rise to antagonistic neuronal populations in hypothalamic feeding circuits. *Nature Medicine* **16**, 403-405 (2010).
16. Bouret, S.G., Draper, S.J. & Simerly, R.B. Formation of Projection Pathways from the Arcuate Nucleus of the Hypothalamus to Hypothalamic Regions Implicated in the Neural Control of Feeding Behavior in Mice. *J. Neurosci.* **24**, 2797-2805 (2004).
17. Grove, K.L., Allen, S., Grayson, B.E. & Smith, M.S. Postnatal development of the hypothalamic neuropeptide Y system. *Neuroscience* **116**, 393-406 (2003).
18. Nilsson, I., Johansen, J.E., Schalling, M., Hokfelt, T. & Fetissov, S.O. Maturation of the hypothalamic arcuate agouti-related protein system during postnatal development in the mouse. *Developmental Brain Research* **155**, 147-154 (2005).

19. Mountjoy, K.G. & Wild, J.M. Melanocortin-4 receptor mRNA expression in the developing autonomic and central nervous systems. *Developmental Brain Research* **107**, 309-314 (1998).
20. Glavas, M.M., Joachim, S.E., Draper, S.J., Smith, M.S. & Grove, K.L. Melanocortinergic Activation by Melanotan II Inhibits Feeding and Increases Uncoupling Protein 1 Messenger Ribonucleic Acid in the Developing Rat. *Endocrinology* **148**, 3279-3287 (2007).
21. Melnick, I., Pronchuck, N., Cowley, M.A., Grove, K.L. & Colmers, W.F. Developmental switch in neuropeptide Y and melanocortin effects in the paraventricular nucleus of the hypothalamus. *Neuron* **56**, 1103-1115 (2007).
22. Grayson, B.E. *et al.* Prenatal development of hypothalamic neuropeptide systems in the nonhuman primate. *Neuroscience* **143**, 975-986 (2006).
23. Yuri Koutcherov, J.r.K.M.K.W.S.A.G.P. Organization of human hypothalamus in fetal development. *The Journal of Comparative Neurology* **446**, 301-324 (2002).
24. Zhang, Y. *et al.* Position cloning of the mouse obese gene and its human homologue. *Nature* **372**, 425 (1994).
25. Kojima, M. *et al.* Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* **402**, 656-660 (1999).
26. Nakazato, M. *et al.* A role for ghrelin in the central regulation of feeding. *Nature* **409**, 194-198 (2001).
27. Halaas, J.L. *et al.* Weight-reducing effects of the plasma protein encoded by the obese gene. *Science* **269**, 543-546 (1995).
28. Pelleymounter, M. *et al.* Effects of the obese gene product on body weight regulation in ob/ob mice. *Science* **269**, 540-543 (1995).
29. Campfield, L.A., Smith, F.J., Guisez, Y., Devos, R. & Burn, P. Recombinant mouse OB protein: evidence for a peripheral signal linking adiposity and central neural networks. *Science* **269**, 546-549 (1995).
30. Ahima, R., Prabakaran, D. & Flier, J. Postnatal leptin surge and regulation of circadian rhythm of leptin by feeding. Implications for energy homeostasis and neuroendocrine function. *J Clin Invest* **101**, 1020-1027 (1998).
31. Ahima, R.S. & Hileman, S.M. Postnatal regulation of hypothalamic neuropeptide expression by leptin: implications for energy balance and body weight regulation. *Regul Pept* **92**, 1-7 (2000).
32. Proulx, K., Richard, D. & Walker, C.-D. Leptin Regulates Appetite-Related Neuropeptides in the Hypothalamus of Developing Rats without Affecting Food Intake. *Endocrinology* **143**, 4683-4692 (2002).
33. Mistry, A., Swick, A. & Romsos, D. Leptin alters metabolic rates before acquisition of its anorectic effect in developing neonatal mice. *Am J Physiol* **277**, R742-747 (1999).
34. Schmidt, I. *et al.* The effect of leptin treatment on the development of obesity in overfed suckling Wistar rats. *International journal of obesity and related metabolic disorders : journal of the International Association for the Study of Obesity* **25**, 1168-1174 (2001).
35. Bouret, S.G., Draper, S.J. & Simerly, R.B. Trophic Action of Leptin on Hypothalamic Neurons That Regulate Feeding. *Science* **304**, 108-110 (2004).
36. Kamitakahara, A., Bouyer, K., Wang, C.H. & Simerly, R. A critical period for the trophic actions of leptin on AgRP neurons in the arcuate nucleus of the hypothalamus. *J Comp Neurol* (2017).
37. Caron, E., Sachot, C., Prevot, V. & Bouret, S.G. Distribution of Leptin-Sensitive Cells in the Postnatal and Adult Mouse Brain. *Journal of Comparative Neurology* **518**, 459-476 (2010).

38. Bouret, S.G., Bates, S.H., Chen, S., Myers, M.G. & Simerly, R.B. Distinct Roles for Specific Leptin Receptor Signals in the Development of Hypothalamic Feeding Circuits. *The Journal of Neuroscience* **32**, 1244-1252 (2012).
39. Bouret, S. & Simerly, R.B. Development of Leptin-Sensitive Circuits. *J Neuroendocrinology* **19**, 575-582 (2007).
40. Tschöp, M., Smiley, D.L. & Heiman, M.L. Ghrelin induces adiposity in rodent. *Nature*, 908-913 (2000).
41. Cowley, M.A. *et al.* The distribution and mechanism of action of ghrelin in the CNS demonstrates a novel hypothalamic circuits regulating energy homeostasis. *Neuron* **37**, 649-661 (2003).
42. Piao, H. *et al.* Ghrelin Stimulates Milk Intake by Affecting Adult Type Feeding Behaviour in Postnatal Rats. *Journal of Neuroendocrinology* **20**, 330-334 (2008).
43. Steculorum SM, C.G., Coupe B, Croizier S, Andrews Z, Jarosch F, Klussmann S, Bouret SG. Ghrelin programs development of hypothalamic feeding circuits. *The Journal of Clinical Investigation* **125**, 846–858 (2015).
44. Tolle, V. *et al.* In vivo and in vitro effects of ghrelin/motilin-related peptide on growth hormone secretion in the rat. *Neuroendocrinology* **73**, 54-61 (2001).
45. Wasinski, F. *et al.* Growth Hormone Receptor Deletion Reduces the Density of Axonal Projections from Hypothalamic Arcuate Nucleus Neurons. *Neuroscience* **434**, 136-147 (2020).
46. Drucker, D.J. Glucagon-like peptides. *Diabetes* **47**, 159-169 (1998).
47. Rozo, A.V. *et al.* Neonatal GLP1R activation limits adult adiposity by durably altering hypothalamic architecture. *Mol Metab* **6**, 748-759 (2017).
48. Li, C. *et al.* Amylin receptor insensitivity impairs hypothalamic POMC neuron differentiation in the male offspring of maternal high-fat diet-fed mice. *Mol Metab* **44**, 101135 (2020).
49. Abegg, K. *et al.* Involvement of Amylin and Leptin in the Development of Projections from the Area Postrema to the Nucleus of the Solitary Tract. *Front Endocrinol (Lausanne)* **8**, 324 (2017).
50. Lutz, T.A. *et al.* Amylin Selectively Signals Onto POMC Neurons in the Arcuate Nucleus of the Hypothalamus. *Diabetes* **67**, 805-817 (2018).
51. Kimura, S. *et al.* The *T/ebp* null mouse: thyroid-specific enhancer-binding protein is essential for the organogenesis of the thyroid, lung, ventral forebrain, and pituitary. *Genes & Development* **10**, 60-69 (1996).
52. Acampora, D. *et al.* Progressive impairment of developing neuroendocrine cell lineages in the hypothalamus of mice lacking the *Orthopedia* gene. *Genes & development* **13**, 2787-2800 (1999).
53. Wang, W. & Lufkin, T. The murine *Otp* homeobox gene plays an essential role in the specification of neuronal cell lineages in the developing hypothalamus. *Dev Biol* **227**, 432-449 (2000).
54. Marín, O., Baker, J., Puelles, L. & Rubenstein, J.L.R. Patterning of the basal telencephalon and hypothalamus is essential for guidance of cortical projections. *Development* **129**, 761-773 (2002).
55. Sussel, L., Marin, O., Kimura, S. & Rubenstein, J.L. Loss of *Nkx2.1* homeobox gene function results in a ventral to dorsal molecular respecification within the basal telencephalon: evidence for a transformation of the pallidum into the striatum. *Development* **126**, 3359-3370 (1999).
56. Yee, C.L., Wang, Y., Anderson, S., Ekker, M. & Rubenstein, J.L.R. Arcuate nucleus expression of *NKX2.1* and *DLX* and lineages expressing these transcription factors in neuropeptide Y+, proopiomelanocortin+, and tyrosine

- hydroxylase+ neurons in neonatal and adult mice. *Journal of Comparative Neurology* **517**, 37-50 (2009).
57. Orquera, D.P. *et al.* The Homeodomain Transcription Factor NKX2.1 Is Essential for the Early Specification of Melanocortin Neuron Identity and Activates Pomc Expression in the Developing Hypothalamus. *J Neurosci* **39**, 4023-4035 (2019).
 58. Nasif, S. *et al.* Islet 1 specifies the identity of hypothalamic melanocortin neurons and is critical for normal food intake and adiposity in adulthood. *Proc Natl Acad Sci U S A* **112**, E1861-1870 (2015).
 59. Lee, B., Lee, S., Lee, S.K. & Lee, J.W. The LIM-homeobox transcription factor Isl1 plays crucial roles in the development of multiple arcuate nucleus neurons. *Development* **143**, 3763-3773 (2016).
 60. Lee, B. *et al.* Dlx1/2 and Otp coordinate the production of hypothalamic GHRH- and AgRP-neurons. *Nat Commun* **9**, 2026 (2018).
 61. Hu, Y. *et al.* Development and characterization of an Otp conditional loss of function allele. *Genesis (New York, N.Y. : 2000)* **58**, e23370 (2020).
 62. Manning, L. *et al.* Regional Morphogenesis in the Hypothalamus: A BMP-Tbx2 Pathway Coordinates Fate and Proliferation through Shh Downregulation. *Developmental Cell* **11**, 873-885 (2006).
 63. Shimogori, T. *et al.* A genomic atlas of mouse hypothalamic development. *Nat Neurosci* **13**, 767-775 (2010).
 64. Lagutin, O.V. *et al.* Six3 repression of Wnt signaling in the anterior neuroectoderm is essential for vertebrate forebrain development. *Genes & Development* **17**, 368-379 (2003).
 65. Geng, X. *et al.* Haploinsufficiency of Six3 fails to activate Sonic hedgehog expression in the ventral forebrain and causes holoprosencephaly. *Dev Cell* **15**, 236-247 (2008).
 66. Lu, F. *et al.* Rax is a selector gene for mediobasal hypothalamic cell types. *J Neurosci* **33**, 259-272 (2013).
 67. Takebayashi, H. *et al.* Dynamic expression of basic helix-loop-helix Olig family members: implication of Olig2 in neuron and oligodendrocyte differentiation and identification of a new member, Olig3. *Mech Dev* **99**, 143-148 (2000).
 68. Zhou, Q., Wang, S. & Anderson, D.J. Identification of a novel family of oligodendrocyte lineage-specific basic helix-loop-helix transcription factors. *Neuron* **25**, 331-343 (2000).
 69. Peng, C.Y., Mukhopadhyay, A., Jarrett, J.C., Yoshikawa, K. & Kessler, J.A. BMP receptor 1A regulates development of hypothalamic circuits critical for feeding behavior. *J Neurosci* **32**, 17211-17224 (2012).
 70. Pelling, M. *et al.* Differential requirements for neurogenin 3 in the development of POMC and NPY neurons in the hypothalamus. *Dev Biol* **349**, 406-416 (2011).
 71. Anthwal, N. *et al.* Conditional deletion of neurogenin-3 using Nkx2.1iCre results in a mouse model for the central control of feeding, activity and obesity. *Dis Model Mech* **6**, 1133-1145 (2013).
 72. McNay, D.E.G., Pelling, M., Claxton, S., Guillemot, F.B. & Ang, S.-L. Mash1 Is Required for Generic and Subtype Differentiation of Hypothalamic Neuroendocrine Cells. *Molecular Endocrinology* **20**, 1623-1632 (2006).
 73. Aujla, P.K., Naratadam, G.T., Xu, L. & Raetzman, L.T. Notch/Rbpjk signaling regulates progenitor maintenance and differentiation of hypothalamic arcuate neurons. *Development* **140**, 3511-3521 (2013).
 74. Crozier, S., Park, S., Maillard, J. & Bouret, S.G. Central Dicer-miR-103/107 controls developmental switch of POMC progenitors into NPY neurons and impacts glucose homeostasis. *Elife* **7** (2018).

75. Quarta, C. *et al.* Functional identity of hypothalamic melanocortin neurons depends on Tbx3. *Nat Metab* **1**, 222-235 (2019).
76. Tessier-Lavigne, M. & Goodman, C.S. The molecular biology of axon guidance. *Science* **274**, 1123-1133 (1996).
77. van der Klaauw, A.A. *et al.* Human Semaphorin 3 Variants Link Melanocortin Circuit Development and Energy Balance. *Cell* **176**, 729-742.e718 (2019).
78. Gervais, M., Labouèbe, G., Picard, A., Thorens, B. & Croizier, S. EphrinB1 modulates glutamatergic inputs into POMC-expressing progenitors and controls glucose homeostasis. *PLoS biology* **18**, e3000680 (2020).
79. Nilsson, I., Johansen, J.E., Schalling, M., H^opfelt, T. & Fetissov, S.O. Maturation of the hypothalamic arcuate agouti-related protein system during postnatal development in the mouse. *Developmental Brain Research* **155**, 147-154 (2005).
80. Klionsky, D.J. Autophagy: from phenomenology to molecular understanding in less than a decade. *Nature Reviews Molecular Cell Biology* **8**, 931-937 (2007).
81. Komatsu, M. *et al.* Impairment of starvation-induced and constitutive autophagy in Atg7-deficient mice. *The Journal of Cell Biology* **169**, 425-434 (2005).
82. Coupe, B. *et al.* Loss of Autophagy in Pro-opiomelanocortin Neurons Perturbs Axon Growth and Causes Metabolic Dysregulation. *Cell Metabolism* **15**, 247-255 (2012).
83. Park, S., Aintablian, A., Coupe, B. & Bouret, S.G. The endoplasmic reticulum stress-autophagy pathway controls hypothalamic development and energy balance regulation in leptin-deficient neonates. *Nat Commun* **11**, 1914 (2020).
84. Fuchs, J.L. & Schwark, H.D. Neuronal primary cilia: a review. *Cell Biol Int* **28**, 111-118 (2004).
85. Carreno, G. *et al.* Hypothalamic sonic hedgehog is required for cell specification and proliferation of LHX3/LHX4 pituitary embryonic precursors. *Development* **144**, 3289-3302 (2017).
86. Lee, C.H. *et al.* Primary cilia mediate early life programming of adiposity through lysosomal regulation in the developing mouse hypothalamus. *Nat Commun* **11**, 5772 (2020).
87. Chen, H., Simar, D. & Morris, M.J. Hypothalamic Neuroendocrine Circuitry is Programmed by Maternal Obesity: Interaction with Postnatal Nutritional Environment. *PLoS ONE* **4**, e6259 (2009).
88. Kirk, S.L. *et al.* Maternal Obesity Induced by Diet in Rats Permanently Influences Central Processes Regulating Food Intake in Offspring. *PLoS ONE* **4**, e5870 (2009).
89. Vogt, M.C. *et al.* Neonatal Insulin Action Impairs Hypothalamic Neurocircuit Formation in Response to Maternal High-Fat Feeding. *Cell* **156**, 495-509 (2014).
90. Park, S., Jang, A. & Bouret, S.G. Maternal obesity-induced endoplasmic reticulum stress causes metabolic alterations and abnormal hypothalamic development in the offspring. *PLoS biology* **18**, e3000296 (2020).
91. Haddad-Tóvolli, R. *et al.* Pro-opiomelanocortin (POMC) neuron transcriptome signatures underlying obesogenic gestational malprogramming in mice. *Mol Metab* **36**, 100963 (2020).
92. Sullivan, E.L. *et al.* Maternal and postnatal high-fat diet consumption programs energy balance and hypothalamic melanocortin signaling in nonhuman primate offspring. *Am J Physiol Regul Integr Comp Physiol* **313**, R169-r179 (2017).
93. Levin, B.E., Dunn-Meynell, A.A., Balkan, B. & Keesey, R.E. Selective breeding for diet-induced obesity and resistance in Sprague-Dawley rats. *Am J Physiol Regul Integr Comp Physiol* **273**, R725-730 (1997).

94. Bouret, S.G. *et al.* Hypothalamic Neural Projections Are Permanently Disrupted in Diet-Induced Obese Rats. *Cell Metabolism* **7**, 179-185 (2008).
95. Horvath, T.L. *et al.* Synaptic input organization of the melanocortin system predicts diet-induced hypothalamic reactive gliosis and obesity. *Proc Natl Acad Sci USA* **107**, 14875-14880 (2010).
96. Dearden, L., Buller, S., Furigo, I.C., Fernandez-Twinn, D.S. & Ozanne, S.E. Maternal obesity causes fetal hypothalamic insulin resistance and disrupts development of hypothalamic feeding pathways. *Mol Metab* **42**, 101079 (2020).
97. Johnson, M.D. *et al.* Early postnatal amylin treatment enhances hypothalamic leptin signaling and neural development in the selectively bred diet-induced obese rat. *Am J Physiol Regul Integr Comp Physiol* **311**, R1032-r1044 (2016).
98. Kennedy, G.C. The development with age of hypothalamic restraint upon the appetite of the rat. . *J Endocrinol* **16**, 9/17 (1957).
99. Widdowson, E.M. & Mc, C.R. Some effects of accelerating growth. I. General somatic development. *Proc R Soc Lond B Biol Sci* **152**, 188-206 (1960).
100. Glavas, M.M. *et al.* Early Overnutrition Results in Early-Onset Arcuate Leptin Resistance and Increased Sensitivity to High-Fat Diet. *Endocrinology* **151**, 1598-1610 (2010).
101. Roberts, B.L., Bennett, C.M., Carroll, J.M., Lindsley, S.R. & Kievit, P. Early overnutrition alters synaptic signaling and induces leptin resistance in arcuate proopiomelanocortin neurons. *Physiol Behav* **206**, 166-174 (2019).
102. Collden, G. *et al.* Neonatal overnutrition causes early alterations in the central response to peripheral ghrelin. *Molecular Metabolism* **4**, 15-24 (2015).
103. Srinivasan, M. *et al.* A high-carbohydrate diet in the immediate postnatal life of rats induces adaptations predisposing to adult-onset obesity. *J Endocrinol* **197**, 565-574 (2008).
104. Plagemann, A. *et al.* Hypothalamic proopiomelanocortin promoter methylation becomes altered by early overfeeding: an epigenetic model of obesity and the metabolic syndrome. *The Journal of Physiology* **587**, 4963-4976 (2009).
105. Kuehnen, P. *et al.* An Alu Element-Associated Hypermethylation Variant of the POMC Gene Is Associated with Childhood Obesity. *PLoS Genet* **8**, e1002543 (2012).
106. Davidowa, H., Li, Y. & Plagemann, A. Altered responses to orexigenic (AGRP, MCH) and anorexigenic (α -MSH, CART) neuropeptides of paraventricular hypothalamic neurons in early postnatally overfed rats. *European Journal of Neuroscience* **18**, 613-621 (2003).

Figures and Figure legends




Figure 1. Important periods of hypothalamic development. The development of a functional hypothalamus occurs in two major phases: the determination of cell numbers, which includes neurogenesis, neuronal migration, and cell fate, and the formation of functional circuits, which includes axon growth and the formation of functional synapses. In rodents, neurogenesis, neuronal migration, and cell fate occur during mid-to-late gestation, while axon growth and synapse formation occur primarily postnatally.




Figure 2. Developmental factors regulating development of the melanocortin system. Hypothalamic development involves cell-intrinsic cellular and molecular factors as well as endocrine signals, genetic and environmental factors.