



HAL
open science

Racah Algebras, the Centralizer

$$Z_n(sl_2)$$

and Its Hilbert–Poincaré Series

Nicolas Crampé, Julien Gaboriaud, Loïc Poulain D’andecy, Luc Vinet

► **To cite this version:**

Nicolas Crampé, Julien Gaboriaud, Loïc Poulain D’andecy, Luc Vinet. Racah Algebras, the Centralizer

$$Z_n(sl_2)$$

and Its Hilbert–Poincaré Series. *Annales Henri Poincaré*, 2022, 23 (7), pp.2657-2682. 10.1007/s00023-021-01152-y . hal-03808212

HAL Id: hal-03808212

<https://hal.science/hal-03808212>

Submitted on 10 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Racah algebras, the centralizer $Z_n(\mathfrak{sl}_2)$ and its Hilbert–Poincaré series

Nicolas Crampé¹, Julien Gaboriaud², Loïc Poulain d’Andecy³, and Luc Vinet⁴

¹Institut Denis-Poisson CNRS/UMR 7013 - Université de Tours - Université d’Orléans,
Parc de Grandmont, 37200 Tours, France.

^{2,4}Centre de Recherches Mathématiques, Université de Montréal,
P.O. Box 6128, Centre-ville Station, Montréal (Québec), H3C 3J7, Canada.

³Laboratoire de mathématiques de Reims UMR 9008, Université de Reims Champagne-Ardenne,
Moulin de la Housse BP 1039, 51100 Reims, France.

E-mail addresses: ¹crampe1977@gmail.com, ²julien.gaboriaud@umontreal.ca,
³loic.poulain-dandecy@univ-reims.fr, ⁴vinet@crm.umontreal.ca

May 5, 2021

Abstract

The higher rank Racah algebra $R(n)$ introduced in [1] is recalled. A quotient of this algebra by central elements, which we call the special Racah algebra $sR(n)$, is then introduced. Using results from classical invariant theory, this $sR(n)$ algebra is shown to be isomorphic to the centralizer $Z_n(\mathfrak{sl}_2)$ of the diagonal embedding of $U(\mathfrak{sl}_2)$ in $U(\mathfrak{sl}_2)^{\otimes n}$. This leads to a first and novel presentation of the centralizer $Z_n(\mathfrak{sl}_2)$ in terms of generators and defining relations. An explicit formula of its Hilbert–Poincaré series is also obtained and studied. The extension of the results to the study of the special Askey–Wilson algebra and its higher rank generalizations is discussed.

Keywords: Racah algebra, centralizer, $U(\mathfrak{sl}_2)$, classical invariant theory, first and second fundamental theorems, Hilbert–Poincaré series.

1 Introduction

This paper clarifies the connection between the higher rank Racah algebra $R(n)$ and the centralizer $Z_n(\mathfrak{sl}_2)$ of the diagonal embedding of $U(\mathfrak{sl}_2)$ in its n -fold tensor product. The central result of the paper is:

A particular quotient of the Racah algebra $R(n)$, which we provide in (4.1), is the centralizer.

In the case $n = 3$, the centralizer $Z_3(\mathfrak{sl}_2)$ contains a number of subalgebras of interest for mathematics and physics. In representations, it is generated by the so-called “intermediate” or “quadratic” Casimir elements [2]. These elements realize the relations of the Racah algebra [3]. One may then wonder if the Racah algebra is the centralizer $Z_3(\mathfrak{sl}_2)$ or if more relations are needed in order to offer a full description of the centralizer by generators and relations.

A similar story repeats itself for the centralizer $Z_n(\mathfrak{sl}_2)$. One observes that the intermediate Casimir elements in the n -fold tensor product of $U(\mathfrak{sl}_2)$ realize the relations of the higher rank Racah algebra [1], but two questions remain: Is the centralizer generated by these intermediate Casimir elements, and are there additional relations needed in order to fully describe the centralizer?

As will be seen, these questions are quite close to questions that appear in classical invariant theory. The issue of finding a generating set (of polynomial functions, for example) is answered by the First Fundamental Theorem and the problem of obtaining defining relations between these generators is answered by the Second Fundamental Theorem.

In this regard, the results that we obtain in Corollary 5.5 and Theorem 5.6 can be seen as non-commutative analogues of the First and Second Fundamental Theorems.

1.1 On the Racah algebra

At this point it would be appropriate to provide some background on the Racah algebra, which arises in numerous areas of mathematics and physics.

A first approach to the Racah algebra is from the theory of orthogonal polynomials. The Racah polynomials are a family of bispectral classical orthogonal polynomials which are characterized by a difference and recurrence operator [4]. These operators obey the quadratic algebraic relations of the Racah algebra; this is actually how the algebra was originally introduced [5] and the reason why it inherited its name. Thus, the representation theory of the Racah algebra involves the eponym polynomials.

The problem of decomposing the tensor product of two irreducible representations of \mathfrak{sl}_2 in a direct sum of irreducible representations is known as the Clebsch–Gordan problem of \mathfrak{sl}_2 . When the three-fold tensor product is considered, there exist two natural decompositions. The question of finding the overlaps between the two associated bases is called the Racah problem of \mathfrak{sl}_2 . The intermediate Casimir elements labelling the two mentioned decompositions realize the Racah algebra [3], and thus the overlaps are found to be given in terms of Racah polynomials. There are also realizations of the Racah algebra in terms of $U(\mathfrak{sl}_2)$ [6–9].

Other instances of the Racah algebra appearing in various contexts include algebraic combinatorics [10, 11], the theory of double affine Hecke algebras [12], and as a symmetry algebra of physical models [13–17]. Its finite-dimensional irreducible modules have also been classified [18].

As is the case for various rich structures that have a lot of applications, the generalization of the Racah algebra is something desirable. A higher rank Racah algebra has been defined in [1] by looking at a 3D superintegrable system that realizes the Racah algebra as its symmetry algebra and then generalizing this system to n dimensions. The algebraic relations between the constants of motion were then computed and used to define abstractly the higher rank Racah algebra $R(n)$. These relations of $R(n)$ are also verified by the intermediate Casimir elements that label various direct sum decompositions of the n -fold tensor product of \mathfrak{sl}_2 irreducible representations [19]. These higher rank Racah algebras have also been observed in physical models [20–22], interpreted in the framework of Howe duality [23], and the study of their relation to multivariate Racah polynomials has been initiated [24–26].

1.2 Outline

The paper is organized as follows. In section 2, the abstract $R(3)$ Racah algebra will be introduced and its quotient by a certain central element will be presented and named the

special Racah algebra $sR(3)$. Then, the higher rank Racah algebra $R(n)$ will be presented in Section 3 and various properties will be highlighted. Section 4 will define the higher rank special Racah algebra $sR(n)$. This algebra is a quotient of $R(n)$ by a number of central elements which will be given precisely. We then come to the main results of the paper. It will be proven in Section 5 that the special Racah algebra is isomorphic to the diagonal centralizer $Z_n(\mathfrak{sl}_2)$ of $U(\mathfrak{sl}_2)$ in its n -fold tensor product. The Hilbert–Poincaré series of the centralizer will be obtained in Section 6 and its rich combinatorial properties will be examined. The PBW basis of the centralizer will then be given for the first few values of n . A conclusion offering some comments about the q -deformation of these results and the connection with the multivariate Racah polynomials will end the paper.

2 The Racah algebra (of rank 1) and a special quotient

We review the definition of the usual Racah algebra (of rank 1 in our terminology), along with the algebraic properties we need for the following. We then give the definition of the special Racah algebra of rank 1, to prepare for the generalization for any rank defined later in the paper.

Definition 2.1. *The Racah algebra $R(3)$ of rank 1 is the associative algebra with generators $P_{11}, P_{12}, P_{13}, P_{22}, P_{23}, P_{33}, F_{123}$, and with the following defining relations for indices i, j, k in $\{1, 2, 3\}$ and all distinct:*

$$P_{ii} \text{ is central,} \tag{2.1a}$$

$$[P_{ij}, P_{jk}] = 2F_{ijk}, \tag{2.1b}$$

$$[P_{jk}, F_{ijk}] = P_{ik}(P_{jk} + P_{jj}) - (P_{jk} + P_{kk})P_{ij}. \tag{2.1c}$$

where $[A, B] = AB - BA$ is the commutator, and P_{ij} and F_{ijk} are defined for any $i, j, k \in \{1, 2, 3\}$ by the requirements:

$$P_{ij} = P_{ji} \quad \text{and} \quad F_{ijk} = -F_{jik} = F_{jki} \quad \text{for any } i, j, k \in \{1, 2, 3\}. \tag{2.2}$$

We say that F_{ijk} is antisymmetric whereas P_{ij} is symmetric. Note that in view of (2.1b), the element F_{123} can be removed from the set of generators of $R(n)$ but it is more convenient to keep it.

Relation (2.1b) is given for any i, j, k . If i, j, k are not ordered, the symmetry properties of P and F are to be used. For example,

$$[P_{23}, P_{13}] = [P_{23}, P_{31}] = 2F_{231} = 2F_{123}, \tag{2.3}$$

where we used the symmetry of P , the antisymmetry of F and relation (2.1b) for $(i, j, k) = (2, 3, 1)$. Similar comments apply to relation (2.1c), which is given for any distinct i, j, k and not only the case $(i, j, k) = (1, 2, 3)$.

Let us introduce the notion of determinant for matrices with non-commuting entries. If A is a $n \times n$ matrix with entries $A_{i,j}$ ($1 \leq i, j \leq n$), we define the symmetrized determinant of A as follows

$$\det A := \frac{1}{n!} \sum_{\rho, \sigma \in S_n} \text{sgn}(\rho) \text{sgn}(\sigma) A_{\rho(1), \sigma(1)} A_{\rho(2), \sigma(2)} \cdots A_{\rho(n), \sigma(n)}, \tag{2.4}$$

where S_n is the permutation group of n elements and $\text{sgn}(\sigma)$ is the signature of σ . For commuting entries, it is the usual definition of the determinant of a matrix. We define also the following 3×3 matrix

$$P_{ijk}^{abc} = \begin{pmatrix} P_{ia} & P_{ib} & P_{ic} \\ P_{ja} & P_{jb} & P_{jc} \\ P_{ka} & P_{kb} & P_{kc} \end{pmatrix}. \quad (2.5)$$

There exist in addition to P_{ii} , other central elements in $R(3)$ given in the following proposition:

Proposition 2.2. *The following elements are central in $R(3)$*

$$Q_3 = P_{12} + P_{13} + P_{23}, \quad (2.6)$$

$$w_{ijk} := F_{ijk}^2 + \frac{1}{2} \det(P_{ijk}^{ijk}) - \frac{1}{3} (\{P_{ij}, P_{ik}\} + \{P_{ij}, P_{jk}\} + \{P_{ik}, P_{jk}\} + P_{ij}P_{kk} + P_{ik}P_{jj} + P_{jk}P_{ii}), \quad (2.7)$$

for $1 \leq i, j, k \leq 3$ distinct and where the anticommutator is defined as $\{A, B\} = AB + BA$. Moreover it is observed that w_{ijk} is symmetric i.e.

$$w_{ijk} = w_{jik} = w_{kji}. \quad (2.8)$$

Proof. To show that an element is central, it is enough to show that it commutes with P_{12}, P_{13}, P_{23} (since the P_{ii} 's are central and $2F_{123} = [P_{12}, P_{23}]$). For Q_3 , this is an easy verification.

The symmetry of w_{ijk} is immediate. So it remains to show the centrality of w_{123} . By symmetry of the algebra under renaming of the indices, only the commutation with one element, say P_{23} , needs to be checked. This is a direct calculation using the defining relations of the algebra. \square

Remark 2.3. *It should be stressed that the element w_{123} is essentially the known Casimir element of the Racah algebra [5].*

The special Racah algebra. The special Racah algebra $sR(3)$ is defined from the Racah algebra $R(3)$ by fixing the value of this non-trivial central element.

Definition 2.4. *The special Racah algebra $sR(3)$ of rank 1 is the quotient of $R(3)$ by*

$$w_{123} = 0. \quad (2.9)$$

Remark 2.5. *This relation (2.9) is akin to the relation that expresses the Casimir element of the Racah algebra in terms of its central elements, see equation (3.4) in [3] for instance.*

Remark 2.6. *It follows from (2.8) that w_{213} and other w_{ijk} obtained from permutations of the indices are null in $sR(3)$.*

The appellation ‘‘special’’ is used in the same way as in [27], where a quotient of the Askey–Wilson algebra by fixing the value of a central element expressed as a determinant was denoted as the ‘‘special Askey–Wilson algebra’’ (this was inspired by the nomenclature of Lie groups).

3 The higher rank Racah algebras

Following [1], we consider the following definition of the Racah algebra (of any rank). Note that we consider it as an abstract algebra defined by generators and relations. It will be clear that for $n = 3$ we recover Definition 2.1 for the rank one Racah algebra $R(3)$.

Definition 3.1. *The Racah algebra $R(n)$ of rank $n-2$ is the associative algebra with generators:*

$$P_{ij}, 1 \leq i \leq j \leq n \quad \text{and} \quad F_{ijk}, 1 \leq i < j < k \leq n, \quad (3.1)$$

and the defining relations are, for all possible indices i, j, k, l, m in $\{1, \dots, n\}$:

$$P_{ii} \text{ is central}, \quad (3.2a)$$

$$[P_{ij}, P_{kl}] = 0 \quad \text{if both } i, j \text{ are distinct from } k, l, \quad (3.2b)$$

$$[P_{ij}, P_{jk}] = 2F_{ijk}, \quad (3.2c)$$

$$[P_{jk}, F_{ijk}] = P_{ik}(P_{jk} + P_{jj}) - (P_{jk} + P_{kk})P_{ij}, \quad (3.2d)$$

$$[P_{kl}, F_{ijk}] = P_{ik}P_{jl} - P_{il}P_{jk}, \quad (3.2e)$$

$$[F_{ijk}, F_{jkl}] = -(F_{ijl} + F_{ikl})P_{jk}, \quad (3.2f)$$

$$[F_{ijk}, F_{klm}] = F_{ilm}P_{jk} - F_{jlm}P_{ik}, \quad (3.2g)$$

where in each relation all indices involved are distinct and P_{ij} and F_{ijk} are defined by:

$$P_{ij} = P_{ji} \quad \text{and} \quad F_{ijk} = -F_{jik} = F_{jki} \quad \text{for any } i, j, k \in \{1, \dots, n\}. \quad (3.3)$$

We say that F_{ijk} (or simply F) is antisymmetric whereas P_{ij} is symmetric. The same comments as for the algebra $R(3)$ after Definition 2.1 apply here. In particular, in view of (3.2c), the elements F_{ijk} can be removed from the set of generators of $R(n)$ but it is more convenient to keep them. Note that to check that a certain element X is central in $R(n)$, it is enough to check that it commutes with all generators P_{ij} .

Remark 3.2. *The form of the defining relations above is very symmetrical, and this is quite useful in practice. Namely, for any permutation π of $\{1, \dots, n\}$, the corresponding renaming of the generators $(P_{ij}, F_{ijk}) \mapsto (P_{\pi(i), \pi(j)}, F_{\pi(i), \pi(j), \pi(k)})$ is an automorphism of the algebra. So when checking a relation in the algebra $R(n)$, it is enough to do it for a chosen set of indices. This property will be used in the proofs.*

Recall that P_{ii} is central in $R(n)$. It is easy to show using (3.2b)-(3.2c) that the following element is central in $R(n)$:

$$Q_n = \sum_{1 \leq i < j \leq n} P_{ij}. \quad (3.4)$$

We now introduce some elements of $R(n)$ that will later play an important part. Recall the definition of $\det(P_{ijk}^{abc})$ (2.5) formulated in the preceding section. We define the following elements in $R(n)$:

$$w_{ijk} := F_{ijk}^2 + \frac{1}{2} \det(P_{ijk}^{ijk}) - \frac{1}{3} (\{P_{ij}, P_{ik}\} + \{P_{ij}, P_{jk}\} + \{P_{ik}, P_{jk}\} + P_{ij}P_{kk} + P_{ik}P_{jj} + P_{jk}P_{ii}), \quad (3.5)$$

$$x_{ijkl} := F_{ijk}F_{jkl} + \frac{1}{2} \det(P_{ijk}^{jkl}) + \frac{1}{2}(F_{ijl} + F_{ikl})P_{jk} - \frac{1}{3}(P_{ij}P_{kl} + P_{ik}P_{jl} + P_{il}P_{jk}), \quad (3.6)$$

$$y_{ijklm} := F_{ijk}F_{klm} + \frac{1}{2} \det(P_{ijk}^{klm}) + \frac{1}{2}(F_{ijl}P_{km} - F_{ijm}P_{kl}), \quad (3.7)$$

$$z_{ijklmp} := F_{ijk}F_{lmp} + \frac{1}{2} \det(P_{ijk}^{lmp}), \quad (3.8)$$

where indices $i, j, k, \ell, m, p \in \{1, \dots, n\}$ are all distinct. Only the element w_{ijk} appears in the Racah algebra $R(3)$ since there are not enough different indices for the other elements.

3.1 The Racah algebra $R(4)$

The algebra $R(3)$ was previously studied in Section 2 so let us consider now the case $n = 4$. According to the definition given above, there are 10 generators P_{ij} of $R(4)$ and 4 generators F_{ijk} . Of the relations (3.2b)–(3.2f) only those involving no more than 4 different indices are necessary here.

We already know that the elements P_{ii} and Q_4 given in (3.4) are central in $R(4)$. The following proposition gives less immediate consequences of the defining relations of $R(4)$, and in particular identifies the elements introduced in (3.5)–(3.6) as central elements.

Proposition 3.3. *The following assertions are true in $R(4)$:*

- For $1 \leq a \leq 4$, the following relations hold:

$$P_{a1}F_{234} - P_{a2}F_{134} + P_{a3}F_{124} - P_{a4}F_{123} = 0. \quad (3.9)$$

- For distinct $i, j, k \in \{1, 2, 3, 4\}$, the elements w_{ijk} are symmetric ($w_{ijk} = w_{jik} = w_{jki}$) and are central in $R(4)$.
- For distinct $i, j, k, \ell \in \{1, 2, 3, 4\}$, the elements x_{ijkl} are symmetric ($x_{\sigma(i)\sigma(j)\sigma(k)\sigma(\ell)} = x_{ijkl}$ for $\sigma \in S_4$) and are central in $R(4)$.

Proof. All these statements are proven by invoking the associativity of the algebra. Here is what is meant by that. Suppose that we have a word CBA , for A, B, C some generators, that we want to reorder into the form ABC . This is done by using the defining relations of the algebra (3.2). We decide, for example, to bring all P 's to the left of the F 's, and to order the F 's and the P 's between themselves in the lexicographical ordering of their indices. There are two ways to proceed: one can first start by reordering the pair (CB) into (BC) + some terms, or one could instead start by reordering the pair (BA) into (AB) + some other terms. We denote symbolically the difference at the end of these two computations by

$$(C(BA) - (CB)A), \quad (3.10)$$

and this must be identically 0 by the associativity of the algebra.

Let us first prove (3.9). In the present case, we shall look at the word $CBA = F_{ij\ell}P_{k\ell}P_{i\ell}$, for i, j, k, ℓ all distinct, and compute

$$\frac{1}{2}((F_{ij\ell}P_{k\ell})P_{i\ell} - F_{ij\ell}(P_{k\ell}P_{i\ell})). \quad (3.11)$$

Using relations (3.2), this can be brought to the form

$$P_{ii}F_{jkl} - P_{ij}F_{ikl} + P_{ik}F_{ijl} - P_{i\ell}F_{ijk}. \quad (3.12)$$

By the argument above, this expression has to be zero. Then, choosing $(i, j, k, \ell) = (1, 2, 3, 4)$, $(2, 3, 4, 1)$, $(3, 4, 1, 2)$ or $(4, 1, 2, 3)$, we recover (3.9) with $a = 1, 2, 3, 4$ respectively.

The proof that w_{ijk} is symmetric and that it commutes with all P_{ab} with $a, b \in \{i, j, k\}$ was already done in Section 2 for the algebra $R(3)$, and is still valid here. Using the symmetry of w_{ijk} and the symmetry of the algebra under renaming of the indices, to prove that w_{ijk} is central, it is enough to prove for example that $[P_{34}, w_{123}] = 0$.

This is done by making use of (3.9) and reducing the calculation to

$$[P_{34}, w_{123}] = ((F_{124} + 2F_{134})F_{123})P_{23} - (F_{124} + 2F_{134})(F_{123}P_{23}) \quad (3.13)$$

which is identically zero by the associativity of the algebra.

For the symmetry of x_{ijkl} , the particular case of $x_{ijkl} = x_{ikjl}$ is immediate using the symmetries of P and F . To complete the proof of the symmetry properties of x , it remains to show that $x_{jkli} = x_{ijkl}$. Using the symmetry of the algebra $R(4)$ under renaming of the indices, it is enough to check that for example $x_{2341} = x_{1234}$. Substituting from the definition of x_{ijkl} , one has

$$\begin{aligned} x_{2341} - x_{1234} &= (F_{134} - F_{123})F_{234} + \frac{1}{2}(\det(P_{134}^{234}) - \det(P_{123}^{234})) \\ &\quad - \frac{1}{2}(P_{34}(F_{123} + F_{124}) + P_{23}(F_{134} + F_{124})). \end{aligned} \quad (3.14)$$

Now, looking at $CBA = F_{234}F_{124}P_{34}$ and making use of (3.9), one computes

$$\begin{aligned} \frac{1}{2}(F_{234}(F_{124}P_{34}) - (F_{234}F_{124})P_{34}) &= (F_{134} - F_{123})F_{234} + \frac{1}{2}(\det(P_{134}^{234}) - \det(P_{123}^{234})) \\ &\quad - \frac{1}{2}(P_{34}(F_{123} + F_{124}) + P_{23}(F_{134} + F_{124})). \end{aligned} \quad (3.15)$$

By the associativity of the algebra (see above), this expression has to be zero. This completes the proof of the symmetry of x_{ijkl} since the right hand sides of (3.14) and (3.15) are the same.

Using the symmetry of x_{ijkl} and the symmetry of the algebra under renaming of the indices, the proof that x_{ijkl} is central reduces to proving that for example $[x_{1234}, P_{23}] = 0$ which is also done by a direct computation using expression (3.17) of x_{1234} . \square

Remark 3.4. *The elements w_{ijk} and x_{ijkl} can be equivalently given by the following formulae*

$$w_{123} = F_{123}^2 - F_{123}P_{13} - P_{12}(P_{13} + P_{23} + P_{33}) + \frac{1}{2} \sum_{\sigma \in S_3} \text{sgn}(\sigma) P_{\sigma(1)1} P_{\sigma(2)2} P_{\sigma(3)3}, \quad (3.16)$$

and

$$x_{1234} = F_{123}F_{234} - F_{123}P_{24} + F_{124}P_{23} + F_{134}P_{23} - P_{14}P_{23} + \frac{1}{2} \sum_{\sigma \in S_3} \text{sgn}(\sigma) P_{\sigma(1)2} P_{\sigma(2)3} P_{\sigma(3)4}. \quad (3.17)$$

3.2 The Racah algebra $R(n)$ for any n

Let now n be any positive integer. We already know that we have in $R(n)$ central elements P_{ii} and Q_n given in (3.4). Building upon all that has been proven up to now, we have the following final proposition about the Racah algebra $R(n)$.

Proposition 3.5. *The following assertions are true in $R(n)$:*

- *The relations below hold for $1 \leq a \leq n$ and $1 \leq i < j < k < \ell \leq n$:*

$$P_{ai}F_{jkl} - P_{aj}F_{ikl} + P_{ak}F_{ijl} - P_{al}F_{ijk} = 0. \quad (3.18)$$

- *For distinct $i, j, k \in \{1, \dots, n\}$, the elements w_{ijk} are symmetric and central in $R(n)$.*
- *For distinct $i, j, k, \ell \in \{1, \dots, n\}$, the elements x_{ijkl} are symmetric and central in $R(n)$.*

- For distinct $i, j, k, \ell, m \in \{1, \dots, n\}$, the elements y_{ijklm} are null in $R(n)$:

$$y_{ijklm} = 0 \quad \text{for all distinct } i, j, k, \ell, m. \quad (3.19)$$

- For distinct $i, j, k, \ell, m, p \in \{1, \dots, n\}$, the elements z_{ijklmp} are null in $R(n)$:

$$z_{ijklmp} = 0 \quad \text{for all distinct } i, j, k, \ell, m, p. \quad (3.20)$$

Proof. If a is equal to i, j, k or ℓ , relation (3.18) only involves 4 indices, and so its validity follows directly from the Proposition 3.3 concerning the algebra $R(4)$. To prove the case when a is different of i, j, k and ℓ , first compare (3.2g) for (i, j, k, ℓ, m) equals to $(1, 2, 5, 3, 4)$ and $(3, 4, 5, 1, 2)$. This leads to the identity

$$P_{15}F_{234} - P_{25}F_{134} + P_{35}F_{124} - P_{45}F_{123} = 0. \quad (3.21)$$

The symmetry of the algebra under renaming of the indices suffices to complete the proof of (3.18).

Regarding the statements about w_{ijk} and building on the verifications made in $R(3)$ and $R(4)$, it remains only to check that w_{ijk} commutes with $P_{\ell m}$ when $\ell, m \notin \{i, j, k\}$. This is immediate since from the defining relations, any two elements (P 's or F 's) with no index in common commute.

Concerning x_{ijkl} , building on the proof of the preceding subsection, it remains to show, for example, that

$$[x_{1234}, P_{45}] = 0. \quad (3.22)$$

This is shown using $y_{ijklm} = 0$ (which is proven below) as well as (3.18). For x_{ijkl} , it remains only to check that it commutes with P_{mp} when $m, p \notin \{i, j, k, \ell\}$, and this is immediate.

To prove that y_{ijklm} is zero, we first look at the case $(i, j, k, \ell, m) = (1, 2, 3, 4, 5)$. Write

$$\frac{1}{2}(F_{245}(F_{123}P_{23}) - (F_{245}F_{123})P_{23}). \quad (3.23)$$

It is seen that this is equal to y_{12345} , by making use of (3.18). Invoking associativity, it follows that $y_{12345} = 0$. Since this can be repeated for all other combinations of distinct indices i, j, k, ℓ, m it is done.

For the nullity of z_{ijklmp} , we invoke again the associativity of the algebra. Looking at

$$\frac{1}{2}((F_{356}F_{234})P_{12} - F_{356}(F_{234}P_{12})) \quad (3.24)$$

it is seen that this is equal to z_{123456} . Thus it follows that

$$z_{123456} = 0. \quad (3.25)$$

A similar reasoning can be repeated for different indices to complete the proof. \square

4 The special Racah algebra $sR(n)$

After the preliminary discussion of the Racah algebra $R(n)$, we are now ready to define the special Racah algebra for any n . This is a generalization to arbitrary rank of the special Racah algebra $sR(3)$ from Section 2.

Definition 4.1. *The special Racah algebra $sR(n)$ of rank $n - 2$ is the quotient of $R(n)$ by all*

$$w_{ijk} = 0, \quad x_{abcd} = 0, \quad (4.1)$$

such that $1 \leq i < j < k \leq n$ and $1 \leq a < b < c < d \leq n$.

Since $sR(n)$ is the algebra involved for the study of the centralizer in the next section, we collect here the generators and defining relations, to give an explicit definition without reference to the Racah algebra (we keep the same name for the generators, which is justified since this is a quotient, this should not lead to any ambiguity).

Definition 4.2 (Equivalent definition). *The special Racah algebra $sR(n)$, of rank $n - 2$ is the associative algebra with generators:*

$$P_{ij}, \quad 1 \leq i \leq j \leq n \quad \text{and} \quad F_{ijk}, \quad 1 \leq i < j < k \leq n. \quad (4.2)$$

To give the defining relations, first we define P_{ij} and F_{ijk} by:

$$P_{ij} = P_{ji} \quad \text{and} \quad F_{ijk} = -F_{jik} = F_{jki} \quad \text{for any } i, j, k \in \{1, \dots, n\}. \quad (4.3)$$

The defining relations are, for all possible indices i, j, k, ℓ, m in $\{1, \dots, n\}$:

$$P_{ii} \quad \text{is central}, \quad (4.4a)$$

$$[P_{ij}, P_{k\ell}] = 0 \quad \text{if both } i, j \text{ are distinct from } k, \ell, \quad (4.4b)$$

$$[P_{ij}, P_{jk}] = 2F_{ijk}, \quad (4.4c)$$

$$[P_{jk}, F_{ijk}] = P_{ik}(P_{jk} + P_{jj}) - (P_{jk} + P_{kk})P_{ij}, \quad (4.4d)$$

$$[P_{k\ell}, F_{ijk}] = P_{ik}P_{j\ell} - P_{i\ell}P_{jk}, \quad (4.4e)$$

$$[F_{ijk}, F_{jkl}] = F_{jkl}P_{ij} - F_{ijk}P_{j\ell} - F_{ik\ell}(P_{jk} + P_{jj}), \quad (4.4f)$$

$$[F_{ijk}, F_{klm}] = F_{ilm}P_{jk} - F_{jlm}P_{ik}, \quad (4.4g)$$

along with

$$F_{ijk}^2 + \frac{1}{2} \det(P_{ijk}^{ijk}) = \frac{1}{3} (\{P_{ij}, P_{ik}\} + \{P_{ij}, P_{jk}\} + \{P_{ik}, P_{jk}\} + P_{ij}P_{kk} + P_{ik}P_{jj} + P_{jk}P_{ii}), \quad (4.4h)$$

for $1 \leq i < j < k \leq n$, and

$$F_{ijk}F_{jkl} + \frac{1}{2} \det(P_{ijk}^{jkl}) = -\frac{1}{2}(F_{ij\ell} + F_{ik\ell})P_{jk} + \frac{1}{3}(P_{ij}P_{k\ell} + P_{ik}P_{j\ell} + P_{i\ell}P_{jk}), \quad (4.4i)$$

for $1 \leq i < j < k < \ell \leq n$.

Consequences of the relations. From the results of the preceding sections, we know that relations (4.4h)-(4.4i) for any distinct i, j, k, ℓ are automatically verified, along with the relations:

$$F_{ijk}F_{klm} + \frac{1}{2} \det(P_{ijk}^{klm}) = \frac{1}{2}(F_{ijm}P_{k\ell} - F_{ij\ell}P_{km}) \quad (4.5a)$$

$$F_{ijk}F_{lmr} + \frac{1}{2} \det(P_{ijk}^{lmr}) = 0, \quad (4.5b)$$

$$P_{ai}F_{jkl} - P_{aj}F_{ik\ell} + P_{ak}F_{ij\ell} - P_{al}F_{ijk} = 0. \quad (4.5c)$$

for distinct $i, j, k, \ell, m, r \in \{1, \dots, n\}$ and for any $a \in \{1, \dots, n\}$. These relations, although satisfied, do not have to be included in the set of defining relations.

Example 4.3. *The special Racah algebra $sR(4)$ of rank 2 is the quotient of $R(4)$ by*

$$w_{123} = 0, \quad w_{124} = 0, \quad w_{134} = 0, \quad w_{234} = 0 \quad \text{and} \quad x_{1234} = 0. \quad (4.6)$$

Remark 4.4. *Some analogues of the relations of $sR(4)$ (excluding the ones of the type (4.5c)) were obtained in a particular realization in [20]. For the higher rank case of $sR(n)$, analogous relations were also observed in a certain realization in [22], once again excluding the ones of the type (4.5c).*

5 Isomorphism between the centralizer $Z_n(\mathfrak{sl}_2)$ and the special Racah algebra $sR(n)$

The goal of this section is to connect the (higher rank) special Racah algebra introduced and characterized in the previous two sections with the centralizer $Z_n(\mathfrak{sl}_2)$ of the diagonal action of $U(\mathfrak{sl}_2)$ in $U(\mathfrak{sl}_2)^{\otimes n}$. This will provide an a posteriori justification for the quotient that was chosen to go from the Racah algebra $R(n)$ to the special Racah algebra $sR(n)$: as will be shown, this quotient is precisely the one that leads to an algebra isomorphic to the centralizer $Z_n(\mathfrak{sl}_2)$.

5.1 Centralizer $Z_n(\mathfrak{sl}_2)$ of the diagonal action $U(\mathfrak{sl}_2)$ into $U(\mathfrak{sl}_2)^{\otimes n}$ and the algebra of polarized traces

We here define the centralizer associated to the Lie algebra \mathfrak{sl}_2 . The generators of \mathfrak{sl}_2 are e_{ij} , $i, j \in \{1, 2\}$ obeying the defining relations

$$[e_{ij}, e_{kl}] = \delta_{jk}e_{il} - \delta_{li}e_{kj}, \quad e_{11} + e_{22} = 0. \quad (5.1)$$

We denote by $U(\mathfrak{sl}_2)$ the universal enveloping algebra of \mathfrak{sl}_2 . Its Casimir element is given by

$$C = e_{11}^2 - e_{11} + e_{12}e_{21}. \quad (5.2)$$

Now consider the tensor product of n copies of $U(\mathfrak{sl}_2)$ and define the following notation for its generators

$$e_{ij}^{(a)} = 1^{\otimes(a-1)} \otimes e_{ij} \otimes 1^{\otimes(n-a)}. \quad (5.3)$$

The diagonal embedding of $U(\mathfrak{sl}_2)$ in its n -fold tensor product is given by

$$\begin{aligned} \delta : U(\mathfrak{sl}_2) &\rightarrow U(\mathfrak{sl}_2)^{\otimes n} \\ e_{ij} &\mapsto \sum_{a=1}^n e_{ij}^{(a)}. \end{aligned} \quad (5.4)$$

There is a natural degree-preserving action of \mathfrak{sl}_2 on $U(\mathfrak{sl}_2)^{\otimes n}$ given by composing the diagonal embedding δ followed by the adjoint action. On the generators, it is given by

$$e_{ij} \cdot e_{kl}^{(a)} = \delta_{jk}e_{il}^{(a)} - \delta_{li}e_{kj}^{(a)}. \quad (5.5)$$

We then define the centralizer $Z_n(\mathfrak{sl}_2)$ of the diagonal embedding of $U(\mathfrak{sl}_2)$ in $U(\mathfrak{sl}_2)^{\otimes n}$ as the kernel of this \mathfrak{sl}_2 action

$$Z_n(\mathfrak{sl}_2) = \{X \in U(\mathfrak{sl}_2)^{\otimes n} \mid g \cdot X = [\delta(g), X] = 0 \quad \forall g \in U(\mathfrak{sl}_2)\} \quad (5.6)$$

or in other words, as the set of elements in $U(\mathfrak{sl}_2)^{\otimes n}$ that commute with the diagonal embedding of $U(\mathfrak{sl}_2)$.

Let us also define the polarized traces (the summation convention is assumed):

$$T^{(a_1, \dots, a_d)} = e_{i_2 i_1}^{(a_1)} e_{i_3 i_2}^{(a_2)} \cdots e_{i_1 i_d}^{(a_d)}, \quad a_1, \dots, a_d \in \{1, \dots, n\}. \quad (5.7)$$

It is seen by a direct computation that these elements are in the centralizer $Z_n(\mathfrak{sl}_2)$.

Remark 5.1. *It is easily checked from the definition (5.7) that $T^{(a_1, a_2)} = T^{(a_2, a_1)}$ and that $T^{(a_1, a_2, a_3)}$ is antisymmetric in its indices a_1, a_2, a_3 , i.e. $T^{(a_1, a_2, a_3)} = T^{(a_2, a_3, a_1)} = -T^{(a_2, a_1, a_3)}$.*

Remark 5.2. *A number of papers in the literature [3, 19] realize the Racah algebra with the so-called ‘‘intermediate Casimir’’ elements C_i, C_{ij} . These elements are given by*

$$C_i = 1^{\otimes(i-1)} \otimes C \otimes 1^{\otimes(n-i)}, \quad C_{ij} = 1^{\otimes(i-1)} \otimes C_{(1)} \otimes 1^{\otimes(j-i-1)} \otimes C_{(2)} \otimes 1^{\otimes(n-j)}, \quad (5.8)$$

where $\Delta(e_{ij}) = e_{ij} \otimes 1 + 1 \otimes e_{ij}$ and we denote $\Delta(C) = C_{(1)} \otimes C_{(2)}$ in Sweedler’s notation. Then the $T^{(i,i)}$ and $T^{(i,j)}$ can be expressed in terms of these intermediate Casimir elements as follows

$$T^{(i,i)} = 2C_i, \quad T^{(i,j)} = C_{ij} - C_i - C_j. \quad (5.9)$$

5.2 Elements of classical invariant theory

We now present results from classical invariant theory about the algebra of polynomial functions on

$$\underbrace{\mathfrak{sl}_2 \times \cdots \times \mathfrak{sl}_2}_{n \text{ factors}} \equiv \mathfrak{sl}_2^n, \quad (5.10)$$

that are invariant under simultaneous conjugations by $SL(2)$. For G elements of $SL(2)$, these actions on a polynomial function of \mathfrak{sl}_2^n are given by:

$$G \cdot f(M_1, \dots, M_n) = f(G^{-1}M_1G, \dots, G^{-1}M_nG), \quad (5.11)$$

for $M_i \in \mathfrak{sl}_2$. The first fundamental theorem of classical invariant theory states that:

Theorem 5.3 (see [28–30] or [31] and references therein). *The algebra $\mathbb{C}[\mathfrak{sl}_2^n]^{inv}$ of polynomial functions on \mathfrak{sl}_2^n that are invariant under simultaneous conjugations by $SL(2)$ elements is generated by the functions*

$$\mathfrak{T}^{(a_1, \dots, a_d)} : (M_1, \dots, M_n) \mapsto Tr(M_{a_1} \cdots M_{a_d}) \quad (5.12)$$

for $M_i \in \mathfrak{sl}_2$, $d \geq 2$ and $a_1, \dots, a_d \in \{1, \dots, n\}$. Moreover, it is sufficient to take $\mathfrak{T}^{(i,j)}$ ($i \leq j$) and $\mathfrak{T}^{(i,j,k)}$ ($i < j < k$) to obtain a generating set.

The generating set of the invariant polynomial functions described in the preceding theorem (the ones of degrees 2 and 3) is not algebraically independent. A set of generators for their ideal of relations is given in the next theorem (second fundamental theorem on these invariants).

Theorem 5.4 (see [32], Theorem 2.3 (ii) or [31], Theorem 3.4 (ii)). *The defining relations for the algebra of polynomial invariant functions are*

$$\begin{aligned} & Tr([M_i, M_j]M_k)Tr([M_p, M_q]M_r) \\ & + 2 \begin{vmatrix} Tr(M_iM_p) & Tr(M_iM_q) & Tr(M_iM_r) \\ Tr(M_jM_p) & Tr(M_jM_q) & Tr(M_jM_r) \\ Tr(M_kM_p) & Tr(M_kM_q) & Tr(M_kM_r) \end{vmatrix} = 0, \end{aligned} \quad (5.13a)$$

$$\begin{aligned} & Tr([M_j, M_k]M_\ell)Tr(M_pM_i) - Tr([M_i, M_k]M_\ell)Tr(M_pM_j) \\ & + Tr([M_i, M_j]M_\ell)Tr(M_pM_k) - Tr([M_i, M_j]M_k)Tr(M_pM_\ell) = 0, \end{aligned} \quad (5.13b)$$

with $i, j, k, \ell, m, n, p, q, r \in \{1, \dots, n\}$.

With the following reasoning that is adapted from [33], we now use Theorem 5.3 to extract information about the algebra of polarized traces and the centralizer.

- The algebra $U(\mathfrak{sl}_2)^{\otimes n}$ is filtered. Take the degree of all generators $e_{ij}^{(a)}$ to be 1, then the associated graded algebra is commutative. Recall the \mathfrak{sl}_2 action (5.5). This induces a natural action on $gr(U(\mathfrak{sl}_2)^{\otimes n})$. Denote the generators of the graded algebra by $e_{ij}^{(a)}$. The induced \mathfrak{sl}_2 -action is given as follows on the generators:

$$e_{ij} \cdot e_{kl}^{(a)} = \delta_{jk}e_{il}^{(a)} - \delta_{li}e_{kj}^{(a)}. \quad (5.14)$$

- The algebra of polynomial functions on \mathfrak{sl}_2^n is the algebra of polynomials $\mathbb{C}[x_{ij}^{(a)}]$, where $x_{ij}^{(a)}$ is the linear form giving the (i, j) coordinate of the a^{th} matrix in the product \mathfrak{sl}_2^n . The simultaneous conjugation action of an element G of $SL(2)$ on a polynomial function of \mathfrak{sl}_2^n (5.11) is given infinitesimally by

$$\epsilon \sum_{k=1}^n f(M_1, \dots, [M_k, g], \dots, M_n) \quad (5.15)$$

for $G = e^{i\epsilon g}$ with $g \in \mathfrak{sl}_2$. Thus, on the generators of polynomials functions $x_{ij}^{(a)}$, this infinitesimal action is

$$e_{ij} \cdot x_{kl}^{(a)} = \delta_{jl}x_{ki}^{(a)} - \delta_{ik}x_{j\ell}^{(a)} \quad (5.16)$$

and can be identified with the induced \mathfrak{sl}_2 action on $gr(U(\mathfrak{sl}_2)^{\otimes n})$ through

$$e_{ij}^{(a)} \leftrightarrow x_{ji}^{(a)}. \quad (5.17)$$

- It follows that under this identification, the invariant functions correspond to the elements of $gr(U(\mathfrak{sl}_2)^{\otimes n})$ in the kernel of the \mathfrak{sl}_2 action, or in other words to the image of the centralizer in $gr(U(\mathfrak{sl}_2)^{\otimes n})$. Moreover, again under this identification, the image in the graded algebra of the polarized trace $T^{(a_1, \dots, a_d)}$ defined in (5.12) is the polynomial function $Tr(M_{a_1} \dots M_{a_d})$.

5.3 The defining relations of $Z_n(\mathfrak{sl}_2)$

Knowing that the generators of the invariant functions (5.12) correspond to the polarized traces (5.7) (see Theorem 5.3), the image of the centralizer in $gr(U(\mathfrak{sl}_2)^{\otimes n})$ is therefore generated by the image of the polarized traces. Now consider an element of degree N in the centralizer. Up to terms of degree $N - 1$, this element can be expressed as a polynomial in $T^{(a_1, \dots, a_d)}$. The same can then be argued for each of the remaining lower degree terms by induction, and thus any element in the centralizer can be expressed as a polynomial in the polarized traces. Since all polarized traces belong in the centralizer, the two algebras thus coincide. So we obtain the analogue of the first fundamental theorem:

Corollary 5.5. *It follows from Theorem 5.3 that the polarized traces $T^{(i,j)}$, $i \leq j$ and $T^{(i,j,k)}$, $i < j < k$ generate the centralizer $Z_n(\mathfrak{sl}_2)$ of the diagonal action of $U(\mathfrak{sl}_2)$ in $U(\mathfrak{sl}_2)^{\otimes n}$.*

Recall that relations (5.13) are a set of defining relations for the image of the centralizer in $gr(U(\mathfrak{sl}_2)^{\otimes n})$. We look for analogous defining relations for the centralizer in $U(\mathfrak{sl}_2)^{\otimes n}$. Once we find the deformations in $U(\mathfrak{sl}_2)^{\otimes n}$ of relations (5.13), we can prove their completeness using for the ideal of relations the same sort of reasoning as before Corollary 5.5. In fact the special Racah algebra was defined such that its set of defining relations gives precisely the complete set of relations for the centralizer.

Theorem 5.6. *With the following identification of the generators:*

$$P_{ab} \mapsto T^{(a,b)} \quad \text{and} \quad F_{ijk} \mapsto -T^{(i,j,k)} \quad (5.18)$$

for i, j, k all distinct, the centralizer $Z_n(\mathfrak{sl}_2)$ is isomorphic to the special Racah algebra $sR(n)$:

$$Z_n(\mathfrak{sl}_2) \cong sR(n). \quad (5.19)$$

In other words, a set of defining relations of $Z_n(\mathfrak{sl}_2)$ is given in (4.4), where P_{ab} and F_{ijk} are replaced by the corresponding polarized traces.

Proof. The defining relations are verified to hold in $U(\mathfrak{sl}_2)^{\otimes n}$ by direct computations. Note that, due to the symmetry under renaming the indices, we only need to make calculations in $U(\mathfrak{sl}_2)^{\otimes n}$ for $n \leq 5$ in degrees less or equal to 6 in the generators.

Under the previous choice of degree for the generators of $U(\mathfrak{sl}_2)^{\otimes n}$ which was $deg(e_{ij}^{(a)}) = 1$, it follows that

$$deg(T^{(i,j)}) = 2, \quad deg(T^{(i,j,k)}) = 3. \quad (5.20)$$

The same degrees are given to the generators of the special Racah algebra $sR(n)$: $deg(P_{ij}) = 2$ and $deg(F_{ijk}) = 3$. This makes it a filtered algebra, and it is straightforward to observe that its associated graded algebra is isomorphic to the algebra of polynomial invariants functions. Indeed, the first set of defining relations (3.2), or equivalently (4.4a)–(4.4g), ensures that the generators all commute in the graded algebra, and then the relations (4.4h)–(4.4i) and (4.5) are mapped to (5.13).

Therefore both algebras related by the morphism in (5.18) have the same associated graded algebras, and so in particular have the same dimensions in each component of the filtration (that is, in each degree). Moreover the morphism is surjective from Corollary 5.5. Consequently, an element in the kernel of the map (in other words, a relation in $Z_n(\mathfrak{sl}_2)$ not implied by the relations of the special Racah algebra), if non-zero, would contradict the equality of dimensions for some degree. \square

Remark 5.7. *It is quite remarkable that we only need to quotient the Racah algebra $R(n)$ by the elements w_{ijk} and x_{ijkl} in order to recover the centralizer for any value of n . Indeed, one could have expected that in order to recover the centralizer for increasing n , we would need to quotient by elements of increasing degree or spanning an increasing number of indices. That this is not the case is quite a surprising simplification.*

6 The Hilbert–Poincaré series of $Z_n(\mathfrak{sl}_2)$

For more information on Hilbert–Poincaré series of graded algebras, we refer to [34]. The Hilbert–Poincaré series contains useful information about a graded, or filtered, algebra. We will illustrate this for the diagonal centralizer $Z_n(\mathfrak{sl}_2)$. We will provide an explicit formula for its Hilbert–Poincaré series, and then use it in conjunction with the defining relations found in Theorem 5.6 to provide bases of $Z_n(\mathfrak{sl}_2)$ for small n .

6.1 An explicit formula

The commutative algebra $\mathbb{C}[\mathfrak{sl}_2^n]^{inv}$ of polynomial functions on \mathfrak{sl}_2^n that are invariant under simultaneous conjugation by $SL(2)$ is a graded algebra: it is the direct sum of the subspaces $\mathbb{C}_k[\mathfrak{sl}_2^n]^{inv}$ of homogeneous invariant polynomial functions of degree k . The Hilbert–Poincaré series records the dimensions of all these subspaces:

$$F_n(t) = \sum_{k \geq 0} \dim(\mathbb{C}_k[\mathfrak{sl}_2^n]^{inv}) t^k. \quad (6.1)$$

The centralizer $Z_n(\mathfrak{sl}_2)$ inherits from $U(\mathfrak{sl}_2)^{\otimes n}$ the structure of a filtered algebra: it is the union of the increasing sequence (in k) of subspaces $Z_n(\mathfrak{sl}_2)_{\leq k}$ of elements of degree less or equal to k (the degree is in the generators of $U(\mathfrak{sl}_2)^{\otimes n}$). The Hilbert–Poincaré series of $Z_n(\mathfrak{sl}_2)$ records the dimensions of the homogeneous subspaces of the associated graded algebra:

$$F_n(t) = \sum_{k \geq 0} \dim(Z_n(\mathfrak{sl}_2)_{\leq k} / Z_n(\mathfrak{sl}_2)_{< k}) t^k, \quad (6.2)$$

and thus, from the discussion in Section 5, is the same as the Hilbert–Poincaré series of the invariant polynomial functions.

Several formulas, using various approaches, have been obtained for the Hilbert–Poincaré series $F_n(t)$ (see references in [31, 35, 36]). The formula presented below seems to be new.

Proposition 6.1. *Let $n \geq 2$ and recall that the rank r is defined by $r = n - 2$. The Hilbert–Poincaré series of $Z_n(\mathfrak{sl}_2)$ is:*

$$F_n(t) = \frac{P_r(t)}{(1 - t^2)^{3(r+1)}}, \quad (6.3)$$

where the numerator is given by:

$$P_r(t) = (1 + t)^r \sum_{k=0}^{2r} (-1)^k a_k t^k, \quad \text{where } \begin{cases} a_{2k} = \binom{r}{k}^2, \\ a_{2k+1} = \binom{r}{k} \binom{r}{k+1}. \end{cases} \quad (6.4)$$

Proof. We take a detour through the graded character of $SL(2)$ on the polynomial functions on \mathfrak{sl}_2^n . The character of $SL(2)$ for a finite-dimensional representation is seen as a Laurent

polynomial in x , given by the trace of the action of the element $\text{Diag}(x, x^{-1})$. For example, for the fundamental representation of $SL(2)$, it is $x + x^{-1}$. For the irreducible representation of dimension $d + 1$, which is the d -symmetrized power of the fundamental representation, the character is thus $\frac{x^{d+1} - x^{-d-1}}{x - x^{-1}}$. Now, it is easy to check that, if we have the character $\chi(x)$ of an arbitrary finite-dimensional representation of $SL(2)$, then the formula:

$$[(1 - x^2)\chi(x)]_0, \quad (6.5)$$

where $[\cdot]_0$ means taking the constant term of a Laurent polynomial, gives the multiplicity of the trivial representation.

After these classical preliminaries, note that the character of the adjoint representation of $SL(2)$ on \mathfrak{sl}_2 is $1 + x^2 + x^{-2}$. On the polynomial function on \mathfrak{sl}_2 , the action of $SL(2)$ preserves the grading, and we record the character of the representation on its graded components as a formal power series in t (also called, the graded character). For each degree, the representation is a symmetrized power of the adjoint representation, so we find that the graded character is:

$$\frac{1}{(1-t)(1-tx^2)(1-tx^{-2})}. \quad (6.6)$$

Equivalently, this is the graded character for the adjoint action on $U(\mathfrak{sl}_2)$. On the polynomial functions \mathfrak{sl}_2^n (or equivalently, on $U(\mathfrak{sl}_2)^{\otimes n}$), the graded character is thus:

$$\frac{1}{(1-t)^n(1-tx^2)^n(1-tx^{-2})^n}. \quad (6.7)$$

Now, in each degree, we look for the dimension of the invariant subspace for the action of $SL(2)$. In other words, we look for the multiplicity of the trivial representation. By what we have recalled above, we obtain that the Hilbert–Poincaré series of $Z_n(\mathfrak{sl}_2)$ is:

$$F_n(t) = \left[\frac{(1-x^2)}{(1-t)^n(1-tx^2)^n(1-tx^{-2})^n} \right]_0. \quad (6.8)$$

Using the expansion $(1-z)^{-n} = \sum_{k \geq 0} \binom{k+n-1}{k} z^k$ and straightforward manipulations, we obtain:

$$F_n(t) = \frac{1}{(1-t)^n} \sum_{k \geq 0} (-1)^k \tilde{a}_k t^k, \quad \text{where } \begin{cases} \tilde{a}_{2k} = \binom{n+k-1}{k}^2, \\ \tilde{a}_{2k+1} = \binom{n+k-1}{k} \binom{n+k}{k+1}. \end{cases} \quad (6.9)$$

Thus the statement of the proposition reduces to the following equality of formal power series:

$$\frac{1}{(1-t)^{r+2}} \sum_{k \geq 0} (-1)^k \tilde{a}_k t^k = \frac{(1+t)^r}{(1-t^2)^{3(r+1)}} \sum_{k \geq 0} (-1)^k a_k t^k. \quad (6.10)$$

To prove this, we multiply the \tilde{a} -series by $(1+t)$ and the a -series by $(1-t)$, and after an application of Pascal rule for binomials, we reach the equivalent formula:

$$\sum_{k \geq 0} (-1)^k \tilde{a}'_k t^k = \frac{1}{(1-t^2)^{2(r+1)}} \sum_{k \geq 0} (-1)^k a'_k t^k, \quad (6.11)$$

where now we have $\begin{cases} \tilde{a}'_{2k} = \binom{r+k+1}{k} \binom{r+k}{k}, \\ \tilde{a}'_{2k+1} = \binom{r+k+1}{k} \binom{r+k+1}{k+1}, \end{cases}$ and $\begin{cases} a'_{2k} = \binom{r}{k} \binom{r+1}{k}, \\ a'_{2k+1} = \binom{r}{k} \binom{r+1}{k+1}. \end{cases}$

This last formula is verified by writing the expansion of the right hand side and checking the equality of the coefficients, making use of the following identity for binomial coefficients [37]:

$$\sum_i \binom{i+a+b}{i} \binom{b}{k-i} \binom{a}{k'-i} = \binom{k'+b}{k} \binom{k+a}{k'}. \quad (6.12)$$

This identity is valid for any a, b, k, k' and we use it for $a = r + 1, b = r$ and $k' \in \{k, k + 1\}$. \square

The exponent $3(r + 1)$ appearing in the denominator of $F_n(t)$ is the Krull, or Gelfand–Kirillov, dimension of the algebra of invariant polynomial functions (see [38],[36]). Here it means that there is a set of $3(r + 1)$ algebraically independent homogeneous elements (a system of parameters) $\theta_1, \dots, \theta_{3(r+1)}$ such that the algebra is a free module of finite dimension over the polynomial subalgebra $\mathbb{C}[\theta_1, \dots, \theta_{3(r+1)}]$. The freeness follows from the property called Cohen–Macaulay, which is ensured here by the Hochster–Roberts theorem from general invariant theory [39]. The form $F_n(t)$ above with the positivity of the numerator $P_r(t)$ (see below) suggests that it might be possible that a system of parameters consists of $3(r + 1)$ elements of degrees 2. If it were to be the case, then $P_r(1)$ would be the dimension of the algebra over $\mathbb{C}[\theta_1, \dots, \theta_{3(r+1)}]$. Moreover, the different monomials in $P_r(t)$ would indicate in which degrees the elements of a basis over $\mathbb{C}[\theta_1, \dots, \theta_{3(r+1)}]$ would have to be found.

Finally, the palindromic property of the numerator $P_r(t)$ in the formula above shows that the Hilbert–Poincaré series satisfies the functional equation:

$$F_n(t^{-1}) = (-1)^{(n-1)} t^{3n} F_n(t). \quad (6.13)$$

This is well-known and related to a property, called being Gorenstein, for the algebra of invariant polynomial functions (see [31] and references therein).

Remark 6.2. *More generally, the Hilbert–Poincaré series of $Z_n(\mathfrak{sl}_2)$ can be defined as a power series in t_1, \dots, t_n if we consider the gradation by the multidegree of $U(\mathfrak{sl}_2)^{\otimes n}$. A slight generalization of the part of the proof up to Formula (6.9) gives the multigraded version of this formula:*

$$F_n(t_1, \dots, t_n) = \frac{1}{(1-t_1)\dots(1-t_n)} \left(\sum_{\substack{\mu \models k \\ \nu \models k}} t_1^{\mu_1+\nu_1} \dots t_n^{\mu_n+\nu_n} - \sum_{\substack{\mu \models k \\ \nu \models k-1}} t_1^{\mu_1+\nu_1} \dots t_n^{\mu_n+\nu_n} \right), \quad (6.14)$$

where $\mu \models k$ means that $\mu = (\mu_1, \dots, \mu_n) \in \mathbb{Z}_{\geq 0}^n$ such that $\mu_1 + \dots + \mu_n = k$. To recover relation (6.9) from this, take $t_1 = \dots = t_n = t$ and use that the number of $\mu \models k$ is $\binom{k+n-1}{k}$.

6.2 Some related combinatorics

We have obtained an expression for the Hilbert–Poincaré series of $Z_n(\mathfrak{sl}_2)$ of the form:

$$F_n(t) = \frac{(1+t)^r Q_r(t)}{(1-t^2)^{3(r+1)}}, \quad \text{where } Q_r(t) = \sum_{k=0}^{2r} (-1)^k a_k t^k, \quad (6.15)$$

and the coefficients a_k are given in the proposition above. It is perhaps not so surprising that the coefficients of the various polynomials involved show some connections with well-studied combinatorial objects of ‘‘Catalan’’ flavor.

The polynomial $Q_r(t)$. The coefficient a_k in the polynomial $Q_r(t)$ counts the number of symmetric Dyck paths of semi-length $2r + 1$ with $k + 1$ peaks (see A088855 in [40]). Their expression with binomial coefficients corresponds to choosing a certain number of peaks and troughs in the first r steps of the paths.

In fact, the polynomial $Q_r(t)$ is a t -deformation of the well-known Catalan number, that is, the value of $Q_r(t)$ at $t = 1$ is the r -th Catalan number:

$$Q_r(1) = c_r = \binom{2r}{r} - \binom{2r}{r+1}. \quad (6.16)$$

Indeed it is not difficult to give a combinatorial proof that the alternating sum of the a_k 's is equal to the Catalan number c_r (the number of Dyck paths of length $2r$). We can also see it as follows. The Catalan number is equal to the constant term of a Laurent polynomial in x :

$$c_r = [(1 - x^2)(x + x^{-1})^{2r}]_0 = [(1 - x^2)(1 + x^2)^r(1 + x^{-2})^r]_0. \quad (6.17)$$

Note that we keep the variable x^2 to stay coherent with the notation used during the proof above. In fact, the first equality expresses that the Catalan number c_r is the multiplicity of the trivial representation in $V^r \otimes (V^*)^r$. The t -deformation is now immediate from this formula for c_r . Indeed, it follows from its explicit expression that the polynomial $Q_r(t)$ is given by:

$$Q_r(t) = [(1 - x^2)(1 + tx^2)^r(1 + tx^{-2})^r]_0. \quad (6.18)$$

In this sense, the polynomial $Q_r(t)$ is a natural t -deformation of the r -th Catalan number.

The numerator $P_r(t)$. The numerator of the Hilbert–Poincaré series of $Z_n(\mathfrak{sl}_2)$ is $P_r(t) = (1 + t)^r Q_r(t)$. We will show explicitly that its coefficients are positive.

First, from what we have said above about $Q_r(t)$, it follows that $P_r(t)$ is a t -deformation of the number $2^r c_r$, that is, its value at $t = 1$ is $P_r(1) = 2^r c_r$. This number counts several classes of combinatorial objects (obtained from objects counted by the Catalan number, see A151374 in [40]). The t -deformation giving $P_r(t)$ can be expressed similarly as before as:

$$P_r(t) = [(1 - x^2)(1 + t)^r(1 + tx^2)^r(1 + tx^{-2})^r]_0. \quad (6.19)$$

Now regrouping the terms with an r -th power, this gives the following expression:

$$\begin{aligned} P_r(t) &= [(1 - x^2)(1 + t^3 + (t + t^2)(1 + x^2 + x^{-2}))^r]_0 \\ &= \sum_{k=0}^r R_k \binom{r}{k} (1 + t^3)^{r-k} (t + t^2)^k, \end{aligned} \quad (6.20)$$

where the positive integer R_k is the Riordan number, one of the closest relatives of the Catalan number, which also admits many combinatorial interpretations (see A005043 in [40]). They are given by either one of the following equalities:

$$R_n = [(1 - x^2)(1 + x^2 + x^{-2})^n]_0 = \sum_{i=0}^n (-1)^{n-i} \binom{n}{i} c_i. \quad (6.21)$$

We actually used the first one in the above calculation, while the second one shows that the Catalan sequence is the binomial transform of the Riordan sequence, and thus allows to recover that $P_r(1) = 2^r c_r$.

The formula (6.20) for $P_r(t)$ has the advantage to show explicitly that it has positive coefficients. So $P_r(t)$ is a t -deformation with positive coefficients of $2^r c_r$ and therefore should be given by an interesting statistics on a certain set of $2^r c_r$ objects.

6.3 PBW basis of $Z_n(\mathfrak{sl}_2)$ for small n

With the help of the Poincaré–Hilbert series obtained and discussed above, we here determine the PBW bases of $Z_n(\mathfrak{sl}_2)$ for $n = 2, 3, 4$. Using Theorem 5.6, we identify systematically $Z_n(\mathfrak{sl}_2)$ with the special Racah algebra, and use the generators and relations in Section 4. Note that from the discussion in Section 5.2, in order to show that a subset of $Z_n(\mathfrak{sl}_2)$ is a spanning set, it is enough to show that the images in the graded algebra of invariant polynomial functions is a spanning set.

The case $n = 2$. The Hilbert–Poincaré series of $Z_2(\mathfrak{sl}_2)$ is:

$$F_2(t) = \frac{1}{(1-t^2)^3}. \quad (6.22)$$

This allows easily to recover that the algebra of invariant polynomial functions, and $Z_2(\mathfrak{sl}_2)$, is a commutative polynomial algebra. A basis of $Z_2(\mathfrak{sl}_2)$ is $P_{11}^a P_{12}^b P_{22}^c$ with $a, b, c \in \mathbb{Z}_{\geq 0}$. Indeed this set is obviously a spanning set, and moreover spans a subspace whose dimensions in each degree are given precisely by the series (6.22). So this set is also linearly independent, and is a basis.

The case $n = 3$. The Hilbert–Poincaré series of $Z_3(\mathfrak{sl}_2)$ is:

$$F_3(t) = \frac{1+t^3}{(1-t^2)^6}. \quad (6.23)$$

This allows to show that the following set is a basis of $Z_3(\mathfrak{sl}_2)$:

$$\{P_{11}^a P_{12}^b P_{13}^c P_{22}^d P_{23}^e P_{33}^f F_{123}^\varepsilon\}, \quad \text{where } a, b, c, d, e, f \in \mathbb{Z}_{\geq 0} \text{ and } \varepsilon \in \{0, 1\}. \quad (6.24)$$

Indeed, such a set is a spanning set since F_{123}^2 can be expressed in terms of the P 's. Comparing with $F_3(t)$, we see directly that this set spans a subspace of the correct dimension in each degree. So this is a basis.

The case $n = 4$. The Hilbert–Poincaré series of $Z_4(\mathfrak{sl}_2)$ is:

$$F_4(t) = \frac{1+t^2+4t^3+t^4+t^6}{(1-t^2)^9}. \quad (6.25)$$

The four sets, for $a, b, c, d, e, f, g, h, i \in \mathbb{Z}_{\geq 0}$,

$$\{F_{123} P_{11}^a P_{12}^b P_{13}^c P_{14}^d P_{22}^e P_{23}^f P_{24}^g P_{33}^h P_{34}^i\}, \quad (6.26a)$$

$$\{F_{124} P_{11}^a P_{12}^b P_{13}^c P_{14}^d P_{22}^e P_{23}^f P_{24}^g P_{34}^h P_{44}^i\}, \quad (6.26b)$$

$$\{F_{134} P_{11}^a P_{12}^b P_{13}^c P_{14}^d P_{23}^e P_{24}^f P_{33}^g P_{34}^h P_{44}^i\}, \quad (6.26c)$$

$$\{F_{234} P_{12}^a P_{13}^b P_{14}^c P_{22}^d P_{23}^e P_{24}^f P_{33}^g P_{34}^h P_{44}^i\}, \quad (6.26d)$$

and, for $a, b, c, d, e, f, g, h, i, j \in \mathbb{Z}_{\geq 0}$ and $aejh = 0$,

$$\{P_{11}^a P_{12}^b P_{13}^c P_{14}^d P_{22}^e P_{23}^f P_{24}^g P_{33}^h P_{34}^i P_{44}^j\} \quad (6.26e)$$

form a basis of $Z_4(\mathfrak{sl}_2)$. To understand this, rewrite its Hilbert–Poincaré series as

$$F_4(t) = \frac{4t^3}{(1-t^2)^9} + \frac{1-t^8}{(1-t^2)^{10}}. \quad (6.27)$$

The first term corresponds to the first 4 sets and the second term corresponds to the fifth ones. These sets are spanning sets since $F_{ijk}F_{mnp}$ can be expressed in terms linear in F by using the relations $w_{ijk} = 0$ and $x_{ijkl} = 0$ and $F_{ijk}P_{\ell\ell}$ (i, j, k, ℓ pairwise distinct) can be expressed in terms of elements of the sets (6.26) by using (3.9). The condition $aejh = 0$ for the fifth set comes from the following fact. Let us define the following 2×2 and 4×4 matrices

$$P_{ij}^{ab} = \begin{pmatrix} P_{ia} & P_{ib} \\ P_{ja} & P_{jb} \end{pmatrix}, \quad P_{ijkl}^{abcd} = \begin{pmatrix} P_{ia} & P_{ib} & P_{ic} & P_{id} \\ P_{ja} & P_{jb} & P_{jc} & P_{jd} \\ P_{ka} & P_{kb} & P_{kc} & P_{kd} \\ P_{la} & P_{lb} & P_{lc} & P_{ld} \end{pmatrix}. \quad (6.28)$$

Recall the definition of the symmetrized determinant (2.4). In $Z_4(\mathfrak{sl}_2)$, the following relation of degree 8 is satisfied by the generators:

$$\begin{aligned} \det(P_{1234}^{1234}) &= -\frac{1}{3} (\det(P_{123}^{124}) - \det(P_{123}^{134}) + \det(P_{123}^{234}) + \det(P_{124}^{134}) - \det(P_{124}^{234}) + \det(P_{134}^{234})) \\ &\quad + \frac{2}{3} (P_{12} \det(P_{34}^{34}) + P_{13} \det(P_{24}^{24}) + P_{14} \det(P_{23}^{23}) \\ &\quad + P_{23} \det(P_{14}^{14}) + P_{24} \det(P_{13}^{13}) + P_{34} \det(P_{12}^{12})). \end{aligned} \quad (6.29)$$

Note that the above relation is not a new relation (it is implied by the defining relations of $sR(4)$ given in (4.4)–(4.5)) and permits to express $P_{11}P_{22}P_{33}P_{44}$ in terms of the elements of the sets (6.26).

7 Conclusion

Classical results about the invariant theory of the polynomials on \mathfrak{sl}_2^m has allowed to provide a description in terms of generators and relations of the diagonal centralizer of \mathfrak{sl}_2 . A precise connection with the higher rank Racah algebra was given. Various questions arise and pave the way to different generalizations.

We would like to emphasize that the natural numbers appearing in the numerator of the Hilbert–Poincaré form very well-known series of integers that have numerous interpretations and appear already in the study of the representation theory of \mathfrak{sl}_2 . This suggests that there should be a further understanding of these numbers.

The classification of the finite irreducible representations of the rank 1 Racah algebra has been done in [12]. Following this, it should be possible to study the finite-dimensional representations of the (special) higher rank Racah algebra $R(n)$ (resp. $sR(n)$). These representations must be closely related to the operators associated to the $(n-2)$ -variable Racah polynomials. Indeed, the difference and recurrence operators characterizing the univariate Racah polynomials satisfy the relations of $R(3)$. The study of the generalization to $(n-2)$ -variable polynomials has been initiated in [25, 26] and it would be interesting to verify if the operators used in this case realize $R(n)$ or $sR(n)$. The Racah algebra appears also as the symmetry algebra of some superintegrable models [13–17]. We trust that the observations and theorems of the present paper will lead to a deeper understanding of this symmetry.

We focused here on the diagonal centralizer of \mathfrak{sl}_2 in the n -fold tensor product of \mathfrak{sl}_2 . Other cases where \mathfrak{sl}_2 is replaced by other algebras are also known. For example the diagonal

centralizer of the oscillator algebra has been studied in [41], the diagonal centralizer of the super Lie algebra $\mathfrak{osp}(1|2)$ is known to be related to the Bannai–Ito algebra [42] and the centralizer of \mathfrak{sl}_3 in its twofold tensor product has been introduced in [33]. An important generalization concerns the quantum group $U_q(\mathfrak{sl}_2)$. Its diagonal centralizer in the 3-fold tensor product has been examined [43] and is associated to the Askey–Wilson algebra (see e.g. [27] for a review). Many attempts [44–46] to generalize this result to n -fold tensor products have yielded relations of this centralizer but certainly did not give all the defining relations. Looking ahead, we are planning to provide a complete set of defining relations, by using some deformation of the defining relations of the special Racah algebra given in this paper.

We have studied the centralizer $Z_n(\mathfrak{sl}_2)$ at an algebraic level. It is however equally important to study this centralizer when each factor in the n -fold tensor product is in a finite-dimensional irreducible representation of \mathfrak{sl}_2 . In the case $n = 3$, a conjecture stating that the centralizer is a quotient of the Racah algebra $R(3)$ was given in [47] (see [48] for the q -deformed case and [49] for $\mathfrak{osp}(1|2)$). This quotient associates the Racah algebra with well-known algebras such as the Temperley–Lieb or Brauer algebras. The generalization of these results to the case of the n -fold tensor product is desirable and the results obtained in the present paper offer a nice starting point. As another follow-up, we plan on finding the explicit quotient that provides a description of these centralizers in representations and to compare them to the recent results reported in [50, 51].

Acknowledgments

N. Crampé and L. Poulain d’Andecy are partially supported by Agence Nationale de la Recherche Projet AHA ANR-18-CE40-0001. J. Gaboriaud held an Alexander-Graham-Bell scholarship from the Natural Sciences and Engineering Research Council of Canada (NSERC) and received scholarships from the ISM and the Université de Montréal. The research of L. Vinet is funded in part by a Discovery Grant from NSERC.

References

- ¹H. De Bie, V. X. Genest, W. van de Vijver, and L. Vinet, “A higher rank Racah algebra and the \mathbb{Z}_2^n Laplace–Dunkl operator”, *Journal of Physics A: Mathematical and Theoretical* **51**, 025203 (2017), arXiv:1610.02638.
- ²G. I. Lehrer and R. B. Zhang, “Strongly multiplicity free modules for Lie algebras and quantum groups”, *Journal of Algebra* **306**, 138–174 (2006).
- ³V. X. Genest, L. Vinet, and A. Zhedanov, “The equitable Racah algebra from three $\mathfrak{su}(1, 1)$ algebras”, *Journal of Physics A: Mathematical and Theoretical* **45**, 025203 (2013), arXiv:1309.3540.
- ⁴R. Koekoek, P. A. Lesky, and R. F. Swarttouw, *Hypergeometric Orthogonal Polynomials and Their q -Analogues*, Springer Monographs in Mathematics (Springer, 2010), 578 pp.
- ⁵Y. A. Granovskii and A. Zhedanov, “Nature of the symmetry group of the $6j$ -symbol”, *Journal of Experimental and Theoretical Physics* **94**, 49–54 (1988).
- ⁶Y. A. Granovskii and A. Zhedanov, “Linear covariance algebra for $SL_q(2)$ ”, *Journal of Physics A: Mathematical and General* **26**, L357 (1993).

- ⁷T. H. Koornwinder, “Askey–Wilson Polynomials as Zonal Spherical Functions on the $SU(2)$ Quantum Group”, *SIAM Journal on Mathematical Analysis* **24**, 795 (1993).
- ⁸S. Bockting-Conrad and H.-W. Huang, “The universal enveloping algebra of \mathfrak{sl}_2 and the Racah algebra”, *Communications in Algebra* **48**, 1022–1040 (2020), arXiv:1907.02135.
- ⁹N. Crampé, D. Shaaban Kabakibo, and L. Vinet, “New realizations of algebras of the Askey–Wilson type in terms of Lie and quantum algebras”, *Reviews in Mathematical Physics* **33**, 2150002 (2021), arXiv:2005.06957.
- ¹⁰P. Terwilliger, “Two linear transformations each tridiagonal with respect to an eigenbasis of the other”, *Linear Algebra and its Applications* **330**, 149 (2001), arXiv:math/0406555.
- ¹¹S. Gao, Y. Wang, and B. Hou, “The classification of Leonard triples of Racah type”, *Linear Algebra and its Applications* **439**, 1834–1861 (2013).
- ¹²H.-W. Huang, “Finite-dimensional modules of the universal Racah algebra and the universal additive DAHA of type (C_1^\vee, C_1) ”, *Journal of Pure and Applied Algebra* **225**, 106653 (2021), arXiv:1906.09160.
- ¹³E. G. Kalnins, J. M. Kress, and W. Miller Jr., “Second-order superintegrable systems in conformally flat spaces. I. Two-dimensional classical structure theory”, *Journal of Mathematical Physics* **46**, 053509 (2005).
- ¹⁴S. Post, “Models of Quadratic Algebras Generated by Superintegrable Systems in 2D”, *Symmetry, Integrability and Geometry: Methods and Applications* **7**, 036–055 (2011), arXiv:1104.0734.
- ¹⁵E. G. Kalnins, W. Miller Jr., and S. Post, “Contractions of 2D 2nd Order Quantum Superintegrable Systems and the Askey Scheme for Hypergeometric Orthogonal Polynomials”, *Symmetry, Integrability and Geometry: Methods and Applications* **9**, 057–084 (2013), arXiv:1212.4766.
- ¹⁶V. X. Genest, L. Vinet, and A. Zhedanov, “Superintegrability in Two Dimensions and the Racah–Wilson Algebra”, *Letters in Mathematical Physics* **104**, 931–952 (2014), arXiv:1307.5539.
- ¹⁷V. X. Genest, L. Vinet, and A. Zhedanov, “The Racah algebra and superintegrable models”, *Journal of Physics: Conference Series* **512**, 012011 (2014), arXiv:1312.3874.
- ¹⁸H.-W. Huang and S. Bockting-Conrad, “Finite-Dimensional Irreducible Modules of the Racah Algebra at Characteristic Zero”, *Symmetry, Integrability and Geometry: Methods and Applications* **16**, 018–034 (2020), arXiv:1910.11446.
- ¹⁹S. Post, “Racah Polynomials and Recoupling Schemes of $\mathfrak{su}(1,1)$ ”, *Symmetry, Integrability and Geometry: Methods and Applications* **11**, 057 (2015), arXiv:1504.03705.
- ²⁰E. G. Kalnins, W. Miller Jr., and S. Post, “Two-Variable Wilson Polynomials and the Generic Superintegrable System on the 3-Sphere”, *Symmetry, Integrability and Geometry: Methods and Applications* **7**, 051 (2011), arXiv:1010.3032.
- ²¹H. De Bie, P. Iliev, and L. Vinet, “Bargmann and Barut-Girardello models for the Racah algebra”, *Journal of Mathematical Physics* **60**, 011701 (2019), arXiv:1811.04203.
- ²²D. Latini, I. Marquette, and Y.-Z. Zhang, “Embedding of the Racah Algebra $R(n)$ and Superintegrability”, *Annals of Physics* **426**, 168397 (2020), arXiv:2010.12822.
- ²³J. Gaboriaud, L. Vinet, S. Vinet, and A. Zhedanov, “The generalized Racah algebra as a commutant”, *Journal of Physics: Conference Series* **1194**, 012034 (2019), arXiv:1808.09518.

- ²⁴J. S. Geronimo and P. Iliev, “Bispectrality of Multivariable Racah–Wilson Polynomials”, *Constructive Approximation* **31**, 417–457 (2010), arXiv:0705.1469.
- ²⁵H. De Bie, P. Iliev, W. van de Vijver, and L. Vinet, “The Racah algebra: An overview and recent results”, (2020), arXiv:2001.11195.
- ²⁶H. De Bie and W. van de Vijver, “A discrete realization of the higher rank Racah algebra”, *Constructive Approximation* **52**, 1–29 (2020), arXiv:1808.10520.
- ²⁷N. Crampé, L. Frappat, J. Gaboriaud, L. Poulain d’Andecy, E. Ragoucy, and L. Vinet, “The Askey–Wilson algebra and its avatars”, *Journal of Physics A: Mathematical and Theoretical* **54**, 063001 (2021), arXiv:2009.14815.
- ²⁸C. Procesi, “The invariant theory of $n \times n$ matrices”, *Advances in Mathematics* **19**, 306–381 (1976).
- ²⁹J. P. Razmyslov, “Trace identities of full matrix algebras over a field of characteristic zero”, *Mathematics of the USSR-Izvestiya* **8**, 727–760 (1974).
- ³⁰K. S. Sibirskii, “Algebraic invariants for a set of matrices”, *Siberian Mathematical Journal* **9**, 115–124 (1968).
- ³¹V. Drensky, “Computing with matrix invariants”, *Mathematica Balkanica* **21**, 141–172 (2007), arXiv:math/0506614.
- ³²V. Drensky, “Defining Relations for the Algebra of Invariants of 2×2 Matrices”, *Algebras and Representation Theory* **6**, 193–214 (2003).
- ³³N. Crampé, L. Poulain d’Andecy, and L. Vinet, “A Calabi–Yau algebra with E_6 symmetry and the Clebsch–Gordan series of $\mathfrak{sl}(3)$ ”, (2020), arXiv:2005.13444.
- ³⁴R. P. Stanley, “Hilbert functions of graded algebras”, *Advances in Mathematics* **28**, 57–83 (1978).
- ³⁵Y. Teranishi, “The ring of invariants of matrices”, *Nagoya Mathematical Journal* **104**, 140–161 (1986).
- ³⁶E. Formanek, “The invariants of $n \times n$ matrices”, in *Invariant Theory*, Lecture Notes in Mathematics, vol 1278 (Springer, Berlin, 1987).
- ³⁷H. W. Gould, “A new symmetrical combinatorial identity”, *Journal of Combinatorial Theory, Series A* **13**, 278–286 (1972).
- ³⁸A. A. Kirillov, “Certain division algebras over a field of rational functions”, *Functional Analysis and Its Applications* **1**, 87–88 (1967).
- ³⁹M. Hochster and J. L. Roberts, “Rings of invariants of reductive groups acting on regular rings are Cohen–Macaulay”, *Advances in Mathematics* **13**, 115–175 (1974).
- ⁴⁰OEIS Foundation Inc., “The On-Line Encyclopedia of Integer Sequences”, <http://oeis.org> (2021).
- ⁴¹N. Crampé, W. van de Vijver, and L. Vinet, “Racah problems for the oscillator algebra, the Lie algebra \mathfrak{sl}_n , and multivariate Krawtchouk polynomials”, *Annales Henri Poincaré* **21**, 3939–3971 (2020), arXiv:1909.12643.
- ⁴²V. X. Genest, L. Vinet, and A. Zhedanov, “The Bannai–Ito polynomials as Racah coefficients of the $sl_{-1}(2)$ algebra”, *Proceedings American Mathematical Society* **142**, 1545 (2012), arXiv:1205.4215.

- ⁴³A. S. Zhedanov, ““Hidden symmetry” of Askey–Wilson polynomials”, *Theoretical and Mathematical Physics* **89**, 1146–1157 (1991).
- ⁴⁴S. Post and A. Walter, “A higher rank extension of the Askey-Wilson Algebra”, (2017), arXiv:1705.01860.
- ⁴⁵H. De Bie, H. De Clercq, and W. van de Vijver, “The Higher Rank q -Deformed Bannai-Ito and Askey-Wilson Algebra”, *Communications in Mathematical Physics* **374**, 277–316 (2020), arXiv:1805.06642.
- ⁴⁶H. De Clercq, “Higher Rank Relations for the Askey-Wilson and q -Bannai-Ito Algebra”, *Symmetry, Integrability and Geometry: Methods and Applications* **15**, 099 (2019), arXiv:1908.11654.
- ⁴⁷N. Crampé, L. Poulain d’Andecy, and L. Vinet, “Temperley-Lieb, Brauer and Racah algebras and other centralizers of $\mathfrak{su}(2)$ ”, (2019), arXiv:1905.06346.
- ⁴⁸N. Crampé, L. Vinet, and M. Zaimi, “Temperley–Lieb, Birman–Murakami–Wenzl and Askey–Wilson algebras and other centralizers of $U_q(\mathfrak{sl}_2)$ ”, (2020), arXiv:2008.04905.
- ⁴⁹N. Crampé, L. Frappat, and L. Vinet, “Centralizers of the superalgebra $\mathfrak{osp}(1|2)$: the Brauer algebra as a quotient of the Bannai–Ito algebra”, *Journal of Physics A: Mathematical and Theoretical* **52**, 424001 (2019), arXiv:1906.03936.
- ⁵⁰N. Crampé and L. Poulain d’Andecy, “Fused braids and centralisers of tensor representations of $U_q(\mathfrak{gl}_N)$ ”, (2020), arXiv:2001.11372.
- ⁵¹S. M. Flores and E. Peltola, “Higher-spin quantum and classical Schur-Weyl duality for \mathfrak{sl}_2 ”, (2020), arXiv:2008.06038.