



**HAL**  
open science

## Photochemistry and Diffusion in Jupiter's Stratosphere

Julianne I. Moses, Thierry Fouchet, Bruno Bézard, Emmanuel Lellouch, G.  
Randall Gladstone

► **To cite this version:**

Julianne I. Moses, Thierry Fouchet, Bruno Bézard, Emmanuel Lellouch, G. Randall Gladstone. Photochemistry and Diffusion in Jupiter's Stratosphere. 34th Annual Lunar and Planetary Science Conference, Mar 2003, League City, Texas, United States. pp.1079. hal-03809277

**HAL Id: hal-03809277**

**<https://hal.science/hal-03809277>**

Submitted on 1 Dec 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**PHOTOCHEMISTRY AND DIFFUSION IN JUPITER'S STRATOSPHERE.** J. I. Moses<sup>1</sup>, T. Fouchet<sup>2</sup>, B. Bézard<sup>3</sup>, E. Lellouch<sup>3</sup>, and G. R. Gladstone<sup>4</sup>, <sup>1</sup>Lunar and Planetary Institute, 3600 Bay Area Blvd., Houston TX 77058-1113 (moses@lpi.usra.edu), <sup>2</sup>Atmospheric, Oceanic and Planetary Physics, Oxford University, Oxford OX1 3PU, England, <sup>3</sup>Observatoire de Paris-Meudon, F-92195 Meudon, France, <sup>4</sup>Space Sciences Dept., Southwest Research Institute, San Antonio TX 78228-0510, USA.

**Introduction:** Infrared spectra from Jupiter exhibit numerous emission features due to higher-order hydrocarbons produced from methane photolysis in the upper stratosphere. Photochemical models to date [e.g., 1,2,3] have done a poor job of reproducing the observed hydrocarbon abundances, especially if one considers the new high-quality observational constraints provided by ISO, the Infrared Space Observatory [see 4,5,6,7]. Recent advances in our knowledge of the stratospheric temperature profile, species abundances, and chemical kinetics data have provided important new model constraints and input parameters, and updated photochemical models are needed in order to explain the observed composition. We use the observational constraints provided by ISO to construct new one-dimensional steady-state models of Jovian stratospheric photochemistry. The models include coupled hydrocarbon and oxygen photochemistry, vertical eddy and molecular diffusion, and radiative transport (including multiple Rayleigh scattering by H<sub>2</sub>, He, and CH<sub>4</sub>). We focus on determining the eddy diffusion coefficient profile in the stratosphere, as this information is currently poorly constrained or contradictory [e.g., 8,9, and Fig. 1]. We also discuss the implications with regard to hydrocarbon photochemistry on all the giant planets.


**Results:** The upper-stratospheric eddy diffusion coefficient derived from observations of Jupiter's 584 Å brightness [e.g., 10] is inconsistent with that derived from the *Voyager* ultraviolet stellar occultation experiment [8] or infrared CH<sub>4</sub> v<sub>3</sub>-band fluorescence data [11]. Therefore, we have created two models (A & B) with very different high-altitude diffusion profiles (see Fig. 1). We also test the eddy diffusion profile used by [1] (Model C). The resulting mixing-ratio profiles for several observed constituents are shown in Fig. 2. Our main conclusions are as follows:

(a) Models A and B both provide a good fit to the ISO C<sub>2</sub>H<sub>2</sub>, C<sub>2</sub>H<sub>6</sub>, CH<sub>3</sub>C<sub>2</sub>H, C<sub>4</sub>H<sub>2</sub>, and C<sub>6</sub>H<sub>6</sub> data. The He profile from Model B is consistent with the 584 Å data, but the Model B CH<sub>4</sub> profile is inconsistent with the methane observations of [8,11]. The Model A CH<sub>4</sub> profile is consistent with [8,11], but the He profile is inconsistent with [10]. A reanalysis of all the above data sets [8,10,11], new UV occultation data (i.e., the *Cassini* flyby), and/or a better determination of the upper at-

mospheric temperature profile as a function of latitude are needed before this issue can be resolved.

(b) Model C and the model of [1] are inconsistent with the C<sub>4</sub>H<sub>2</sub> upper limit and the C<sub>2</sub>H<sub>6</sub>/C<sub>2</sub>H<sub>2</sub> ratio inferred from ISO data [4], suggesting that the eddy diffusion coefficient adopted in these models is too large at pressures of 1-100 mbar and/or the hydrocarbon chemistry from [1] is incorrect or incomplete.

(c) The large abundance of benzene on Jupiter and the other giant planets [e.g., 5] most probably results from the long dissociation lifetime of the C<sub>6</sub>H<sub>6</sub> molecule. Upon absorption of a UV photon, electronically excited C<sub>6</sub>H<sub>6</sub> can be collisionally quenched at lower stratospheric pressures before it dissociates.



**Figure 1.** The eddy diffusion coefficients adopted for Model A (solid line), Model B (dashed line), and Gladstone *et al.* [1] and Model C (dotted line) compared with various observational constraints (individual data points).

**Figure 2 (next page):** Mixing ratio profiles from our models compared with various infrared and ultraviolet observations (line styles as labeled in Fig. 2a).

**References:** [1] Gladstone, G. R. *et al.* (1996) *Icarus* **119**, 1. [2] Romani, P. N. (1996) *Icarus* **122**, 233. [3] Lee, A. Y. T. *et al.* (2000) *JGR* **105**, 20207. [4] Fouchet, T. *et al.* (2000) *Astron. Astrophys.* **355**, L13. [5] Bézard, B. *et al.* (2001) *Icarus* **154**, 492. [6] Drossart, P. *et al.* (1999) [7] Encenaz, T. *et al.* (1996) *Astron. Astrophys.* **315**, L397. [8] Yelle, R. V. *et al.* (1996) *JGR* **101**, 2149. [9] Festou, M. C. *et al.* (1981) *JGR* **86**, 5715. [10] Vervack, R. J., Jr. *et al.* (1995). *Icarus* **114**, 163. [11] Drossart, P. *et al.* (1999), *In The Universe as Seen by ISO*, p. 169, ESA SP-427.

## PHOTOCHEMISTRY AND DIFFUSION IN JUPITER'S STRATOSPHERE: J. I. Moses et al.

