



HAL
open science

Receptor Tyrosine Kinase KIT: A New Look for an Old Receptor

Julie Ledoux, Luba Tchertanov

► **To cite this version:**

Julie Ledoux, Luba Tchertanov. Receptor Tyrosine Kinase KIT: A New Look for an Old Receptor. Bioinformatics and Biomedical Engineering 2022, pp.133 - 137, 2022, 10.1007/978-3-031-07802-6_11 . hal-03809900

HAL Id: hal-03809900

<https://hal.science/hal-03809900>

Submitted on 11 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Metadata of the chapter that will be visualized in SpringerLink

Book Title	Bioinformatics and Biomedical Engineering	
Series Title		
Chapter Title	Receptor Tyrosine Kinase KIT: A New Look for an Old Receptor	
Copyright Year	2022	
Copyright HolderName	Springer Nature Switzerland AG	
Corresponding Author	Family Name	Ledoux
	Particle	
	Given Name	Julie
	Prefix	
	Suffix	
	Role	
	Division	
	Organization	Université Paris-Saclay, ENS Paris-Saclay, Centre Borelli
	Address	4 Avenue des Sciences, 91190, Gif-sur-Yvette, France
	Email	julie.ledoux@ens-paris-saclay.fr
Author	Family Name	Tchertanov
	Particle	
	Given Name	Luba
	Prefix	
	Suffix	
	Role	
	Division	
	Organization	Université Paris-Saclay, ENS Paris-Saclay, Centre Borelli
	Address	4 Avenue des Sciences, 91190, Gif-sur-Yvette, France
	Email	
Abstract	<p>RTK KIT regulates a variety of crucial cellular processes via its cytoplasmic (CD) domain composed of the tyrosine kinase domain crowned by highly flexible domains - juxtamembrane region, kinase insert domain and C-tail, key recruitment regions for downstream signalling proteins. We generated 3D models of the full-length CD attached to the transmembrane helix to prepare a structural basis for characterization of interactions of native KIT and its oncogenic mutant D816V with signalling proteins (KIT INTERACTOME). Generic models of native KIT in inactive state and constitutively activated KIT mutant D816V were studied by molecular dynamics simulation under conditions mimicking the natural environment of KIT. With the accurate atomistic description of the multidomain KIT dynamics, we explained the role of somatic mutation D816V on structural and dynamical properties of RTK KIT focusing on its intrinsic (intra-domain) and extrinsic (inter-domain) disorder. Conformational ensembles of native and mutated KIT were represented through free energy landscapes. Strongly coupled movements within each domain and between distant domains of KIT prove the functional interdependence of these regions, described as allosteric regulation, a phenomenon widely observed in many proteins.</p>	
Keywords (separated by '-')	<p>Receptor tyrosine kinase - RTK - Full-length KIT cytoplasmic region - Intrinsically disordered regions - Somatic mutation d816v - Phosphotyrosine - Modelling - Molecular dynamics - Conformational transition - Allosteric regulation and deregulations - Free energy landscape</p>	



Receptor Tyrosine Kinase KIT: A New Look for an Old Receptor

Julie Ledoux^(✉) and Luba Tchertanov

Université Paris-Saclay, ENS Paris-Saclay, Centre Borelli, 4 Avenue des Sciences,
91190 Gif-sur-Yvette, France
julie.ledoux@ens-paris-saclay.fr

Abstract. RTK KIT regulates a variety of crucial cellular processes via its cytoplasmic (CD) domain composed of the tyrosine kinase domain crowned by highly flexible domains - juxtamembrane region, kinase insert domain and C-tail, key recruitment regions for downstream signalling proteins. We generated 3D models of the full-length CD attached to the transmembrane helix to prepare a structural basis for characterization of interactions of native KIT and its oncogenic mutant D816V with signalling proteins (KIT INTERACTOME). Generic models of native KIT in inactive state and constitutively activated KIT mutant D816V were studied by molecular dynamics simulation under conditions mimicking the natural environment of KIT. With the accurate atomistic description of the multidomain KIT dynamics, we explained the role of somatic mutation D816V on structural and dynamical properties of RTK KIT focusing on its intrinsic (intra-domain) and extrinsic (inter-domain) disorder. Conformational ensembles of native and mutated KIT were represented through free energy landscapes. Strongly coupled movements within each domain and between distant domains of KIT prove the functional interdependence of these regions, described as allosteric regulation, a phenomenon widely observed in many proteins.

[AQ1](#)

Keywords: Receptor tyrosine kinase · RTK · Full-length KIT cytoplasmic region · Intrinsically disordered regions · Somatic mutation d816v · Phosphotyrosine · Modelling · Molecular dynamics · Conformational transition · Allosteric regulation and deregulations · Free energy landscape

Receptor tyrosine kinases (RTKs) control various signalling pathways in cells. Their remarkable conformational plasticity enables the specific recognition of many molecules such as ligands, substrates or proteins. In solution, RTKs are at equilibrium between different conformations ranging from an inactive auto-inhibited state to a fully active state. Ligand-induced activation of RTKs leads to recruitment and activation of multiple downstream signalling proteins which alter the expression of genes governing cell physiology [10]. Explicit elucidation of signalling events is an important and unresolved problem in cell biology. The initiation of these cascade-like processes involves different domains of

RTK, each performs specific actions, finely concerted by a regulated allosteric mechanism controlling all functional biological processes [4, 12]. Dysregulation of RTK-controlled signalling pathways, prompted generally by gain-of-function mutations, underlies abnormal cell development leading to tumorigenesis.

Focusing on the RTK KIT, an important target in oncology [3, 11], we will discuss this RTK as a key regulator of intracellular signalling mediated by regions possessing multiple phosphorylation sites: juxtamembrane region (JMR), kinase insert domain (KID), activation loop (A-loop) and C-terminal tail. Since these regions are very flexible or disordered, their properties are not yet well understood. As KIT is a multi-functional protein, its different regions regulate catalytic processes and/or events that activate and control the signalling cascade. To complete the functions required in more than one region, these regions should be directly or collaterally coupled.

We generated the 3D model of the inactive full-length native KIT (KIT WT) attached to the transmembrane helix (Fig. 1), to (i) prepare a structural basis for the characterisation of interconnections between functional regions - JMR, tyrosine kinase (TK) domain, KID and C-tail -, and (ii) establish the interactions of KIT with its signalling proteins. Then, we investigated this model by molecular dynamics (MD) simulation in conditions mimicking its natural environment. We suggested that such atomistic description of KIT will fully elucidate structural and dynamical properties of its different functional regions. To the best of our knowledge, we have presented for the first time a model of a full-length cytoplasmic region of an RTK KIT attached to a transmembrane helix and its molecular dynamics simulations under conditions that mimic its natural environment [9].

Analysis of the simulation data (three 2- μ s MD trajectories) put in evidence that the multidomain RTK KIT is a modular protein consisting of a quasi-stable TK domain crowned by at least four intrinsically disordered (ID) regions - JMR, KID, A-loop and C-tail. These ID regions belong to two types - the very elongated (extended) and poorly folded regions (JMR, A-loop and C-tail), and the globule-like (collapsed) KID having a high level of the helical structures. KIT ID regions contain transient structures (helices and β -strands) and their local architecture displays various degrees of compaction and elongation. Therefore, the structure of each ID domain of KIT represents a very complex mixture of a broad variety of differently folded conformations which describe a reversible folding-unfolding process, specific for each ID domain. In particular, the KID, composed of transient helices linked by coils, is the most disordered domain in respect to other KIT domains, but shows a globule-like shape stabilised by non-covalent interactions [5, 7].

Also, the inherently disordered KID shows different positions derived from two types of motions - linear (translation) and angular (rotation) displacement - regarding the stable TK domain. The elongated regions (JMR, A-loop and C-tail) show rather local disorder as evidenced by alternating positions of their short segments relative to the stable TK domain.




Fig. 1. The 3D model of full-length KIT in native and constitutively active states attached to the transmembrane helix. Modelling and study by MD simulations (top panel) analysed to (i) characterise the global motions and coupling, (ii) to represent the MD conformations as the free energy landscape, (iii) to describe the relative positions of the KIT functional regions and the position of phosphotyrosine residues.

The two-level disorder (intrinsic and extrinsic) provides high conformational variability of KIT and supplies the high adaptability of JMR, KID and C-tail required for the scaffolding (docking sites) and recruitment of different protein partners of KIT and accomplishes the tight regulation of cellular processes. Consequently, the overall structure of KIT represents a continuous spectrum of conformations with a different degree and depth of disorder, thereby generating a complex protein structural space. It is partially reflected by free energy landscapes (FEL) lacking a unique global deep minimum as typically observed in ordered proteins. Such energy landscapes, with two local minima joined by a ‘flattened plateau’ containing the intermediate conformations, show that KIT is extremely sensitive to different environmental changes (e.g. phosphorylation) that can alter its FEL in different ways. We suggest that JMR, displaced from its packed auto-inhibited position upon the SCF-induced activation of KIT, will achieve higher levels of disorder, and therefore a higher level of adaptability for the recruitment of signalling proteins. We suggest that JMR, displaced from its packed auto-inhibited position upon the SCF-induced activation of KIT, will achieve higher levels of disorder, and therefore a higher level of adaptability for the recruitment of signalling proteins.

Since ID domains are multiple in RTK KIT, does the disorder/order of one domain depend on the disorder/order of other remote regions? As evidenced by the cross-correlations, the highly coupled motions of distant regions of KIT suggest their functional dependence, classified as allosteric regulation, phenomenon

largely observed in many proteins. In particular, the coupling motions within the TK domain reflect the allosteric regulation of kinase function which is well-described for different non-receptor and receptor tyrosine kinases [1]. The coupled/uncoupled motions of A-loop and JMR were described through their allosteric communication in the wild-type KIT and oncogenic mutants [2,6], although this characterization was made using limited structural data with only partially resolved JMR.

To characterise function-related mutation-induced effects in KIT, we studied KID D816V mutant represented by the most realistic the full-length model [8]. Even through KID D816V structure is highly similar to native KIT, motions of these two proteins are quite different. Indeed, KID D816V mutant shows significantly increasing coupling of intra- and inter-domain motions. Moreover, KIT communication pathways network is strongly different in two proteins. Such multipanel characterisation has explained more explicitly the role of each region in maintaining KIT inactive and constitutively active state and/or as a signalling support for the phosphotyrosines-containing regions, and has established relationships between them.

This dynamic model of allosterically regulated KIT in two states is the first step for the reconstruction of its (i) INTERACTOME, composed of a set of KIT complexes with its signalling proteins, and its (ii) DYNASOME, constituting of an ensemble of KIT intermediate conformations before, over and after post-transduction processes upon its physiological and pathological context. We also suggested that KIT in inactive state encodes all properties of the active protein and post-transduction events. Such hypothesis echoes high dependences of the INTERACTOME to the DYNASOME.

References

1. Ahuja, L.G., Aoto, P.C., Kornev, A.P., Veglia, G., Taylor, S.S.: Dynamic allostery-based molecular workings of kinase: peptide complexes. *Proc. Natl. Acad. Sci.* **116**(30), 15052–15061 (2019). <https://doi.org/10.1073/pnas.1900163116>. <http://www.pnas.org/lookup/doi/10.1073/pnas.1900163116>
2. Allain, A., Chauvot de Beauchêne, I., Langenfeld, F., Guarracino, Y., Laine, E., Tchertanov, L.: Allosteric pathway identification through network analysis: from molecular dynamics simulations to interactive 2D and 3D graphs. *Faraday Discuss.* **169**, 303–321 (2014). <https://doi.org/10.1039/C4FD00024B>. <http://xlink.rsc.org/?DOI=C4FD00024B>
3. Gilreath, J., Tchertanov, L., Deininger, M.: Novel approaches to treating advanced systemic mastocytosis. *Clin. Pharmacol. Adv. Appl.* **11**, 77–92 (2019). <https://doi.org/10.2147/CPAA.S206615>. <https://www.dovepress.com/novel-approaches-to-treating-advanced-systemic-mastocytosis-peer-reviewed-article-CPAA>
4. Gunasekaran, K., Ma, B., Nussinov, R.: Is allostery an intrinsic property of all dynamic proteins? *Proteins Struct. Function Bioinform.* **57**(3), 433–443 (2004). <https://doi.org/10.1002/prot.20232>. <https://onlinelibrary.wiley.com/doi/10.1002/prot.20232>

5. Inizan, F., Hanna, M., Stolyarchuk, M., Chauvot de Beauchêne, I., Tchertanov, L.: The first 3D model of the full-length KIT cytoplasmic domain reveals a new look for an old receptor. *Sci. Rep.* **10**(1), 5401 (2020). <https://doi.org/10.1038/s41598-020-62460-7>. <http://www.nature.com/articles/s41598-020-62460-7>
6. Laine, E., Auclair, C., Tchertanov, L.: Allosteric communication across the native and mutated KIT receptor tyrosine kinase. *PLoS Comput. Biol.* **8**(8), e1002661 (2012). <https://doi.org/10.1371/journal.pcbi.1002661>. <https://dx.plos.org/10.1371/journal.pcbi.1002661>
7. Ledoux, J., Trouvé, A., Tchertanov, L.: Folding and intrinsic disorder of the receptor tyrosine kinase KIT insert domain seen by conventional molecular dynamics simulations. *Int. J. Mol. Sci.* **22**(14), 7375 (2021). <https://doi.org/10.3390/ijms22147375>. <https://www.mdpi.com/1422-0067/22/14/7375>
8. Ledoux, J., Trouvé, A., Tchertanov, L.: Does cancerogenic mutation D816V influence modular receptor tyrosine kinase KIT intrinsic disorder? Paper in preparation (2022)
9. Ledoux, J., Trouvé, A., Tchertanov, L.: The inherent coupling of intrinsically disordered regions in the multidomain receptor tyrosine kinase KIT. *Int. J. Mol. Sci.* **23**(3), 1589 (2022). <https://doi.org/10.3390/ijms23031589>. <https://www.mdpi.com/1422-0067/23/3/1589>
10. Lemmon, M.A., Schlessinger, J.: Cell signaling by receptor tyrosine kinases. *Cell* **141**(7), 1117–1134 (2010). <https://doi.org/10.1016/j.cell.2010.06.011>. <https://linkinghub.elsevier.com/retrieve/pii/S0092867410006653>
11. Pham, D.D.M., Guhan, S., Tsao, H.: KIT and melanoma: biological insights and clinical implications. *Yonsei Med. J.* **61**(7), 562 (2020). <https://doi.org/10.3349/ymj.2020.61.7.562>. <https://eymj.org/DOIx.php?id=10.3349/ymj.2020.61.7.562>
12. Wodak, S.J., et al.: Allostery in its many disguises: from theory to applications. *Structure* **27**(4), 566–578 (2019). <https://doi.org/10.1016/j.str.2019.01.003>. <https://linkinghub.elsevier.com/retrieve/pii/S0969212619300036>

Author Queries

Chapter 11

Query Refs.	Details Required	Author's response
AQ1	This is to inform you that corresponding author has been identified as per the information available in the Copyright form.	