



HAL
open science

Advances in construction and modeling of functional neural circuits in vitro

Siu Yu A. Chow, Huaruo Hu, Tatsuya Osaki, Timothee Levi, Yoshiho Ikeuchi

► **To cite this version:**

Siu Yu A. Chow, Huaruo Hu, Tatsuya Osaki, Timothee Levi, Yoshiho Ikeuchi. Advances in construction and modeling of functional neural circuits in vitro. *Neurochemical Research*, 2022, 47, pp.2529-2544. 10.1007/s11064-022-03682-1 . hal-03810108

HAL Id: hal-03810108

<https://hal.science/hal-03810108v1>

Submitted on 11 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Advances in construction and modeling of functional neural circuits *in vitro*

Siu Yu A. Chow^{1,2,3} · Huaruo Hu^{1,2} · Tatsuya Osaki^{1,2,3} · Timothée Levi^{1,3,4} · Yoshiho Ikeuchi^{1,2,3}

Received: 31 March 2022 / Revised: 26 June 2022 / Accepted: 8 July 2022 / Published online: 9 August 2022
© The Author(s) 2022

Abstract

Over the years, techniques have been developed to culture and assemble neurons, which brought us closer to creating neuronal circuits that functionally and structurally mimic parts of the brain. Starting with primary culture of neurons, preparations of neuronal culture have advanced substantially. Development of stem cell research and brain organoids has opened a new path for generating three-dimensional human neural circuits. Along with the progress in biology, engineering technologies advanced and paved the way for construction of neural circuit structures. In this article, we overview research progress and discuss perspective of *in vitro* neural circuits and their ability and potential to acquire functions. Construction of *in vitro* neural circuits with complex higher-order functions would be achieved by converging development in diverse major disciplines including neuroscience, stem cell biology, tissue engineering, electrical engineering and computer science.

Introduction

During development *in vivo*, cells can flawlessly generate the whole human body. One of the highlights of embryogenesis is the formation of the brain, which is achieved by the most complex choreography and orchestration of cells. In addition to cellular processes that occur during formation of other organs (e.g. proliferation, differentiation, and migration), diverse types of neurons extend and connect their dendrites and axons, and go through maturation process, to generate complex networks of neural circuits. The functions

of neurons as the key elements of the brain are to propagate action potential within the cells and to transmit and receive activity through synapses. Neurons can be cultured *in vitro*, go through maturation process, retain their elementary functions as neurons, and exhibit characteristics of diverse cell types. Therefore, it would be natural to hope and be optimistic for the formation of functional neural circuits *in vitro*. However, construction of functional neural circuits *in vitro* could be considered as one of the most difficult tasks that have been sought after for a long time.

Functions of neural circuits could be appreciated from two perspectives. Considering neurons as elements for neural circuits, abilities or characteristics of neurons emerged by forming networks could be considered as functions of the circuits. Considering neural circuits as sub-systems of the brains, activity or response that provide mechanistic processes for the brain performance could be functions of the circuits as well. With these criteria, *in vitro* neural circuits have demonstrated their diverse mechanistic functions including complex oscillation activity and network plasticity. It is fair to assume that more unique and higher-order functions could be achieved by constructing *in vitro* neural circuits with certain structures that are organized to acquire

✉ Yoshiho Ikeuchi
yikeuchi@iis.u-tokyo.ac.jp

¹ Institute of Industrial Science, The University of Tokyo, Tokyo, Japan

² Department of Chemistry and Biotechnology, School of Engineering, The University of Tokyo, Tokyo, Japan

³ Institute for AI and Beyond, The University of Tokyo, Tokyo, Japan

⁴ IMS laboratory, CNRS UMR 5218, University of Bordeaux, Talence, France




Fig. 1 Levels of functionality and organization of neurons

signal processing capability. Ultimately, neural circuits with desired functions might be generated by connecting neurons just as a computer assembled with semiconductor and other elements.

Attempts to construct neural circuits with certain functions directly link with efforts to understand minimal structural requirements of the circuits which execute mechanistic functions within the brain. If a particular function of the brain is executed by a specific neural circuit, in theory, the same function would be reproduced by an *in vitro* neural circuit that precisely mimics the functional *in vivo* circuit structure. The neural circuit structures in the brain have been explored in detail by classical anatomical approaches and recent connectome studies [1, 2]. The functions of the neural circuits *in vivo* have been revealed rapidly by various approaches including electrophysiology, fMRI, EEG, electrode array, calcium imaging and optogenetics [3–6]. As structures and functions of the brain circuits are being clarified, the need for *in vitro* neural circuit models that can investigate the circuits' fundamental mechanisms increased correspondingly.

Neurological diseases indicated that disruption of various circuits triggers diverse functional defects of the brain. It is essential to create *in vitro* disease models that precisely recapitulate the symptomatic regions or circuits of the brain to understand their pathophysiological mechanisms. Correspondingly, *in vitro* disease modeling provide important insights into functionality of circuits by understanding their deregulation phenotypes in relation to the disease symptoms.

Conventional two-dimensional culture of neurons generate limited functionality due to random orientation and interaction of neurons. Efforts have been made to mimic neural circuits with neurons cultured *in vitro* to provide structural basis for functionalization of neurons. Various engineering techniques have been developed to control positioning, connection, and assembly of living neural cells that are fragile, unstable and hard to control. In parallel, stem cell research

enabled formation of three-dimensional brain-like tissues (brain organoids) from human stem cells, which mimic brain structures including layers of subtypes of neurons. Brain organoids showed great potentials of creating neural circuit models that resembles the brain structure and functionality more than two-dimensional cell cultures. Research of neural circuit formation is currently dynamically changing and advancing through convergence of engineering and biology and incorporation of other diverse research fields. Could higher-order functions emerge and be recognized in the *in vitro* neural circuits by extending the current efforts to control and assemble neural cells? We review the progress and perspectives toward building functional neural circuits in this review.

Individual neurons exhibit their “elementary functions” including generation of action potentials (left). The neurons form circuits and gain “mechanistic functions” including generation of complex network activity (middle). The brain exhibit “higher-order functions” by coordinating activity of the neural circuits (right).

Culture of neurons

Primary neuron culture

The foundation for the construction of a functional neural circuit *in vitro* relies on the culture of neurons capable of performing their elementary functions (Fig. 1). Dissociated neuronal culture of rat neurons was first developed over 40 years ago [7]. Isolated primary rodent neurons can develop and acquire morphologies and characteristics resembling neurons *in vivo*. Diverse types of neurons from different regions of the brain can be cultured [8–12]. The cultured neurons mature *in vitro* and exhibit ability to form synapses and induce action potentials. Over the years, primary rodent neuronal cultures have been widely applied as a model

system and significantly facilitated our understanding of neuron physiology and functions, particularly in the field of morphogenesis and synaptogenesis. Defined media and growth factors have been established for the primary neuron cultures [13, 14], which paved the way for further development of cell and tissue culture methods. Importantly, the *in vitro* cultured neurons can produce periodic synchronized burst activity [15], which could be considered as a relatively simple yet important network function, and has been proposed to model sleep/wake cycles [16, 17].

Stem cells

Primary culture of rodent neural stem cells (NSCs) have facilitated understanding of their asymmetric/symmetric division and differentiation *in vitro* [18]. Culture of NSCs as free-floating aggregates, known as neurospheres, was developed as an efficient method to maintain and expand NSCs [19, 20], illustrating the importance of cell-cell interaction for the growth of NSCs. Primary cultured neurons must be isolated from animals for each experiment. Also, primary NSCs cannot proliferate infinitely. Therefore, cell lines capable of differentiating into neurons have been used as alternative cell models, such as Neuro2A (mouse line) and SH-SY5Y (human line) [21–23]. While these cell lines present the advantage of permanent proliferation, they have insufficient differentiation potential to produce mature neurons. Although human primary NSCs have also been cultured [24, 25], difficulty in obtaining human specimens highlights the need for alternative methods to acquire human neurons.

Embryonic stem (ES) cells isolated from the inner cell mass of fertilized eggs can be differentiated into diverse cell types including neurons [26–32]. In 1998, human ES cell lines were generated and provided a major advance in the investigation of human neurons [33–36]. However, ethical debates restricted the usage of ES cells derived from human embryos. The research of human neurons was then boosted by the revolutionary invention of human induced pluripotent stem cells (iPS cells), in which somatic cells are reprogrammed to acquire pluripotency. Somatic cells could become pluripotent stem cells by overexpression of only four transcription factors (OCT3/4, KLF4, SOX2, c-MYC). Mouse iPS cells were first reported in 2006, followed by human iPS cells in 2007 [37, 38]. Human iPS cells can be produced from any individual, which is ideal for personalized regenerative medicines, disease models, and drug development. Methods for differentiating iPS cells into various neural cells have been established [39–42]. In parallel, methods to directly reprogram somatic cells into induced neurons (iNs) have also been developed and utilized widely [43]. These reliable and powerful culture methods provided

access to human neurons *in vitro* for cellular and molecular neuroscience and other fields of study.

Slice culture

While the above-mentioned neurons are randomly oriented in the culture, *in vivo* neural circuits are complex yet well organized. To extract and obtain the functional neural circuits, acute slices of the brain from animals are prepared. Acute slices of rodent hippocampus are the most studied *ex vivo* neural circuits and have significantly advanced the study of synaptic plasticity [44–47]. Brain slices of multiple regions (e.g., cortex and thalamus) have also been developed, thereby providing researchers opportunities to investigate both local microcircuits and the communication between distant target regions [48–50]. They can exhibit oscillatory activity similar to the neural circuits *in vivo* [47, 51], indicating that the neuronal circuits are at least partially functional in the slices. Organotypic brain slices can be cultured and maintained *ex vivo* for a couple of weeks, which allows researchers to observe temporal (developmental) changes of the cells. However, it is challenging to keep the circuit structure perfect for an extended period of culture, since the cells tend to migrate within and out from the tissue. Furthermore, a major limitation of brain slice methods is that it is not applicable to humans, which restricts the potential for drug development and human brain studies.

Engineered circuits in two-dimensional culture of neurons

The cultured neurons have the property of spontaneously extending neurites and forming synapses with other neurons. If no physical restriction is applied to the neurites, they will extend in random directions, form synapses between accidentally encountered neurons, and create circuits without architectural plans. Synapses can even be formed between the dendrites and axons extended from the same cell [52].

Towards establishing organized neural circuits in two-dimensional culture, spatial control of neurons and their processes is one of the most critical techniques. Since the Campenot chamber was first introduced in 1977, various microfluidic culture chips have been developed and utilized to control neurites. Typically, these devices allow separation between cell bodies and axons through control of neurite outgrowth by size. Microslits in a wall separating multiple culture chambers are designed to be wide enough for axons (approximately 0.5 μm in width) to pass through, whereas cell bodies (several tens of μm wide) cannot pass through. Thus, neurons plated into one chamber stay inside of the chamber, but axons grown from the neurons pass through the microslits and extend into another chamber.

While porous membranes can be also used for separation of axons from cell bodies, microslit chips are superior for more elaborate designs aimed at circuit formation. By using the devices, neurons can be connected via axons within the microgroove to create circuit structures. Furthermore, more advanced microfluidic devices have been applied not only for connecting two types of neurons but also for connecting more than three types of neurons to achieve co-culture of multiple cell types [53]. Techniques have been developed to control the direction of synaptic connections by devising the shape of microgrooves and compartmentalized chambers. For example, by arranging a short microslit for dendrites to pass through on one side, and a long microslit for axons on the other side, the directionality of synapses can be modeled [54]. Also, other microfluidic studies demonstrated the control of axonal outgrowth with asymmetric channels and gradient generator (e.g. Netrin-1 gradient) to regulate axonal outgrowth direction [55, 56].

It is also possible to direct neuron outgrowth by micropatterning and microstructures on the culture surface [57–61]. Application of technologies that dynamically change the patterns on the culture surface by light or heat have been invented as well [62–65]. Additionally, more advanced axon-controlling methods such as microrobot and microplates have been proposed [66–68]. Advancements in technologies to control neurons will allow us to design and create neural circuits.

Over the years, various technologies have been developed one after another in two-dimensional neuron cultures, and related ongoing research is steadily progressing towards the establishment of neural circuits with organized structures. The development of these circuits has made it possible to study neural plasticity and generate simple oscillatory activities *in vitro*, but improvements are still required to generate more complicated activities and acquire higher-order functions.

Three-dimensional culture of neurons

Assembly of three-dimensional neuronal tissues

Two-dimensional cultures of neurons contributed tremendously to the understanding of neuronal physiology, taking advantage of their accessibility for observation and ease of manipulation. Neuron culture systems and engineering techniques in three dimensions could advance the modeling of more complex neural circuits (Fig. 2).

As mentioned earlier, primary NSCs are often cultured in three-dimensional as neurospheres. The three-dimensional environment facilitates cell-cell interactions within the tissues and exposure to extracellular matrix produced by

themselves, which mimic the *in vivo* environment and promote cellular maturation. More importantly, neurons could form synapses and generate unorganized circuits within the tissues, although the internal structure of neurospheres do not necessarily mimic the brain.

Inter-regional connections are critical for coordination and information processing of the brain. To achieve this, methods to integrate the neurospheres as building blocks have been developed [69]. Neural networks were built with three-dimensional neuronal tissues mutually connected by axons [70]. Importantly, these studies demonstrated that neurons could extend their axons out from the tissues, with their cell bodies remaining in the tissue even in the *in vitro* culture. The physical segregation between axons and cell bodies is naturally achieved *in vivo*, but often cells migrate out of tissues *in vitro*, impeding the creation of neural circuits. This indicates that three-dimensional assembly of cells within the neurospheres provides strong adhesion between cells, serving as a robust structural foundation to build circuit-like structures.

Brain organoids

Stem cell research independently propelled three-dimensional tissue culture technologies. Dr. Yoshiki Sasai's group first demonstrated that cultured three-dimensional embryoid bodies of mouse ES cells can spontaneously form neural tube-like neuroepithelial structures with inner lumens [71]. The neuroepithelia can differentiate and form cortical layer structures, recapitulating early *in vivo* cortical embryogenesis, which was not typically observed in the neurospheres that are mentioned in the previous section. This marked an important turning point for *in vitro* neuroscience research by demonstrating that stem cells can self-organize to produce complex and organized three-dimensional tissues.

Similarly, human iPS cells and ES cells in three-dimensional aggregates can undergo differentiation and self-organization that result in tissues with brain-like structures [72]; such brain-like tissues are called human brain organoids. The development of brain organoids opened new opportunities to model human brain development *in vitro*. In addition, brain organoids derived from patient iPS cells or genetically engineered disease-model human stem cells can enhance our understanding of disease pathogenesis and facilitate the search for new drugs.

What distinguishes brain organoids from previous human neuron models is the structures of cellular assemblies e.g. spontaneously organized layers, similar to the *in vivo* human brain development. This could be extremely beneficial in the establishment of complex neural circuit constructs. Also, a population of neural stem cells named outer radial glia (oRG) cells that are present in human brains




Fig. 2 Development of *in vitro* neuronal cultures with advanced structures

has been identified in the organoids, indicating that organoids are good models of the human brain.

Various regions of our brain are mostly originated from a neural tube, in which each region gradually segments and acquires its identity. Formation of the brain regions are controlled by morphogens that are differently expressed along the anterior-posterior and dorso-ventral axes of the neural tube and the body. Organoids resembling different brain regions can be created by mimicking and reproducing the corresponding developmental processes through treatment with extrinsic factors [73]. Brain organoid has evolved as an excellent technology that successfully simulates the early stages of brain development and generates region-specific neuron subtypes.

Strikingly, organoids cultured for long periods of time (several months) have been reported to exhibit complex oscillatory activities, with the activity patterns similar to those of premature infants' brainwaves [74]. Brain organoids cultured for over a year have shown signs of post-natal stages between 250 and 300 days following culture [75]. In addition, from the perspective of neural circuit formation, complex macroscopic circuits generation is challenging as interregional circuit cannot form spontaneously or randomly within conventional organoids despite after extended duration of culture.

Conventional two-dimensional culture of neurons form random networks in a dish. Bioengineering techniques to generate organized networks mimicking local circuit structures have been developed. This could be further incorporated into a network of networks that mimic macroscopic circuit structures. Neurons could be also cultured in a three-dimensional aggregate known as a neurosphere. The brain organoids bearing advanced three-dimensional internal structures can be generated by spontaneous differentiation

and maturation of stem cells in a spheroid. The brain organoids could be further used to develop advanced structures including fused and connected organoids.

Brain organoids as disease models

Developmental brain diseases

In the past, disease pathogenesis of the human brain was largely explored through post-mortem patient biopsy, which provides only snapshots of the disease. Among various types of disease models that allow researchers to actively investigate disease pathogenesis and develop treatment, brain organoids demonstrate their unique characteristics of mimicking the human brain.

In 2013, Lancaster et al. reported the differentiation of cerebral organoids from iPS cells derived from microcephaly patients, marking the first time in which brain organoids were used to model disease phenotypes [72]. The patient-derived cerebral organoids were smaller in size and displayed premature neural differentiation than non-patient cerebral organoids. Brain organoids also contain a diverse cellular composition. By understanding the cellular heterogeneity of organoids, cell-type specific phenotypes of neurological disorders can be uncovered. For example, a recent paper modeled the neurodevelopmental phenotypes of tuberous sclerosis (TSC) [76]. Single-cell RNA-seq analyses identified a sub-group of progenitor cells in TSC patient-derived organoids that could give rise to both cortical tubers and tumors.

The unique ability of brain organoids to self-organize and mature to form layered structures led early studies using patient-derived brain organoids to reveal critical physical

phenotypes *in vitro*. As our brain's higher-order capacity depends on the functionality of neural circuits and their coordination, most neurological diseases are not limited to structural and cellular alterations of the brain, but also functional changes at a network level. As emerging evidence indicates that brain organoids contain intrinsically complex networks, the potential of brain organoids to study network disruptions in disease states has come under the spotlight. Sun *et al.* modeled Angelman Syndrome, a neurodevelopmental disease characterized by developmental delays and seizures, with CRISPR-mediated UBE3A knock-out (KO) human ES cells [77]. The KO organoids exhibited seizure-like hyperactivity and synchronous network activity that was reversed by the big potassium channel blocker paxilline. Paulsen *et al.* looked at risk genes involved in autism spectrum disorder (ASD), including SUV420H1. SUV420H1 mutant organoids displayed reduced network burst frequency and duration [78]. Correspondingly, the SUV420H1 mutant organoids exhibited disruption of developmental cell type specification. More advanced organoid disease models were established for Timothy Syndrome (TS). Hypersynchronous activity was observed in TS patient-derived dorsal forebrain organoids, which was more prominent when fused with ventral forebrain organoids (a.k.a. assembloids; Please see below "Multi-regional organoids" section) [79]. Another assembloid model for Rett Syndrome exhibited hypersynchrony, recurring epileptiform-appearing spikes and high frequency oscillations, despite no obvious deficit in cytoarchitecture [80]. Gamma oscillation in mutant assembloid was partially rescued by the administration of Pifithrin- α , a p53 inhibitor recently shown to be able to rescue premature aging induced by MeCP2 mutation, highlighting the potential of brain organoids in drug testing [80].

Neurodegenerative diseases and neurological damages

Other than neurological diseases that happen early in development, brain organoids have also shown promise in the study of late-onset, neurodegenerative diseases (NDDs). These diseases uniquely affect specific regions of the nervous system, but have the common hallmark of aggregated protein accumulation, linked to neuronal damage and their loss. NDDs cause dysfunctions of certain neural circuits that underly their distinct symptoms, making it appropriate to investigate functionality of representative neural circuits. In fact, correspondingly, recent studies suggest that aggregated proteins propagate along specific neural networks, which further suggest importance of neural circuits as disease models for understanding their pathological mechanisms. To date, various neurodegenerative disease models have been established, including the use of cerebral organoids for

Alzheimer's disease (AD) [81, 82], midbrain organoids for Parkinson's Disease (PD) [83, 84] and motor neuron organoids for Amyotrophic Lateral Sclerosis (ALS) [85, 86].

In addition to investigation on protein aggregation and cellular death, the functionality of NDD organoid models is gradually coming to light. For instance, APP mutant and PSEN1 mutant cerebral organoids were studied as AD models [87]. The authors observed increased synchronous bursts in both of the AD models, mimicking hyperexcitability and hypersynchronization observed in early stages of AD mouse models and patients [88, 89]. Chen *et al.* proposed an alternative model for pathogenesis of AD through mimicking blood-brain barrier leakage by serum exposure [90]. The exposed organoids showed synaptic loss and reduced network activity as characterized by a decrease in number of spikes, bursts and synchronization. When investigating cerebral organoid network activity in a model of amyotrophic lateral sclerosis/frontotemporal dementia (ALS/FTD), the authors found that the connectivity was comparable between mutant and control organoids despite neuronal loss in mutant organoids, demonstrating neuronal plasticity [91].

A study by Foliaki *et al.* investigated the electrophysiological properties of cerebral organoid models generated with cells from donors representing Creutzfeldt Jakob disease and PD along with Down syndrome. In early stages (3–4 months), the control and the disease model organoids exhibited no difference in activity. Interestingly, by 6–10 months, the disease model organoids demonstrated slower burst rates and longer periods between network events, as well as neural oscillation changes [92]. Since PD primarily affects dopaminergic neurons, midbrain organoids have been used to study PD. A recent study observed decreased firing rate and number of spikes that was evident following the maturation of PD-model midbrain organoids with SNCA mutation [93]. Studies on the network functionality of NDD *in vitro* are increasing, and further development is expected to uncover pathogenesis mechanisms of NDDs and to serve as platforms for drug development.

Besides the diseases with genetic causes, models have also been developed for diseases caused by damages induced by external triggers including traumatic brain injury, viral infection and malaria [94–97]. Substance abuse is also modeled with brain organoids [98, 99]. Functional analyses of network activity upon drug applications have been demonstrated [100]. Analyses on circuit functions would be useful for a deeper understanding on the chronic and acute effects of the external damages in the models.

Psychiatric diseases

Psychiatric diseases exhibit diverse symptoms that affect higher-order brain functions. Our understanding of the etiology and pathophysiology underlying these disorders remain limited. With no definite biomarkers or anatomical changes, diagnosis of psychiatric diseases largely relies on clinical symptoms such as behavioral and cognitive impairments. *In vivo* studies of patients revealed neural circuitry alterations that often span multiple brain regions. A deeper understanding of developmental progression and early network changes associated with psychiatric diseases can possibly result in earlier intervention and more treatment options. Due to the complexity of psychiatric diseases, it is extremely difficult to model the disorders *in vitro*. Cellular or molecular analyses have not been very effective to decipher the higher-order dysfunctions and it is now clear that analyses at a network level are necessary to study psychiatric disease. Fortunately, recent progress in technology and cultures has shown promises toward understanding psychiatric diseases *in vitro*.

Three-dimensional brain organoids have opened new opportunities in studying the developmental progression of psychiatric diseases. Several papers reported abnormalities in neurogenesis and neurodevelopment using organoid models of schizophrenia and bipolar disorder, including reduced neural progenitor cell proliferation, disrupted neural differentiation, and altered excitatory-inhibitory balance [101–104].

Recent studies have attempted to explore the functional alterations in organoids modelling psychiatric disorders. Schizophrenia patient-derived organoids and bipolar disorder patient-derived organoids exhibited diminished response to stimuli, although the baseline electrical activity did not differ from the control [105, 106]. These studies suggest that functional study on neural circuits could provide important insights into deregulation mechanisms of neural circuits underlying the disease symptoms, and the emerging field could advance further by developing better models and analyses.

In this last decade, we have seen major advancements in brain organoids and disease models using them. Their complexity has offered unprecedented opportunities in uncovering the functional aspects of neurological diseases at a cellular and network level. More advanced models combined with diverse technologies would provide a deeper understanding of not only the diseases but also our normal brain functions and mechanisms.

Brain organoid engineering

Multi-regional organoids

Inter-regional circuit models are crucial in understanding brain functions and mechanisms. Beyond the models of local and regional circuitry, researchers have developed more advanced strategies for modeling neural circuits structured at a relatively larger scale. Importantly, developmental interaction between brain regions is essential to form proper structures of the brain, and the interaction facilitate formation of neural networks. To model the inter-regional interaction, multiple approaches have been proposed with their effectiveness demonstrated.

The first approach is to induce developmental body axes for generating multiple regions within an organoid along the axes. During brain development, the dorsal forebrain produces excitatory neurons and becomes the cerebral cortex. Meanwhile, the ventral forebrain becomes ganglion eminence and produces inhibitory neurons which migrate tangentially to the dorsal forebrain and are incorporated into the cerebral cortex. To reproduce this process *in vitro*, a method of locally inducing ventral differentiation within an organoid has been proposed [107]. By incorporating cells that secrete the ventralization factor, SHH, an organoid gains the dorsoventral axis and forms both dorsal and ventral sides within an organoid. Local expression of SHH has been demonstrated by using light-activatable Cre recombinase system [108]. Toward this purpose, light-activatable HH signaling modulator compounds have been developed [109, 110]. The first approach is to induce developmental body axes for generating multiple regions within an organoid along the axes. Interestingly, spontaneous formation of dorsoventral axis within an organoid has been achieved by culturing the organoid on a gel and exposing the organoid to SHH or equivalent compounds [111].

Another type of approach is direct fusion of multiple organoids that models different regions of the brain. Various types of brain organoids are fused to model interaction between different pairs of regions. Such fused organoids are widely recognized as “assembloids” [112–115]. This approach is also effective in modeling the dorsal-ventral interaction of forebrain [112], in which GABAergic inhibitory neurons generated from the ventral region migrate to the dorsal organoids. Furthermore, an assembloid generated by fusing three organoids including the cerebrum, motor nerves, and muscles, has been reported, demonstrating its applicability to model complex neural circuits [116]. The assembloid approach provides a simple and powerful approach to model connections of multiple regions beyond the body axis and regardless of their native distance in the nervous system.

Construction of the brain organoid circuits

Though the above mentioned approaches have shown feasibility in modeling the interactions between multiple regions and creating circuit structures, none of them generate an axonal tract that appears to be the signature structure in the brain and serves as a critical hub for the inter-regional connections. In the brain, axonal tracts in and between cerebral hemispheres form the white matter. While distal cortical regions are connected through the white matter (extrinsic pathway) in the human brains, the corresponding axons in the mouse brains are either fewer or located inside the grey matter (intrinsic pathway) [117]. This structural difference of the axonal tracts might lead to functional differences of the brains between the species, given that axonal tracts could serve as a structural mechanism to support the massive information-processing and complex coordination in human brains. To understand the significance of this structural difference, and to accurately model the human macroscopic neural circuits, it is essential to create a model circuit in which human brain organoids are connected via axonal tracts.

The first step towards reconstructing a brain circuit that mimics the distal connections of different brain regions is the establishment of an *in vitro* axon tract model that contains self-assembled bundles of axons. With this in mind, we have previously developed a method to fabricate motor nerve organoids in microfluidic devices [118]. In this method, axons extended from the organoid is physically guided by a wall of a culture vessel into a thin microchannel in which axons extend unidirectionally and spontaneously assemble into a bundle through axo-axonal interaction. Based on this technique, a method was established to connect two cerebral organoids with an axon bundle by using the same culture vessel (microdevice) [119]. Human iPS cell-derived cerebral organoids are placed in two small chambers at the both ends of the microchannel. The organoids extend their axons into a microchannel between them, and eventually the axons reciprocally reach the cerebral organoid that has been placed on the other end of the microchannel. The two interconnected cerebral organoids can be obtained as one tissue, mimicking a cortico-cortical connection in the brain. We call these connected organoids “connectoids”. The inter-organoid connection could be also formed by connecting two organoids with reciprocally extending axons within a tube of gel [120, 121].

The difference between assembloids and connectoids lies in the axon bundles which can be useful for understanding how axonal tracts assemble and form. In addition to the importance of the axon bundle in structural and developmental studies, functionality of the tissue could be dramatically changed by the structure. Using multi-electrode array

(MEA), the activities of connectoids were observed to be significantly more complex and active than those of single cerebral organoid and cortico-cortical assembloids [122], suggesting importance of macroscopic structure and architecture of neural circuits *in vitro*.

Perspective: developing functions of neural circuits *in vitro*

In vitro neural circuits at a thrilling intersection of multiple disciplines

As described, organoid models of neural circuits are quickly advancing through the application of bioengineering technologies on biologically generated tissues, in parallel to the advances in two-dimensional neural circuits, where recent improvements largely depend on bioengineering techniques (Fig. 2). The neurons with their elementary functions acquire the ability to exhibit assembly and orchestration of neural activity as a circuit, which recapitulates the mechanistic functions underlying the higher-order functions of the brain (Fig. 1). The examples of bioengineered neural networks highlight the progress of *in vitro* neural cultures as a result of fruitful collaboration between stem cell biology and tissue engineering (Fig. 3). This is truly exciting since the two fields of study share the same ultimate goal to build and learn from organs, although different approaches have been applied. Combining the strength of the two fields certainly boosts a new (yet anticipated) stream of research in neural circuit modeling *in vitro*. Biology-driven approaches for generating neural culture and tissues, which heavily rely on cellular self-organization processes, could in theory generate tissues precisely mimicking the brain as the cells are programmed to generate the whole organ. However, self-assembly often suffers from a lack of environmental context which is supplied in embryos. Obviously making the whole embryo would be ethically unfeasible for the study of neural circuit models. Instead, a dash of engineering techniques could go a long way to supply missing guidance and components to tissues, and thus obtain models of interest at the level of desired complexity and structures. Other technologies also became essential to model *in vitro* neural circuits. To create model tissues in diverse context including diseases, genome editing and other molecular genetic approaches are becoming as important as usage of patient-derived cells. To understand activity of neurons and their functions, it is becoming essential to incorporate other disciplines including artificial intelligence (AI) and electrical engineering, which will be described below. The merging interest and participation from various fields of study on *in vitro* neural circuits create a unique opportunity for the




Fig. 3 Development of advanced functional *in vitro* neural circuits

researchers to explore new ways to code and decode neural activity patterns along with developing the circuits at the same time. This could construct a new platform and scientific community to probe fundamental mechanisms of neural circuits underlying higher-order functions of the brain.

Stem cell biology and tissue engineering technologies converge into organoid engineering technologies. By incorporating other technologies including genome editing, patient-derived cells, artificial intelligence, and electrical engineering would generate advanced functional *in vitro* models.

Structure-function relationship

A demonstration of *in vitro* circuits of three snail neurons that generate oscillation is one of the simplest and most sophisticated examples of *in vitro* network executing clear functions [123]. Since then, advanced methods to create complex and ordered neural circuits have been developed in the hope for the biomimetic circuits to be more functional. Various approaches proved that the structured and organized circuits could have more functional capacity than random circuits in both two-dimensional and three-dimensional conditions.

Random neural circuits *in vitro* often generate simple periodic burst activity, which by itself could be considered as a functional signature of the neural circuits. The periodic bursts could be generated by strong recruitment of individual neurons to fire together with other neurons to form synchronous burst activity, due to their tight and strong synaptic connections and uniformness of the circuits. This could be also considered as a hyperactive (possibly epileptic) pattern in which too many neurons fire together and hence its information coding capability is hindered, which indicate that too strong and too uniform connections limit functionality

of neural circuits. Adequate amount of inhibitory neurons and astrocytes are also critical to maintain complex activity patterns of the networks. In contrast to the random network, regional circuits in the brain are connected with other regions with their axons projected between the regions, allowing information to flow from one region to another, and the whole brain to function coordinately. Additionally, the axonal projections could serve as structural constraint for the circuit to protect them from excessive connection and thus epileptic seizures. The axonal connections also could provide time delay as important factor for functionality of computation and oscillation of neural circuits. One could speculate that the structural constraint can provide convergence points for information and neural activity, whereas the cortical regional networks as divergence segments, which could serve as an important principle for information processing. Thus, neural circuits that contains regional circuits that are inter-connected with axons could be useful for assessing the importance of the convergence/divergence structure motif. Due to structural and functional complexity, the connected organoids could be an ideal platform to assess these hypotheses and functional importance of the inter-regional connections and to formulate a fundamental structural requirement for maximizing functionality of neural circuits. Two-dimensional cultures with the macroscopic separation and connections between segmented neural circuits have demonstrated promising structural contribution to their network activity [124, 125]. Once we understand the basic requirement to increase functionality of *in vitro* neural circuits, more unique brain region- or input- specific functions should be modeled by incorporating more organoids in numbers and types into the circuits.

Unlocking and understanding higher-order functionality of *in vitro* neural circuits

In vitro neural circuits have been demonstrated to exhibit functionality including synchronized bursts, oscillation, network propagation, and neural avalanches. These activity patterns generated as neural circuits could be considered as mechanistic functions that provide foundation for higher-order functions of the brain. One of the important questions is whether higher-order functions could emerge in the neural circuits that exhibit mechanistic functionality. This question leads to another question of whether higher-order functionality could be appreciated by observers, even if the neural circuits have the capability of executing higher-order functions in itself. To answer these questions, it is essential to provide a system to relate input to the neural circuits and output from the circuits (Fig. 4). Since most higher-order functions require plasticity of neurons to execute, a closed loop stimulation system would be required to assess these important questions. In this system, neurons receive inputs and generate output signals which would be recorded and instantly processed by computers or electrical components and then it generate the input sequences for the circuits. In a closed loop setup, neural circuits are connected to virtual space where they can interact with the hypothetical external environment. To achieve this, two technological parts make up these closed-loop systems: the interface and the processing. The interface between the biological component (typically, analog signal) and the physical component (typically digital signal) is important to achieve bidirectional communication over analog-digital conversion. These interfaces includes both recording of neuronal activity and stimulation equipment of neural activity [126]. The neuronal recording must be spatiotemporally precise and needs to be captured from a wide range of culture area. Most of the systems are either electrical, in particular by using networks of electrodes (MEA [127, 128]), or optical, in particular by the use of optogenetics and fluorescent activity indicators [129]. The processing part is also an important step for bio-hybrid systems [130, 131]. Indeed, a key feature of these systems is the real-time processing and decoding of neural signals with super low latency (< 1-100 msec) to drive feedback for modulation or replacement of neural function. To this end, enhancing biohybrid systems with artificial neural networks (ANN) is the emerging strategy to achieve complex interactions between biological and artificial neurons [132]. These ANNs can be used for detection and classification but also to reproduce the biological neural network for replacement [125, 132, 133]. One approach to achieve this is a neuromorphic system allowing bidirectional communication in real time [134, 135]. Recent studies demonstrated that *in vitro* neural circuits could be incorporated into computational




Fig. 4 Computer-assisted understanding and generation of higher-order functions of neural circuits

recurrent neural network (RNN) or a “reservoir” to achieve tasks, suggesting that employing the mechanistic structure for artificial intelligence could be a powerful approach for understanding higher-order functionality of neural circuits [136]. Another notable example is that *in vitro* neural circuits have been investigated to achieve responsive game task [137], in which “free energy principle” theory has been examined. They utilized noise to destabilize the network to manipulate the neural circuit activity. With these approaches, researchers demonstrated maneuvers of objects or robots to achieve tasks, suggesting that *in vitro* neural circuits are capable of processing information and executing higher-order functions. These highlights that *in vitro* neural circuits could be ideal for testing theories and methods to unlock the ability of neural circuits and execute higher-order functions. Higher-order functionality of the biological *in vitro* neural circuits are still limited at the current stage but combining closed loop systems with circuit models equipped with more sophisticated structures would pave the way to unleash advanced higher-order functions, and lead to better understanding on the basic principles of neural network mechanisms of the brain. *In vitro* neural circuits could be ideal for both exploring methods to unlock higher-order functions and inventing new methods to effectively process information inspired by the neural circuits.

Do neurons have higher-order functions? We could investigate this by real-time interaction with them through a system with continuous input and output that provides virtual environments.

Concluding notes

Current limitations

Human stem cell-derived brain organoids provide numerous advantages to model brain functions and structure. However, currently there are a few major limitations. One of the potential concerns is the variability between organoids. Organoids often differ in their structures and activity patterns even within the same batch of organoids from the same iPS cell line, resulting in difficulty to control their quality. Cell-line-to-cell-line variability also exists. Relatively long time required for maturation and production of some of the cell types could also be a limitation. Human brain organoids need a few hundreds of days for their maturation, which is critical for producing some of the late-emerging cell types including upper layer neurons and astrocytes. Also, the brain organoids usually lack major cell types and structures, including those for the blood and immune systems. By resolving these issues, the brain organoids would have better ability to model the human brain.

Considering the functions of brain organoids, one of the most critical limitations of the current organoids is the lack of input and output as stated above. This is critical not only for understanding their hidden functional capability, but also for creating functional circuits within or with the organoids through activity-dependent synapse formation and organization mechanisms. Another limitation is the recording techniques. Although high density electrode arrays and wide-field calcium imaging techniques enable recording of neuronal activity, it is still difficult to record from all neurons in the complex tissue with substantial three-dimensional size.

Engineering techniques are rapidly developing for brain organoids, but there are many technical limitations to overcome. For example, it is currently difficult to build complex three-dimensional structures that are compatible with long-term culture and observation of the organoids. It is also difficult to establish stable long-term gradient of morphogens to establish body axes within organoids. While numerous technical challenges exist, we believe that new technologies in biology and engineering will continue to be invented to advance the construction and modeling of functional neural circuits.

Conclusions

In this review, we overviewed research progress and discussed perspective of *in vitro* neural circuits and their ability and potential to acquire functions. The field is quickly advancing and changing as exciting discoveries and inventions emerge almost every week. We apologize that many

important and interesting studies are not mentioned in this review due to the space limitation. We hope that this review helps the readers to know current advances in the field and to feel the potential for future developments.

Acknowledgements We thank all Ikeuchi lab members for critical discussion for the manuscript. This work was supported in part by a Grant-in-Aid for Challenging Research (Pioneering) from the JSPS (20K20643), a Grant-in-Aid for Transformative Research Areas (B) (20H05786), AMED-CREST (JP20gm1410001), and the Institute for AI and Beyond.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Fornito A, Zalesky A, Breakspear M. The connectomics of brain disorders. *Nat Rev Neurosci*. 2015;16(3):159–72. doi: <https://doi.org/10.1038/nrn3901>.
2. Sporns O. The human connectome: origins and challenges. *Neuroimage*. 2013;80:53–61. doi: <https://doi.org/10.1016/j.neuroimage.2013.03.023>.
3. Bassett DS, Sporns O. Network neuroscience. *Nat Neurosci*. 2017;20(3):353–64. doi: <https://doi.org/10.1038/nn.4502>.
4. Zhao S, Ting JT, Atallah HE, Qiu L, Tan J, Gloss B, et al. Cell type-specific channelrhodopsin-2 transgenic mice for optogenetic dissection of neural circuitry function. *Nat Methods*. 2011;8(9):745–52. doi: <https://doi.org/10.1038/nmeth.1668>.
5. Obien ME, Deligkaris K, Bullmann T, Bakkum DJ, Frey U. Revealing neuronal function through microelectrode array recordings. *Front Neurosci*. 2014;8:423. doi: <https://doi.org/10.3389/fnins.2014.00423>.
6. Smith SM, Vidaurre D, Beckmann CF, Glasser MF, Jenkinson M, Miller KL, et al. Functional connectomics from resting-state fMRI. *Trends Cogn Sci*. 2013;17(12):666–82. doi: <https://doi.org/10.1016/j.tics.2013.09.016>.
7. Banker GA, Cowan WM. Rat hippocampal neurons in dispersed cell culture. *Brain Res*. 1977;126(3):397–42. doi: [https://doi.org/10.1016/0006-8993\(77\)90594-7](https://doi.org/10.1016/0006-8993(77)90594-7).
8. Gaven F, Marin P, Claeysen S. Primary culture of mouse dopaminergic neurons. *J Vis Exp*. 2014(91):e51751. doi: <https://doi.org/10.3791/51751>.
9. Graber DJ, Harris BT. Purification and culture of spinal motor neurons from rat embryos. *Cold Spring Harb Protoc*. 2013;2013(4):319–26. doi: <https://doi.org/10.1101/pdb.prot074161>.
10. Lin YT, Chen JC. Dorsal Root Ganglia Isolation and Primary Culture to Study Neurotransmitter Release. *J Vis Exp*. 2018(140). doi: <https://doi.org/10.3791/57569>.

11. Mao L, Wang JQ. Primary striatal neuronal culture. *Methods Mol Med.* 2003;79:379–86. doi: <https://doi.org/10.1385/1-59259-358-5:379>.
12. Beaudoin GM, Lee SH, Singh D, Yuan Y, Ng YG, Reichardt LF, et al. Culturing pyramidal neurons from the early postnatal mouse hippocampus and cortex. *Nat Protoc.* 2012;7(9):1741–54. doi: <https://doi.org/10.1038/nprot.2012.099>.
13. Brewer GJ, Torricelli JR, Evege EK, Price PJ. Optimized survival of hippocampal neurons in B27-supplemented Neurobasal, a new serum-free medium combination. *J Neurosci Res.* 1993;35(5):567–76. doi: <https://doi.org/10.1002/jnr.490350513>.
14. Brewer GJ. Serum-free B27/neurobasal medium supports differentiated growth of neurons from the striatum, substantia nigra, septum, cerebral cortex, cerebellum, and dentate gyrus. *J Neurosci Res.* 1995;42(5):674–83. doi: <https://doi.org/10.1002/jnr.490420510>.
15. Bacci A, Verderio C, Pravettoni E, Matteoli M. Synaptic and intrinsic mechanisms shape synchronous oscillations in hippocampal neurons in culture. *Eur J Neurosci.* 1999;11(2):389–97. doi: <https://doi.org/10.1046/j.1460-9568.1999.00440.x>.
16. Saberi-Moghadam S, Simi A, Setareh H, Mikhail C, Tafti M. In vitro Cortical Network Firing is Homeostatically Regulated: A Model for Sleep Regulation. *Sci Rep.* 2018;8(1):6297. doi: <https://doi.org/10.1038/s41598-018-24339-6>.
17. Yokoi R, Okabe M, Matsuda N, Odawara A, Karashima A, Suzuki I. Impact of Sleep-Wake-Associated Neuromodulators and Repetitive Low-Frequency Stimulation on Human iPSC-Derived Neurons. *Front Neurosci.* 2019;13:554. doi: <https://doi.org/10.3389/fnins.2019.00554>.
18. Stemple DL, Anderson DJ. Isolation of a stem cell for neurons and glia from the mammalian neural crest. *Cell.* 1992;71(6):973–85. doi: [https://doi.org/10.1016/0092-8674\(92\)90393-q](https://doi.org/10.1016/0092-8674(92)90393-q).
19. Reynolds BA, Weiss S. Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system. *Science.* 1992;255(5052):1707–10. doi: <https://doi.org/10.1126/science.1553558>.
20. Reynolds BA, Weiss S. Clonal and population analyses demonstrate that an EGF-responsive mammalian embryonic CNS precursor is a stem cell. *Dev Biol.* 1996;175(1):1–13. doi: <https://doi.org/10.1006/dbio.1996.0090>.
21. Ross RA, Spengler BA, Biedler JL. Coordinate morphological and biochemical interconversion of human neuroblastoma cells. *J Natl Cancer Inst.* 1983;71(4):741–7.
22. Seeds NW, Gilman AG, Amano T, Nirenberg MW. Regulation of axon formation by clonal lines of a neural tumor. *Proc Natl Acad Sci U S A.* 1970;66(1):160–7. doi: <https://doi.org/10.1073/pnas.66.1.160>.
23. Rathjen FG, Schachner M. Immunocytological and biochemical characterization of a new neuronal cell surface component (L1 antigen) which is involved in cell adhesion. *EMBO J.* 1984;3(1):1–10.
24. Palmer TD, Schwartz PH, Taupin P, Kaspar B, Stein SA, Gage FH. Cell culture. Progenitor cells from human brain after death. *Nature.* 2001;411(6833):42–3. doi: <https://doi.org/10.1038/35075141>.
25. Stein JL, de la Torre-Ubieta L, Tian Y, Parikshak NN, Hernandez IA, Marchetto MC, et al. A quantitative framework to evaluate modeling of cortical development by neural stem cells. *Neuron.* 2014;83(1):69–86. doi: <https://doi.org/10.1016/j.neuron.2014.05.035>.
26. Nichols J, Smith A. The origin and identity of embryonic stem cells. *Development.* 2011;138(1):3–8. doi: <https://doi.org/10.1242/dev.050831>.
27. Evans MJ, Kaufman MH. Establishment in culture of pluripotent cells from mouse embryos. *Nature.* 1981;292(5819):154–6. doi: <https://doi.org/10.1038/292154a0>.
28. Martin GR. Isolation of a pluripotent cell line from early mouse embryos cultured in medium conditioned by teratocarcinoma stem cells. *Proc Natl Acad Sci U S A.* 1981;78(12):7634–8. doi: <https://doi.org/10.1073/pnas.78.12.7634>.
29. Bibel M, Richter J, Schrenk K, Tucker KL, Staiger V, Korte M, et al. Differentiation of mouse embryonic stem cells into a defined neuronal lineage. *Nat Neurosci.* 2004;7(9):1003–9. doi: <https://doi.org/10.1038/nn1301>.
30. Wichterle H, Lieberam I, Porter JA, Jessell TM. Directed differentiation of embryonic stem cells into motor neurons. *Cell.* 2002;110(3):385–97. doi: [https://doi.org/10.1016/S0092-8674\(02\)00835-8](https://doi.org/10.1016/S0092-8674(02)00835-8).
31. Kawasaki H, Mizuseki K, Nishikawa S, Kaneko S, Kuwana Y, Nakanishi S, et al. Induction of midbrain dopaminergic neurons from ES cells by stromal cell-derived inducing activity. *Neuron.* 2000;28(1):31–40. doi: [https://doi.org/10.1016/S0896-6273\(00\)00083-0](https://doi.org/10.1016/S0896-6273(00)00083-0).
32. Bain G, Kitchens D, Yao M, Huettner JE, Gottlieb DI. Embryonic stem cells express neuronal properties in vitro. *Dev Biol.* 1995;168(2):342–57. doi: <https://doi.org/10.1006/dbio.1995.1085>.
33. Thomson JA, Itskovitz-Eldor J, Shapiro SS, Waknitz MA, Swiergiel JJ, Marshall VS, et al. Embryonic stem cell lines derived from human blastocysts. *Science.* 1998;282(5391):1145–7. doi: <https://doi.org/10.1126/science.282.5391.1145>.
34. Zhang SC, Wernig M, Duncan ID, Brüstle O, Thomson JA. In vitro differentiation of transplantable neural precursors from human embryonic stem cells. *Nat Biotechnol.* 2001;19(12):1129–33. doi: <https://doi.org/10.1038/nbt1201-1129>.
35. Eguizabal C, Aran B, Chuva de Sousa Lopes SM, Geens M, Heindryckx B, Panula S, et al. Two decades of embryonic stem cells: a historical overview. *Hum Reprod Open.* 2019;2019(1):hoy024. doi: <https://doi.org/10.1093/hropen/hoy024>.
36. Reubinoff BE, Itsykson P, Turetsky T, Pera MF, Reinhartz E, Itzik A, et al. Neural progenitors from human embryonic stem cells. *Nat Biotechnol.* 2001;19(12):1134–40. doi: <https://doi.org/10.1038/nbt1201-1134>.
37. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell.* 2006;126(4):663–76. doi: <https://doi.org/10.1016/j.cell.2006.07.024>.
38. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, et al. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell.* 2007;131(5):861–72. doi: <https://doi.org/10.1016/j.cell.2007.11.019>.
39. Tao Y, Zhang SC. Neural Subtype Specification from Human Pluripotent Stem Cells. *Cell Stem Cell.* 2016;19(5):573–86. doi: <https://doi.org/10.1016/j.stem.2016.10.015>.
40. Parr CJC, Yamanaka S, Saito H. An update on stem cell biology and engineering for brain development. *Mol Psychiatry.* 2017;22(6):808–19. doi: <https://doi.org/10.1038/mp.2017.66>.
41. McComish SF, Caldwell MA. Generation of defined neural populations from pluripotent stem cells. *Philos Trans R Soc Lond B Biol Sci.* 2018;373(1750). doi: <https://doi.org/10.1098/rstb.2017.0214>.
42. Harschnitz O, Studer L. Human stem cell models to study host-virus interactions in the central nervous system. *Nat Rev Immunol.* 2021;21(7):441–53. doi: <https://doi.org/10.1038/s41577-020-00474-y>.
43. Yang N, Ng YH, Pang ZP, Südhof TC, Wernig M. Induced neuronal cells: how to make and define a neuron. *Cell Stem Cell.* 2011;9(6):517–25. doi: <https://doi.org/10.1016/j.stem.2011.11.015>.
44. Nicoll RA. A Brief History of Long-Term Potentiation. *Neuron.* 2017;93(2):281–90. doi: <https://doi.org/10.1016/j.neuron.2016.12.015>.

45. Frey U, Morris RG. Synaptic tagging and long-term potentiation. *Nature*. 1997;385(6616):533–6. doi: <https://doi.org/10.1038/385533a0>.
46. Schwartzkroin PA, Wester K. Long-lasting facilitation of a synaptic potential following tetanization in the in vitro hippocampal slice. *Brain Res*. 1975;89(1):107–19. doi: [https://doi.org/10.1016/0006-8993\(75\)90138-9](https://doi.org/10.1016/0006-8993(75)90138-9).
47. Rhee JK, Iwamoto Y, Baker BJ. Visualizing Oscillations in Brain Slices With Genetically Encoded Voltage Indicators. *Front Neuroanat*. 2021;15:741711. doi: <https://doi.org/10.3389/fnana.2021.741711>.
48. Llano DA, Slater BJ, Lesicko AM, Stebbings KA. An auditory colliculothalamocortical brain slice preparation in mouse. *J Neurophysiol*. 2014;111(1):197–207. doi: <https://doi.org/10.1152/jn.00605.2013>.
49. Inagaki HK, Chen S, Ridder MC, Sah P, Li N, Yang Z, et al. A midbrain-thalamus-cortex circuit reorganizes cortical dynamics to initiate movement. *Cell*. 2022;185(6):1065–81.e23. doi: <https://doi.org/10.1016/j.cell.2022.02.006>.
50. Sato H, Fukutani Y, Yamamoto Y, Tataru E, Takemoto M, Shimamura K, et al. Thalamus-derived molecules promote survival and dendritic growth of developing cortical neurons. *J Neurosci*. 2012;32(44):15388–402. doi: <https://doi.org/10.1523/JNEUROSCI.0293-12.2012>.
51. Naggar I, Stewart M, Orman R. High Frequency Oscillations in Rat Hippocampal Slices: Origin, Frequency Characteristics, and Spread. *Front Neurol*. 2020;11:326. doi: <https://doi.org/10.3389/fneur.2020.00326>.
52. Ikeda K, Bekkers JM. Autapses. *Curr Biol*. 2006;16(9):R308. doi: <https://doi.org/10.1016/j.cub.2006.03.085>.
53. Park J, Koito H, Li J, Han A. Microfluidic compartmentalized co-culture platform for CNS axon myelination research. *Biomed Microdevices*. 2009;11(6):1145–53. doi: <https://doi.org/10.1007/s10544-009-9331-7>.
54. Lenoir S, Genoux A, Agasse F, Saudou F, Humbert S. Recreating mouse cortico-hippocampal neuronal circuit in microfluidic devices to study BDNF axonal transport upon glucocorticoid treatment. *STAR Protoc*. 2021;2(1):100382. doi: <https://doi.org/10.1016/j.xpro.2021.100382>.
55. Gladkov A, Pigareva Y, Kutyina D, Kolpakov V, Bukatin A, Mukhina I, et al. Design of Cultured Neuron Networks in vitro with Predefined Connectivity Using Asymmetric Microfluidic Channels. *Sci Rep*. 2017;7(1):15625. doi: <https://doi.org/10.1038/s41598-017-15506-2>.
56. Taylor AM, Menon S, Gupton SL. Passive microfluidic chamber for long-term imaging of axon guidance in response to soluble gradients. *Lab Chip*. 2015;15(13):2781–9. doi: <https://doi.org/10.1039/c5lc00503e>.
57. Hardelauf H, Waide S, Sinaisake J, Jacob P, Hausher V, Schöbel N, et al. Micropatterning neuronal networks. *Analyst*. 2014;139(13):3256–64. doi: <https://doi.org/10.1039/c4an00608a>.
58. Kfoury G, El Habbaki V, Malaeb W, Weaver S, Momotenko D, Mhanna R. Alginate Sulfate Substrates Control Growth Factor Binding and Growth of Primary Neurons: Toward Engineered 3D Neural Networks. *Adv Biosyst*. 2020;4(7):e2000047. doi: <https://doi.org/10.1002/adbi.202000047>.
59. Liu W, Fu W, Sun M, Han K, Hu R, Liu D, et al. Straightforward neuron micropatterning and neuronal network construction on cell-repellent polydimethylsiloxane using microfluidics-guided functionalized Pluronic modification. *Analyst*. 2021;146(2):454–62. doi: <https://doi.org/10.1039/d0an02139c>.
60. Xie J, Liu W, MacEwan MR, Bridgman PC, Xia Y. Neurite outgrowth on electrospun nanofibers with uniaxial alignment: the effects of fiber density, surface coating, and supporting substrate. *ACS Nano*. 2014;8(2):1878–85. doi: <https://doi.org/10.1021/nn406363j>.
61. Froeter P, Huang Y, Cangellaris OV, Huang W, Dent EW, Gillette MU, et al. Toward intelligent synthetic neural circuits: directing and accelerating neuron cell growth by self-rolled-up silicon nitride microtube array. *ACS Nano*. 2014;8(11):11108–17. doi: <https://doi.org/10.1021/nn504876y>.
62. Hong N, Nam Y. Thermoplasmonic neural chip platform for in situ manipulation of neuronal connections in vitro. *Nat Commun*. 2020;11(1):6313. doi: <https://doi.org/10.1038/s41467-020-20060-z>.
63. Kaehr B, Allen R, Javier DJ, Currie J, Shear JB. Guiding neuronal development with in situ microfabrication. *Proc Natl Acad Sci U S A*. 2004;101(46):16104–8. doi: <https://doi.org/10.1073/pnas.0407204101>.
64. Suzuki I, Sugio Y, Jimbo Y, Yasuda K. Stepwise pattern modification of neuronal network in photo-thermally-etched agarose architecture on multi-electrode array chip for individual-cell-based electrophysiological measurement. *Lab Chip*. 2005;5(3):241–7. doi: <https://doi.org/10.1039/b406885h>.
65. Yamamoto H, Okano K, Demura T, Hosokawa Y, Masuhara H, Tani T, et al. guidance of individual neuronal processes by wet femtosecond-laser processing of self-assembled monolayers. *Appl Phys Lett*. 2011;99(16):163701. doi: <https://doi.org/10.1063/1.3651291>.
66. Kim E, Jeon S, An HK, Kianpour M, Yu SW, Kim JY, et al. A magnetically actuated microrobot for targeted neural cell delivery and selective connection of neural networks. *Sci Adv*. 2020;6(39). doi: <https://doi.org/10.1126/sciadv.abb5696>.
67. Yoshida S, Teshima T, Kuribayashi-Shigetomi K, Takeuchi S. Mobile Microplates for Morphological Control and Assembly of Individual Neural Cells. *Adv Healthc Mater*. 2016;5(4):415–20. doi: <https://doi.org/10.1002/adhm.201500782>.
68. Yoshida S, Kato-Negishi M, Takeuchi S. Assembly and Connection of Micropatterned Single Neurons for Neuronal Network Formation. *Micromachines (Basel)*. 2018;9(5). doi: <https://doi.org/10.3390/mi9050235>.
69. Kato-Negishi M, Morimoto Y, Onoe H, Takeuchi S. Millimeter-sized neural building blocks for 3D heterogeneous neural network assembly. *Adv Healthc Mater*. 2013;2(12):1564–70. doi: <https://doi.org/10.1002/adhm.201300052>.
70. Kato-Negishi M, Tsuda Y, Onoe H, Takeuchi S. A neurospheroid network-stamping method for neural transplantation to the brain. *Biomaterials*. 2010;31(34):8939–45. doi: <https://doi.org/10.1016/j.biomaterials.2010.08.008>.
71. Eiraku M, Watanabe K, Matsuo-Takasaki M, Kawada M, Yone-mura S, Matsumura M, et al. Self-organized formation of polarized cortical tissues from ESCs and its active manipulation by extrinsic signals. *Cell Stem Cell*. 2008;3(5):519–32. doi: <https://doi.org/10.1016/j.stem.2008.09.002>.
72. Lancaster MA, Renner M, Martin CA, Wenzel D, Bicknell LS, Hurler ME, et al. Cerebral organoids model human brain development and microcephaly. *Nature*. 2013;501(7467):373–9. doi: <https://doi.org/10.1038/nature12517>.
73. Tanaka Y, Park IH. Regional specification and complementation with non-neuroectodermal cells in human brain organoids. *J Mol Med (Berl)*. 2021;99(4):489–500. doi: <https://doi.org/10.1007/s00109-021-02051-9>.
74. Trujillo CA, Gao R, Negraes PD, Gu J, Buchanan J, Preissl S, et al. Complex Oscillatory Waves Emerging from Cortical Organoids Model Early Human Brain Network Development. *Cell Stem Cell*. 2019;25(4):558–69.e7. doi: <https://doi.org/10.1016/j.stem.2019.08.002>.
75. Gordon A, Yoon SJ, Tran SS, Makinson CD, Park JY, Andersen J, et al. Long-term maturation of human cortical organoids matches key early postnatal transitions. *Nat Neurosci*. 2021;24(3):331–42. doi: <https://doi.org/10.1038/s41593-021-00802-y>.

76. Eichmüller OL, Corsini NS, Vértesy Á, Morassut I, Scholl T, Gruber VE, et al. Amplification of human interneuron progenitors promotes brain tumors and neurological defects. *Science*. 2022;375(6579):eabf5546. doi: <https://doi.org/10.1126/science.abf5546>.
77. Sun AX, Yuan Q, Fukuda M, Yu W, Yan H, Lim GGY, et al. Potassium channel dysfunction in human neuronal models of Angelman syndrome. *Science*. 2019;366(6472):1486–92. doi: <https://doi.org/10.1126/science.aav5386>.
78. Paulsen B, Velasco S, Kedaigle AJ, Pigioli M, Quadrato G, Deo AJ, et al. Autism genes converge on asynchronous development of shared neuron classes. *Nature*. 2022;602(7896):268–73. doi: <https://doi.org/10.1038/s41586-021-04358-6>.
79. Birey F, Li MY, Gordon A, Thete MV, Valencia AM, Revah O, et al. Dissecting the molecular basis of human interneuron migration in forebrain assembloids from Timothy syndrome. *Cell Stem Cell*. 2022;29(2):248–64.e7. doi: <https://doi.org/10.1016/j.stem.2021.11.011>.
80. Samarasinghe RA, Miranda OA, Buth JE, Mitchell S, Ferando I, Watanabe M, et al. Identification of neural oscillations and epileptiform changes in human brain organoids. *Nat Neurosci*. 2021;24(10):1488–500. doi: <https://doi.org/10.1038/s41593-021-00906-5>.
81. Gonzalez C, Armijo E, Bravo-Alegria J, Becerra-Calixto A, Mays CE, Soto C. Modeling amyloid beta and tau pathology in human cerebral organoids. *Mol Psychiatry*. 2018;23(12):2363–74. doi: <https://doi.org/10.1038/s41380-018-0229-8>.
82. Zhao J, Fu Y, Yamazaki Y, Ren Y, Davis MD, Liu CC, et al. APOE4 exacerbates synapse loss and neurodegeneration in Alzheimer's disease patient iPSC-derived cerebral organoids. *Nat Commun*. 2020;11(1):5540. doi: <https://doi.org/10.1038/s41467-020-19264-0>.
83. Kim H, Park HJ, Choi H, Chang Y, Park H, Shin J, et al. Modeling G2019S-LRRK2 Sporadic Parkinson's Disease in 3D Midbrain Organoids. *Stem Cell Reports*. 2019;12(3):518–31. doi: <https://doi.org/10.1016/j.stemcr.2019.01.020>.
84. Zagare A, Barmba K, Smajic S, Smits LM, Grzyb K, Grünwald A, et al. Midbrain organoids mimic early embryonic neurodevelopment and recapitulate LRRK2-p.Gly2019Ser-associated gene expression. *Am J Hum Genet*. 2022;109(2):311–27. doi: <https://doi.org/10.1016/j.ajhg.2021.12.009>.
85. Osaki T, Uzel SGM, Kamm RD. Microphysiological 3D model of amyotrophic lateral sclerosis (ALS) from human iPSC-derived muscle cells and optogenetic motor neurons. *Sci Adv*. 2018;4(10):eaat5847. doi: <https://doi.org/10.1126/sciadv.aat5847>.
86. Kemmerer ZA, Robinson KP, Schmitz JM, Manicki M, Paulson BR, Jochem A, et al. UbiB proteins regulate cellular CoQ distribution in *Saccharomyces cerevisiae*. *Nat Commun*. 2021;12(1):4769. doi: <https://doi.org/10.1038/s41467-021-25084-7>.
87. Ghatak S, Dolatabadi N, Trudler D, Zhang X, Wu Y, Mohata M, et al. Mechanisms of hyperexcitability in Alzheimer's disease hiPSC-derived neurons and cerebral organoids vs isogenic controls. *Elife*. 2019;8. doi: <https://doi.org/10.7554/eLife.50333>.
88. Palop JJ, Chin J, Roberson ED, Wang J, Thwin MT, Bien-Ly N, et al. Aberrant excitatory neuronal activity and compensatory remodeling of inhibitory hippocampal circuits in mouse models of Alzheimer's disease. *Neuron*. 2007;55(5):697–711. doi: <https://doi.org/10.1016/j.neuron.2007.07.025>.
89. Vossel KA, Beagle AJ, Rabinovici GD, Shu H, Lee SE, Naasan G, et al. Seizures and epileptiform activity in the early stages of Alzheimer disease. *JAMA Neurol*. 2013;70(9):1158–66. doi: <https://doi.org/10.1001/jamaneurol.2013.136>.
90. Chen X, Sun G, Tian E, Zhang M, Davtyan H, Beach TG, et al. Modeling Sporadic Alzheimer's Disease in Human Brain Organoids under Serum Exposure. *Adv Sci (Weinh)*. 2021;8(18):e2101462. doi: <https://doi.org/10.1002/advs.202101462>.
91. Szebényi K, Wenger LMD, Sun Y, Dunn AWE, Limegrover CA, Gibbons GM, et al. Human ALS/FTD brain organoid slice cultures display distinct early astrocyte and targetable neuronal pathology. *Nat Neurosci*. 2021;24(11):1542–54. doi: <https://doi.org/10.1038/s41593-021-00923-4>.
92. Foliaki ST, Schwarz B, Groveman BR, Walters RO, Ferreira NC, Orrù CD, et al. Neuronal excitatory-to-inhibitory balance is altered in cerebral organoid models of genetic neurological diseases. *Mol Brain*. 2021;14(1):156. doi: <https://doi.org/10.1186/s13041-021-00864-w>.
93. Modamio J, Saraiva C, Giro GG, Nickels SL, Jarazo J, Antony P, et al. Synaptic decline precedes dopaminergic neuronal loss in human midbrain organoids harboring a triplication of the SNCA gene. *bioRxiv*. 2021:2021.07.15.452499. doi: <https://doi.org/10.1101/2021.07.15.452499>.
94. Harbuzariu A, Pitts S, Cespedes JC, Harp KO, Nti A, Shaw AP, et al. Modelling heme-mediated brain injury associated with cerebral malaria in human brain cortical organoids. *Sci Rep*. 2019;9(1):19162. doi: <https://doi.org/10.1038/s41598-019-55631-8>.
95. Ramirez S, Mukherjee A, Sepulveda S, Becerra-Calixto A, Bravo-Vasquez N, Gherardelli C, et al. Modeling Traumatic Brain Injury in Human Cerebral Organoids. *Cells*. 2021;10(10). doi: <https://doi.org/10.3390/cells10102683>.
96. Krenn V, Bosone C, Burkard TR, Spanier J, Kalinke U, Calistri A, et al. Organoid modeling of Zika and herpes simplex virus 1 infections reveals virus-specific responses leading to microcephaly. *Cell Stem Cell*. 2021. doi: <https://doi.org/10.1016/j.stem.2021.03.004>.
97. Song E, Zhang C, Israelow B, Lu-Culligan A, Prado AV, Skriabine S, et al. Neuroinvasion of SARS-CoV-2 in human and mouse brain. *J Exp Med*. 2021;218(3). doi: <https://doi.org/10.1084/jem.20202135>.
98. Dang J, Tiwari SK, Agrawal K, Hui H, Qin Y, Rana TM. Glial cell diversity and methamphetamine-induced neuroinflammation in human cerebral organoids. *Mol Psychiatry*. 2021;26(4):1194–207. doi: <https://doi.org/10.1038/s41380-020-0676-x>.
99. Lee CT, Chen J, Kindberg AA, Bendriem RM, Spivak CE, Williams MP, et al. CYP3A5 Mediates Effects of Cocaine on Human Neocorticalgenesis: Studies using an In Vitro 3D Self-Organized hPSC Model with a Single Cortex-Like Unit. *Neuropsychopharmacology*. 2017;42(3):774–84. doi: <https://doi.org/10.1038/npp.2016.156>.
100. Sirenko O, Parham F, Dea S, Sodhi N, Biesmans S, Mora-Castilla S, et al. Functional and Mechanistic Neurotoxicity Profiling Using Human iPSC-Derived Neural 3D Cultures. *Toxicol Sci*. 2019;167(1):58–76. doi: <https://doi.org/10.1093/toxsci/kfy218>.
101. Kathuria A, Lopez-Lengowski K, Jagtap SS, McPhie D, Perlis RH, Cohen BM, et al. Transcriptomic Landscape and Functional Characterization of Induced Pluripotent Stem Cell-Derived Cerebral Organoids in Schizophrenia. *JAMA Psychiatry*. 2020;77(7):745–54. doi: <https://doi.org/10.1001/jamapsychiatry.2020.0196>.
102. Kathuria A, Lopez-Lengowski K, Vater M, McPhie D, Cohen BM, Karmacharya R. Transcriptome analysis and functional characterization of cerebral organoids in bipolar disorder. *Genome Med*. 2020;12(1):34. doi: <https://doi.org/10.1186/s13073-020-00733-6>.
103. Srikanth P, Lagomarsino VN, Muratore CR, Ryu SC, He A, Taylor WM, et al. Shared effects of DISC1 disruption and elevated WNT signaling in human cerebral organoids. *Transl Psychiatry*. 2018;8(1):77. doi: <https://doi.org/10.1038/s41398-018-0122-x>.
104. Sawada T, Chater TE, Sasagawa Y, Yoshimura M, Fujimori-Tonou N, Tanaka K, et al. Developmental excitation-inhibition imbalance underlying psychoses revealed by single-cell analyses of discordant twins-derived cerebral organoids. *Mol*

- Psychiatry. 2020;25(11):2695–711. doi: <https://doi.org/10.1038/s41380-020-0844-z>.
105. Stachowiak EK, Benson CA, Narla ST, Dimitri A, Chuye LEB, Dhiman S, et al. Cerebral organoids reveal early cortical maldevelopment in schizophrenia-computational anatomy and genomics, role of FGFR1. *Transl Psychiatry*. 2017;7(11):6. doi: <https://doi.org/10.1038/s41398-017-0054-x>.
 106. Notaras M, Lodhi A, Dündar F, Collier P, Sayles NM, Tilgner H, et al. Schizophrenia is defined by cell-specific neuropathology and multiple neurodevelopmental mechanisms in patient-derived cerebral organoids. *Mol Psychiatry*. 2021. doi: <https://doi.org/10.1038/s41380-021-01316-6>.
 107. Cederquist GY, Ascioia JJ, Tchieu J, Walsh RM, Cornacchia D, Resh MD, et al. Specification of positional identity in forebrain organoids. *Nat Biotechnol*. 2019;37(4):436–44. doi: <https://doi.org/10.1038/s41587-019-0085-3>.
 108. De Santis R, Etoc F, Rosado-Olivieri EA, Brivanlou AH. Self-organization of human dorsal-ventral forebrain structures by light induced SHH. *Nat Commun*. 2021;12(1):6768. doi: <https://doi.org/10.1038/s41467-021-26881-w>.
 109. Misawa R, Minami T, Okamoto A, Ikeuchi Y. A Light-Inducible Hedgehog Signaling Activator Modulates Proliferation and Differentiation of Neural Cells. *ACS Chem Biol*. 2020;15(6):1595–603. doi: <https://doi.org/10.1021/acschembio.0c00195>.
 110. Misawa R, Minami T, Okamoto A, Ikeuchi Y. Light-inducible control of cellular proliferation and differentiation by a Hedgehog signaling inhibitor. *Bioorg Med Chem*. 2021;38:116144. doi: <https://doi.org/10.1016/j.bmc.2021.116144>.
 111. Zheng Y, Xue X, Resto-Irizarry AM, Li Z, Shao Y, Zhao G, et al. Dorsal-ventral patterned neural cyst from human pluripotent stem cells in a neurogenic niche. *Sci Adv*. 2019;5(12):eaax5933. doi: <https://doi.org/10.1126/sciadv.aax5933>.
 112. Bagley JA, Reumann D, Bian S, Lévi-Strauss J, Knoblich JA. Fused cerebral organoids model interactions between brain regions. *Nat Methods*. 2017;14(7):743–51. doi: <https://doi.org/10.1038/nmeth.4304>.
 113. Xiang Y, Tanaka Y, Patterson B, Kang YJ, Govindaiah G, Roselaar N, et al. Fusion of Regionally Specified hPSC-Derived Organoids Models Human Brain Development and Interneuron Migration. *Cell Stem Cell*. 2017;21(3):383–98.e7. doi: <https://doi.org/10.1016/j.stem.2017.07.007>.
 114. Xiang Y, Tanaka Y, Cakir B, Patterson B, Kim K-Y, Sun P, et al. hESC-Derived Thalamic Organoids Form Reciprocal Projections When Fused with Cortical Organoids. *Cell Stem Cell*. 2019;24(3):487–97.e7. doi: <https://doi.org/10.1016/j.stem.2018.12.015>.
 115. Birey F, Andersen J, Makinson CD, Islam S, Wei W, Huber N, et al. Assembly of functionally integrated human forebrain spheroids. *Nature*. 2017;545(7652):54–9. doi: <https://doi.org/10.1038/nature22330>.
 116. Andersen J, Revah O, Miura Y, Thom N, Amin ND, Kelley KW, et al. Generation of Functional Human 3D Cortico-Motor Assembloids. *Cell*. 2020;183(7):1913–29.e26. doi: <https://doi.org/10.1016/j.cell.2020.11.017>.
 117. Watakabe A, Hirokawa J. Cortical networks of the mouse brain elaborate within the gray matter. *Brain Struct Funct*. 2018;223(8):3633–52. doi: <https://doi.org/10.1007/s00429-018-1710-5>.
 118. Kawada J, Kaneda S, Kirihara T, Maroof A, Levi T, Eggan K, et al. Generation of a Motor Nerve Organoid with Human Stem Cell-Derived Neurons. *Stem Cell Reports*. 2017;9(5):1441–9. doi: <https://doi.org/10.1016/j.stemcr.2017.09.021>.
 119. Kirihara T, Luo Z, Chow SYA, Misawa R, Kawada J, Shibata S, et al. A Human Induced Pluripotent Stem Cell-Derived Tissue Model of a Cerebral Tract Connecting Two Cortical Regions. *iScience*. 2019;14:301–11. doi: <https://doi.org/10.1016/j.isci.2019.03.012>.
 120. Cullen DK, Gordián-Vélez WJ, Struzyna LA, Jgamadze D, Lim J, Wofford KL, et al. Bundled Three-Dimensional Human Axon Tracts Derived from Brain Organoids. *iScience*. 2019;21:57–67. doi: <https://doi.org/10.1016/j.isci.2019.10.004>.
 121. Adewole DO, Struzyna LA, Burrell JC, Harris JP, Nemes AD, Petrov D, et al. Development of optically controlled “living electrodes” with long-projecting axon tracts for a synaptic brain-machine interface. *Sci Adv*. 2021;7(4). doi: <https://doi.org/10.1126/sciadv.aay5347>.
 122. Osaki T, Ikeuchi Y. Advanced Complexity and Plasticity of Neural Activity in Reciprocally Connected Human Cerebral Organoids. *bioRxiv*. 2021:2021.02.16.431387. doi: <https://doi.org/10.1101/2021.02.16.431387>.
 123. Syed NI, Bulloch AG, Lukowiak K. In vitro reconstruction of the respiratory central pattern generator of the mollusk *Lymnaea*. *Science*. 1990;250(4978):282–5. doi: <https://doi.org/10.1126/science.2218532>.
 124. Yamamoto H, Moriya S, Ide K, Hayakawa T, Akima H, Sato S, et al. Impact of modular organization on dynamical richness in cortical networks. *Sci Adv*. 2018;4(11):eaau4914. doi: <https://doi.org/10.1126/sciadv.aau4914>.
 125. Buccelli S, Bornat Y, Colombi I, Ambroise M, Martines L, Pasquale V, et al. A Neuromorphic Prosthesis to Restore Communication in Neuronal Networks. *iScience*. 2019;19:402–14. doi: <https://doi.org/10.1016/j.isci.2019.07.046>.
 126. Broccard FD, Joshi S, Wang J, Cauwenberghs G. Neuromorphic neural interfaces: from neurophysiological inspiration to biohybrid coupling with nervous systems. *Journal of Neural Engineering*. 2017;14(4):41002-. doi: <https://doi.org/10.1088/1741-2552/aa67a9>.
 127. Angotzi GN, Boi F, Lecomte A, Miele E, Malerba M, Zucca S, et al. SiNAPS: An implantable active pixel sensor CMOS-probe for simultaneous large-scale neural recordings. *Biosensors and Bioelectronics*. 2019;126:355–64.
 128. Forro C, Caron D, Angotzi GN, Gallo V, Berdondini L, Santoro F, et al. Electrophysiology read-out tools for brain-on-Chip biotechnology. *Micromachines*. 2021;12(2):124.
 129. Mosbacher Y, Khojatee F, Goldin M, Kanner S, Malakay Y, Silva M, et al. Toward neuroprosthetic real-time communication from in silico to biological neuronal network via patterned optogenetic stimulation. *Scientific Reports*. 2020;10(1):7512-. doi: <https://doi.org/10.1038/s41598-020-63934-4>.
 130. Vassanelli S, Mahmud M. Trends and Challenges in Neuroengineering: Toward “Intelligent” Neuroprostheses through Brain-“Brain Inspired Systems” Communication. *Frontiers in Neuroscience*. 2016;10:438-. doi: <https://doi.org/10.3389/fnins.2016.00438>.
 131. Keren H, Partzsch J, Marom S, Mayr CG. A Biohybrid Setup for Coupling Biological and Neuromorphic Neural Networks. *Frontiers in Neuroscience*. 2019;13:432-. doi: <https://doi.org/10.3389/fnins.2019.00432>.
 132. Sharifshazileh M, Burelo K, Sarnthein J, Indiveri G. An electronic neuromorphic system for real-time detection of high frequency oscillations (HFO) in intracranial EEG. *Nature communications*. 2021;12(1):1–14.
 133. Joucla S, Ambroise M, Levi T, Lafon T, Chauvet P, Saighi S, et al. Generation of Locomotor-Like Activity in the Isolated Rat Spinal Cord Using Intraspinal Electrical Microstimulation Driven by a Digital Neuromorphic CPG. *Frontiers in Neuroscience*. 2016;10:67-. doi: <https://doi.org/10.3389/fnins.2016.00067>.
 134. George R, Chiappalone M, Giugliano M, Levi T, Vassanelli S, Partzsch J, et al. Plasticity and Adaptation in Neuromorphic Biohybrid Systems. *Iscience*. 2020:101589-.

135. Christensen DV, Dittmann R, Linares-Barranco B, Sebastian A, Gallo ML, Redaelli A, et al. Roadmap on Neuromorphic Computing and Engineering. arXiv preprint arXiv:210505956. 2021.
136. Yada Y, Yasuda S, Takahashi H. Physical reservoir computing with FORCE learning in a living neuronal culture. *Applied Physics Letters*. 2021;119(17):173701-.
137. Kagan BJ, Kitchen AC, Tran NT, Parker BJ, Bhat A, Rollo B, et al. In vitro neurons learn and exhibit sentience when embodied in a simulated game-world. *bioRxiv*. 2021.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.