



**HAL**  
open science

# Electron ionization mass fragmentation of monounsaturated long-chain alpha,omega-dicarboxylic acid trimethylsilyl derivatives.

Jean-Francois Rontani, Claude Aubert

► **To cite this version:**

Jean-Francois Rontani, Claude Aubert. Electron ionization mass fragmentation of monounsaturated long-chain alpha,omega-dicarboxylic acid trimethylsilyl derivatives.. 2022. hal-03810241v1

**HAL Id: hal-03810241**

**<https://hal.science/hal-03810241v1>**

Preprint submitted on 11 Oct 2022 (v1), last revised 19 Oct 2022 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## Electron ionization mass fragmentation of monounsaturated long-chain $\alpha,\omega$ -dicarboxylic acid trimethylsilyl derivatives

Journal:	<i>Rapid Communications in Mass Spectrometry</i>
Manuscript ID	RCM-22-0068.R1
Wiley - Manuscript type:	Letter to the Editor
Date Submitted by the Author:	28-Apr-2022
Complete List of Authors:	RONTANI, Jean-François; M.I.O., Aix-Marseille University AUBERT, Claude; Laboratoire de Pharmacocinétique et Toxicocinétique (EA 3286)
Keywords:	EIMS, monounsaturated dicarboxylic acids, TMS derivatives, Mass fragmentation, Neutral losses
Abstract:	<p><b>RATIONALE</b> Long-chain monounsaturated <math>\alpha,\omega</math>-dicarboxylic acids are important components of the cuticle and suberized cell walls of plants. Identification and quantification of these compounds, which are widespread in environmental samples, can be challenging.</p> <p><b>METHODS</b> The EI-MS fragmentation pathways of long-chain monounsaturated <math>\alpha,\omega</math>-dicarboxylic acid TMS derivatives were investigated. These pathways were deduced by low-energy CID-GC-EI-MS/MS and accurate mass measurement.</p> <p><b>RESULTS</b> Low-energy CID-MS/MS analyses and accurate mass measurement enabled us to elucidate EI-MS fragmentations of long-chain monounsaturated <math>\alpha,\omega</math>-dicarboxylic acid TMS derivatives. These fragmentations involves successive losses of neutral molecules of TMSOH and CO initiated by transfer of allylic hydrogen atom to the ionized ester groups, which are thus totally lacking in EI mass spectra of saturated homologues.</p> <p><b>CONCLUSIONS</b> EI-MS fragmentations of long-chain monounsaturated <math>\alpha,\omega</math>-dicarboxylic acids TMS derivatives exhibit specific fragment ions that can be very useful for the quantification in MRM mode of trace amounts of these compounds in environmental samples.</p>

SCHOLARONE™  
Manuscripts

## RATIONALE

Long-chain monounsaturated  $\alpha,\omega$ -dicarboxylic acids are important components of the cuticle and suberized cell walls of plants. Identification and quantification of these compounds, which are widespread in environmental samples, can be challenging.

## METHODS

The EI-MS fragmentation pathways of long-chain monounsaturated  $\alpha,\omega$ -dicarboxylic acid TMS derivatives were investigated. These pathways were deduced by low-energy CID-GC-EI-MS/MS and accurate mass measurement.

## RESULTS

Low-energy CID-MS/MS analyses and accurate mass measurement enabled us to elucidate EI-MS fragmentations of long-chain monounsaturated  $\alpha,\omega$ -dicarboxylic acid TMS derivatives. These fragmentations involves successive losses of neutral molecules of TMSOH and CO initiated by transfer of allylic hydrogen atom to the ionized ester groups, which are thus totally lacking in EI mass spectra of saturated homologues.


## CONCLUSIONS

EI-MS fragmentations of long-chain monounsaturated  $\alpha,\omega$ -dicarboxylic acids TMS derivatives exhibit specific fragment ions that can be very useful for the quantification in MRM mode of trace amounts of these compounds in environmental samples.

Dear Editor,


***Electron ionization mass fragmentation of monounsaturated long-chain  $\alpha,\omega$ -dicarboxylic acid trimethylsilyl derivatives***

Electron ionization (EI) mass spectral fragmentation of  $\alpha,\omega$ -dicarboxylic acid trimethylsilyl derivatives was previously investigated.<sup>1,2</sup> The mass spectra of these compounds exhibit intense fragment ions  $\mathbf{a}^+$ ,  $\mathbf{b}^{+\bullet}$  and  $\mathbf{c}^+$  at  $m/z$  217,  $m/z$  204 and  $M^{+\bullet} - 131$  ( $m/z$  327 in the case of octadecanedioic acid TMS derivative) (Fig. 1A). Fragment ions  $\mathbf{a}^+$  and  $\mathbf{b}^{+\bullet}$  are formed via an initial trimethylsilyl transfer between the two ester groups (Scheme 1),<sup>2</sup> which generally occurs through cyclic transition states,<sup>3</sup> and is thus favored in the case of long-chain diesters where intramolecular reaction is nearly strainless.<sup>4</sup> In the other hand, the formation of the fragment ion  $\mathbf{c}^+$  involves the loss of a radical  $\bullet\text{CH}_2\text{-COOTMS}$ .<sup>2</sup>



**FIGURE 1.** EI mass spectra of octadecanedioic acid (A) and octadec-9-enedioic (B) TMS derivatives

Octadec-9-enedioic acid trimethylsilyl derivative exhibits EI mass spectrum (Fig. 1B) quite distinct from this of its saturated counterpart (Fig. 1A), with weaker fragment ions  $\mathbf{a}^+$  and  $\mathbf{b}^{+\bullet}$ , lack of ion  $\mathbf{c}^+$  and the presence of intense additional ions at  $m/z$  366 ( $\mathbf{d}^{+\bullet}$ ),  $m/z$  276 ( $\mathbf{e}^{+\bullet}$ ) and  $m/z$  248 ( $\mathbf{f}^{+\bullet}$ ) corresponding to  $[\mathbf{M} - 90]^{+\bullet}$ ,  $[\mathbf{M} - 180]^{+\bullet}$  and  $[\mathbf{M} - 208]^{+\bullet}$ , respectively.



SCHEME 1

Measurements of accurate masses of ions  $\mathbf{d}^{+\bullet}$ ,  $\mathbf{e}^{+\bullet}$  and  $\mathbf{f}^{+\bullet}$  in TOF mass spectrum of octadec-9-enedioic TMS derivative revealed elemental compositions of  $\text{C}_{21}\text{H}_{38}\text{O}_3\text{Si}$  ( $m/z$  366.2579),  $\text{C}_{18}\text{H}_{28}\text{O}_2$  ( $m/z$  276.2086) and  $\text{C}_{17}\text{H}_{28}\text{O}$  ( $m/z$  248.2133), respectively. Moreover, low energy collision induced dissociation (CID/MS/MS) analyses with precursor ions at  $m/z$  366 ( $\mathbf{d}^{+\bullet}$ ) and  $m/z$  276 ( $\mathbf{e}^{+\bullet}$ ) allowed to demonstrate that ion  $\mathbf{e}^{+\bullet}$  is formed from ion  $\mathbf{d}^{+\bullet}$  and ion  $\mathbf{f}^{+\bullet}$  from ion  $\mathbf{e}^{+\bullet}$ . On the basis of these results, two subsequent losses of a neutral molecule of trimethylsilanol were proposed to explain the formation of ions  $\mathbf{d}^{+\bullet}$  and  $\mathbf{e}^{+\bullet}$  from the molecular ion (Scheme 1). These losses are initiated by the transfer of an allylic hydrogen atom to the ionized OTMS groups and result to the formation of cyclic ketonic structures (Scheme 1). A loss of a neutral molecule of CO, which is a characteristic of six and higher-membered cyclic ketones,<sup>5</sup> was proposed to explain the conversion of ion  $\mathbf{e}^{+\bullet}$  to ion  $\mathbf{f}^{+\bullet}$ . Due to the lack of allylic hydrogen atoms in saturated  $\alpha,\omega$ -dicarboxylic acid trimethylsilyl derivatives, ions  $\mathbf{d}^{+\bullet}$ ,  $\mathbf{e}^{+\bullet}$  and  $\mathbf{f}^{+\bullet}$  are totally lacking in EI mass spectra of these compounds (Fig. 1A). Mass-to-charge ratios and relative abundances of ions  $\mathbf{a}^+$ ,  $\mathbf{b}^{+\bullet}$ ,  $\mathbf{d}^{+\bullet}$ ,  $\mathbf{e}^{+\bullet}$  and  $\mathbf{f}^{+\bullet}$  in EI mass spectra of  $\text{C}_{14}$ - $\text{C}_{21}$  monounsaturated  $\alpha,\omega$ -dicarboxylic acid TMS derivatives are summarized in Table 1.

**Table 1.** Relative abundances of the fragment ions **a-f** in EI mass spectra of homologous monounsaturated  $\alpha,\omega$ -dicarboxylic acid di-TMS derivatives.

Diacids	$\mathbf{a}^+$	$\mathbf{b}^{+\bullet}$	$\mathbf{d}^{+\bullet}$ [M – TMSOH] $^{+\bullet}$	$\mathbf{e}^{+\bullet}$ [M – 2TMSOH] $^{+\bullet}$	$\mathbf{f}^{+\bullet}$ [M – 2TMSOH – CO] $^{+\bullet}$
$\text{C}_{14:1}$	217(5.3) <sup>a</sup>	204(6.4)	310(3.8)	220(6.7)	192(5.1)
$\text{C}_{15:1}$	217(7.0)	204(6.9)	324(3.1)	234(7.3)	206(5.6)
$\text{C}_{16:1}$	217(6.7)	204(7.0)	338(3.8)	248(15.8)	220(4.5)
$\text{C}_{17:1}$	217(6.3)	204(4.7)	352(4.0)	262(12.8)	234(3.6)
$\text{C}_{18:1}$	217(9.0)	204(8.4)	366(6.8)	276(23.7)	248(5.6)
$\text{C}_{19:1}$	217(6.7)	204(6.5)	380(4.6)	290(13.1)	262(3.7)
$\text{C}_{20:1}$	217(7.0)	204(6.9)	394(4.3)	304(11.6)	276(3.8)
$\text{C}_{21:1}$	217(5.8)	204(7.3)	408(4.0)	318(9.3)	290(2.5)

<sup>a</sup>  $m/z$ (relative abundance)

Long chain monounsaturated  $\alpha,\omega$ -dicarboxylic acids are important components of the cuticle and suberized cell walls of plants<sup>6,7</sup>. These compounds play a role in controlling solute and water transport across cell walls and provide a barrier against pathogens<sup>8</sup>. The transitions  $\mathbf{d}^{+\bullet} \rightarrow \mathbf{e}^{+\bullet}$ ,  $\mathbf{d}^{+\bullet} \rightarrow \mathbf{f}^{+\bullet}$  and  $\mathbf{e}^{+\bullet} \rightarrow \mathbf{f}^{+\bullet}$  should be very useful for the characterization in MRM mode of trace amounts of TMS derivatives of these compounds in environmental samples.

## ACKNOWLEDGEMENTS

Financial support from the Centre National de la Recherche Scientifique (CNRS) and the Aix-Marseille University is gratefully acknowledged. Thanks are due to the FEDER OCEANOMED (N° 1166-39417) for the funding of the instruments employed and to an anonymous reviewer for his useful and constructive comments.

*Jean-François Rontani<sup>1\*</sup>*

*Claude Aubert<sup>2</sup>*

<sup>1</sup> *Aix Marseille Univ, Université de Toulon, CNRS, IRD, MIO UM 110, 13288 Marseille, France.*

<sup>2</sup> *Laboratoire de Pharmacocinétique et Toxicocinétique (EA 3286), Faculté de Pharmacie, 13385 Marseille, France.*

\*Correspondence to:

*Jean-François Rontani, Aix Marseille University, Campus de Luminy, Bâtiment 26P, MIO UM 110, 13288 Marseille, France.*

*Email: [jean-francois.rontani@mio.osupytheas.fr](mailto:jean-francois.rontani@mio.osupytheas.fr)*

## REFERENCES

1. Draffan GH, Stillwell RN, McCloskey JA. Electron impact-induced rearrangement of trimethylsilyl groups in long chain compounds. *Org Mass Spectrom* 1968; 1(5): 669-685.
2. Rontani J-F, Aubert C. Trimethylsilyl transfer during electron ionization mass spectral fragmentation of some  $\omega$ -hydroxycarboxylic and  $\omega$ -dicarboxylic acid trimethylsilyl derivatives and the effect of chain length. *Rapid Commun Mass Spectrom* 2004; 18(17): 1889-1895.
3. Harvey DJ, Vouros P. Mass spectrometric fragmentation of trimethylsilyl and related alkylsilyl derivatives. *Mass Spectrom Rev* 2020; 39(1-2): 105-211.

- 1
  - 2
  - 3
  - 4
  - 5
  - 6
  - 7
  - 8
  - 9
  - 10
  - 11
  - 12
  - 13
  - 14
  - 15
  - 16
  - 17
  - 18
  - 19
  - 20
  - 21
  - 22
  - 23
  - 24
  - 25
  - 26
  - 27
  - 28
  - 29
  - 30
  - 31
  - 32
  - 33
  - 34
  - 35
  - 36
  - 37
  - 38
  - 39
  - 40
  - 41
  - 42
  - 43
  - 44
  - 45
  - 46
  - 47
  - 48
  - 49
  - 50
  - 51
  - 52
  - 53
  - 54
  - 55
  - 56
  - 57
  - 58
  - 59
  - 60
4. Casadei MA, Galli C, Mandolini L. Ring-closure reactions. 22. Kinetics of cyclization of diethyl (omega-bromoalkyl) malonates in the range of 4-to 21-membered rings. Role of ring strain. *J Am Chem Soc* 1984; 106(4): 1051-1056.
5. Sparkman OD, Penton Z., Kitson FG. Gas chromatography and mass spectrometry: a practical guide. Academic press. 2011; pp. 345–349.
6. Kolattukudy PE, Kronman K, Poulouse AJ. Determination of structure and composition of suberin from the roots of carrot, parsnip, rutabaga, turnip, red beet, and sweet potato by combined gas-liquid chromatography and mass spectrometry. *Plant Physiol* 1984; 55(3): 567-573.
7. Razeq FM, Kosma DK, França D, Rowland O, Molina I. Extracellular lipids of *Camelina sativa*: Characterization of cutin and suberin reveals typical polyester monomers and unusual dicarboxylic fatty acids. *Phytochemistry* 2021; 184: 112665.
8. Pollard M, Beisson F, Li Y, Ohlrogge JB. Building lipid barriers: biosynthesis of cutin and suberin. *Trends Plant Sci* 2008; 13: 236–246.