



# **A phenotyping procedure for imaging root hair on whole root systems reveal responses to nutrient availability and inoculation with PGPR-like bacteria**

**Ikram Madani, Thanyakorn Rongsawat, Volker Baecker, Anne-Aliénor Véry,  
Hervé Sentenac, Jean-Benoît Peltier**

## **► To cite this version:**

Ikram Madani, Thanyakorn Rongsawat, Volker Baecker, Anne-Aliénor Véry, Hervé Sentenac, et al.. A phenotyping procedure for imaging root hair on whole root systems reveal responses to nutrient availability and inoculation with PGPR-like bacteria. 11th Symposium of the International Society of Root Research, May 2021, the University of Missouri Columbia, Missouri, US, United States. hal-03812295

**HAL Id: hal-03812295**

**<https://hal.science/hal-03812295>**

Submitted on 12 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A phenotyping procedure for imaging root hairs on whole root system reveals responses to nutrient availability and to inoculation with PGPR-like bacteria

I. Madani<sup>1</sup>, T. Rongsawat<sup>1</sup>, V. Baecker<sup>2</sup>, A.A. Véry<sup>1</sup>, H. Sentenac<sup>1</sup>, J.B. Peltier<sup>1</sup>

<sup>1</sup>UMR B&PMP, CNRS-INRAE-Supagro-Univ Montpellier, 2 place Pierre Viala – 34060 Montpellier Cedex 2, France


<sup>2</sup>Montpellier Ressources Imagerie, BioCampus Montpellier, CNRS, INSERM, University of Montpellier

## Introduction

Breeding new cultivars allowing to reduce the ecological costs of conventional agriculture, including fertilization and irrigation, is a major objective of the research in plant sciences [1]. Root hairs at the root-soil interface provide a major contribution to nutrient and water uptake [2]. Different methodologies for root architecture phenotyping have been described in the literature and implemented on different platforms at the international level [3]. However, these methodologies generally do not take into account the development of root hair zones at the root surface, in particular because of the small size of root hairs, which makes them difficult to distinguish


### Our objectives are :

- To develop a methodology for phenotyping root development and elongation of root hairs.
- To investigate the effect of abiotic (water and nutrient availability) and biotic (inoculation with bacterial strains) treatments on wheat (cv. Oued Zenati) root hair development .



## The methodologies developed

### A- Root hair phenotyping system ( Rhizobox-like system)




A: Front view of rhizobox

B: Vertical distribution of rhizobox components

C: The rhizobox system and the irrigation system

The rhizobox is scanned periodically with high resolution scanner


### B- A developed program to analyse root surface area




Analyze Complex Root Tool (ACRT) program<sup>4</sup>

## Wheat root and root hair response to low N with or without inoculation with PGPR-like bacteria

### C- Root area analysed per distance and depth using ACRT program



### D- Root and root hair surface area analysed using winRHIZO™ software



## Results

Inoculation with PGPR bacteria resulted an increase in total root hair surface area

## Conclusion and perspectives

The developed methodology allows for instance to reveal root hair developmental responses to low N inoculated *versus* low N non inoculated conditions. The results obtained indicate that *Bradyrhizobium japonicum* ORS285 strain can interact with the wheat cultivar (Oued Zenati) by increasing the root hair surface area.

Such analyses pave the way to screenings of collections of microorganisms (PGPR) in different abiotic conditions (nutrient availability) and to comparisons between wheat genotypes recapitulating wheat domestication in term of root and root hair responses to abiotic stress and efficient interaction with PGPR bacteria.

### References :

- [1] J.P. Lynch (2007) Roots of the Second Green Revolution. Aust. J. Bot. 55: 493–512
- [2] T. Rongsawat *et al.* (2020) Looking for root hairs to overcome poor soils. Trends Plant Sci. 26: 83-94
- [3] J.A. Atkinson *et al.* (2019) Uncovering the hidden half of plants using new advances in root phenotyping. Curr Opin Biotechnol. 55: 1-8
- [4] V.Baecker, [https://github.com/MontpellierRessourcesImagerie/imagej\\_macros\\_and\\_scripts/wiki/Analyze\\_Complex\\_Roots\\_Tool](https://github.com/MontpellierRessourcesImagerie/imagej_macros_and_scripts/wiki/Analyze_Complex_Roots_Tool)