



HAL
open science

Utility Maximization Problem with Uncertainty and a Jump Setting

Achraf Tamtalini, Anis Matoussi, Sarah Kaakai

► **To cite this version:**

Achraf Tamtalini, Anis Matoussi, Sarah Kaakai. Utility Maximization Problem with Uncertainty and a Jump Setting. 2022. hal-03813812v2

HAL Id: hal-03813812

<https://hal.science/hal-03813812v2>

Preprint submitted on 17 Oct 2022 (v2), last revised 25 Nov 2022 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Utility Maximization Problem with Uncertainty and a Jump Setting*

Achraf Tamtalini [†] Anis Matoussi [‡] Sarah Kaakai [§]

Abstract

We study a robust utility maximization problem in the unbounded case with a general penalty term and information including jumps. We focus on time consistent penalties and we prove that there exists an optimal probability measure solution of the robust problem. Then, we characterize the dynamic value process of our stochastic control problem as the unique solution of a Quadratic-Exponential BSDE.

Keywords: Utility maximization, Robustness, Quadratic BSDEs with jumps, Time-consistent penalties, Bellman Martingale Optimality principle.

Introduction

One of the major problems in asset pricing is the valuation in incomplete markets. In such markets, the decision maker/agent could use the well known utility maximization approach and the literature is particularly rich on the subject (see for example [Rouge and El Karoui \(2000\)](#), [Hu et al. \(2005\)](#), [Morlais \(2009\)](#) and [Carmona \(2008\)](#) among many others). However, in many cases, the decision maker does not know the probability distribution (also called prior or model) governing the stochastic nature of the problem she/he is facing. Thus, before solving the utility maximization problem, the decision maker is faced with an intermediate

*Acknowledgements: The authors research is part of the ANR project DREAMeS (ANR-21-CE46-0002) and benefited from the support of the "Chair Risques Emergents en Assurance" under the aegis of Fondation du Risque, a joint initiative by Le Mans University and Covéa.

[†]Laboratoire Manceau de Mathématiques & FR CNRS N° 2962, Institut du Risque et de l'Assurance, Le Mans University

[‡]Laboratoire Manceau de Mathématiques & FR CNRS N° 2962, Institut du Risque et de l'Assurance, Le Mans University

[§]Laboratoire Manceau de Mathématiques & FR CNRS N° 2962, Institut du Risque et de l'Assurance, Le Mans University

problem of choosing an “optimal” probability. This type of problems are called robust utility maximization problems or utility maximization problems under model uncertainty. In the mathematical finance literature, we can find two types of approaches to solve robust utility maximization problems. The first one relies on convex duality methods which are presented in [Quenez \(2004\)](#), [Gundel \(2005\)](#), [Schied \(2007\)](#) and [Schied and Wu \(2005\)](#). The second one, which we will follow in this article, is based on a stochastic control approach and the powerful tool of BSDEs.

In this article, this uncertainty is captured by considering a set of plausible probability measures that will be penalized through a penalty functional. This penalty functional will measure the distance between any plausible probability Q and the reference/historical one denoted P . In [Anderson et al. \(2003\)](#) and [Hansen et al. \(2006\)](#) for example, a hedging problem was addressed by using the classical entropic penalty under a Markovian setting and hence Hamilton-Jacobi-Bellman (HJB) equations were derived in order to characterize the optimal strategies. The authors in [Skiadas \(2003\)](#) have followed the same point of view and obtained a BSDE that coincides with the one describing a stochastic differential utility (see also [Duffie and Epstein \(1992\)](#) and [Duffie and Skiadas \(1994\)](#) for more about stochastic differential utilities).

More recently, [Bordigoni et al. \(2007\)](#) addressed a robust problem in a more general setting which is non Markovian by using stochastic control techniques. More precisely, they studied the following robust maximization problem:

$$\sup_{\pi} \inf_{Q \in \mathcal{Q}} \mathbb{U}(\pi, Q) \tag{0.1}$$

where π runs through a set of strategies and $Q \in \mathcal{Q}$ through a set of models. The simplest case corresponds to the case where the set \mathcal{Q} is the singleton $\{P\}$ and $\mathbb{U}(\pi, P)$ is simply the P -expected utility from a non bounded terminal wealth and consumption/investment portfolio. The term $\mathbb{U}(\pi, Q)$ is the sum of Q -expected utility and an entropic penalization term. The set \mathcal{Q} is assumed to have certain properties and usually does not need to be specified in any detail. Their work is cast in the case of a continuous filtration and the first minimization problem was solved by proving the existence of a unique optimal probability Q^* . They also characterized the value process of the stochastic control problem as the unique solution of a Quadratic BSDE (QBSDE). In the same spirit, [Faidi et al. \(2013\)](#) studied the same problem using two type of penalties: the first one is the f -divergence penalties in the general framework of a continuous filtration and the second one is time-consistent penalty and studied in the context of a Brownian filtration. For the latter, they also characterized the value process as the unique solution of a QBSDE.

In this paper, we study the first minimization problem in (0.1) in the case of discontinuous filtration (where the information includes jumps) using time consistent penalties. Note that in our framework, the relative entropic penalty, as we will see further, is a special case of the class of time-consistent penalties. We first start by showing that the minimization

problem in (0.1) is well posed and we prove the existence of the optimal probability Q^* using a Komlós-type argument. Second, we prove that the value process of the minimization problem is described by a class of Quadratic-Exponential BSDE with jumps (QEBSDEJ) with unbounded terminal condition. We stress that for a given unbounded terminal condition, the study of Quadratic BSDEs is a difficult problem, see for instance Briand and Hu (2006), Briand and Hu (2008) and Barrieu and El Karoui (2013) in the continuous framework and we emphasize that adding jumps to our optimization problem involves significant difficulties in solving the related BSDEs. Karoui et al. (2016) have obtained existence result for this new class of BSDEs with jumps with unbounded terminal condition. However, they have showed uniqueness only in the bounded case. In this paper, we use the convexity of the generator of our BSDE to show the uniqueness of solution of the BSDE by extending the work of Briand and Hu (2008) in the Brownian setting.

The paper is structured as follows. Section 1 establishes the general framework, in which we assume the existence of a stochastic basis carrying a Brownian motion and a compensated integer-valued random measure that possesses a weak predictable representation property. In section 2, we give a number of estimates for subsequent use. We then prove with the help of Komlós theorem that there exists an optimal probability Q^* . Finally, in section 3, we treat our optimization problem from a stochastic control point of view, and show, thanks to Bellman Optimality Principle, that the corresponding value process is the unique solution of a QEBSDEJ.

1 Framework of the optimization problem

1.1 Setting and notations

This section sets out the notation and the assumptions that will be assumed to hold in the sequel. Let $(\Omega, \mathcal{F}, \mathbb{F}, P)$ be a filtered probability space with a finite time horizon $T < \infty$ and a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$ satisfying the usual conditions of right continuity and completeness, in which all semimartingales are taken to have right continuous paths with left limits. We assume that that \mathcal{F}_0 is trivial and $\mathcal{F} = \mathcal{F}_T$. On this stochastic basis, let W be a d -dimensional standard Brownian motion and let $\mu(dt, dx) = (\mu(w, dt, dx) | w \in \Omega)$ denote an integer-random valued measure on $([0, T] \times E, \mathcal{B}([0, T]) \otimes \mathcal{E})$ with compensator $\nu := \nu^P(w, dt, dx)$ under P , where $E := \mathbb{R}^d \setminus \{0\}$ is equipped with its Borel σ -field $\mathcal{E} := \mathcal{B}(E)$.

On $(\tilde{\Omega}, \tilde{\mathcal{F}}) := (\Omega \times [0, T] \times E, \mathcal{F} \otimes \mathcal{B}[0, T] \otimes \mathcal{E})$, we define the measure $P \times \nu$ by

$$P \times \nu(\tilde{B}) = E \left[\int_{[0, T] \times E} \mathbb{1}_{\tilde{B}}(w, t, x) \nu(w, dt, dx) \right], \quad \tilde{B} \in \tilde{\mathcal{F}}. \quad (1.1)$$

We denote by $\tilde{\mathcal{P}} := \mathcal{P} \otimes \mathcal{E}$ where \mathcal{P} is the predictable σ -field on $\Omega \times [0, T]$. We say that a function on $\tilde{\Omega}$ is *predictable* if it is $\tilde{\mathcal{P}}$ -measurable.

We will assume that the compensator ν is absolutely continuous with respect to the $\lambda \otimes dt$ with a density ξ :

$$\nu(w, dt, dx) = \xi_t(w, x)\lambda(dx)dt, \quad (1.2)$$

where λ is a σ -finite measure on (E, \mathcal{E}) , that satisfies the following condition: $\int_E 1 \wedge |x|^2 \lambda(dx) < \infty$ and the density ξ is $\tilde{\mathcal{P}}$ -measurable, positive and bounded:

$$0 < \xi_t(w, x) \leq C_\nu < \infty, \quad P \times \lambda(dx) \times dt - a.e. \quad \text{for some constant } C_\nu. \quad (1.3)$$

Note that, thanks to (1.2), we have that $\nu(\{t\} \times E) = 0$ for all t , and $\nu([0, T] \times E) \leq C_\nu T \lambda(E)$. For ψ a predictable function on $\tilde{\Omega}$, we define its integral process with respect to μ as:

$$(\psi \star \mu)_t := \begin{cases} \int_{[0, t] \times E} \psi_t(x) \mu(w, ds, dx) & \text{if } \int_{[0, t] \times E} |\psi_t(w, x)| \mu(w, ds, dx) < \infty, \\ + \infty & \text{otherwise.} \end{cases} \quad (1.4)$$

In the same way, we define the integral process with respect to ν .

Let $\tilde{\mu}$ be the compensated measure of μ

$$\tilde{\mu}(w, dt, dx) = \mu(w, dt, dx) - \nu(w, dt, dx). \quad (1.5)$$

To alleviate the notations, we will omit the dependence on w in the different stochastic quantities. In the following, we recall some properties that can be found in [Becherer \(2006\)](#) or [Jacod and Shiryaev \(2013\)](#). First, for any predictable function ψ , the process $\psi \star \nu$ is a predictable process whereas $\psi \star \mu$ is an optional process. We recall that $E[|\psi| \star \mu_T] = E[|\psi| \star \nu_T]$. If $(|\psi|^2 \star \mu)^{1/2}$ is locally integrable, then ψ is integrable with respect to $\tilde{\mu}$ and $\psi \star \tilde{\mu}$ is defined as the purely discontinuous local martingale (under P) with jump process $\int_E U \mu(\{t\}, dx)$. If the increasing process $|\psi| \star \mu$ (or equivalently, $|\psi| \star \nu$) is locally integrable, then again, ψ is integrable with respect to $\tilde{\mu}$ and is the purely discontinuous local martingale as in the first case and we have $\psi \star \tilde{\mu} = \psi \star \mu - \psi \star \nu$. Finally, if the process $|\psi|^2 \star \nu$ is integrable, then U is integrable with respect to $\tilde{\mu}$ and $\tilde{Z} \star \tilde{\mu}$ is a square integrable martingale, purely discontinuous, with predictable quadratic variation $\langle \psi \star \tilde{\mu} \rangle = |\psi|^2 \star \nu$. These properties and their proofs can be found in Section II.1.d of [Jacod and Shiryaev \(2013\)](#).

We will assume that W and $\tilde{\mu}$ satisfy the following weak representation property with respect to P and \mathcal{F} : Every local martingale M with respect to (\mathcal{F}, P) admits the following decomposition:

$$M_t = M_0 + (\eta \cdot W)_t + (\psi \star \tilde{\mu})_t := M_0 + \int_0^t \eta_s dW_s + (\psi \star \tilde{\mu})_t, \quad \forall t \geq 0, \quad P - a.s. \quad (1.6)$$

where η is a progressively measurable process and ψ a predictable process such that

$$\int_0^T |\eta_s|^2 ds < \infty, \quad (|\psi|^2 \star \nu)_T < \infty, \quad P - a.s. \quad (1.7)$$

We introduce the following spaces:

- L^{exp} is the space of all \mathcal{F}_T -measurable random variables X such that:

$$E[\exp(\gamma|X|)] < \infty, \quad \forall \gamma > 0.$$

- $\mathcal{D}_0^{\text{exp}}$ is the space of progressively measurable processes $(X_t)_{0 \leq t \leq T}$ with

$$E \left[\exp \left(\gamma \operatorname{ess\,sup}_{0 \leq t \leq T} |X_t| \right) \right] < \infty, \quad \forall \gamma > 0.$$

- $\mathcal{D}_1^{\text{exp}}$ is the space of progressively measurable processes $(X_t)_{0 \leq t \leq T}$ with

$$E \left[\exp \left(\gamma \int_0^T |X_s| ds \right) \right] < \infty, \quad \forall \gamma > 0.$$

- $\mathcal{H}_\lambda^{2,p}$ is the space of predictable processes ψ such that

$$\left(E \left[\int_0^T |\psi|_{s,\lambda}^2 ds \right]^{\frac{p}{2}} \right)^{\frac{1}{p}},$$

where

$$|\psi|_{s,\lambda}^2 := \int_E |\psi_s(x)|^2 \xi_s(x) \lambda(dx).$$

- $H^{2,p}$ is the set of all predictable processes η such that

$$E \left[\left(\int_0^T |\eta_s|^2 ds \right)^{\frac{p}{2}} \right]^{\frac{1}{p}} < \infty.$$

1.2 The optimization problem

For every probability $Q \ll P$ on \mathcal{F}_T , we denote by $D = (D_t)_{0 \leq t \leq T}$ its Radon-Nikodym density with respect to P , that is,

$$D_t = E \left[\frac{dQ}{dP} \middle| \mathcal{F}_t \right], \quad t \geq 0.$$

D is a càdlàg nonnegative P -martingale. Let $\tau_n := \inf\{t \geq 0, D_t \leq 1/n\}$ and consider the local martingale $M_t^n = \int_0^{t \wedge \tau_n} D_s^{-1} dD_s$. Thanks to the weak representation property, there exist two predictable processes (η_s^n) and (ψ_s^n) , $s \leq \tau_n$, such that, $\int_0^{t \wedge \tau_n} |\eta_s^n|^2 ds < \infty$ and $(|\psi|^2 \star \nu)_{t \wedge \tau_n} < \infty$ and

$$D_{t \wedge \tau_n} = \mathcal{E}((\eta^n \cdot W) + (\psi^n \star \tilde{\mu}))_{t \wedge \tau_n}, \quad t \geq 0, \quad P - a.s.$$

Consistency requires that we should have $\eta_t^n = \eta_t^{n+1}$ $dt \otimes dP$ -a.e and $\psi_t^n(x) = \psi_t^{n+1}(x)$ $\nu(dx) \otimes dt \otimes dP$ -a.e on $\{t \leq \tau_n \wedge T\}$. By the fact that $\tau_n \nearrow \infty$ Q -a.s., we obtain the existence of Q -a.s. defined predictable processes η and ψ such that,

$$D_t = \mathcal{E}((\eta \cdot W)_t + (\psi \star \tilde{\mu})_t), \quad t \geq 0, \quad Q - a.s. \quad (1.8)$$

where $\int_0^T |\eta_s|^2 ds < \infty$ and $(|\psi|^2 \star \nu)_T < \infty$ $Q - a.s.$. Note that since for all $t \in [0, T]$, $D_t > 0$ Q -a.s., then we must have for every $t \in [0, T]$, $\psi_t(x) > -1$ $dQ \times \nu(dx, dt)$ -a.e. and we can rewrite (D_t) as in the following:

$$D_t = \exp\left((\eta \cdot W)_t + (\psi \star \tilde{\mu})_t - \frac{1}{2} \int_0^t |\eta_s|^2 ds + ((\ln(1 + \psi) - \psi) \star \mu)_t\right) \quad Q - a.s. \quad (1.9)$$

We now introduce the following time consistent penalty for a probability $Q \ll P$ on \mathcal{F}_T :

$$\gamma_t(Q) := E_Q \left[\int_t^T r(s, w, \eta_s, \psi_s) ds \middle| \mathcal{F}_t \right], \quad (1.10)$$

where $r : [0, T] \times \Omega \times \mathbb{R}^d \times L^2(E, \lambda) \rightarrow [0, +\infty]$ is suitable measurable function that is convex and lower-semicontinuous in (η, ψ) and such that $r(t, 0, 0) = 0$. Note that, since r is non-negative, r is minimal at $\eta = 0$ and $\psi = 0$ and this corresponds to the probabilistic model P . Therefore, the reference probability has the highest plausibility. In the following, we will consider probabilities $Q \in \mathcal{Q}^f$ where

$$\mathcal{Q}^f = \{Q \ll P, \gamma_0(Q) < \infty\}. \quad (1.11)$$

In order to solve the stochastic control problem with BSDEs, we need to impose some regularity and growth conditions on the penalty function. In a Brownian setting, [Faidi et al. \(2013\)](#) assumed the penalty function to be bounded from below by the relative entropy. In the same way, we will assume that there exists $\tilde{K}_2, \tilde{K}_1 > 0$ such that,

$$\gamma_0(Q) \geq -\tilde{K}_2 + \tilde{K}_1 H(Q|P).$$

Let f be the function defined as follow:

$$f(x) = \begin{cases} (1+x) \log(1+x) - x, & \text{if } x \geq -1; \\ \infty & \text{otherwise;} \end{cases} \quad (1.12)$$

For the latter inequality to be verified, a sufficient condition on r will be the following:

(\mathcal{A}_r) There exists $K_1, K_2 > 0$ such that for all $w \in \Omega$, $t \in [0, T]$, $\eta \in \mathbb{R}^d$ and $\psi \in L^2(\mathcal{E}, \lambda; \mathbb{R})$, we have,

$$r(t, w, \eta, \psi) \geq -K_2 + K_1 \left(\frac{|\eta|^2}{2} + \int_E f(\psi(x)) \xi_t(w, x) \lambda(dx) \right).$$

The following proposition shows that the entropic penalty can be retrieved with a special choice for r . A detailed proof is given in the Appendix 4.

Proposition 1.1. *Let $r(t, \eta, \psi) = \frac{1}{2}|\eta|^2 + \int_E f(\psi(x))\xi_t(x)\lambda(dx)$ and $Q \in \mathcal{Q}_f$. Then, the penalty function corresponds to the relative entropy, that is*

$$\gamma_0(Q) \stackrel{(\Delta)}{=} E_Q \left[\int_0^T \left(\frac{|\eta_t|^2}{2} + \int_E f(\psi_t(x))\xi_t(x)\lambda(dx) \right) dt \right] = H_0(Q|P),$$

where,

$$H_t(Q|P) = E_Q \left[\log \left(\frac{dQ}{dP} \right) \middle| \mathcal{F}_t \right]. \quad (1.13)$$

Moreover, we have for a general r verifying (A_r) ,

$$H(Q|P) \leq \frac{\gamma_0(Q)}{K_1} + \frac{TK_2}{K_1}. \quad (1.14)$$

In particular, $H(Q|P)$ is finite for all $Q \in \mathcal{Q}_f$.

Remark 1.2. *Let $r^*(t, z, \tilde{z}) = \sup_{\eta \in \mathbb{R}^d, \psi \in L^2(\lambda)} (z \cdot \eta + \int_E \tilde{z}(x)\psi(x)\lambda(dx) - r(t, \eta, \psi))$ the Fenchel conjugate r . Assumption (A_r) implies that, for $w \in \Omega$, $z \in \mathbb{R}^d$, $t \in [0, T]$ and $\tilde{z} \in L^2(\mathcal{E}, \lambda; \mathbb{R})$,*

$$r^*(t, w, z, \tilde{z}) \leq K_2 + \frac{|z|^2}{2K_1} + K_1 \int_E f^* \left(\frac{\tilde{z}(x)}{K_1 \xi_t(x)} \right) \xi_t(x)\lambda(dx), \quad (1.15)$$

where $f^*(x) := e^x - x - 1$ is the Fenchel conjugate of the function f .

Now, given a positive adapted process δ , we define the discounting process:

$$S_t^\delta := \exp \left(- \int_0^t \delta_s ds \right), \quad 0 \leq t \leq T,$$

and the auxiliary quantities,

$$\begin{aligned} \mathcal{U}_{t,T}^\delta &:= \alpha \int_t^T \frac{S_s^\delta}{S_t^\delta} U_s ds + \bar{\alpha} \frac{S_T^\delta}{S_t^\delta} \bar{U}_T, \quad 0 \leq t \leq T, \quad \alpha, \bar{\alpha} \geq 0, \\ \mathcal{R}_{t,T}^\delta(Q) &:= \int_t^T \frac{S_s^\delta}{S_t^\delta} r(s, \eta_s, \psi_s) ds, \quad 0 \leq t \leq T. \end{aligned}$$

Now we consider the cost functional

$$c(w, Q) := \mathcal{U}_{0,T}^\delta(w) + \beta \mathcal{R}_{0,T}^\delta(Q)(w), \quad (1.16)$$

which consists of two terms. The first one is a discounted utility term that is the sum of a final utility \bar{U}_T and a continuous utility with utility rate (U_s) . For instance, (U_s) can be seen as the utility coming from investing/consuming and \bar{U}_T as the utility coming from the

terminal wealth. The second term is simply the penalty term and measure the “distance” between the probability Q and the reference probability P . The parameter β might be viewed as measuring the degree of confidence of the reference probability P . The higher β is, the more confident we are in P , with the limiting case $\beta \uparrow \infty$ (respectively $\beta \downarrow \infty$) corresponding to full degree of confidence (respectively distrust).

Our objective is to solve the following optimization problem:

$$\text{Minimize the functional } Q \mapsto \Gamma(Q) := E_Q[c(\cdot, Q)], \quad (1.17)$$

over the set \mathcal{Q}_f . To guarantee the well-posedness of the problem, we will assume the following:

- (\mathcal{A}_u) i. The discounting process is bounded by some constant $\|\delta\|_\infty$;
- ii. The process U belongs to D_1^{exp} ;
- iii. The terminal utility \bar{U}_T belongs to L^{exp} .

Remark 1.3. Under assumption (\mathcal{A}_u), we have

$$E \left[\exp \left(\lambda \int_0^T |U_s| ds + \mu |\bar{U}_T| \right) \right] < \infty, \quad \forall \lambda, \mu \in \mathbb{R}^+. \quad (1.18)$$

Indeed, using the convexity of the exponential function, we get,

$$\begin{aligned} E \left[\exp \left(\lambda \int_0^T |U_s| ds + \mu \bar{U}_T \right) \right] &= E \left[\exp \left(\frac{1}{2} \times 2\lambda \int_0^T |U_s| ds + \frac{1}{2} \times 2\mu \bar{U}_T \right) \right] \\ &\leq E \left[\frac{1}{2} \exp \left(2\lambda \int_0^T |U_s| ds \right) + \frac{1}{2} \exp (2\mu |\bar{U}_T|) \right] \\ &= \frac{1}{2} E \left[\exp \left(2\lambda \int_0^T |U_s| ds \right) \right] + \frac{1}{2} E [\exp (2\mu |\bar{U}_T|)] < \infty, \end{aligned}$$

where the finiteness of the two last expectations is due to assumption (\mathcal{A}_u).

2 Some helpful estimates and existence of optimal probability

2.1 Auxiliary estimates

The main objective of this section is to prove the existence of an optimal probability Q^* that minimizes the functional Γ . To achieve this, we start by proving some useful auxiliary estimates. We will adapt the steps in [Bordigoni et al. \(2007\)](#) and the inequalities therein into our setting.

Proposition 2.1. Under assumption (\mathcal{A}_r) and (\mathcal{A}_u), there exists a constant $C \in (0, \infty)$

which depends only on $\alpha, \bar{\alpha}, \beta, \delta, T, U, \bar{U}_T$ such that

$$\Gamma(Q) \leq E_Q[|c(\cdot, Q)|] \leq C(1 + \gamma_0(Q)), \quad \text{for all } Q \in \mathcal{Q}_f. \quad (2.1)$$

In particular, this shows that $\Gamma(Q)$ is well defined and finite for every $Q \in \mathcal{Q}_f$.

Proof. The first inequality is obvious. As for the second, denoting $\mathbb{U} := \alpha \int_0^T |U_t| dt + \bar{\alpha} |\bar{U}_T|$, we have for $Q \in \mathcal{Q}_f$, using the fact that, $0 \leq S_t^\delta \leq 1$,

$$E_Q[|c(\cdot, Q)|] \leq E_Q \left[\alpha \int_0^T |U_t| dt + \bar{\alpha} |\bar{U}_T| \right] + \beta E_Q \left[\int_0^T r(t, \eta_t, \psi_t) dt \right] \quad (2.2)$$

$$= E_Q[\mathbb{U}] + \beta \gamma_0(Q). \quad (2.3)$$

Fenchel inequality applied to $x \mapsto x \log(x)$, gives

$$xy \leq \frac{1}{\lambda} (x \log(x) + e^{\lambda y - 1}), \quad \forall (x, y, \lambda) \in \mathbb{R}_+^* \times \mathbb{R}_+^* \times \mathbb{R}^*. \quad (2.4)$$

Therefore, using this inequality with $\lambda = 1$, we get,

$$\begin{aligned} E_Q[\mathbb{U}] &= E \left[\frac{dQ}{dP} \mathbb{U} \right] \\ &\leq E \left[\frac{dQ}{dP} \log \left(\frac{dQ}{dP} \right) \right] + E[e^{\mathbb{U}-1}] \\ &= H(Q|P) + E[e^{\mathbb{U}-1}] \\ &\leq \frac{\gamma_0(Q)}{K_1} + \frac{TK_2}{K_1} + e^{-1} E[e^{\mathbb{U}}], \end{aligned}$$

where we used (1.14) in the last inequality. Going back to (2.2), we obtain,

$$E_Q[|c(\cdot, Q)|] \leq \left(\beta + \frac{1}{K_1} \right) \gamma_0(Q) + \left(\frac{TK_2}{K_1} + e^{-1} E[e^{\mathbb{U}}] \right),$$

where the term $E[e^{\mathbb{U}}]$ is finite as pointed out in remark 1.3. We then conclude by setting $C := \max \left(\beta + \frac{1}{K_1}, \frac{TK_2}{K_1} + e^{-1} E[e^{\mathbb{U}}] \right)$. \square

The next result shows that Γ is bounded from below by $\gamma_0(Q)$. This will be very useful for proving the existence of an optimal probability.

Proposition 2.2. *Assume (A_r) and (A_u) hold. Then, there exists $C \in (0, \infty)$ depending on $\alpha, \bar{\alpha}, \beta, \delta, T, U, \bar{U}_T$ such that for all $Q \in \mathcal{Q}_f$*

$$\gamma_0(Q) \leq C(1 + \Gamma(Q)). \quad (2.5)$$

In particular, we have $\inf_{Q \in \mathcal{Q}_f} \Gamma(Q) > -\infty$

Proof. Using the same notation as in the proof of the previous proposition, we have,

$$E_Q[\mathcal{U}_{0,T}^\delta] \geq -E_Q[\mathbb{U}].$$

For every $\lambda \in \mathbb{R}^*$, using the inequality (2.4), we get,

$$\begin{aligned} E_Q[\mathbb{U}] &\leq \frac{1}{\lambda} H(Q|P) + \frac{1}{\lambda} E[e^{\lambda \mathbb{U}-1}] \\ &\leq \frac{1}{\lambda} \left(\frac{\gamma_0(Q)}{K_1} + \frac{TK_2}{K_1} \right) + \frac{e^{-1}}{\lambda} E[e^{\lambda \mathbb{U}}], \end{aligned}$$

where we used (1.14) in the last inequality. On the other hand, since the discounting process is bounded from above, we have

$$E_Q[\mathcal{R}_{0,T}] \stackrel{(\Delta)}{=} E_Q \left[\int_0^T S_t^\delta r(t, \eta_t, \psi_t) dt \right] \geq e^{-\|\delta\|_\infty T} E_Q \left[\int_0^T r(t, \eta_t, \psi_t) dt \right] = e^{-\|\delta\|_\infty T} \gamma_0(Q).$$

Combining the two previous inequalities leads to the following,

$$\Gamma(Q) = E_Q[\mathcal{U}_{0,T}^\delta + \beta \mathcal{R}_{0,T}^\delta] \geq \left(\beta e^{-\|\delta\|_\infty T} - \frac{1}{\lambda K_1} \right) \gamma_0(Q) - \frac{TK_2}{\lambda K_1} - \frac{e^{-1}}{\lambda} E[e^{\lambda \mathbb{U}}]$$

Choosing λ large enough such that $\mu := \beta e^{-\|\delta\|_\infty T} - \frac{1}{\lambda K_1} > 0$, we get the desired result by setting $C := \frac{1}{\mu} \max \left(1, \frac{TK_2}{\lambda K_1} + \frac{e^{-1}}{\lambda} E[e^{\lambda \mathbb{U}}] \right)$. \square

The following is a direct consequence of the previous proposition and inequality (1.14).

Corollary 2.3. *Under assumptions (\mathcal{A}_r) and (\mathcal{A}_u) , there exists $K \in (0, \infty)$ such that for every $Q \in \mathcal{Q}_f$, we have the following*

$$H(Q|P) \leq K(1 + \Gamma(Q)). \quad (2.6)$$

In the same spirit of the proof of the above proposition, we have the following estimate that is crucial in proving the existence of an optimal probability $Q^* \in \mathcal{Q}_f$.

Lemma 2.4. *For any $\lambda > 0$ and any measurable set $A \in \mathcal{F}_T$, we have for every $Q \in \mathcal{Q}_f$*

$$E_Q[|\mathcal{U}_{0,T}^\delta| \mathbb{1}_A] \leq \frac{\gamma_0(Q)}{\lambda K_1} + \frac{TK_2}{\lambda K_1} + \frac{e^{-1}}{\lambda} E \left[\mathbb{1}_A \exp(\lambda \alpha \int_0^T |U_s| ds + \lambda \bar{\alpha} |\bar{U}_T|) \right]. \quad (2.7)$$

Proof. Using inequality (2.4), we have for every $\lambda > 0$ and $Q \in \mathcal{Q}_f$,

$$\frac{dQ}{dP} |\mathcal{U}_{0,T}^\delta| \mathbb{1}_A \leq \frac{1}{\lambda} \left(\frac{dQ}{dP} \log \left(\frac{dQ}{dP} \right) + e^{-1} e^{\lambda \mathbb{U}} \right) \mathbb{1}_A.$$

Taking the expectation under P and using (1.14), we consequently get, $E_Q[|\mathcal{U}_{0,T}^\delta| \mathbb{1}_A] \leq \frac{\gamma_0(Q)}{\lambda K_1} + \frac{TK_2}{\lambda K_1} + \frac{e^{-1}}{\lambda} E[\mathbb{1}_A \exp(\lambda \mathbb{U})]$. \square

2.2 Existence of optimal probability

In this subsection, we prove the existence of an optimal probability $Q^* \in \mathcal{Q}_f$ using a standard Komlós-type argument, but before let us show two important properties of the functionals Γ and γ_0 . We will introduce the following Linderberg condition on sequences of martingales converging almost surely to 0. This technical assumption is needed to prove the lower-semicontinuity of γ_0 :

(\mathcal{A}_L) Every sequence (M_n) of locally square integrable martingales with the representation $dM_t^n = \eta_t^n dW_t + \int_E \psi_t^n(x) \tilde{\mu}(dt, dx)$ converging P -a.s. to 0 for each $t \in [0, T]$, verifies the following Linderberg condition:

$$\forall \epsilon \in (0, 1], \int_0^T \int_E |\psi_t^n(x)|^2 \mathbb{1}_{|\psi_t^n(x)| \geq \epsilon} \nu(dt, dx) \xrightarrow{n \rightarrow \infty} 0, \text{ in } P - \text{Probability}$$

Proposition 2.5. *Under the assumption (\mathcal{A}_L), we have the following:*

1. \mathcal{Q}_f is a convex set and the functional $Q \in \mathcal{Q}_f \mapsto \Gamma(Q)$ is convex.
2. γ_0 is lower-semicontinuous for $L^1(P)$ convergence.

Proof. 1. Let $\lambda \in (0, 1)$, $Q, \tilde{Q} \in \mathcal{Q}_f$ and $Q^\lambda := \lambda Q + (1 - \lambda)\tilde{Q}$. Let D and \tilde{D} denote the corresponding density processes and $(\eta, \psi), (\tilde{\eta}, \tilde{\psi})$ the associated processes via (1.8). Consider the following processes:

$$\begin{aligned} \eta_t^\lambda &:= \frac{\lambda D_{t-} \eta_t + (1 - \lambda) \tilde{D}_{t-} \tilde{\eta}_t}{D_{t-}^\lambda} \mathbb{1}_{D_{t-}^\lambda > 0}, \\ \psi_t^\lambda &:= \frac{\lambda D_{t-} \psi_t + (1 - \lambda) \tilde{D}_{t-} \tilde{\psi}_t}{D_{t-}^\lambda} \mathbb{1}_{D_{t-}^\lambda > 0}, \end{aligned}$$

where $D^\lambda := \lambda D + (1 - \lambda)\tilde{D}$ is the density process of Q^λ with respect to P . It is easy to see that the density D^λ satisfies the following SDE:

$$dD_t^\lambda = D_{t-}^\lambda \left(\eta_t^\lambda dW_t + \int_E \psi_t^\lambda(x) \tilde{\mu}(dt, dx) \right), t \in [0, T], Q^\lambda - a.s.$$

Hence, using the convexity assumption of r , we have,

$$\begin{aligned} \gamma_0(Q^\lambda) &= E_{Q^\lambda} \left[\int_0^T r(t, \eta_t^\lambda, \psi_t^\lambda) dt \right] \\ &\leq E_{Q^\lambda} \left[\int_0^T \left(\frac{\lambda D_{t-}}{D_{t-}^\lambda} r(t, \eta_t, \psi_t) + \frac{(1 - \lambda) \tilde{D}_{t-}}{D_{t-}^\lambda} r(t, \tilde{\eta}_t, \tilde{\psi}_t) \right) \mathbb{1}_{D_{t-}^\lambda > 0} dt \right] \\ &= E \left[D_T^\lambda \int_0^T \left(\frac{\lambda D_{t-}}{D_{t-}^\lambda} r(t, \eta_t, \psi_t) + \frac{(1 - \lambda) \tilde{D}_{t-}}{D_{t-}^\lambda} r(t, \tilde{\eta}_t, \tilde{\psi}_t) \right) \mathbb{1}_{D_{t-}^\lambda > 0} dt \right]. \end{aligned}$$

Using Fubini's Theorem to interchange integral and expectation followed by conditioning on \mathcal{F}_t and the martingale property of the density process D^λ , yields,

$$\gamma_0(Q^\lambda) \leq E \left[\int_0^T D_t^\lambda \left(\frac{\lambda D_{t^-}}{D_{t^-}^\lambda} r(t, \eta_t, \psi_t) + \frac{(1-\lambda) \tilde{D}_{t^-}}{D_{t^-}^\lambda} r(t, \tilde{\eta}_t, \tilde{\psi}_t) \right) \mathbb{1}_{D_{t^-}^\lambda > 0} dt \right].$$

Since D^λ is right continuous, the set $\{t \in [0, T], D_t^\lambda \neq D_{t^-}^\lambda\}$ is countable. Therefore, we have,

$$\begin{aligned} \gamma_0(Q^\lambda) &= E \left[\int_0^T D_{t^-}^\lambda \left(\frac{\lambda D_{t^-}}{D_{t^-}^\lambda} r(t, \eta_t, \psi_t) + \frac{(1-\lambda) \tilde{D}_{t^-}}{D_{t^-}^\lambda} r(t, \tilde{\eta}_t, \tilde{\psi}_t) \right) \mathbb{1}_{D_{t^-}^\lambda > 0} dt \right] \\ &\leq E \left[\int_0^T \lambda D_{t^-} r(t, \eta_t, \psi_t) + (1-\lambda) \tilde{D}_{t^-} r(t, \tilde{\eta}_t, \tilde{\psi}_t) dt \right] \\ &= \lambda \gamma_0(Q) + (1-\lambda) \gamma_0(\tilde{Q}) < \infty. \end{aligned}$$

We have showed then that \mathcal{Q}_f is convex. The convexity of the functional Γ follows readily by using the same arguments used above.

2. Let (Q^n) be a sequence of probability measures that converges to Q in $L^1(\Omega, P)$, i.e., $D_T^n \rightarrow D_T$ in $L^1(\Omega, P)$ where D^n and D the corresponding densities processes. Let (η^n, ψ^n) and (η, ψ) (resp.) be the processes given by (1.8) of D^n and D (resp.). Since we know that D_T^n converges to D_T in $L^1(P)$, the maximal Doob's inequality

$$P\left(\sup_{0 \leq t \leq T} |D_t^n - D_t| \geq \epsilon\right) \leq \frac{1}{\epsilon} E[|D_T^n - D_T|], \quad \forall \epsilon > 0,$$

implies that $(\sup_{0 \leq t \leq T} |D_t^n - D_t|)$ converges to 0 in P -probability. By passing by a subsequence, we can assume that $(\sup_{0 \leq t \leq T} |D_t^n - D_t|)$ converges to 0 $P - a.s.$

We denote $M_t^n := \sup_{0 \leq s \leq t} |D_s^n - D_s|$ and introduce the following stopping time $\tau_n := \inf\{t \in [0, T], M_t^n \geq 1\} \wedge T$. We have $M_{\tau_n}^n \leq M_{\tau_n}^n + |D_{\tau_n}^n - D_{\tau_n}|$ and by taking expectation in the latter we get

$$E[M_{\tau_n}^n] \leq E[M_{\tau_n}^n] + E[|D_{\tau_n}^n - D_{\tau_n}|]. \quad (2.8)$$

Recall that $M_T^n \xrightarrow{n \rightarrow \infty} 0$ and since $(M_t^n)_t$ is a nondecreasing process we have $M_{\tau_n}^n \leq M_T^n$ so that $M_{\tau_n}^n \xrightarrow{n \rightarrow \infty} 0$. We also have by the definition of the stopping time τ_n that $M_{\tau_n}^n \leq 1$. Hence, by the dominated convergence theorem, we obtain that

$$E[M_{\tau_n}^n] \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (2.9)$$

Furthermore,

$$\begin{aligned} E[|D_{\tau_n}^n - D_{\tau_n}|] &= E[|E[D_T^n | \mathcal{F}_{\tau_n}] - E[D_T | \mathcal{F}_{\tau_n}]|] = E[|E[D_T^n - D_T | \mathcal{F}_{\tau_n}]|] \\ &\leq E[E[|D_T^n - D_T| | \mathcal{F}_{\tau_n}]] = E[|D_T^n - D_T|] \xrightarrow{n \rightarrow \infty} 0. \end{aligned} \quad (2.10)$$

Combining (2.8), (2.9) and (2.10), we deduce that $M_{\tau_n}^n$ converges to 0 in $L^1(P)$. Then, by Burkholder-Davis-Gundy's inequality, we get that $[D^n - D]_{\tau_n}^{\frac{1}{2}}$ converges to 0 in $L^1(P)$ and a fortiori in P -probability. Now, as $[D^n - D]_T = [D^n - D]_{\tau_n} \mathbb{1}_{\tau_n=T} + [D^n - D]_T \mathbb{1}_{\tau_n < T}$, then for every $\epsilon > 0$,

$$\begin{aligned} P([D^n - D]_T \geq \epsilon) &\leq P([D^n - D]_{\tau_n} \mathbb{1}_{\tau_n=T} \geq \epsilon) + P([D^n - D]_T \mathbb{1}_{\tau_n < T} \geq \epsilon) \\ &\leq P([D^n - D]_{\tau_n} \geq \epsilon) + P(\tau_n < T), \end{aligned}$$

and

$$P(\tau_n < T) = P(\exists t \in [0, T] \text{ s.t. } M_t^n \geq 1) \leq P(M_T^n \geq 1) \xrightarrow{n \rightarrow \infty} 0.$$

So, we get that $[D^n - D]_T$ converges to 0 in P -probability. On the other hand, since $D_t^n - D_t \rightarrow 0$, thanks to the assumption (\mathcal{A}_L) , we get from Corollary 1 in Shiriyayev (1981) that $\langle D^n - D \rangle_T$ converges to 0 in P -probability and by passing to a subsequence while keeping the same notation, we may say that $\langle D^n - D \rangle_T$ converges to 0 $P - a.s.$. But, we know that,

$$\langle D^n - D \rangle_T = \int_0^T |D_{t-}^n \eta_t^n - D_{t-} \eta_t|^2 dt + \int_0^T \int_E |D_{t-}^n \psi_t^n(x) - D_{t-} \psi_t(x)|^2 \nu(dt, dx).$$

Therefore, we immediately obtain that $D_{t-}^n \eta_t^n \rightarrow D_{t-} \eta_t$ $dP \times dt - a.e.$ and $dP \times dt - a.e.$, $D_{t-}^n \psi_t^n(x) \rightarrow D_{t-} \psi_t(x)$ in $L^2(E, \lambda)$. Next, we will show that $\gamma_0(Q) \leq \liminf_{n \rightarrow \infty} \gamma_0(Q^n)$. Assume by way of contradiction that $\gamma_0(Q) > l := \liminf_{n \rightarrow \infty} \gamma_0(Q^n)$. By passing to a subsequence, we may assume that $\gamma_0(Q^n) \rightarrow l$. Let $\zeta := \inf\{t \in [0, T], D_t = 0\}$ and $\zeta_n := \inf\{t \in [0, T], D_t^n = 0\}$. Since $D_t^n = 0$ on $\{t > \zeta_n\}$, we must have $\zeta \leq \liminf_{n \rightarrow \infty} \zeta_n$. Hence, for $\epsilon := \frac{\gamma_0(Q) - l}{2}$, there is $k \in \mathbb{N}$ such that for $T_k := \zeta \wedge \{\zeta_k, \zeta_{k+1}, \dots\}$, we have

$$\begin{aligned} \gamma_0(Q) &= E_Q \left[\int_0^T r(t, \eta_t, \psi_t) dt \right] = E \left[\int_0^\zeta D_{t-} r(t, \eta_t, \psi_t) dt \right] \\ &\leq E \left[\int_0^{T_k} D_{t-} r(t, \eta_t, \psi_t) dt \right] + \epsilon \\ &= E \left[\int_0^{T_k} g(t, D_{t-}, D_{t-} \eta_t, D_{t-} \psi_t) dt \right] + \epsilon, \end{aligned}$$

where $g(t, x, y, z) := xr(t, \frac{y}{x}, \frac{z}{x})$. Clearly, since r is lower-semicontinuous in (η, ψ) , we

get that also g is also lower-semicontinuous. Hence, by Fatou's lemma, we obtain

$$\begin{aligned} E \left[\int_0^{T_k} g(t, D_{t-}, D_{t-}\eta_t, D_{t-}\psi_t) dt \right] &\leq \liminf_{n \geq k} E \left[\int_0^{T_k} g(t, D_{t-}^n, D_{t-}^n \eta_t, D_{t-}^n \psi_t) dt \right] \\ &\leq \liminf_{n \geq k} E \left[\int_0^{\zeta_k} g(t, D_{t-}^n, D_{t-}^n \eta_t^n, D_{t-}^n \psi_t^n) dt \right] \\ &= \liminf_{n \geq k} \gamma_0(Q^n) = l, \end{aligned}$$

so that we have $\gamma_0(Q) \leq l + \epsilon < \gamma_0(Q)$ which is a contradiction. \square

In the next theorem, we show the existence of an optimal probability $Q^* \in \mathcal{Q}_f$.

Theorem 2.6. *Assume (A_r) , (A_u) and (A_L) hold. Then there exists a probability measure Q^* minimizing $Q \mapsto \Gamma(Q)$ over \mathcal{Q}_f .*

Proof. Let Q^n a minimizing sequence in \mathcal{Q}_f such that

$$\Gamma(Q^n) \searrow \inf_{Q \in \mathcal{Q}_f} \Gamma(Q),$$

and we denote by D^n the corresponding density process. Since we have $D_T^n \geq 0$, it follows from Komlós' lemma that there exists a sequence denoted \bar{D}_T^n with such that $\bar{D}_T^n \in \text{conv}(D_T^n, Z_T^{n+1}, \dots)$ for each $n \in \mathbb{N}$ and (\bar{D}_T^n) converges $P - a.s.$ to a random variable \bar{D}_T^∞ . Now, we will show that \bar{D}_T^∞ is associated with a probability measure \bar{Q}^∞ . First, we have \bar{D}_T^∞ is nonnegative as the $P - a.s.$ limit of the nonnegative sequence $(\bar{D}_T^n)_n$. Second, since \mathcal{Q}_f is convex, each \bar{D}_T^n is associated with a probability measure $\bar{Q}^n \in \mathcal{Q}_f$. Now, thanks to the convexity of Γ and the fact that $(\Gamma(Q^n))_n$ is decreasing, we have the following,

$$\Gamma(\bar{Q}^n) \leq \sup_{m \geq n} \Gamma(Q^m) = \Gamma(Q^n) \leq \Gamma(Q^1). \quad (2.11)$$

Consequently, using (2.6), we get,

$$\sup_{n \in \mathbb{N}} E[\bar{D}_T^n \log(\bar{D}_T^n)] = \sup H(\bar{Q}^n | P) \leq K(1 + \sup_{n \in \mathbb{N}} \Gamma(\bar{Q}^n)) \leq K(1 + \Gamma(Q^1)) < \infty.$$

By Vallée-Poussin's criterion, the sequence (\bar{D}_T^n) is P - uniformly integrable and therefore converges to \bar{D}_T^∞ in $L^1(P)$. Hence, we have, $E[\bar{D}_T^\infty] = \lim_{n \rightarrow \infty} E[\bar{D}_T^n] = 1$ since $E[\bar{D}_T^n] = 1$ for all $n \in \mathbb{N}$. This shows that \bar{D}_T^∞ can be associated with a probability \bar{Q}^∞ on \mathcal{F}_T such that $d\bar{Q}^\infty = \bar{D}_T^\infty dP$. Our next step is to prove that this probability \bar{Q}^∞ belongs to \mathcal{Q}_f . By Proposition 2.5, we know that γ_0 is lower-semicontinuous with respect to $L^1(\Omega, P)$ convergence. Therefore, we get since $\bar{D}_T^n \xrightarrow{L^1} \bar{D}_T^\infty$,

$$\gamma_0(\bar{Q}^\infty) \leq \liminf_{n \rightarrow \infty} \gamma_0(\bar{Q}^n).$$

But, thanks to (2.5), we know that $\gamma_0(Q) \leq C(1 + \Gamma(Q))$. Consequently, we obtain that,

$$\liminf_{n \rightarrow \infty} \gamma_0(\bar{Q}^n) \leq C(1 + \sup_{n \in \mathbb{N}} \Gamma(\bar{Q}^n)).$$

The RHS of the last inequality is finite thanks to (2.11). We then conclude that $\gamma_0(\bar{Q}^\infty) < \infty$, i.e., $\bar{Q}^\infty \in \mathcal{Q}_f$. It remains to show that \bar{Q}^∞ is optimal. Note that using the same arguments in the proof of Proposition 2.5, the function $Q \mapsto E_Q[\mathcal{R}_{0,T}^\delta(Q)] = E_Q[\int_0^T S_t^\delta r(t, \eta_t, \psi_t) dt]$ is lower-semicontinuous for $L^1(\Omega, P)$ convergence and therefore we get immediately that,

$$E_{\bar{Q}^\infty}[\mathcal{R}_{0,T}^\delta(\bar{Q}^\infty)] \leq \liminf_{n \rightarrow \infty} E_{\bar{Q}^n}[\mathcal{R}_{0,T}^\delta(\bar{Q}^n)].$$

We denote $\bar{Y}^n := \bar{D}_T^n \mathcal{U}_{0,T}^\delta$ and $\bar{Y}^\infty := \bar{D}_T^\infty \mathcal{U}_{0,T}^\delta$. If we prove that we also have

$$E[\bar{Y}^\infty] \leq \liminf_{n \rightarrow \infty} E[\bar{Y}^n], \tag{2.12}$$

then we would have

$$\begin{aligned} \Gamma(\bar{Q}^\infty) &= E[\bar{Y}^\infty] + E_{\bar{Q}^\infty}[\mathcal{R}_{0,T}^\delta(\bar{Q}^\infty)] \\ &\leq \liminf_{n \rightarrow \infty} E[\bar{Y}^n] + \liminf_{n \rightarrow \infty} E_{\bar{Q}^n}[\mathcal{R}_{0,T}^\delta(\bar{Q}^n)] \\ &\leq \liminf_{n \rightarrow \infty} E[\bar{Y}^n] + E_{\bar{Q}^n}[\mathcal{R}_{0,T}^\delta(\bar{Q}^n)] \\ &= \liminf_{n \rightarrow \infty} \Gamma(\bar{Q}^n) = \inf_{Q \in \mathcal{Q}_f} \Gamma(Q), \end{aligned}$$

which proves that indeed \bar{Q}^∞ is optimal. Although \bar{Y} is linear in \bar{D}_T^n , we cannot use Fatou's lemma since the term $\mathcal{U}_{0,T}^\delta$ has no lower bound. To remediate this, we introduce the following:

$$\tilde{R}_m := \mathcal{U}_{0,T}^\delta \mathbb{1}_{\mathcal{U}_{0,T}^\delta \geq -m} \geq -m, \quad m \in \mathbb{N}.$$

Hence, we have for $n \in \mathbb{N} \cup \{\infty\}$,

$$\bar{Y}^n = \bar{D}_T^n \mathcal{U}_{0,T}^\delta = \bar{D}_T^n \tilde{R}_m + \bar{D}_T^n \mathcal{U}_{0,T}^\delta \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}.$$

Because now \tilde{R}_m is bounded below by $-m$, we can use Fatou's lemma to get,

$$E[\bar{D}^\infty \tilde{R}_m] \leq \liminf_{n \rightarrow \infty} E[\bar{D}^n \tilde{R}_m].$$

Consequently, by adding and subtracting $E[\bar{D}_T^n \mathcal{U}_{0,T}^\delta \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}]$, we obtain,

$$\begin{aligned} E[\bar{Y}^\infty] &\leq \liminf_{n \rightarrow \infty} E[\bar{D}^n \tilde{R}_m] + E[\bar{D}_T^\infty \mathcal{U}_{0,T}^\delta \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}] \\ &\leq \liminf_{n \rightarrow \infty} E[\bar{Y}^n] + 2 \sup_{n \in \mathbb{N} \cup \{\infty\}} E[\bar{D}_T^n |\mathcal{U}_{0,T}^\delta| \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}]. \end{aligned}$$

The desired inequality (2.12) will follow once we prove that

$$\lim_{m \rightarrow \infty} \sup_{n \in N \cup \{\infty\}} E[\mathcal{D}_T^n | \mathcal{U}_{0,T}^\delta | \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}] = 0,$$

and this is where we use Lemma 2.4. Indeed, thanks to this lemma, we have,

$$\begin{aligned} E[\mathcal{D}_T^n | \mathcal{U}_{0,T}^\delta | \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}] &= E_{\overline{Q}^n} [| \mathcal{U}_{0,T}^\delta | \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}] \leq \frac{\gamma_0(\overline{Q}^n)}{\lambda K_1} + \frac{TK_2}{\lambda K_1} + \frac{e^{-1}}{\lambda} E[\exp(\lambda \mathbb{U}) \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}] \\ &\leq \frac{C(1 + \Gamma(\overline{Q}^n))}{\lambda K_1} + \frac{TK_2}{\lambda K_1} + \frac{e^{-1}}{\lambda} E[\exp(\lambda \mathbb{U}) \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}]. \end{aligned}$$

Using (2.11), we deduce that

$$E[\mathcal{D}_T^n | \mathcal{U}_{0,T}^\delta | \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}] \leq \frac{C(1 + \max(\Gamma(Q^1), \Gamma(\overline{Q}^\infty)))}{\lambda K_1} + \frac{TK_2}{\lambda K_1} + \frac{e^{-1}}{\lambda} E[\exp(\lambda \mathbb{U}) \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}].$$

By the dominated convergence theorem, the third term in the RHS of the previous inequality goes to 0 as $m \rightarrow \infty$. Hence, we for all $\lambda > 0$, we have

$$\lim_{m \rightarrow \infty} E[\mathcal{D}_T^n | \mathcal{U}_{0,T}^\delta | \mathbb{1}_{\mathcal{U}_{0,T}^\delta < -m}] \leq \frac{C(1 + \max(\Gamma(Q^1), \Gamma(\overline{Q}^\infty)))}{\lambda K_1} + \frac{TK_2}{\lambda K_1}.$$

Sending λ to ∞ , we finally obtain the desired result. \square

3 Related BSDE with jumps

This section is devoted to the study of the dynamic value process V associated to the optimization problem (1.17) using stochastic control techniques. More precisely, we prove that the dynamic process is the unique solution of a certain QEBSDEJ. This extends the previous work by Schroder and Skiadas (1999), Skiadas (2003) and Lazrak and Quenez (2003).

We first introduce some notations that we will use below. Let \mathcal{S} denote the set of all stopping time τ with values in $[0, T]$ and \mathcal{D} the space of all density processes D^Q with $Q \in \mathcal{Q}_f$. We also define,

$$\begin{aligned} \mathcal{D}(Q, \tau) &:= \{Q' \in \mathcal{Q}_f, Q' = Q \text{ on } \mathcal{F}_\tau\}, \\ \Gamma(Q, \tau) &:= E_Q[c(\cdot, Q) | \mathcal{F}_\tau]. \end{aligned}$$

As in Karoui (1981), we define the minimal conditional cost at time τ by

$$J(Q, \tau) := Q - \operatorname{ess\,inf}_{Q' \in \mathcal{D}(Q, \tau)} \Gamma(Q', \tau).$$

For $Q \in \mathcal{Q}_f$ and $\tau \in \mathcal{S}$, we now define the value of the control problem starting at time τ instead of 0 and assuming one has used the model Q up to time τ ,

$$\begin{aligned}\tilde{V}(Q', \tau) &:= E_{Q'}[\mathcal{U}_{\tau, T}^\delta | \mathcal{F}_\tau] + \beta E_{Q'}[\mathcal{R}_{\tau, T}^\delta(Q') | \mathcal{F}_\tau], \\ V(Q, \tau) &:= Q - \operatorname{ess\,inf}_{Q' \in \mathcal{D}(Q, \tau)} \tilde{V}(Q', \tau).\end{aligned}$$

The following martingale optimality principle is a consequence of Theorems 1.15, 1.17 and 1.21 in Karoui (1981). It is the analogue of Proposition 3.4 in Faidi et al. (2013) in a Brownian setting but the proofs also hold in our setting with obvious modifications.

Proposition 3.1. *Under (\mathcal{A}_u) and (\mathcal{A}_r) , we have:*

- The family $\{J(Q, \tau) | \tau \in \mathcal{S}, Q \in \mathcal{Q}_f\}$ is a submartingale system, that is for any $Q \in \mathcal{Q}_f$ and stopping times $\sigma \leq \tau$, we have,

$$E_Q[J(Q, \tau) | \mathcal{F}_\sigma] \geq J(Q, \sigma) \quad Q - a.s. \quad (3.1)$$

- $\hat{Q} \in \mathcal{Q}_f$ is optimal if and only if the family $\{J(\hat{Q}, \tau) | \tau \in \mathcal{S}\}$ is a \hat{Q} -martingale system which means that for any stopping times $\sigma \leq \tau$

$$E_{\hat{Q}}[J(\hat{Q}, \tau) | \mathcal{F}_\sigma] = J(\hat{Q}, \sigma) \quad \hat{Q} - a.s.$$

- For each $Q \in \mathcal{Q}_f$, there exists an adapted RCLL process $J^Q = (J_t^Q)_{t \in [0, T]}$ which is a right closed Q -submartingale such that for every stopping time τ

$$J_\tau^Q = J(Q, \tau) \quad Q - a.s.$$

Before stating the BSDE verified by the value process V , we will need to define a strong order relation on the set of increasing processes defined below.

Definition 3.2. *Let X and Y two increasing processes. We say that $X \preceq Y$ if the process $Y - X$ is increasing.*

Theorem 3.3. *Assume assumptions (\mathcal{A}_r) , (\mathcal{A}_u) and (\mathcal{A}_L) hold. If the optimal probability \bar{Q}^∞ in Theorem 2.6 is equivalent to P , then there exists Z and \tilde{Z} such that (V, Z, \tilde{Z}) is solution in $D_0^{\text{exp}} \times \mathcal{H}^{2,p} \times \mathcal{H}_\lambda^{2,p}$ of the following BSDE:*

$$\begin{cases} dV_t = \left(\delta_t V_t - \alpha U_t + \beta r^* \left(t, \frac{Z_t}{\beta}, \xi_t \frac{\tilde{Z}_t}{\beta} \right) \right) dt - Z_t dW_t - \int_E \tilde{Z}_t(x) \tilde{\mu}(dx, dt), \\ V_T = \bar{\alpha} \bar{U}_T. \end{cases} \quad (3.2)$$

Proof. We will split the proof into three steps: First, we will prove that the value process V is a P -special martingale, that is it can be decomposed as $V = V_0 + M^V + A^V$, where M^V

is a local martingale that can be written as $M^V = (Z \cdot W) + (\tilde{Z} \star \tilde{\mu})$ and A^V a predictable finite variation process. Then, we show that (V, Z, \tilde{Z}) is a solution of the BSDE. Finally, we prove that (V, Z, \tilde{Z}) is in the required spaces.

Step 1: First, note that since we assumed that $\bar{Q}^\infty \sim P$, then,

$$\inf_{Q \in \mathcal{Q}_f} \Gamma(Q) = \inf_{Q \in \mathcal{Q}_f^e} \Gamma(Q),$$

where $\mathcal{Q}_f^e := \{Q \in \mathcal{Q}_f, Q \sim P\}$ and we define $\mathcal{D}^e(Q, \tau)$ accordingly. Hence, we will restrict our attention to probabilities $Q \in \mathcal{Q}_f^e$ and all essential infimums can be taken with respect to P in the expression of $V(Q, \tau)$ and $J(Q, \tau)$, i.e.,

$$\begin{aligned} J(Q, \tau) &= P - \operatorname{ess\,inf}_{Q' \in \mathcal{D}^e(Q, \tau)} \Gamma(Q', \tau), \\ V(Q, \tau) &= P - \operatorname{ess\,inf}_{Q' \in \mathcal{D}^e(Q, \tau)} \tilde{V}(Q', \tau). \end{aligned}$$

By Bayes' formula and the definition of $\mathcal{R}_{\tau, T}(Q')$, it is easy to see that $\tilde{V}(Q', \tau)$ depends only on the values of the density process D' of Q' on $[\tau, T]$ and is therefore independent of Q . Hence, we can denote $V(Q, \tau)$ by $V(\tau)$. From the definition of $\mathcal{R}_{t, T}^\delta(Q')$ and $\mathcal{U}_{t, T}^\delta$, we have

$$\begin{aligned} \mathcal{R}_{0, T}^\delta(Q') &= \int_0^\tau S_t^\delta r(t, q'_t, \psi'_t) dt + S_\tau^\delta \mathcal{R}_{\tau, T}^\delta(Q'), \\ \mathcal{U}_{0, T}^\delta &= \alpha \int_0^\tau S_t U_t dt + \mathcal{U}_{\tau, T}^\delta. \end{aligned}$$

Comparing $V(\tau)$ and $J(Q, \tau)$ yields for $Q \in \mathcal{Q}_f^e$ with density process $D^Q = \mathcal{E}((\eta \cdot W) + (\psi \star \tilde{\mu}))$,

$$J(Q, \tau) = S_\tau^\delta V(\tau) + \alpha \int_0^\tau S_t^\delta U_t dt + \beta \int_0^\tau S_t^\delta r(t, \eta_t, \psi_t) dt, \quad P - a.s. \quad (3.3)$$

From the martingale optimality principle in Proposition 3.1, there exists an adapted RCLL process denoted $J^Q = (J_t^Q)_{t \in [0, T]}$ such that $J_\tau^Q = J(Q, \tau)$, $Q - a.s.$ From (3.3), we deduce that we can choose an adapted RCLL process $(V_t)_{t \in [0, T]}$ such that $V_\tau = V(\tau)$, $P - a.s.$ for all $\tau \in \mathcal{S}$. We can then rewrite (3.3) as for every $Q \in \mathcal{Q}_f^e$,

$$J_t^Q = S_t^\delta V_t + \alpha \int_0^t S_s^\delta U_s ds + \beta \int_0^t S_s^\delta r(s, \eta_s, \psi_s) dt, \quad dt \times dP - a.e. \quad (3.4)$$

As the probability $P \in \mathcal{Q}_f^e$ corresponds to $\eta = 0$ and $\psi = 0$ and $r(t, 0, 0) = 0$, we get in particular for $Q = P$ in (3.4) that $J^P = S^\delta V + \alpha \int_0^\cdot S_s^\delta U_s ds$. By Proposition 3.1, J^P is a P -submartingale and thus we deduce that V is a P -special semimartingale, i.e. its canonical decomposition can be written as

$$V = V_0 + M^V + A^V, \quad (3.5)$$

where M^V is a local martingale and A^V is a predictable finite variation process. By the weak representation assumption, the local martingale M^V can be written as:

$$M^V = -(Z \cdot W) - (\tilde{Z} \star \tilde{\mu}).$$

Step 2: We now prove that (V, Z, \tilde{Z}) is a solution of QEBSDEJ in (3.2). Plugging (3.5) into (3.4) yields

$$dJ_t^Q = -\delta_t S_t^\delta V_t dt + \alpha S_t U_t dt + S_t^\delta \left(\beta r(t, \eta_t, \psi_t) dt - Z_t dW_t - \int_E \tilde{Z}_t(x) \tilde{\mu}(dx, dt) + dA_t^V \right).$$

For each $Q \in \mathcal{Q}_f^e$, we have, $D^Q = \mathcal{E}((\eta \cdot W) + (\psi \star \tilde{\mu}))$, P -a.s. and by Girsanov's theorem, we have, $dW_t^Q = dW_t - \eta_t dt$ is a Q Brownian motion and $\nu^Q(dx, dt) = (1 + \psi_t(x))\nu(dt, dx)$ is the compensation of μ under Q . Rewriting the dynamic of J^Q , we obtain,

$$dJ_t^Q = -\delta_t S_t^\delta V_t dt + \alpha S_t U_t dt + S_t^\delta \left(\beta r(t, \eta_t, \psi_t) dt - Z_t \eta_t dt - \int_E \tilde{Z}_t(x) \psi_t(x) \nu(dx, dt) + dA_t^V \right) - S_t^\delta \left(Z_t dW_t^Q + \int_E \tilde{Z}_t(x) \tilde{\mu}^Q(dx, dt) \right). \quad (3.6)$$

But, we know thanks to Proposition 3.1, that for every $Q \in \mathcal{Q}_f^e$, J^Q is a Q -submartingale and $J^{\bar{Q}^\infty}$ is a \bar{Q}^∞ -martingale. This means that we should have,

$$dA_t^V \geq Z_t \eta_t dt + \int_E \tilde{Z}_t(x) \psi_t(x) \nu(dx, dt) - \beta r(t, \eta_t, \psi_t) dt + \delta_t V_t dt - \alpha S_t^\delta U_t dt, \quad dt \times dQ - a.e.$$

$$dA_t^V = Z_t \bar{\eta}_t^\infty dt + \int_E \tilde{Z}_t(x) \bar{\psi}_t^\infty(x) \nu(dx, dt) - \beta r(t, \bar{\eta}_t^\infty, \bar{\psi}_t^\infty) + \delta_t V_t dt - \alpha S_t^\delta U_t dt, \quad dt \times d\bar{Q}^\infty - a.e.$$

Note that the above inequality and equality are verified $dt \times dP - a.e.$ since $Q \in \mathcal{Q}_f^e$ and by the assumption that $\bar{Q}^\infty \in \mathcal{Q}_f^e$, in which case they become equivalent to,

$$dA_t^V \geq \operatorname{ess\,sup}_{\eta_t \in \mathbb{R}^d, \psi_t \in L^2(\lambda)} \left(Z_t \eta_t dt + \int_E \tilde{Z}_t(x) \psi_t(x) \nu(dx, dt) - \beta r(t, \eta_t, \psi_t) dt \right) + \delta_t V_t dt - \alpha S_t^\delta U_t dt, \quad dt \times dP - a.e. \quad (3.7)$$

$$dA_t^V = Z_t \bar{\eta}_t^\infty dt + \int_E \tilde{Z}_t(x) \bar{\psi}_t^\infty(x) \nu(dx, dt) - \beta r(t, \bar{\eta}_t^\infty, \bar{\psi}_t^\infty) + \delta_t V_t dt - \alpha S_t^\delta U_t dt, \quad dt \times dP - a.e. \quad (3.8)$$

By denoting

$$r^*(t, z, \tilde{z}) = \sup_{\eta \in \mathbb{R}^d, \psi \in L^2(\lambda)} (z \cdot \eta + \int_E z(x) \psi(x) \lambda(dx) - r(t, \eta, \psi)),$$

the Fenchel conjugate of r , equation (3.7) implies that $dt \times dP - a.e.$,

$$\begin{aligned}
 dA_t^V &= \operatorname{ess\,sup}_{\eta_t \in \mathbb{R}^d, \psi_t \in L^2(\lambda)} \left(Z_t \eta_t dt + \int_E \tilde{Z}_t(x) \psi_t(x) \nu(dx, dt) - \beta r(t, \eta_t, \psi_t) dt \right) + \delta_t V_t dt - \alpha S_t^\delta U_t dt \\
 &= \beta r^*(t, \frac{1}{\beta} Z_t, \frac{1}{\beta} \xi_t \tilde{Z}_t) + \delta_t V_t dt - \alpha S_t^\delta U_t dt \\
 &= Z_t \bar{\eta}_t^\infty dt + \int_E \tilde{Z}_t(x) \bar{\psi}_t^\infty(x) \nu(dx, dt) - \beta r(t, \bar{\eta}_t^\infty, \bar{\psi}_t^\infty) + \delta_t V_t dt - \alpha S_t^\delta U_t dt.
 \end{aligned} \tag{3.9}$$

This shows in particular that

$$\left(\frac{Z_t}{\beta}, \frac{\tilde{Z}_t}{\beta} \xi_t \right) \in \partial r(t, \bar{\eta}_t^\infty, \bar{\psi}_t^\infty), \quad dt \times dP - a.e. \tag{3.10}$$

Going back to equation (3.5) and replacing the finite variation process A^V by its expression in (3.9), it follows that (V, Z, \tilde{Z}) is solution of the following equation,

$$\begin{cases} dV_t = \left(\delta_t V_t - \alpha U_t + \beta r^* \left(t, \frac{Z_t}{\beta}, \xi_t \frac{\tilde{Z}_t}{\beta} \right) \right) dt - Z_t dW_t - \int_E \tilde{Z}_t(x) \tilde{\mu}(dx, dt), \\ V_T = \bar{\alpha} \bar{U}_T. \end{cases}$$

Step 3: In this step, we show that the $(V, Z, \tilde{Z}) \in D_0^{\text{exp}} \times \mathcal{H}_\lambda^{2,p} \times \mathcal{H}^{2,2}$. $V \in D_0^{\text{exp}}$ follows as in Faidi et al. (2013). As for Z and \tilde{Z} , the proof will lean on some exponential transform. We introduce the following processes defined for $t \in [0, T]$ as:

$$\begin{aligned}
 Y_t^- &= -CV_t + C \int_0^t (\alpha |U_s| + K_2 \beta) ds + C \int_0^t \delta_s |V_s| ds, \\
 Y_t^+ &= CV_t + C \int_0^t (\alpha |U_s| + K_2 \beta) ds + C \int_0^t \delta_s |V_s| ds, \\
 K_t^- &= \exp(Y_t^-), \quad K_t^+ = \exp(Y_t^+),
 \end{aligned}$$

where $C = \frac{1}{K_1 \beta}$. For any $p \geq 1$, we have

$$\begin{aligned}
 \sup_{t \in [0, T]} (K_t^\pm)^p &= \sup_{t \in [0, T]} \exp(pY_t^\pm) \leq \sup_{t \in [0, T]} \exp(pC|V_t| + pC \int_0^t (\alpha |U_s| + K_2 \beta) ds + pC \int_0^t \delta_s |V_s| ds) \\
 &\leq \exp(pC \sup_{t \in [0, T]} |V_t| + pC \alpha \int_0^T |U_s| ds + pC K_2 \beta T + pC \|\delta\|_\infty T \sup_{t \in [0, T]} |V_t|).
 \end{aligned}$$

Since $V \in D_0^{\text{exp}}$ and $U \in D_1^{\text{exp}}$, from the above inequality we deduce that $\sup_{t \in [0, T]} K_t^\pm \in L^p(\Omega)$.

We turn our attention to the process Y^- . Using (3.2), the process Y^- verifies:

$$\begin{aligned}
 dY_t^- &= -CdV_t + C(\alpha|U_t| + K_2\beta)dt + C\delta_t|V_t|dt \\
 &= C \left(\delta_t(|V_t| - V_t) + \alpha(|U_t| + U_t) + K_2\beta - \beta r^* \left(t, \frac{Z_t}{\beta}, \xi_t \frac{\tilde{Z}_t}{\beta} \right) \right) dt + CZ_t dW_t + \int_E C\tilde{Z}_t(x)\tilde{\mu}(dt, dx) \\
 &= \left(CK_2\beta - C\beta r^* \left(t, \frac{Z_t}{\beta}, \xi_t \frac{\tilde{Z}_t}{\beta} \right) + \frac{|CZ_t|^2}{2} + \int_E f^*(C\tilde{Z}_t(x))\xi_t(x)\lambda(dx) \right) dt + C\delta_t(|V_t| - V_t)dt \\
 &\quad + C\alpha(|U_t| + U_t)dt + CZ_t dW_t - \frac{|CZ_t|^2}{2}dt + \int_E C\tilde{Z}_t(x)\tilde{\mu}(dt, dx) - \int_E f^*(C\tilde{Z}_t(x))\xi_t(x)\lambda(dx)dt \\
 &= dI_t^- + dL_t^-,
 \end{aligned}$$

where

$$\begin{cases}
 dI_t^- = \left(C\delta_t(|V_t| - V_t) + C\alpha(|U_t| + U_t) + CK_2\beta + C\beta r^* \left(t, \frac{Z_t}{\beta}, \xi_t \frac{\tilde{Z}_t}{\beta} \right) + \frac{|CZ_t|^2}{2} \right) dt, \\
 \quad + \int_E f^*(C\tilde{Z}_t(x))\xi_t(x)\lambda(dx)dt \\
 dL_t^- = CZ_t dW_t - \frac{|CZ_t|^2}{2}dt + \int_E C\tilde{Z}_t(x)\tilde{\mu}(dt, dx) - \int_E f^*(C\tilde{Z}_t(x))\xi_t(x)\lambda(dx)dt.
 \end{cases}$$

Thanks to inequality given in (1.15), we have the following:

$$-C\beta r^* \left(t, \frac{Z_t}{\beta}, \xi_t \frac{\tilde{Z}_t}{\beta} \right) + CK_2\beta + \frac{|CZ_t|^2}{2} + \int_E f^*(C\tilde{Z}_t(x))\xi_t(x)\lambda(dx) \geq 0, \quad dt \times dP - a.e.$$

It is also easy to see, by the definition of Doléans-Dade's exponential, that $\exp(L_t^-) = \mathcal{E}(M^-)_t$ where, $dM_t^- = CZ_t dW_t + \int_E (e^{C\tilde{Z}_t(x)} - 1)\tilde{\mu}(dt, dx)$. Therefore, we obtain,

$$K_t^- \stackrel{(\Delta)}{=} \exp(Y_t^-) = \exp(V_0) \exp(I_t^-) \exp(L_t^-) = \exp(V_0) \exp(I_t^-) \mathcal{E}(M^-)_t.$$

Using the integration by part formula, we get, $dK_t^- = K_t^- (dM_t^- + dI_t^-)$, which implies, that the predictable quadratic variation of K^- verifies, $d\langle K^- \rangle_t = (K_t^-)^2 d\langle M^- \rangle_t$ and as a consequence, $d\langle M^- \rangle_t = \frac{1}{(K_t^-)^2} d\langle K^- \rangle_t$. Hence,

$$\langle M^- \rangle_T \leq \sup_{t \in [0, T]} \left(\frac{1}{(K_t^-)^2} \right) \times \langle K^- \rangle_T. \quad (3.11)$$

Now, we need to have an estimate for $\langle K^- \rangle$ in order to get one for $\langle M^- \rangle$. Itô's formula yields,

$$d(K_t^-)^2 = 2K_t^- dK_t^- + d[K^-]_t = 2(K_t^-)^2 (dM_t^- + dI_t^-) + d[K^-]_t.$$

Taking a sequence of stopping times (T_n) such that for each $n \in \mathbb{N}$, $(\int_0^{t \wedge T_n} 2(K_s^-)^2 dM_s^-)_t$ is a uniformly integrable martingale and integrating the above equation between a stopping

time $\sigma \leq T$ and $T \wedge T_n$, we get,

$$[K^-]_{T \wedge T_n} - [K^-]_\sigma = (K_{T \wedge T_n}^-)^2 - (K_\sigma^-)^2 - 2 \int_\sigma^{T \wedge T_n} (K_t^-)^2 (dM_t^- + dI_t^-).$$

Since $\int_0^{T_n} (K_t^-)^2 dI_t^- \geq 0$, by taking conditional expectations on both sides, we obtain,

$$E[\langle K^- \rangle_{T \wedge T_n} - \langle K^- \rangle_\sigma | \mathcal{F}_\sigma] = E[[K^-]_{T \wedge T_n} - [K^-]_\sigma | \mathcal{F}_\sigma] \leq E[(K_{T \wedge T_n}^-)^2 | \mathcal{F}_\sigma] \leq E[\sup_{t \in [0, T]} (K_t^-)^2 | \mathcal{F}_\sigma].$$

Finally, passing to the limit as $n \rightarrow +\infty$ and using the Monotone Convergence theorem, we have,

$$E[\langle K^- \rangle_T - \langle K^- \rangle_\sigma | \mathcal{F}_\sigma] \leq E[\sup_{t \in [0, T]} (K_t^-)^2 | \mathcal{F}_\sigma].$$

Now, since for every $p \geq 1$, $\sup_{t \in [0, T]} K_t^- \in L^p(\Omega)$, it follows from Garcia and Neveu Lemma (see for example Lemma 4.3 in [Barrieu and El Karoui \(2013\)](#) or [Neveu \(1972\)](#)) that

$$E[\langle K^- \rangle_T^p] < \infty, \quad \forall p \geq 1. \quad (3.12)$$

With the same arguments used to show that $\sup_{t \in [0, T]} K_t^- \in L^p(\Omega)$, we have also that $\sup_{t \in [0, T]} \frac{1}{K_t^-} \in L^p(\Omega)$ for any $p \geq 1$. From (3.11) and (3.12) together with Cauchy-Schwartz inequality, we deduce that

$$E[\langle M^- \rangle_T^p] < \infty, \quad \forall p \geq 1. \quad (3.13)$$

As for the process Y^+ , it verifies, $dY_t^+ = dI_t^+ + dL_t^+$ where,

$$\begin{cases} dI_t^+ = \left(C\delta_t(|V_t| + V_t) + C\alpha(|U_t| - U_t) + CK_2\beta + C\beta r^* \left(t, \frac{Z_t}{\beta}, \xi_t \frac{\tilde{Z}_t}{\beta} \right) + \frac{|CZ_t|^2}{2} \right) dt \\ \quad + \int_E f^*(-C\tilde{Z}_t(x))\xi_t(x)\lambda(dx)dt, \\ dL_t^+ = -CZ_t dW_t - \frac{|CZ_t|^2}{2} dt - \int_E C\tilde{Z}_t(x)\tilde{\mu}(dt, dx) - \int_E f^*(-C\tilde{Z}_t(x))\xi_t(x)\lambda(dx)dt. \end{cases}$$

As r^* and f^* are positive functions, the process I_t^+ is increasing and as previously, by easy calculations, we can see that $\exp(L_t^+) = \mathcal{E}(M^+)_t$ where $dM_t^+ = -CZ_t dW_t + \int_E (e^{-C\tilde{Z}_t(x)} - 1)\tilde{\mu}(dt, dx)$. Going through the same lines as with Y^- , we obtain,

$$E[\langle M^+ \rangle_T^p] < \infty, \quad \forall p \geq 1. \quad (3.14)$$

But, expressing the expression of predictable quadratic variation of M^+ and M^- , we get,

$$\begin{aligned} E \left[\left(\int_0^T |CZ_t|^2 dt + \int_0^T \int_E (e^{C\tilde{Z}_t(x)} - 1)^2 \nu(dt, dx) \right)^p \right] &< \infty, \quad p \geq 1, \\ E \left[\left(\int_0^T |CZ_t|^2 dt + \int_0^T \int_E (e^{-C\tilde{Z}_t(x)} - 1)^2 \nu(dt, dx) \right)^p \right] &< \infty, \quad p \geq 1. \end{aligned}$$

This implies from one hand that,

$$E \left[\left(\int_0^T |Z_t|^2 dt \right)^p \right] < \infty, \quad p \geq 1,$$

and from the other hand, using the fact that $|y|^2 \leq 2(|e^y - 1|^2 + |e^{-y} - 1|^2)$, $y \in \mathbb{R}$, we get that,

$$E \left[\left(\int_0^T \int_E |\tilde{Z}_t(x)|^2 \nu(dt, dx) \right)^p \right] < \infty, \quad p \geq 1.$$

In conclusion, we have showed that $Z \in \mathcal{H}^{2,p}$ and $\tilde{Z} \in \mathcal{H}_\lambda^{2,p}$. \square

In the next proposition, we establish a comparison theorem for the class of BSDEs in (3.2). For two random variables, we write $A \leq B$ if $A \leq BP - a.s.$ and for two processes X and Y , we write $X \leq Y$ if $\forall t \in [0, T]$, $X_t \leq Y_t P - a.s.$ Finally, we write $(A, X) \leq (B, Y)$ if $A \leq B$ and $X \leq Y$.

Proposition 3.4. *Assume that for $k = 1, 2$, (V^k, Z^k, \tilde{Z}^k) is a solution of the BSDE (3.2) in $D_0^{\text{exp}} \times \mathcal{H}^{2,p} \times \mathcal{H}_\lambda^{2,p}$ associated with (U^k, \bar{U}_T^k) . If $(U^1, \bar{U}_T^1) \leq (U^2, \bar{U}_T^2)$, then,*

$$\forall t \in [0, T], \quad V_t^1 \leq V_t^2 \quad P - a.s.$$

Proof. In general, establishing comparison theorems for BSDE is obtained through an estimate of the quantity $((V^1 - V^2)^+)^2$. Here, in order to take advantage of the convexity of the finite variation part of the BSDE, we will rather estimate $V^1 - \theta V^2$ for each $\theta \in (0, 1)$. Similar idea was used in [Briand and Hu \(2008\)](#) for the continuous case.

Let $\theta \in (0, 1)$ and $V^\theta = V^1 - \theta V^2$. We define accordingly Z^θ , \tilde{Z}^θ , U^θ and \bar{U}_T^θ . From (3.2), the dynamics of the process V^θ discounted are given by

$$\begin{aligned} dS_t^\delta V_t^\theta &= S_t^\delta \left[-\alpha U_t^\theta + \beta \left(r^* \left(t, \frac{Z_t^1}{\beta}, \xi_t \frac{\hat{Z}_t^1}{\beta} \right) - \theta r^* \left(t, \frac{Z_t^2}{\beta}, \xi_t \frac{\hat{Z}_t^2}{\beta} \right) \right) \right] dt \\ &\quad - S_t^\delta Z_t^\theta dW_t - \int_E S_t^\delta \tilde{Z}_t^\theta(x) \tilde{\mu}(dx, dt) \\ &= S_t^\delta (-\alpha U_t^\theta + \beta(r^{*,1} - \theta r^{*,2})) dt - S_t^\delta Z_t^\theta dW_t - \int_E S_t^\delta \tilde{Z}_t^\theta(x) \tilde{\mu}(dx, dt), \end{aligned}$$

where, to alleviate the notations, we have denoted, $r^{*,i} = r^*(t, \frac{Z_t^i}{\beta}, \xi_t \frac{\hat{Z}_t^i}{\beta})$. Now, since r^* is

convex, the term $r^{*,1} - \theta r^{*,2}$ can be bounded from above. Indeed,

$$\begin{aligned} r^{*,1} &= r^* \left(t, \frac{Z_t^1}{\beta}, \xi_t \frac{\widehat{Z}_t^1}{\beta} \right) \leq \theta r^* \left(t, \frac{Z_t^2}{\beta}, \xi_t \frac{\widehat{Z}_t^2}{\beta} \right) + (1-\theta) r^* \left(t, \frac{Z_t^\theta}{\beta(1-\theta)}, \xi_t \frac{\widetilde{Z}_t^\theta}{\beta(1-\theta)} \right) \\ &:= \theta r^{*,2} + (1-\theta) r^{*,\theta}. \end{aligned} \quad (3.15)$$

Moreover, thanks to (1.15), we have

$$r^{*,\theta} \leq K_2 + \frac{|Z_t^\theta|^2}{2K_1\beta^2(1-\theta)^2} + K_1 \int_E f^* \left(\frac{\widetilde{Z}_t^\theta(x)}{K_1\beta(1-\theta)} \right) \xi_t(x) \lambda(dx).$$

Using this last inequality in (3.15), we get that,

$$\beta(r^{*,1} - \theta r^{*,2}) \leq K_2\beta(1-\theta) + \frac{|Z_t^\theta|^2}{2K_1\beta(1-\theta)} + K_1\beta(1-\theta) \int_E f^* \left(\frac{\widetilde{Z}_t^\theta(x)}{K_1\beta(1-\theta)} \right) \xi_t(x) \lambda(dx). \quad (3.16)$$

To get rid of the quadratic and exponential terms in the inequality above, we will use an exponential change of variables. More precisely, let c be a negative constant (to be specified later), and set $P_t = \exp(cS_t^\delta V_t^\theta)$, $Q_t = cS_t^\delta P_{t-} Z_t^\theta$ and $\widetilde{Q}_t = cS_t^\delta P_{t-} \widetilde{Z}_t^\theta$. Using Itô's formula, we deduce,

$$\begin{aligned} dP_t &= P_{t-} \left[cd\langle S_t^\delta V_t^\theta \rangle + \frac{c^2}{2} d\langle S_t^\delta V_t^\theta \rangle_t + \int_E f^*(-cS_t^\delta \widetilde{Z}_t^\theta(x)) \mu(dx, dt) \right] \\ &= cP_{t-} \left[S_t^\delta (-\alpha U_t^\theta + \beta(r^{*,1} - \theta r^{*,2})) dt - S_t^\delta Z_t^\theta dW_t - \int_E S_t^\delta \widetilde{Z}_t^\theta(x) \widetilde{\mu}(dx, dt) \right. \\ &\quad \left. + \frac{c}{2} |S_t^\delta Z_t^\theta|^2 dt + \frac{1}{c} \int_E f^*(-cS_t^\delta \widetilde{Z}_t^\theta(x)) \mu(dx, dt) \right] \\ &= cS_t^\delta P_{t-} \left[-\alpha U_t^\theta + \beta(r^{*,1} - \theta r^{*,2}) + \frac{c}{2} S_t^\delta |Z_t^\theta|^2 + \frac{1}{cS_t^\delta} \int_E f^*(-cS_t^\delta \widetilde{Z}_t^\theta(x)) \xi_t(x) \lambda(dx) \right] dt \\ &\quad - Q_t dW_t - \int_E \widetilde{Q}_t(x) \widetilde{\mu}(dx, dt) + P_{t-} \int_E f^*(-cS_t^\delta \widetilde{Z}_t^\theta(x)) \widetilde{\mu}(dx, dt) \\ &:= G_t dt - Q_t dW_t - \int_E \widetilde{Q}_t(x) \widetilde{\mu}(dx, dt) + P_{t-} \int_E f^*(-cS_t^\delta \widetilde{Z}_t^\theta(x)) \widetilde{\mu}(dx, dt). \end{aligned}$$

Thanks to equation (3.16), the G_t term is bounded from above,

$$\begin{aligned} G_t &\leq cS_t^\delta P_{t-} \left[-\alpha U_t^\theta + K_2\beta(1-\theta) + \frac{|Z_t^\theta|^2}{2} \left(\frac{1}{K_1\beta(1-\theta)} + cS_t^\delta \right) \right. \\ &\quad \left. + \int_E \left(K_1\beta(1-\theta) f^* \left(\frac{\widetilde{Z}_t^\theta(x)}{K_1\beta(1-\theta)} \right) - \frac{1}{-cS_t^\delta} f^*(-cS_t^\delta \widetilde{Z}_t^\theta(x)) \right) \xi_t(x) \lambda(dx) \right] \\ &:= cS_t^\delta P_{t-} \left[-\alpha U_t^\theta + K_2\beta(1-\theta) + \frac{|Z_t^\theta|^2}{2} \left(\frac{1}{K_1\beta(1-\theta)} + cS_t^\delta \right) \right. \\ &\quad \left. + \int_E \left(h(K_1\beta(1-\theta), \widetilde{Z}_t^\theta(x)) - h\left(\frac{-1}{cS_t^\delta}, \widetilde{Z}_t^\theta(x)\right) \right) \xi_t(x) \lambda(dx) \right], \end{aligned} \quad (3.17)$$

where $h : \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ defined as $h(x, z) := x f^*(z/x) = x e^{z/x} - x - z$. We need to choose c such that the term next to $|Z_t^\theta|^2$ is negative, that choose c such that,

$$\frac{1}{K_1 \beta (1 - \theta)} \leq -c S_t^\delta.$$

Since $S_t^\delta \geq e^{\|\delta\|_\infty T}$, it is sufficient to set $c(\theta) := -\frac{e^{-\|\delta\|_\infty T}}{K_1 \beta (1 - \theta)}$. Computing the derivative of h with respect to x , we get, $\partial_x h(x, z) = e^{z/x} - (z/x) e^{z/x} - 1$. Studying the sign of the function $x \rightarrow e^x - x e^x - 1$ by calculating its derivative, we obtain that $e^x - x e^x - 1 \leq 0, \forall x \in \mathbb{R}$. Therefore, we deduce that $\partial_x h(x, z) \leq 0, \forall x \in \mathbb{R}$, that h is decreasing. Hence, going back to (3.17), we get that,

$$G_t \leq c(\theta) S_t^\delta P_{t-} (-\alpha U_t^\theta + K_2 \beta (1 - \theta)) \leq S_t^\delta P_{t-} e^{-\|\delta\|_\infty T} \left(\frac{\alpha}{K_1 \beta} U_t^1 - \frac{K_2}{K_1} \right), \quad (3.18)$$

where we have used, in the second inequality, the fact that,

$$U_t^\theta = U_t^1 - \theta U_t^2 = \theta (U_t^1 - U_t^2) + (1 - \theta) U_t^1 \leq (1 - \theta) U_t^1.$$

Finally, denoting $D_t = \exp\left(-e^{-\|\delta\|_\infty T} \int_0^t S_s^\delta \left(\frac{\alpha}{K_1 \beta} U_t^1 - \frac{K_2}{K_1}\right) ds\right)$, and introducing $P_t^D := D_t P_t$, $Q_t^D := Q_t D_t$ and $\tilde{Q}_t^D := \tilde{Q}_t D_t$. Using again Itô's formula, for any stopping time $0 \leq t \leq \tau \leq T$,

$$P_t^D \geq P_\tau^D + \int_t^\tau Q_s^D dW_s + \int_t^\tau \tilde{Q}_s^D(x) \tilde{\mu}(dx, ds) - \int_t^\tau \int_E P_{s-} D_s f^*(-c S_s^\delta \tilde{Z}_s^\theta(x)) \tilde{\mu}(dx, ds).$$

Considering a localizing sequence of stopping time τ_n , such that the local martingales, in the above inequality, stopped in τ_n are martingales, we obtain,

$$P_t \geq E \left[P_{\tau_n} \exp\left(-e^{-\|\delta\|_\infty T} \int_t^{\tau_n} S_s \left(\frac{\alpha}{K_1 \beta} U_t^1 - \frac{K_2}{K_1}\right) ds\right) \middle| \mathcal{F}_t \right].$$

In view of the integrability assumptions on U^1 and on V , by the dominated convergence theorem, we can deduce that,

$$P_t \geq E \left[P_T \exp\left(-e^{-\|\delta\|_\infty T} \int_t^T S_s \left(\frac{\alpha}{K_1 \beta} U_t^1 - \frac{K_2}{K_1}\right) ds\right) \middle| \mathcal{F}_t \right].$$

But by definition of P , $P_T = \exp(c(\theta) S_T^\delta V_T^\theta) = \exp(c(\theta) S_T^\delta (\bar{U}_T^1 - \theta \bar{U}_T^2))$, and because $\bar{U}_T^1 \leq \bar{U}_T^2$ and $c(\theta)$ is negative, we get,

$$c(\theta) S_T^\delta V_T^\theta \geq -\frac{e^{-\|\delta\|_\infty T}}{K_1 \beta} S_T^\delta \bar{U}_T^1.$$

Therefore, we have,

$$P_t \geq E \left[\exp \left(-e^{-\|\delta\|_\infty T} \left(\frac{S_T^\delta}{K_1 \beta} \bar{U}_T^1 + \int_t^T S_s \left(\frac{\alpha}{K_1 \beta} U_t^1 - \frac{K_2}{K_1} \right) ds \right) \right) \middle| \mathcal{F}_t \right],$$

which implies that,

$$V_t^\theta \leq -\frac{K_1 \beta (1 - \theta) e^{\|\delta\|_\infty T}}{S_t^\theta} \ln E \left[\exp \left(-e^{-\|\delta\|_\infty T} \left(\frac{S_T^\delta}{K_1 \beta} \bar{U}_T^1 + \int_t^T S_s \left(\frac{\alpha}{K_1 \beta} U_t^1 - \frac{K_2}{K_1} \right) ds \right) \right) \middle| \mathcal{F}_t \right].$$

Taking the limit when $\theta \nearrow 1$, we finally get,

$$V_t^1 \leq V_t^2.$$

□

The following corollary is a direct consequence of the comparison result above.

Corollary 3.5. *Under assumptions (A_r) and (A_u) , the BSDE (3.2) has a unique solution $(V, Z, \tilde{Z}) \in D_0^{\text{exp}} \times \mathcal{H}^{2,p} \times \mathcal{H}_\lambda^{2,p}$.*

4 Appendix

Lemma 4.1. *Let $r(t, \eta, \psi) = \frac{1}{2}|\eta|^2 + \int_E f(\psi(x)) \xi_t(x) \lambda(dx)$ and $Q \in \mathcal{Q}_f$. Then, the following processes,*

$$M_t = \int_0^t \eta_s dW_s^Q, \quad M'_t = \int_0^t \int_E \log(1 + \psi_s(x)) \tilde{\mu}^Q(ds, dx),$$

are Q -martingales.

Proof. Since $Q \in \mathcal{Q}_f$, we have,

$$E_Q \left[\int_0^T r(t, \eta_t, \psi_t) dt \right] = E_Q \left[\int_0^T \left(\frac{1}{2} |\eta_t|^2 + \int_E f(\psi_t(x)) \xi_t(x) \lambda(dx) \right) dt \right] < \infty. \quad (4.1)$$

In particular, $E_Q \left[\int_0^T \frac{1}{2} |\eta_t|^2 dt \right] < \infty$, which implies that M is Q -martingale. Now, we prove that M' is also a Q -martingale. First, note that,

$$f(x) = (1+x) \log(1+x) - x \geq \frac{1}{6} (1+x) \log^2(1+x) \geq 0, \quad \text{for } -1 \leq x \leq e^2 - 1.$$

Hence, as the RHS of (4.1) is finite, we get that

$$\log(1 + \psi_s(x)) \mathbb{1}_{\psi_s(x) \leq e^2 - 1} \in L^2(dQ \times \nu^Q(ds, dx)). \quad (4.2)$$

Moreover, for $x > e^2 - 1$, we have,

$$(1+x) \log(1+x) \leq 2((1+x) \log(1+x) - x).$$

Again, as the RHS of (4.1) is finite, we get that,

$$\log(1 + \psi_s(x)) \mathbb{1}_{\psi_s(x) > e^{2-1}} \in L^1(dQ \times \nu^Q(ds, dx)). \quad (4.3)$$

From (4.2) and (4.3), and using Theorem 1.8(i) in [Jacod and Shiryaev \(2013\)](#), we obtain that M' is Q -martingale. \square

Proposition 4.2. *Let $r(t, \eta, \psi) = \frac{1}{2}|\eta|^2 + \int_E f(\psi(x))\xi_t(x)\lambda(dx)$ and $Q \in \mathcal{Q}_f$. Then, we have,*

$$H(Q|P) = E_Q \left[\int_0^T r(t, \eta_t, \psi_t) dt \right] = E_Q \left[\int_0^T \left(\frac{1}{2}|\eta_t|^2 + \int_E f(\psi_t(x))\xi_t(x)\lambda(dx) \right) dt \right]. \quad (4.4)$$

Proof. Let $Q \in \mathcal{Q}_f$ with corresponding (η, ψ) . We introduce the following sequence of processes $(\psi_m)_{m \in \mathbb{N}^*}$ defined as:

$$\psi_{m,s}(x) = \psi_s(x) \mathbb{1}_{\psi_s(x) \leq m} \mathbb{1}_{|x| \geq 1/m}.$$

It is clear that $\psi_m \in L^2(dQ \times \nu^Q(ds, dx))$. Developing the logarithm of Radon-Nikodym derivative of Q w.r.t P gives Q -a.s.:

$$\begin{aligned} \log \left(\frac{dQ}{dP} \right) &= \lim_{m \rightarrow \infty} \log \left\{ \mathcal{E} \left(\int_0^T \eta_s dW_s + \int_0^T \int_E \psi_{m,s}(x) \tilde{\mu}(ds, dx) \right) \right\} \\ &= \lim_{m \rightarrow \infty} \left\{ \int_0^T \eta_s dW_s - \frac{1}{2} \int_0^T |\eta_s|^2 ds + \int_0^T \int_E \psi_{m,s}(x) \tilde{\mu}(ds, dx) \right. \\ &\quad \left. + \int_0^T \int_E (\log(1 + \psi_{m,s}(x)) - \psi_{m,s}(x)) \mu(ds, dx) \right\} \\ &= \lim_{m \rightarrow \infty} \left\{ \int_0^T \eta_s dW_s^Q + \frac{1}{2} \int_0^T |\eta_s|^2 ds + \int_0^T \int_E \psi_{m,s}(x) \tilde{\mu}^Q(ds, dx) \right. \\ &\quad \left. + \int_0^T \int_E (\log(1 + \psi_{m,s}(x)) - \psi_{m,s}(x)) \tilde{\mu}^Q(ds, dx) \right. \\ &\quad \left. + \int_0^T \int_E [\psi_{m,s}(x)\psi_s(x) + (1 + \psi_s(x)) (\log(1 + \psi_{m,s}(x)) - \psi_{m,s}(x))] \nu(ds, dx) \right\} \\ &= \lim_{m \rightarrow \infty} \left\{ \int_0^T \eta_s dW_s^Q + \frac{1}{2} \int_0^T |\eta_s|^2 ds + \int_0^T \int_E \log(1 + \psi_{m,s}(x)) \tilde{\mu}^Q(ds, dx) \right. \\ &\quad \left. + \int_0^T \int_E [(1 + \psi_s(x)) \log(1 + \psi_{m,s}(x)) - \psi_{m,s}(x)] \nu(ds, dx) \right\}, \end{aligned} \quad (4.5)$$

where we used from the second to the third inequality that, by the definition of the process ψ_m , $(1 + \psi_s)(\log(1 + \psi_{m,s}) - \psi_{m,s}) \in L^1(\nu(ds, dx))$ and $\psi_m \psi \in L^1(\nu(ds, dx))$. In particular, $\int_0^t \int_E (\log(1 + \psi_{m,s}(x)) - \psi_{m,s}(x)) \tilde{\mu}^Q(ds, dx)$ is a well defined. Lemma 4.1 above insures that the following processes:

$$M_t := \int_0^t \eta_s dW_s^Q, \quad M'_t := \int_0^t \int_E \log(1 + \psi_s(x)) \tilde{\mu}^Q(ds, dx), \quad M'_{m,t} := \int_0^t \int_E \log(1 + \psi_{m,s}(x)) \tilde{\mu}^Q(ds, dx),$$

are Q -martingales. Moreover, we also have that $M'_{m,T}$ converges to M'_T in $L^1(Q)$. Indeed, decomposing M' and M'_m as in Lemma 4.1 in the following way,

$$M'_{m,t} = \widehat{M}_{m,t} + \widetilde{M}_{m,t},$$

where,

$$\widehat{M}_{m,t} := \int_0^t \int_E \log(1 + \psi_{s,m}(x)) \mathbb{1}_{\psi_{s,m}(x) \leq e^2 - 1}, \quad \widetilde{M}_{m,t} := \int_0^t \int_E \log(1 + \psi_{s,m}(x)) \mathbb{1}_{\psi_{s,m}(x) > e^2 - 1}.$$

We have the positive (resp. negative) part of $\log(1 + \psi_{m,s}(x)) \mathbb{1}_{\psi_{s,m}(x) \leq e^2 - 1}$ increases (resp. decreases) to the positive (resp. negative) part of $\log(1 + \psi_s(x)) \mathbb{1}_{\psi_s(x) \leq e^2 - 1}$ as m goes to infinity and by the monotone convergence we deduce that $\widehat{M}_{m,T}$ converges to \widehat{M}_T in $L^1(Q)$. Using the same argument, we have also $\widetilde{M}_{m,T}$ converges to \widetilde{M}_T in $L^1(Q)$ as m goes to infinity. Hence, we obtain that $M'_{m,T}$ converges to M'_T as $m \rightarrow \infty$. By passing to a subsequence, we may assume that $M'_{m,T}$ converges to M'_T Q -a.s. Finally, by Monotone convergence theorem, the last term in (4.5) also converges to $\int_0^T \int_E f(\psi_s(x)) \nu(ds, dx)$. Consequently, (4.5) becomes,

$$\begin{aligned} \log\left(\frac{dQ}{dP}\right) &= \int_0^T \eta_s dW_s^Q + \frac{1}{2} \int_0^T |\eta_s|^2 ds + \int_0^T \int_E \log(1 + \psi_s(x)) \widetilde{\mu}^Q(ds, dx) \\ &\quad + \int_0^T \int_E f(\psi_s(x)) \nu(ds, dx). \end{aligned}$$

Taking the expectation under Q and using the fact that M and M' are martingales yields (4.4). \square

References

- Anderson, E. W., Hansen, L. P., and Sargent, T. J. (2003). A quartet of semigroups for model specification, robustness, prices of risk, and model detection. *Journal of the European Economic Association*, 1(1):68–123.
- Barrieu, P. and El Karoui, N. (2013). Monotone stability of quadratic semimartingales with applications to unbounded general quadratic bsdes. *The Annals of Probability*, 41(3B):1831–1863.
- Becherer, D. (2006). Bounded solutions to backward sdes with jumps for utility optimization and indifference hedging. *The Annals of Applied Probability*, 16(4):2027–2054.
- Bordigoni, G., Matoussi, A., and Schweizer, M. (2007). A stochastic control approach to a robust utility maximization problem. In *Stochastic analysis and applications*, pages 125–151. Springer.

-
- Briand, P. and Hu, Y. (2006). Bsde with quadratic growth and unbounded terminal value. Probability Theory and Related Fields, 136(4):604–618.
- Briand, P. and Hu, Y. (2008). Quadratic bsdes with convex generators and unbounded terminal conditions. Probability Theory and Related Fields, 141(3):543–567.
- Carmona, R. (2008). Indifference pricing: theory and applications. Princeton University Press.
- Duffie, D. and Epstein, L. G. (1992). Stochastic differential utility. Econometrica: Journal of the Econometric Society, pages 353–394.
- Duffie, D. and Skiadas, C. (1994). Continuous-time security pricing: A utility gradient approach. Journal of Mathematical Economics, 23(2):107–131.
- Faidi, W., Matoussi, A., and Mni, M. (2013). Robust utility maximization problem with a general penalty term. Preprint.
- Gundel, A. (2005). Robust utility maximization for complete and incomplete market models. Finance and Stochastics, 9(2):151–176.
- Hansen, L. P., Sargent, T. J., Turmuhambetova, G., and Williams, N. (2006). Robust control and model misspecification. Journal of Economic Theory, 128(1):45–90.
- Hu, Y., Imkeller, P., and Müller, M. (2005). Utility maximization in incomplete markets. The Annals of Applied Probability, 15(3):1691–1712.
- Jacod, J. and Shiryaev, A. (2013). Limit theorems for stochastic processes, volume 288. Springer Science & Business Media.
- Karoui, N. E. (1981). Les aspects probabilistes du contrôle stochastique. In École d’été de Probabilités de Saint-Flour IX-1979, pages 73–238. Springer.
- Karoui, N. E., Matoussi, A., and Ngupeyou, A. (2016). Quadratic exponential semimartingales and application to bsdes with jumps. arXiv preprint arXiv:1603.06191.
- Lazrak, A. and Quenez, M. C. (2003). A generalized stochastic differential utility. Mathematics of operations research, 28(1):154–180.
- Morlais, M.-A. (2009). Quadratic bsdes driven by a continuous martingale and applications to the utility maximization problem. Finance and Stochastics, 13(1):121–150.
- Neveu, J. (1972). Martingales à temps discret, volume 1. Masson Paris.
- Quenez, M.-C. (2004). Optimal portfolio in a multiple-priors model. In Seminar on Stochastic Analysis, Random Fields and Applications IV, pages 291–321. Springer.
-

- Rouge, R. and El Karoui, N. (2000). Pricing via utility maximization and entropy. Mathematical Finance, 10(2):259–276.
- Schied, A. (2007). Optimal investments for risk-and ambiguity-averse preferences: a duality approach. Finance and Stochastics, 11(1):107–129.
- Schied, A. and Wu, C.-T. (2005). Duality theory for optimal investments under model uncertainty.
- Schroder, M. and Skiadas, C. (1999). Optimal consumption and portfolio selection with stochastic differential utility. Journal of Economic Theory, 89(1):68–126.
- Shiryayev, A. N. (1981). Martingales: recent developments, results and applications. International Statistical Review/Revue Internationale de Statistique, pages 199–233.
- Skiadas, C. (2003). Robust control and recursive utility. Finance and Stochastics, 7(4):475–489.