



What Can We Learn from Synaptic Connectivity Maps about Cerebellar Internal Models?

Ludovic Spaeth, Philippe Isope

► To cite this version:

Ludovic Spaeth, Philippe Isope. What Can We Learn from Synaptic Connectivity Maps about Cerebellar Internal Models?. The Cerebellum, 2022, <10.1007/s12311-022-01392-6>. <hal-03815818>

HAL Id: hal-03815818

<https://hal.science/hal-03815818v1>

Submitted on 14 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



What Can We Learn from Synaptic Connectivity Maps about Cerebellar Internal Models?

Ludovic Spaeth^{1,2} · Philippe Isope¹

Accepted: 5 March 2022
© The Author(s) 2022

Abstract

The cerebellum is classically associated with fine motor control, motor learning, and timing of actions. However, while its anatomy is well described and many synaptic plasticity have been identified, the computation performed by the cerebellar cortex is still debated. We, here, review recent advances on how the description of the functional synaptic connectivity between granule cells and Purkinje cells support the hypothesis that the cerebellum stores internal models of the body coordinates. We propose that internal models are specific of the task and of the locomotor context of each individual.

Keywords cerebellum · synaptic transmission · sensorimotor adaptation · internal models

Introduction

One of the major role of the cerebellum in sensorimotor adaptation is to learn to predict the future in order to adjust or correct the course of our actions [1–3]. To do this, many studies demonstrated that the cerebellum stores internal models of the motor apparatus, which combine motor commands with the current body state in order to predict an expected sensory feedback [4–7]. The comparison between the actual and the expected sensory feedback defines a prediction error that allows for correction, re-adjustment of the internal models, and ultimately promotion of behavioral adaptation. At the microcircuit level, the adapted behavior would correspond to modified network dynamics underpinned by a re-organization of the excitatory and inhibitory synaptic inputs at different cerebellar connections. Synergies of synaptic plasticity can ensure re-adaptation to a constantly changing environment. Therefore, the description of the functional synaptic organization (i.e., the connectivity maps) under normal or perturbed conditions should help understand how cerebellar internal models enable sensorimotor

adaptation. After introducing the topography of cerebellar climbing and mossy fiber inputs, we will discuss how connectivity maps between granule cells (GC) and Purkinje cells (PC) can shed light on internal models implementation in the cerebellum.

Climbing fibers define sagittal functional microzones

The cerebellar cortex receives two major excitatory inputs, the climbing (CF) and the mossy fibers (MF), which are both topographically organized [8–11]. A given group of neighboring PCs is contacted by CFs originating in a discrete region of the inferior olive and conveying information from the same receptive fields, defining a functional unit called microzone [12–15]. Since olivary cells are coupled by gap junctions and fire together, this pathway leads to synchronized activation of PCs from the same microzone [16]. Microzones were therefore defined as parasagittal zones of the cerebellar cortex composed of a group of PCs, the molecular layer interneurons (MLI) in the vicinity and the GCs located underneath [8, 9, 17]. Since PCs project to a specific group of nuclear neurons from which a subgroup project back to olivary cells, microzones constitute the cortical part of an olivo-cerebellar module processing information related to a defined area of the body [8]. Microzone boundaries can be identified by a family of neurochemical markers, which are specifically expressed in sagittal bands of

✉ Philippe Isope
philippe.isope@inci-cnrs.unistra.fr

¹ Institut des Neurosciences Cellulaires et Intégratives, CNRS, Université de Strasbourg, 67084 Strasbourg, France

² Present Address: Dominick P Purpura Department of Neuroscience, Albert Einstein College of Medicine, Bronx, NY, USA




Fig. 1 GC-PC connectivity maps. **A**, left. PCs (green) are recorded using patch-clamp and filled with biocytin for reconstruction and positioning (zebrin bands labeled in red). RuBi-Glutamate is uncaged using patterned blue light illumination, while GC-PC postsynaptic currents are recorded. The maximum of each GC column is used to determine a 1D pattern of connectivity for each PC (see C). Note the patchy organization of connected sites. **Right**, examples of excitatory postsynaptic currents evoked by glutamate photolysis in corresponding sites. **B**, correlation between postsynaptic currents recorded in neighboring PCs indicating that they have similar connectivity maps. **C**, median of the 1D connectivity maps (z-scored) across animals ($n=18$), illustrating that some specific GC layer columns are connected to midline PCs in many animals. Z-score < 3 corresponds to silent sites (i.e., illuminated GC layer of $41 \times 41 \times \sim 100 \mu\text{m}$ eliciting

current $< 15 \text{ pA}$ in the recorded PC). Note local and distal hotspots of GC-PC connections. **A**, **B** and **C** Adapted from [23]. **D**, Strategy to map synaptic connections in neural networks and correlate it to animal behavior. Panel 1, animals training in different locomotor contexts. Panel 2, extraction of behavioral features related to sensorimotor adaptation for each mouse (e.g., distance traveled in a wheel or gait symmetry in a corridor). Panel 3, GC-PC synaptic connectivity maps description in a specific area of the cerebellum involved in the behavior. Panel 4, determination of the structure of these maps using mathematical algorithms from graph theory. Panel 5, the structure of the maps and the behavioral parameters of each mouse are used to build a classifier that can predict animal behavior from the synaptic map. The classifier is validated by testing its efficiency on a different set of data. Adapted from [24]

PCs [15, 17] (Fig. 1A). When these markers are labeled, the cerebellar cortex looks like a zebra displaying positive and negative bands; hence, they are collectively called zebrins [17]. Many studies have also demonstrated that microzones have anatomical and functional regional differences leading to a large diversity of information processing in individual microzones [18–22]. Therefore, an exhaustive functional description of the synaptic communication between microzones is necessary to understand how they collectively influence motor coordination.

The mossy fiber to Purkinje cell pathway: an anatomy favoring internal model implementation

As opposed to the strict sagittal orientation of the CF inputs, MF inputs, which contact GCs, project to many different GC layer areas in a given lobule sending the same information into multiple microzones [11, 25]. Indeed, in vivo electrophysiological micro-mappings of the GC layer demonstrated that MF receptive fields in the GC layer have a fractured and redundant somatotopy [26]. Furthermore, many pre-cerebellar nuclei carrying distinct modalities project MFs to the GC layer of the same microzone [27, 11]. In the anterior lobe, for example, lobule IV and V receive MFs from the spinocerebellar tract (e.g., DSCT, dorsal spinocerebellar tract) and the external cuneate conveying proprioceptive information from the hindlimbs and the forelimbs, respectively, as well

as from the pontine nuclei, which carry information from the cerebral cortex such as motor plans and motor commands [28]. Therefore, individual GCs can integrate multimodal information [26, 29] but see [30] that combine both motor plans or commands and sensory feedback. GC axons, the parallel fibers (PF), which travel across multiple microzones, make excitatory synapses onto both local and distant PCs and molecular interneurons (MLI) located at millimeters from the incoming microzone [31, 32]. Within a given lobule, PFs likely control the PC population code via direct excitation and indirect inhibition in a wide array of microzones [33–35]. Moreover, many experiments demonstrated that the GC-PC connection is a major site of information storage in the cerebellar cortex through CF-dependent plasticity: GC-PC synapses are depressed when CF and PFs are active in conjunction and potentiated when only PFs are activated [36, 37].

The organization of MF inputs into the GC layer expands encoding possibilities and is in agreement with the long-standing Marr–Albus–Ito hypothesis, suggesting that expansion recoding in the GC layer enables pattern separation of similar MF inputs [38–42]. Moreover, the convergence of sensorimotor inputs from the spinal cord and cortico-pontine inputs suggests that the GC layer can compute motor commands as well as sensory context or feedback, a prerequisite configuration for the implementation of forward internal models. Indeed, in vivo PC recordings during movement have shown that modulation of PC discharge can lead or lag movement onset [2, 43, 44]. Recent experiments have demonstrated that in an operant task in which monkeys have to follow a cursor on a screen PC discharge can predict the position error of the hand by a few hundred milliseconds ahead and compute in return the visual feedback shortly after the movement [6] (see also [3, 45]). Since the modulation of PC discharge relies on the combined activity of GCs and MLIs, these results suggest that the cerebellar cortex stores an internal model of this skilled movement. In a different task, in which mice move a handle to get a reward, GCs encode a predicted reward, suggesting that internal models are not only sensorimotor engrams [46]. Therefore, understanding how the cerebellar cortex encode sensorimotor information and store internal models requires determining how the GC-(MLI)-PC pathway is functionally organized.

Interrogating the spatial organization of the GC-PC connection

Glutamate uncaging combined with patch-clamp recordings can be used on acute brain slices [47–49]. This approach allows the systematic excitation of single or small groups of presynaptic cells [50, 51] while monitoring the resulting synaptic inputs in the target cell soma (Fig. 1A). Uncaged

glutamate excites mostly dendritic or somatic receptors preventing the direct excitation of neighboring axons as opposed to electrical or optogenetic stimulation. When glutamate uncaging is combined with a system enabling specific and localized restricted illumination (laser scanning or patterned light illumination), the spatial organization of the presynaptic cells as well as their synaptic weights on the target cell can be reconstructed [23, 24, 52, 53] (Fig. 1A, B). Hence, each recording yields a single connectivity map. This method is particularly well adapted to the study of the GC-PC connectivity maps as PCs from identified microzones can be targeted using transgenic mice expressing fluorescent zebrin markers [54]. Therefore, the use of caged-glutamate photostimulation and whole-cell patch clamp of PCs in identified zebrin bands allows the description of specific connectivity maps across different animals and eventually different behavioral conditions [23, 24].

Spatially organized GC-PC connectivity maps as an implementation of internal models

Many in vitro and in vivo experiments demonstrated that while direct stimulation of PF beams can activate PCs along the beam, GC or MF activation actually modulate specific groups of PCs [55–58]. In vivo recordings of tactile receptive fields showed that in a given microzone GC layer, Golgi cells and MLIs are excited by the same area of the body, suggesting that incoming MF inputs lead to the activation of most cell types [14, 56, 59]. Conversely, PCs are excited by specific MF inputs targeting GCs in distant microzones, illustrating a functional selection of GC-PC synapses [23, 60]. Indeed, in two seminal studies, Jorntell and Ekerot [59, 61] demonstrated in vivo that PCs sharing the same PF beam are excited by different and specific receptive fields, but upon repeated high frequency PF stimulation PCs respond to all. Altogether, these results suggest the existence of a priori nonfunctional—or silent—GC-PC synapses (e.g., depressed by the CF-dependent LTD) that are potentiated—or awakened—via activity-dependent plasticity. This hypothesis was confirmed in vitro, using paired recordings and photostimulation studies in acute slices, showing that while PCs are always connected to local (i.e., underneath) GCs, a selection of distant hotspots of connected GC patches is interleaved with GCs making silent or nonfunctional synapses [23, 24, 62] (Fig. 1A, C). Silent synapses were also recently observed during development and in adult rodents [24, 63].

By repeating photostimulation and establishing excitatory GC-PC and GC-MLI connectivity maps in an identified set of microzones across many mice, several other operational rules were described [23, 24]: (1) neighboring PCs have a similar patchy connectivity map defining clusters of

PCs spatially organized (Fig. 1A, B). While PC clusters are related to zebrin band, they overlap with zebrin boundaries, suggesting that a finer functional organization may coexist with zebrin band topography [17]. (2) MLIs from an identified microzone have different connectivity maps than local PCs, but they are similarly organized with local and distal patches of connected GCs. (3) Although functional connectivity is highly variable between animals, some specific areas of the synaptic maps are conserved between individuals, highlighting specific association between microzones through GC-PC synapses (Fig. 1C). (4) Connectivity maps are plastic and can be modified by protocols of PF stimulation, which lead to a reorganization of the map with potentiated or depressed GC sites, suggesting that they are sensitive to activity of the MF-GC-PC network. GC-PC maps were also recorded after mice underwent locomotor adaptation in different locomotor context [24]: a transient locomotor impairment (i.e., recovery following unilateral alteration of the sciatic nerve) or a training in a wheel for three weeks. These specific locomotor contexts induced both a sustainable increase in overall synaptic weights of GC-PC connections synapses and a specific reorganization of the connectivity maps. Moreover, using mathematical algorithms based on graph theory, we demonstrated that a synaptic connectivity map from a given mouse is correlated with the unique behavioral performance of this mouse in a simple task such as walking on a corridor or running in a wheel (Fig. 1D). These results suggest that GC-PC connectivity maps are specific engrams (i.e., internal models) of behavioral adaptation, allowing encoding of movement features in a context-specific manner throughout individual motor learning.

Patchy organization of GC-PC maps and “sparse” coding in the cerebellar cortex

Altogether, these operational rules underlying microzonal communication in the cerebellar cortex are likely at the core of cerebellar computation [8]. The description of synaptic connectivity maps in cerebellar lobules could shed light on the combination of microzones that control a given adapted behavior. We postulate that local MF inputs in recipient microzones set a minimal framework of activated PCs while the patchy organization of GC-PC synaptic maps allows the spatial and temporal combination of an extended array of microzones by PFs. Such organization may reconcile Marr–Albus–Ito [38] and adaptive filter [39, 64] theoretical models with the recent studies describing GC activity using *in vivo* two-photon imaging [44, 65, 66]. In these theoretical models, a sparse coding strategy at the MF-GC synapses is a prerequisite for the expansion-recoding hypothesis and

few GCs should be activated by a specific combination of MF inputs.

This assumption might appear in contradiction with the dense activation of GCs observed using two-photon imaging of genetically encoded calcium indicators in behaving mice or zebrafishes [44, 65, 66]. They demonstrated that large areas of the GC layer seem simultaneously active at specific moment of the task and correlated with sensorimotor or predictive reward inputs. Anatomical, physiological, and technical issues may explain this apparent discrepancy. MFs carrying information from a given body part target specific areas of the GC layer making many rosettes in a restricted area. This arrangement favors the aforementioned patchy organization of the GC-PC functional connectivity and may explain why two-photon imaging identified patches of dense GC activity. Nevertheless, since the majority of GC-PC synapses are silent [23, 24, 62], GC-PC connectivity can still be sparse. Moreover, short-term plasticity at the MF-GC synapses expands temporal integration in GCs from tens to hundreds of milliseconds [67], suggesting that in a given GC patch targeted by a combination of multimodal MFs, GC discharges may be split into a wide range of temporal patterns yielding GC population decorrelation. As explained in [40], this mechanisms may enable pattern separation even if GCs are densely activated. In two-photon GCaMP6 imaging experiments, these patterns cannot be temporally discriminated and are merged into a single giant patch of simultaneously activated GCs. Heterogeneity of short-term plasticity was also described recently at the GC-MLI and GC-MLI-PC connection [33, 68], indicating that the temporal expansion performed by the GC layer is amplified in the molecular layer. Therefore, the spatial organization of the MF-GC-PC connections combined with the spread in the temporal integration window enables pattern discrimination and PCs to encode thousands of different patterns of information as suggested in the Marr–Albus–Ito hypothesis [69].

To conclude, patchy connectivity maps at the GC-MLI-PC connections may underlie the communication between specific microzones in the cerebellar cortex. Graph network properties of GC-PC connectivity maps suggest that context-dependent and task-specific internal models are encoded at the GC-PC synapses. Next step will be to describe how the feedforward inhibition influences connectivity maps and intermodular communication. Unraveling the rules governing the coordination between microzones will shed light on how the cerebellar cortex controls the output stage of the cerebellum. As an appealing perspective, it will be important to describe how the MF-GC-MLI pathway correlates microzones using genetically encoded voltage sensitive indicators combined with two-photon imaging. Knowing the combination of microzones and their sequence of activation in a specific behavior will help designing and refining cerebellar stimulation protocols to compensate for cerebellar dysfunction in pathological models [70].

Acknowledgements This work was supported by the Centre National pour la Recherche Scientifique (CNRS), the Université de Strasbourg, the Agence Nationale pour la Recherche (ANR-2015-CeMod, ANR-2019-MultiMod, ANR-2019-NetOnTime) and by the Fondation pour la Recherche Médicale to PI (#DEQ20140329514) and by the Fondation pour la Recherche Médicale to LS (#FDT201825005172).

Declarations

Conflict of interest The authors have no relevant financial or non-financial interests to disclose. The authors have no competing interests to declare that are relevant to the content of this article. The authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript. The authors have no financial or proprietary interests in any material discussed in this article.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Bastian AJ. Learning to predict the future: the cerebellum adapts feedforward movement control. *Curr Opin Neurobiol*. 2006;16(6):645–9. <https://doi.org/10.1016/j.conb.2006.08.016>
- Thach WT, Goodkin HP, Keating JG. The cerebellum and the adaptive coordination of movement. *Annu Rev Neurosci*. 1992;15:403–42. <https://doi.org/10.1146/annurev.ne.15.030192.002155>
- Shadmehr R, Smith MA, Krakauer JW. Error correction, sensory prediction, and adaptation in motor control. *Annu Rev Neurosci*. 2010;33:89–108. <https://doi.org/10.1146/annurev-neuro-060909-153135>
- Wolpert DM, Miall RC, Kawato M. Internal models in the cerebellum. *Trends Cogn Sci*. 1998;2(9):338–47. [https://doi.org/10.1016/S1364-6613\(98\)01221-2](https://doi.org/10.1016/S1364-6613(98)01221-2)
- Blakemore SJ, Wolpert DM, Frith CD. Central cancellation of self-produced tickle sensation. *Nat Neurosci*. 1998;1(7):635–40. <https://doi.org/10.1038/2870>
- Streng ML, Popa LS, Ebner TJ. Modulation of sensory prediction error in Purkinje cells during visual feedback manipulations. *Nat Commun*. 2018;9(1):1099. <https://doi.org/10.1038/s41467-018-03541-0>
- Brooks JX, Carriot J, Cullen KE. Learning to expect the unexpected: rapid updating in primate cerebellum during voluntary self-motion. *Nat Neurosci*. 2015;18(9):1310–7. <https://doi.org/10.1038/nn.4077>
- Apps R, Hawkes R, Aoki S, Bengtsson F, Brown AM, Chen G et al. Cerebellar Modules and Their Role as Operational Cerebellar Processing Units: A Consensus paper. *Cerebellum*. 2018;17(5):654–82. <https://doi.org/10.1007/s12311-018-0952-3>
- Ruigrok TJH. Ins and outs of cerebellar modules. *Cerebellum*. 2011;10(3):464–74. <https://doi.org/10.1007/s12311-010-0164-y>
- Voogd J. Cerebellar zones: a personal history. *Cerebellum*. 2011;10(3):334–50. <https://doi.org/10.1007/s12311-010-0221-6>
- Shinoda Y, Sugihara I, Wu HS, Sugiyuchi Y. The entire trajectory of single climbing and mossy fibers in the cerebellar nuclei and cortex. *Prog Brain Res*. 2000;124:173–86. [https://doi.org/10.1016/S0079-6123\(00\)24015-6](https://doi.org/10.1016/S0079-6123(00)24015-6)
- Oscarsson O. Functional units of the cerebellum - sagittal zones and microzones. *Trends Neurosci*. 1979;2(June):143–5. [https://doi.org/10.1016/0166-2236\(79\)90057-2](https://doi.org/10.1016/0166-2236(79)90057-2)
- Voogd J, Ruigrok TJH. The organization of the corticonuclear and olivocerebellar climbing fiber projections to the rat cerebellar vermis: the congruence of projection zones and the zebrin pattern. *J Neurocytol*. 2004;33(1):5–21. <https://doi.org/10.1023/B:NEUR.0000029645.72074.2b>
- Apps R, Garwicz M. Anatomical and physiological foundations of cerebellar information processing. *Nat Rev Neurosci*. 2005;6(4):297–311. <https://doi.org/10.1038/nrn1646>
- Sugihara I, Shinoda Y. Molecular, topographic, and functional organization of the cerebellar cortex: a study with combined aldolase C and olivocerebellar labeling. *J Neurosci*. 2004;24(40):8771–85. <https://doi.org/10.1523/JNEUROSCI.1961-04.2004>
- De Zeeuw CI, Hoebeek FE, Bosman LWJ, Schonewille M, Witter L, Koekkoek SK. Spatiotemporal firing patterns in the cerebellum. *Nat Rev Neurosci*. 2011;12(6):327–44. <https://doi.org/10.1038/nrn3011>
- Apps R, Hawkes R. Cerebellar cortical organization: a one-map hypothesis. *Nat Rev Neurosci*. 2009;10(9):670–81. <https://doi.org/10.1038/nrn2698>
- Cerminara NL, Lang EJ, Sillitoe R V., Apps R. Redefining the cerebellar cortex as an assembly of non-uniform Purkinje cell microcircuits. *Nat Rev Neurosci*. 2015;16(2):79–93. <https://doi.org/10.1038/nrn3886>
- Zhou H, Lin Z, Voges K, Ju C, Gao Z, Bosman LWJ et al. Cerebellar modules operate at different frequencies. *Elife*. 2014;3:e02536. <https://doi.org/10.7554/eLife.02536.001>
- Xiao J, Cerminara NL, Kotsurovsky Y, Aoki H, Burroughs A, Wise AK et al. Systematic Regional Variations in Purkinje Cell Spiking Patterns. *PLoS One*. 2011;4(9):e105633. <https://doi.org/10.1371/journal.pone.0105633>
- Witter L, De Zeeuw CI. Regional functionality of the cerebellum. *Curr Opin Neurobiol*. 2015;33:150–5. <https://doi.org/10.1016/j.conb.2015.03.017>
- De Zeeuw CI. Bidirectional learning in upbound and downbound microzones of the cerebellum. *Nat Rev Neurosci*. 2021;22(2):92–110. <https://doi.org/10.1038/s41583-020-00392-x>
- Valera AM, Binda F, Pawlowski SA, Dupont J-L, Casella J-F, Rothstein JD et al. Stereotyped spatial patterns of functional synaptic connectivity in the cerebellar cortex. *Elife*. 2016;5:e09862. <https://doi.org/10.7554/eLife.09862.001>
- Spaeth L, Bahuguna J, Gagneux T, Dorgans K, Sugihara I, Poulain B et al. Cerebellar connectivity maps embody individual adaptive behavior in mice. *Nat Commun*. 2022;13(1):580. <https://doi.org/10.1038/s41467-022-27984-8>
- Sillitoe R V, Fu Y, Watson C (2012) *Cerebellum in Mouse Nerv Syst*, Academic Press. 2012. p. 360–97. <https://doi.org/10.1016/B978-0-12-369497-3.10011-1>
- Shambes GM, Gibson JM, Welker W. Fractured somatotopy in granule cell tactile areas of rat cerebellar hemispheres revealed by micromapping. *Brain Behav Evol*. 1978;15(2):94–140. <https://doi.org/10.1159/000123774>
- Ji Z, Hawkes R. Topography of Purkinje cell compartments and mossy fiber terminal fields in lobules II and III of the rat

- cerebellar cortex: pinocerebellar and cuneocerebellar projections. *Neuroscience*. 1994;61:935–54.
28. Biswas MS, Luo Y, Sarpong GA, Sugihara I. Divergent projections of single pontocerebellar axons to multiple cerebellar lobules in the mouse. *J Comp Neurol*. 2019;527:1966–85.
 29. Bengtsson F, Jörntell H. Sensory transmission in cerebellar granule cells relies on similarly coded mossy fiber inputs. *Proc Natl Acad Sci U S A*. 2009;106(7):2389–94. <https://doi.org/10.1073/pnas.0808428106>
 30. Huang C-C, Sugino K, Shima Y, Guo C, Bai S, Mensh BD et al. Convergence of pontine and proprioceptive streams onto multimodal cerebellar granule cells. *Elife*. 2013;2:e00400. <https://doi.org/10.7554/eLife.00400.001>
 31. Pichitpornchai C, Rawson JA, Rees S. Morphology of parallel fibres in the cerebellar cortex of the rat: an experimental light and electron microscopic study with biocytin. *J Comp Neurol*. 2014;342(2):206–20. <https://doi.org/10.1002/cne.903420205>
 32. Harvey R, Napper R. Quantitative studies on the mammalian cerebellum. *Prog Neurobiol*. 1991;36(6):437–63. [https://doi.org/10.1016/0304-0082\(91\)90012-P](https://doi.org/10.1016/0304-0082(91)90012-P)
 33. Mittmann W, Koch U, Häusser M. Feed-forward inhibition shapes the spike output of cerebellar Purkinje cells. *J Physiol*. 2005;563(Pt 2):369–78. <https://doi.org/10.1113/jphysiol.2004.075028>
 34. Jelitai M, Puggioni P, Ishikawa T, Rinaldi A, Duguid I. Dendritic excitation-inhibition balance shapes cerebellar output during motor behaviour. *Nat Commun*. 2016;7:13722. <https://doi.org/10.1038/ncomms13722>
 35. Grangeray-Vilmin A, Valera AM, Kumar A, Isope P. Short-Term Plasticity Combines with Excitation-Inhibition Balance to Expand Cerebellar Purkinje Cell Dynamic Range. *J Neurosci*. 2018;38(22):5153–67.
 36. Ito M. Bases and implications of learning in the cerebellum-adaptive control and internal model mechanism. *Prog Brain Res*. 2005;148:95–109. [https://doi.org/10.1016/S0079-6123\(04\)48009-1](https://doi.org/10.1016/S0079-6123(04)48009-1)
 37. Jörntell H, Hansel C. Synaptic memories upside down: bidirectional plasticity at cerebellar parallel fiber-Purkinje cell synapses. *Neuron*. 2006;52(2):227–38. <https://doi.org/10.1016/j.neuron.2006.09.032>
 38. Marr D. A theory of cerebellar cortex. *J Physiol*. 1969;202(2):437–70. <https://doi.org/10.1113/jphysiol.1969.sp008820>
 39. Albus JS. A theory of cerebellar function. *Math Biosci*. 1971;10(1–2):25–61. [https://doi.org/10.1016/0025-5564\(71\)90051-4](https://doi.org/10.1016/0025-5564(71)90051-4)
 40. Ito M. Cerebellar circuitry as a neuronal machine. *Prog Neurobiol*. 2006;78(3–5):272–303. <https://doi.org/10.1016/j.pneurobio.2006.02.006>
 41. Dean P, Porrill J, Ekerot C-F, Jörntell H. The cerebellar microcircuit as an adaptive filter: experimental and computational evidence. *Nat Rev Neurosci*. 2010;11(1):30–43. <https://doi.org/10.1038/nrn2756>
 42. Cayco-Gajic NA, Silver RA. Re-evaluating Circuit Mechanisms Underlying Pattern Separation. *Neuron*. 2019;101(4):584–602. <https://doi.org/10.1016/j.neuron.2019.01.044>
 43. Chen S, Augustine GJ, Chadderton P. The cerebellum linearly encodes whisker position during voluntary movement. *Elife*. 2016;5:1–16. <https://doi.org/10.7554/eLife.10509.001>
 44. Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R. Encoding of action by the Purkinje cells of the cerebellum. *Nature*. 2015;526(7573):439–42. <https://doi.org/10.1038/nature15693>
 45. Tseng Y-w., Diedrichsen J, Krakauer JW, Shadmehr R, Bastian AJ. Sensory Prediction Errors Drive Cerebellum-Dependent Adaptation of Reaching. *J Neurophysiol*. 2007;98(1):54–62. <https://doi.org/10.1152/jn.00266.2007>
 46. Wagner MJ, Kim TH, Savall J, Schnitzer MJ, Luo L. Cerebellar granule cells encode the expectation of reward. *Nature*. 2017;544(7648):96–100. <https://doi.org/10.1038/nature21726>
 47. Callaway EM, Katz LC. Photostimulation using caged glutamate reveals functional circuitry in living brain slices. *Proc Natl Acad Sci U S A*. 1993;90(16):7661–5. <https://doi.org/10.1073/pnas.90.16.7661>
 48. Katz LC, Dalva MB. Scanning laser photostimulation: a new approach for analyzing brain circuits. *J Neurosci Methods*. 1994;54(2):205–18. [https://doi.org/10.1016/0165-0270\(94\)90194-5](https://doi.org/10.1016/0165-0270(94)90194-5)
 49. Fino E, Araya R, Peterka DS, Salierno M, Etchenique R, Yuste R. RuBi-Glutamate: Two-Photon and Visible-Light Photoactivation of Neurons and Dendritic spines. *Front Neural Circuits*. 2009;3:2. <https://doi.org/10.3389/neuro.04.002.200>
 50. Shepherd GMG. Circuit Mapping by Ultraviolet Uncaging of Glutamate. *Cold Spring Harb Protoc*. 2012;(9):pdb.prot070664–pdb.prot070664. <https://doi.org/10.1101/pdb.prot070664>
 51. Callaway EM (2002) Cell type specificity of local cortical connections. *J Neurocytol*. 2002;31:231–237. <https://doi.org/10.1023/a:1024165824469>
 52. Spaeth L, Gagneux T, Dorgans K, Valera A, Sugihara I, Isope P. Mapping Synaptic Connectivity in the Cerebellar Cortex Using Uncaging. In: Sillitoe RV, editor. *Meas Cerebellar Funct. Humana: Ne*. Springer Nature; 2022. p. 171–85. https://doi.org/10.1007/978-1-0716-2026-7_9
 53. Walter JT, Dizon M-J, Khodakhah K. The functional equivalence of ascending and parallel fiber inputs in cerebellar computation. *J Neurosci*. 2009;29(26):8462–73. <https://doi.org/10.1523/JNEUROSCI.5718-08.2009>
 54. Fujita H, Aoki H, Ajioka I, Yamazaki M, Abe M, Oh-Nishi A et al. Detailed expression pattern of aldolase C (Aldoc) in the cerebellum, retina and other areas of the CNS studied in Aldoc-Venus knock-in mice. *PLoS One*. 2014;9(1):e86679. <https://doi.org/10.1371/journal.pone.0086679>
 55. Cramer SW, Gao W, Chen G, Ebner T. Reevaluation of the beam and radial hypotheses of parallel fiber action in the cerebellar cortex. *J Neurosci*. 2013;33(28):11412–24. <https://doi.org/10.1523/JNEUROSCI.0711-13.2013>
 56. Cohen D, Yarom Y. Unravelling cerebellar circuitry: an optical imaging study. *Prog Brain Res*. 2000;124:107–14. [https://doi.org/10.1016/S0079-6123\(00\)24011-9](https://doi.org/10.1016/S0079-6123(00)24011-9)
 57. Bower JM, Woolston DC. Congruence of spatial organization of tactile projections to granule cell and Purkinje cell layers of cerebellar hemispheres of the albino rat: vertical organization of cerebellar cortex. *J Neurophysiol*. 2003;49(3):745–66. <https://doi.org/10.1152/jn.1983.49.3.745>
 58. Brown IE, Bower JM. Congruence of mossy fiber and climbing fiber tactile projections in the lateral hemispheres of the rat cerebellum. *J Comp Neurol*. 2001;429(1):59–70. [https://doi.org/10.1002/1096-9861\(2000101\)429:1%3C59::aid-cne5%3E3.0.co;2-3](https://doi.org/10.1002/1096-9861(2000101)429:1%3C59::aid-cne5%3E3.0.co;2-3)
 59. Ekerot CF, Jörntell H. Parallel fibre receptive fields of Purkinje cells and interneurons are climbing fibre-specific. *Eur J Neurosci*. 2001;13(7):1303–10. <https://doi.org/10.1046/j.0953-816x.2001.01499.x>
 60. Ekerot C-F, Jörntell H. Parallel fiber receptive fields: a key to understanding cerebellar operation and learning. *Cerebellum*. 2003;2(2):101–9. <https://doi.org/10.1080/14734220309411>
 61. Jörntell H, Ekerot C-F. Reciprocal Bidirectional Plasticity of Parallel Fiber Receptive Fields in Cerebellar Purkinje Cells and Their Afferent Interneurons. *Neuron*. 2002;34(5):797–806. [https://doi.org/10.1016/S0896-6273\(02\)00713-4](https://doi.org/10.1016/S0896-6273(02)00713-4)
 62. Isope P, Barbour B. Properties of Unitary Granule {Cell→Purkinje} Cell Synapses in Adult Rat Cerebellar Slices. *J Neurosci*. 2002;22(22):9668–78. <https://doi.org/10.1523/JNEUROSCI.22-22-09668.2002>

63. Ho S, Lajaunie R, Lerat M, Le M, Crépel V, Loulier K et al. A stable proportion of Purkinje cell inputs from parallel fibers are silent during cerebellar maturation. *Proc Natl Acad Sci U S A*. 2021;118(45). <https://doi.org/10.1073/pnas.2024890118>
64. Fujita M. Adaptive filter model of the cerebellum. *Biol Cybern*. 1982;45(3):195–206. <https://doi.org/10.1007/BF00336192>
65. Giovannucci A, Badura A, Deverett B, Najafi F, Pereira TD, Gao Z, et al. Cerebellar granule cells acquire a widespread predictive feedback signal during motor learning. *Nat Neurosci*. 2017;20(5):727–34. <https://doi.org/10.1038/nn.4531>
66. Knogler LD, Markov DA, Dragomir EI, Štíh V, Portugues R. Sensorimotor Representations in Cerebellar Granule Cells in Larval Zebrafish Are Dense, Spatially Organized, and Non-temporally Patterned. *Curr Biol*. 2017;27(9):1288–302. <https://doi.org/10.1016/j.cub.2017.03.029>
67. Chabrol FP, Arenz A, Wiechert MT, Margrie TW, DiGregorio DA. Synaptic diversity enables temporal coding of coincident multi-sensory inputs in single neurons. *Nat Neurosci*. 2015;18(5):718–27. <https://doi.org/10.1038/nn.3974>
68. Dorgans K, Demais V, Bailly Y, Poulain B, Isope P, Doussau F. Short-term plasticity at cerebellar granule cell to molecular layer interneuron synapses expands information processing. *Elife*. 2019;8:1–24. <https://doi.org/10.7554/eLife.41586.001>
69. Brunel N, Hakim V, Isope P, Nadal J-P, Barbour B. Optimal information storage and the distribution of synaptic weights: perceptron versus Purkinje cell. *Neuron*. 2004;43(5):745–57. <https://doi.org/10.1016/j.neuron.2004.08.023>
70. Miterko LN, Lin T, Zhou J, van der Heijden ME, Beckinghausen J, White JJ et al. Neuromodulation of the cerebellum rescues movement in a mouse model of ataxia. *Nat Commun*. 2021;12(1):1295. <https://doi.org/10.1038/s41467-021-21417-8>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.