



HAL
open science

On the regularity of conical Calabi-Yau potentials

Tran-Trung Nghiem

► **To cite this version:**

| Tran-Trung Nghiem. On the regularity of conical Calabi-Yau potentials. 2022. hal-03817877

HAL Id: hal-03817877

<https://hal.science/hal-03817877v1>

Preprint submitted on 17 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ON THE REGULARITY OF CONICAL CALABI-YAU POTENTIALS

TRAN-TRUNG NGHIEM

ABSTRACT. Using pluripotential theory on degenerate Sasakian manifolds, we show that a locally bounded conical Calabi-Yau potential on a Fano cone is actually smooth on the regular locus. This work is motivated by a similar result obtained by R. Berman in the case where the cone is toric. Our proof is purely pluripotential and independent of any extra symmetry imposed on the cone.

1. INTRODUCTION

1.1. Background and motivation. The problem of finding Kähler-Einstein metrics has been central in the development of Kähler geometry, leading to the solution by Chen-Donaldson-Sun of the celebrated Yau-Tian-Donaldson conjecture [CDS15a, CDS15b, CDS15c]. While the problem is well understood on compact Kähler manifolds, or more generally compact Kähler varieties [EGZ09, Li22], the non-compact case is still relatively open. In the pioneering work of Martelli-Sparks-Yau [MSY08], the existence of conical Calabi-Yau metrics (alias Ricci-flat Kähler cone metrics) on toric varieties with an isolated singularity is shown to be equivalent to a volume minimization principle for Euclidean convex cones. This principle still holds for mildly singular toric varieties as proved by Berman [Ber20]. A more systematic study of polarized affine varieties with an isolated singularity was done by Collins and Székelyhidi [CS19], generalizing the work of Chen-Donaldson-Sun to the context of Kähler cones, or equivalently, Sasakian manifolds.

A Sasakian manifold is a compact Riemannian manifold such that the metric cone over it is Kähler. Sasakian manifolds can be viewed as odd-dimensional analogs of compact Kähler manifolds since they have a natural transverse Kähler structure on an intrinsic horizontal distribution. The existence of Ricci-flat Kähler cone metrics on a Kähler cone is in fact equivalent to the existence of Sasaki-Einstein metrics on the link, which boils down to a Kähler-Einstein-like problem on the transverse structure.

The existence of a (singular) Kähler-Einstein metrics is equivalent to solving a (degenerate) complex Monge-Ampère equation. An interesting problem to ask is the regularity of a singular Kähler-Einstein metric on the smooth locus. In the present paper, we are concerned with the regularity problem on a class of mildly singular affine varieties called *Fano cones*.

In order to state the main result, let us first give some preliminaries on Fano cones and conical Calabi-Yau potentials. Recall that a normal variety is called \mathbb{Q} -Gorenstein if a multiple of its canonical line bundle is Cartier. The action of a complex torus T on Y is said to be *good* if it is effective and has a unique fixed point contained in any orbit closure.

Definition 1.1. *A cone Y is a normal affine variety endowed with the good action of a complex torus $T \simeq (\mathbb{C}^*)^k$. We say that Y is a Fano cone if it is \mathbb{Q} -Gorenstein with klt singularities. The unique fixed point of Y , denoted by 0_Y , is called the vertex of Y .*

Let $\mathcal{M} := \text{Hom}(T, \mathbb{C}^*) \simeq \mathbb{Z}^k$ be the weight lattice and $\mathcal{N} := \mathcal{M}^* = \text{Hom}(\mathbb{C}^*, T)$ the coweight lattice. The ring of regular functions of Y admits a decomposition in to T -modules

$$\mathbb{C}[Y] = \bigoplus_{\alpha \in \Gamma} R_\alpha, \quad \Gamma := \{\alpha \in \mathcal{M}, R_\alpha \neq 0\}$$

where R_α is the T -module with weight α . Let $\mathcal{M}_{\mathbb{R}} := \mathcal{M} \otimes \mathbb{R}$ and $\mathcal{N}_{\mathbb{R}} := \mathcal{N} \otimes \mathbb{R}$. The set Γ is an affine semi-group of finite type which generates a strictly convex polyhedral cone $\sigma^\vee \subset \mathcal{M}_{\mathbb{R}}$. Equivalently, the dual cone σ in $\mathcal{N}_{\mathbb{R}}$ is polyhedral of maximal dimension k . This results from the assumption that Y has a unique fixed point lying in the closure of every T -orbit (cf. [AH06]). The interior of σ is then non-empty and coincides with its relative interior :

$$\text{Int}(\sigma) = \{\xi \in \mathcal{N}_{\mathbb{R}}, \langle \alpha, \xi \rangle > 0, \forall \alpha \in \Gamma\}$$

Definition 1.2. *The interior of the cone σ is called the Reeb cone of Y . An element $\xi \in \text{Int}(\sigma)$ is called a Reeb vector. A Fano cone decorated with a Reeb vector (Y, ξ) is said to be a polarized Fano cone. We say that (Y, ξ) is quasi-regular if $\xi \in \mathcal{N}_{\mathbb{Q}}$, and otherwise irregular if $\xi \notin \mathcal{N}_{\mathbb{Q}}$.*

The closure inside $\text{Aut}(Y)$ of the one-parameter subgroup generated by the infinitesimal action of ξ is a compact torus $T_\xi \subset T_c$, where $T_c \simeq (\mathbb{S}^1)^k$ is a maximal compact subtorus of T . If ξ is quasi-regular then $T_\xi \simeq \mathbb{S}^1$, but if it is irregular then $T_\xi \simeq (\mathbb{S}^1)^m$, $k \geq m > 1$. Equivalently, in the quasi-regular (resp. irregular) case, the holomorphic vector field associated to ξ generates an action of \mathbb{C}^* (resp. $(\mathbb{C}^*)^k$). It can be shown that in the quasi-regular case, the quotient $(Y \setminus \{0_Y\})/\mathbb{C}^*$ is a Fano orbifold (see [Kol04, Paragraph 42]). Note however that in the irregular case, the quotient by $(\mathbb{C}^*)^k$ is only well-defined as an algebraic space (cf. [Kol97]). For more details on Fano cones, the reader may consult for example [LLX20], [DS17] and references therein.

Given a Fano cone (Y, T) , by Sumihiro's theorem (see [Sum74, Theorem 1, Lemma 8]), there exists an embedding $Y \subset \mathbb{C}^N$ such that T corresponds to a diagonal subgroup of $GL_N(\mathbb{C}^N)$ acting linearly. Given an embedding $Y \subset \mathbb{C}^N$, we say that a function f is plurisubharmonic (psh for short) on Y if it is locally the restriction to Y of a psh function on the ambient space \mathbb{C}^N .

Definition 1.3. A ξ -radial function (or ξ -conical potential) $r^2 : Y \rightarrow \mathbb{R}_{>0}$ is a psh function on Y that is invariant under the action of ξ and 2-homogeneous under $-J\xi$, namely

$$\mathcal{L}_\xi r^2 = 0, \quad \mathcal{L}_{-J\xi} r^2 = 2r^2$$

on Y_{reg} .

If Y is a \mathbb{Q} -Gorenstein cone, then for $m > 0$ large enough, mK_Y is a Cartier divisor and naturally linearized by the T -action. Moreover, there exists a T -invariant non-vanishing holomorphic section $s \in mK_Y$ and a volume form dV_Y such that

$$dV_Y = \left(i^{(n+1)^2} m s \wedge \bar{s} \right)^{1/m}$$

where $n+1 = \dim_{\mathbb{C}} Y$. To simplify the notation, by an abuse of language we will sometimes say that s is a “multivalued” section of K_Y and simply write $dV_Y = i^{(n+1)^2} s \wedge \bar{s}$.

A *canonical volume form* dV_Y on Y is a volume form that is $(2n+2)$ -homogeneous under the action of $r\partial r$, namely

$$\mathcal{L}_{r\partial r} dV_Y = 2(n+1)dV_Y$$

on Y_{reg} .

The \mathbb{Q} -Gorenstein and klt singularities assumptions on Y guarantee that there exists a unique canonical volume form on Y up to a constant, see [MSY08], [CS19].

A $(1,1)$ -Kähler current ω on a polarized Fano cone (Y, ξ) is said to be a ξ -Kähler cone current if there exists a locally bounded ξ -radial function such that

$$\omega = dd^c r^2$$

This is well-defined thanks to the local theory of Bedford-Taylor [BT76]. If moreover the function r^2 satisfies the Calabi-Yau condition

$$(1) \quad \omega^{n+1} = (dd^c r^2)^{n+1} = dV_Y$$

in the pluripotential sense, then r^2 is said to be a (singular) *conical Calabi-Yau potential*.

Definition 1.4. We say that a Kähler cone current $\omega = dd^c r^2$ is a *conical Calabi-Yau metric* if the function r^2 is a singular conical Calabi-Yau potential which is smooth on the regular locus of Y .

The motivation for studying these metrics on Fano cones actually has its origin in the compact Fano case. Concretely, Fano cones arise as metric tangent cones of the Gromov-Hausdorff limit of a Fano manifolds sequence [DS17]. If each term of the sequence is moreover Kähler-Einstein, then the Fano cone admits conical Calabi-Yau metrics. As discussed in [Ber20, Section 4] (see also Remark 4.10), it is expected that a singular conical Calabi-Yau potential restricts to a smooth function on the regular locus of Y . Our goal in this article is to give an affirmative answer to this problem.

Theorem 1.5. *Let (Y, ξ) be a polarized Fano cone and r^2 be a singular ξ -conical Calabi-Yau potential on Y . Then r^2 is smooth on the regular locus of Y . In particular, the curvature form of r^2 is a well-defined conical Calabi-Yau metric.*

Such smoothness result is well-known for singular Kähler-Einstein metrics on compact Kähler varieties [EGZ09], [BEGZ10], [BBE⁺19, Lemma 3.6]. In the non-compact setting, when the cone has a unique singularity at the vertex, the Sasakian link is smooth, so the conical metric is automatically smooth outside the vertex. For toric Fano cones with non-isolated singularities, a regularity property was obtained by Berman [Ber20] by using the toric symmetry to reformulate the problem in terms of real Monge-Ampère equations. As discussed in [Ber20, Remark 4.10], the only places where the toric structure was used were the L^∞ -estimate and uniqueness of the Monge-Ampère equation. Although it is possible to generalize the same approach to a larger class of highly symmetric varieties, such as horospherical varieties, we provide a proof closer to the pluripotential spirit and independent of any symmetry other than the given effective torus action. It is an interesting problem to ask if we can weaken the regularity assumption of the solution.

1.2. **Organization.** The organization of the article is as follows.

- In Section 2, we give a quick review of the structure of degenerate Sasakian manifolds. We then gather results in pluripotential theory on these manifolds based on the work of Guedj-Zeriahi [GZ05] and He-Li [HL21]. We also introduce *extremal functions* associated to a Reeb-invariant Borel set on a degenerate Sasakian manifold, which seems to be new in the literature. These objects were not studied in [HL21] in all generality (but see [HL21, Prop. 3.17, Thm. 3.1] for results concerning weighted global extremal functions). The capacity-extremal function comparison is crucial in the proof of the uniform estimate.
- Section 3 is devoted to the proof of our main result. The general strategy is based on [EGZ09], [BEGZ10], [BBE⁺19] and [Ber20]. Let us give a brief explanation. After taking a resolution of singularities, the conical Calabi-Yau problem is translated by pullback to a Calabi-Yau problem on a degenerate Sasakian manifold.

Our key theorem is the uniform L^∞ -estimate of a family of solutions, which relies on a domination-by-capacity property (cf. Prop. 3.7). This, combined with a transverse Yau-Aubin inequality, allows us to obtain a Laplacian estimate of the family, which implies regularity of the solution.

- In Section 4, we provide a proof for the transverse version of Yau-Aubin inequality, which is used in the Laplacian estimate.

Acknowledgements. This article is part of a thesis supervised by Thibaut Delcroix and Marc Herzlich. I wish to thank Vincent Guedj, Eleonora Di Nezza, and Tat-Dat To for their generosity as well as many helpful discussions and remarks. Thanks are also due to the hospitality of the Vietnam Institute for Advanced Study in Mathematics (VIASM), where this work first begun.

2. PLURIPOTENTIAL THEORY ON SASAKIAN MANIFOLDS

2.1. Structure of Sasakian manifolds. In this section, we introduce the notion of *degenerate Sasakian manifolds*. These are compact manifolds having all the essential properties of a Sasakian manifold, except that the form $d\eta$ is not positive-definite, hence does not define a transverse Kähler structure. Still, we assume that a degenerate Sasakian manifold has a transverse Kähler structure, but that the basic Kähler form is not induced by the contact form.

Degenerate Sasakian manifolds arise as the link of the resolution of Fano cones (see Lem. 3.1). The reader should compare this setting to the Kähler situation: a resolution of a Kähler space is still Kähler, but the Kähler structure of the resolution is not the pullback of the Kähler structure on the base.

We refer the reader to [BG08] for a detailed treatment of almost contact structures and Sasakian manifolds.

Let S be a compact differentiable manifold of dimension $2n+1$. A *contact structure* on S is the data of a 1-form η on S such that $\eta \wedge (d\eta)^n \neq 0$. The manifold S is then said to be a *contact manifold*. On a contact manifold, there exists a unique vector field ξ , called the *Reeb vector field*, such that $\eta(\xi) = 1, \mathcal{L}_\xi \eta = 0$. The distribution $\mathcal{D} := \ker(\eta)$ is called the *horizontal distribution* of S .

Definition 2.1. *An almost contact structure is given by (S, ξ, η, Φ) , where η is a contact form, ξ the corresponding Reeb vector field, and Φ a $(1, 1)$ -tensor of TS such that:*

$$\Phi^2 = -Id + \xi \otimes \eta, \quad d\eta(\Phi \cdot, \Phi \cdot) = d\eta, \quad d\eta(\cdot, \Phi \cdot) > 0$$

In particular, $\Phi|_{\mathcal{D}}$ is an almost complex structure.

A degenerate almost contact structure is the same as an almost contact structure, except that $d\eta(\cdot, \Phi \cdot)$ is only semipositive, i.e. $d\eta(\cdot, \Phi \cdot) \geq 0$.

Definition 2.2. *A degenerate metric contact structure is a degenerate almost contact structure (S, ξ, η, Φ) endowed with a Riemannian metric g satisfying*

$$g(\Phi \cdot, \Phi \cdot) = g(\cdot, \cdot) - \eta \otimes \eta$$

Such a metric is said to be compatible.

A (degenerate) almost contact structure is said to be *normal* if the horizontal distribution \mathcal{D} is integrable. A form α on S is said to be *basic* if

$$\mathcal{L}_\xi \alpha = i_\xi \alpha = 0$$

Definition 2.3. A degenerate Sasakian manifold (S, ξ, η, ω_B) is a normal degenerate contact structure with a transverse Kähler metric defined by a basic positive-definite $(1, 1)$ -form ω_B .

Let g_B be the Riemannian metric associated to ω_B . A degenerate Sasakian manifold admits a Riemannian metric, defined by

$$g_S := \eta \otimes \eta + g_B,$$

which restricts to a transverse Kähler metric on \mathcal{D} , but the latter is in general different from the semipositive form induced by the contact form. In particular, a degenerate Sasakian manifold has a degenerate metric contact structure.

Remark. In [BG08], a Sasakian manifold is defined as a *normal metric contact structure*. In our paper, one should distinguish between a *metric contact structure* and an *degenerate metric contact structure*. Both are almost contact structures with a compatible metric, but the metric of the former is exactly $d\eta(Id \otimes \Phi)$, while the latter has a compatible metric $g_S \neq d\eta(Id \otimes \Phi)$.

Many properties of Sasakian manifolds still hold on their degenerate counterparts. For example, on a degenerate Sasakian manifold, we still have a cover by *local foliation charts*, coming from the foliation \mathcal{F}_ξ by the Reeb vector field ξ on S .

Definition 2.4. The foliation atlas on a degenerate Sasakian manifold is defined as a collection of charts (U_α, Φ_α) covering S with diffeomorphisms:

$$\begin{aligned} \Phi_\alpha : W_\alpha \times]-t, t[&\rightarrow U_\alpha \\ (z, x) &\longrightarrow (\varphi_\alpha(z), \tau_\alpha(z, x)) \end{aligned}$$

such that:

- The open interval $]-t, t[\subset \mathbb{R}$ has coordinate x . Here, t can be taken to be independent of α .
- For all α , $W_\alpha \simeq B_\delta(0)$ is the ball of radius $\delta > 0$ centered in $0 \in \mathbb{C}^n$ with coordinates $z = (z_1, \dots, z_n)$. Moreover, the transition map $\varphi_{\alpha\beta} := \varphi_\alpha \circ \varphi_\beta^{-1}$ from $W_\alpha \cap W_\beta$ to itself is holomorphic. In practice, we usually take $\delta = 1$.

Each chart (U_α, Φ_α) is called a foliation chart, and each W_α is said to be a transverse chart (or transverse neighborhood).

In a foliation chart U_α , we may identify ξ with ∂_x and a point $p \in S$ can be written as $p = (z_1, \dots, z_n, x)$.

Let Ω_B^k be the sheaf of basic k -forms on S . Since the exterior differential d on S preserves basic forms, it descends to the *basic exterior differential* $d_B := d|_{\Omega_B^k}$. We then have a subcomplex $\Omega_B(\mathcal{F}_\xi)$ of the de Rham complex, and the corresponding *basic cohomology* H_B^* . The integrable complex structure

on \mathcal{D} leads to the decompositions

$$d_B = \partial_B + \bar{\partial}_B, \quad \Omega_B^k = \bigoplus_{p+q=k} \Omega_B^{p,q}$$

as well as the basic Dolbeault complex and the corresponding cohomologies $H_B^{p,q}$. We then say that a basic function is *transversely holomorphic* if it vanishes under $\bar{\partial}_B$. The Kähler structure on \mathcal{D} induces the decomposition in basic cohomologies as in the classic Hodge theory:

$$H_B^k = \bigoplus_{p+q=k} H_B^{p,q}$$

In short, usual Kähler properties still hold for a Kähler leaf space. We refer the reader to [EKA90] for proofs.

2.2. Quasipsh functions and capacities. We present here some results concerning intrinsic capacities on degenerate Sasakian manifolds, following the lines of Guedj-Zeriahi [GZ05], slightly generalizing the work of He-Li [HL21]. Apart from a subtlety in the definition of capacity, there are generally no supplementary difficulties compared to the case of a classic Sasakian manifold studied by He and Li.

Let (S, ξ, η, ω_B) be a degenerate Sasakian manifold of dimension $(2n+1)$, where ω_B a basic Kähler form on S , while $\theta := d\eta$ is smooth, semipositive and *big*; the latter meaning:

$$0 < \text{vol}_\theta(S) := \int_S \theta^n \wedge \eta < +\infty$$

Let $g_S := \eta \otimes \eta + g_B$ be the corresponding Riemannian metric on S . We denote by

$$\mu_{\omega_B} := \omega_B^n \wedge \eta$$

the volume form on S associated to g_S .

Definition 2.5. *By a ξ -invariant object (function, set, etc.), we mean that the object is invariant under the action of the compact torus T_ξ generated by ξ .*

By a function in $L^1(S)$, we mean a function being L^1 with respect to the measure μ_{ω_B} on S .

A (p, q) -*transverse current* is a collection $\{(W_\alpha, T_\alpha)\}$ where W_α is a transverse neighborhood and T_α a current of bidegree (p, q) on W_α such that

$$\varphi_{\alpha\beta}^* T_\beta|_{W_\alpha \cap W_\beta} = T_\alpha|_{W_\alpha \cap W_\beta}$$

The current T is said to be closed (resp. positive) if each T_α is closed (resp. positive) on W_α . Recall that a basic function on S is a ξ -invariant function. A *basic psh function* u on U_α is a basic, upper-semicontinuous function on U_α such that $u|_{W_\alpha}$ is a classical psh function. In particular, u is locally integrable.

Definition 2.6. We say that a function $u: S \rightarrow \mathbb{R} \cup \{-\infty\}$ is basic θ -psh if u is locally the sum of a basic smooth function and a basic psh function, such that

$$(\theta + d_B d_B^c u)|_{\mathcal{D}} \geq 0$$

in the sense of transverse currents. We will denote by $PSH(S, \xi, \theta)$ the set of basic θ -psh functions. If $u \in PSH(S, \xi, \theta)$, we put $\theta_u := \theta + dd^c u$.

In particular, a θ -psh function is ξ -invariant, upper-semicontinuous and $L^1(S)$. A Sasakian analogue of the Bedford-Taylor theory was developed by van Coevering [vC18] in the case where θ is Kähler and u is a θ -psh bounded function on S . Let us give some details of the construction.

Let $u \in PSH(S, \xi, \theta) \cap L^\infty(S)$ and T a transverse closed positive current on S . Since θ is a closed and basic $(1, 1)$ -form, θ_u defines a transverse $(1, 1)$ -current. After perhaps resizing the transverse neighborhood W_α , there exists a local ξ -invariant potential v such that $\theta = dd^c v$. We then define on each W_α

$$\theta_u \wedge T := dd^c((v + u).T)$$

This allows one to define inductively $\theta_u^k \wedge T$ on each W_α . Passing to the foliation chart $U_\alpha = W_\alpha \times]-t, t[$, the Monge-Ampère operator of u is defined as

$$\theta_u^n \wedge dx$$

where we identify the contact form η with dx in the local coordinate of $]-t, t[$. One can check that this definition is independent of the foliation chart. We will denote the (sasakian) Monge-Ampère measure of u by

$$\text{MA}_\theta(u) := \theta_u^n \wedge \eta$$

In particular, $\text{MA}_\theta(u)$ is a ξ -invariant Radon measure, which has the following continuity property.

Proposition 2.7. [vC18, Theorem 2.3.1] The sasakian Monge-Ampère operator is continuous for monotone convergence. In other words, if $(u_k)_{k \in \mathbb{N}} \subset PSH(S, \xi, \theta) \cap L^\infty(S)$ increases (or decreases) towards u , then $\text{MA}_\theta(u_k) \rightarrow \text{MA}_\theta(u)$ in the sense of measures.

If u is bounded, then by supposing $u \geq 0$ and noting that u^2 is basic and psh, one can define the transverse closed positive current:

$$du \wedge d^c u \wedge T := \frac{1}{2} dd^c u^2 \wedge T - u dd^c \wedge T$$

As in the (transverse) Kähler case, we have for all $u \in PSH(S, \xi, \theta) \cap L^\infty(S)$

$$\int_S \theta_u^n \wedge \eta = \text{vol}_\theta(S)$$

i.e. a locally bounded θ -psh function is of full mass.

We record the following regularization property for a later use:

Lemma 2.8. Given $u \in PSH(S, \xi, \omega_B)$, there exists a sequence $(u_k)_{k \in \mathbb{N}} \subset PSH(S, \xi, \omega_B) \cap C^\infty(S)$ decreasing to u .

Proof. We use the regularization procedure as in [Ber19, Theorem 3.3]. First, for a smooth basic function f and $\beta > 0$, consider the basic Calabi-Yau-type problem on S :

$$(\omega_B + d_B d_B \varphi_\beta)^n \wedge \eta = e^{\beta(\varphi_\beta - f)} \omega_B^n \wedge \eta$$

A solution φ_β verifying $\sup \varphi_\beta = 0$ exists and is unique (cf. [EKA90, 3.5.5]). We will denote by $P_\beta(f)$, $\beta > 0$ the unique solution.

Now let

$$P_{\omega_B}(f)(p) := \sup \{ \varphi(p), \varphi \leq f, \varphi \in PSH(S, \xi, \omega_B) \},$$

This function belongs to $PSH(S, \xi, \omega_B)$ (cf. [HL21, Proposition 3.17]). Consider

$$P'_{\omega_B}(f)(p) := \sup \{ \varphi(p), \varphi \leq f, \varphi \in PSH(S, \xi, \omega_B) \cap C^\infty(S) \}$$

Since u is u.s.c. and basic, it is a decreasing limit of a sequence of smooth basic functions (f_j) . We assert that the sequence $(v_j)_{j \in \mathbb{N}} := (P'_{\omega_B}(f_j))_{j \in \mathbb{N}}$, which consists of basic functions, decreases to u . Indeed, since P'_{ω_B} is a decreasing operator, (v_j) is a decreasing sequence and $f_j \geq v_j \geq u$ by construction. Since $f_j \searrow u$, for all x and $\varepsilon > 0$, there exists j_0 such that for all $j \geq j_0$:

$$u(x) \leq v_j(x) \leq f_j(x) \leq u(x) + \varepsilon$$

hence $v_j(x)$ decreases to $u(x)$.

Arguing as in [Ber19, Proposition 2.3], one can show that the sequence of basic ω_B -psh functions $v_{j,\beta} := P_\beta(f_j)$ converges uniformly to v_j as $\beta \rightarrow \infty$, hence for appropriate $\varepsilon_j \rightarrow 0$, the sequence

$$u_j := v_{j,\beta(j)} + \varepsilon_j$$

which consists of smooth basic ω_B -psh functions, decreases to u . \square

We also have the *comparison principle* for θ -psh functions in the degenerate Sasakian context.

Proposition 2.9. For all $u, v \in PSH(S, \xi, \theta) \cap L^\infty(S)$,

$$\int_{\{v < u\}} MA_\theta(u) \leq \int_{\{v < u\}} MA_\theta(v)$$

Proof. We first prove the following *maximum principle*:

$$1_{\{v < u\}} MA_\theta(\max(u, v)) = 1_{\{v < u\}} MA_\theta(u)$$

It is enough to prove the equality on a foliation chart U_α . First remark that since u, v are both basic, on U_α they depend only on the z -coordinates,

hence $U_\alpha \cap \{v < u\} =]-t, t[\times \{z \in W_\alpha, v < u\}$. Since $\text{MA}_\theta(u)$ is ξ -invariant, it restricts to $\theta_u^n \wedge dx$ on U_α . The equality is then equivalent to

$$1_{]-t, t[\times \{z \in W_\alpha, v < u\}} \theta_{\max(u, v)}^n \wedge dx = 1_{]-t, t[\times \{z \in W_\alpha, v < u\}} \theta_u^n \wedge dx$$

on each foliation chart. By contracting with $\xi = \partial_x$, this is exactly the classical local maximum principle for θ -psh functions.

It follows from the maximum principle that

$$\begin{aligned} \int_{\{v < u\}} \text{MA}_\theta(u) &= \int_S 1_{\{v < u\}} \text{MA}_\theta(\max(u, v)) \\ &= \text{vol}_\theta(S) - \int_{\{v \geq u\}} \text{MA}_\theta(\max(u, v)) \\ &\leq \int_S \text{MA}_\theta(v) - \int_{\{v > u\}} \text{MA}_\theta(\max(u, v)) = \int_{\{v \leq u\}} \text{MA}_\theta(v) \end{aligned}$$

By arguing the same way with $u - \varepsilon$ and v , we obtain

$$\int_{\{v < u - \varepsilon\}} \text{MA}_\theta(u) \leq \int_{\{v \leq u - \varepsilon\}} \text{MA}_\theta(v) \leq \int_{\{v < u\}} \text{MA}_\theta(v)$$

The proof is now concluded by remarking that $\{v < u - \varepsilon\}$ increases to $\{v < u\}$. \square

We record the following result for a later use.

Proposition 2.10. Let $U = B_1(0) \times]-t, t[$ be a foliation chart on S . For every $\varphi \in PSH(S, \xi, \theta) \cap L^\infty(S)$, there exists a unique $\tilde{\varphi} \in PSH(S, \xi, \theta) \cap L^\infty(S)$ such that

$$\text{MA}_\theta(\tilde{\varphi}) = 0 \text{ on } U, \tilde{\varphi} = \varphi \text{ on } S \setminus U, \tilde{\varphi} \geq \varphi \text{ on } S$$

Moreover, if $\varphi_1 \leq \varphi_2$, then $\tilde{\varphi}_1 \leq \tilde{\varphi}_2$.

Proof. The proof is a direct consequence of the local Dirichlet problem on a degenerate Sasakian manifold. The problem can be solved in exactly the same way as in the classical case by remarking that for a basic function u in a foliation chart (z_1, \dots, z_n, x) .

$$(d_B d_B^c u)^n \wedge \eta = \det \left(\frac{\partial^2 u}{\partial z_i \partial \bar{z}_j} \right) \bigwedge_{k=1}^n \frac{i}{2} dz_k \wedge d\bar{z}_k \wedge dx = 0 \iff \det(u_{i\bar{j}}) = 0$$

Hence the local Dirichlet problem on a degenerate Sasakian manifold becomes the classical Dirichlet problem (see [BT76], [BT82] for a proof). \square

Proposition 2.11. Let $(\varphi_j)_{j \in \mathbb{N}} \subset PSH(S, \xi, \theta)^\mathbb{N}$.

1. There exists a constant $C = C(\mu_{\omega_B}, \theta)$ such that for all $u \in PSH(S, \xi, \theta)$:

$$-C + \sup_S u \leq \int_S u d\mu_{\omega_B} \leq \text{vol}_{\omega_B}(S) \sup_S u$$

2. If (φ_j) is uniformly bounded on S , then either (φ_j) converges locally uniformly to $-\infty$, or (φ_j) is relatively compact in $L^1(S)$.

3. If $\varphi_j \rightarrow \varphi$ in $L^1(S)$, then φ coincides almost-everywhere with a function $\varphi^* \in PSH(S, \xi, \theta)$. Moreover,

$$\sup_S \varphi^* = \lim_{j \rightarrow +\infty} \sup_S \varphi_j$$

4. The family

$$\mathcal{F}_0 := \{\varphi \in PSH(S, \xi, \theta), \sup \varphi = 0\}$$

is a compact subset of $PSH(S, \xi, \theta)$.

Proof. For 1), we can adapt the strategy in [HL21, Prop. 3.3] to the non-degenerate Sasakian case. Let us sketch the arguments. We only need to prove the first inequality in the statement (the second one is trivial). Assuming without loss of generality that $\sup_S u = 0$, the inequality then reduces to

$$\int_S u d\mu_{\omega_B} \geq -C$$

There exists two finite covering of S by foliation charts $V_\alpha \subset U_\alpha$ such that $V_\alpha \simeq B_1(0) \times]-t, t[$ is relatively compact in $U_\alpha \simeq B_4(0) \times]-2t, 2t[$. To prove the desired result, it is enough to show that

$$\int_{V_\alpha} u d\mu_{\omega_B} \geq -C_\alpha$$

where $C_\alpha = C_\alpha(\theta)$. But on V_α , this is equivalent to

$$\int_{B_1(0) \times]-t, t[} u d\mu_{z,x} = 2t \int_{B_1(0)} u(z) d\mu_z \geq -C_\alpha$$

where $d\mu_{z,x}$ and $d\mu_z$ are respectively the measures $\omega_B^n \wedge \eta$ and ω_B^n on V_α and $B_1(0)$. Let φ_α be a local potential of θ on $B_4(0)$ (φ_α exists by the $\partial_B \bar{\partial}_B$ -lemma). The function $\varphi_\alpha + u$ is independent of x and psh in $B_4(0)$. By upper-semicontinuity, u attains its local supremum $u(p_1) = 0$ at $p_1 = (z_1, 0) \in B_4(0)$. By the submean inequality on $B_2(z_1) \subset B_4(0)$,

$$(\varphi_\alpha + u)(z_1, 0) = \varphi_\alpha(z_1, 0) \leq \frac{1}{\mu_z(B_2(z_1))} \int_{B_2(z_1)} (\varphi_\alpha + u)(z, 0) d\mu_z$$

Since $u \leq 0$ and $B_1(0) \subset B_2(z_1)$, this completes our proof.

2) is a consequence of 1) (cf. [HL21, Proposition 3.4]).

3) is a consequence of the local result for psh functions (see e.g. [GZ17, Theorem 1.46 (2)]). Indeed, by assumption, on each foliation chart $U_\alpha \simeq B_1(0) \times]-t, t[$, we have $\varphi_j \rightarrow \varphi$ in $L^1_{\text{loc}}(U_\alpha)$. In particular, $\varphi_j \rightarrow \varphi$ in $L^1_{\text{loc}}(B_1(0))$ as psh functions.

4) is a direct consequence of 2) and 3). \square

The following is a Chern-Levine-Nirenberg-type inequality.

Lemma 2.12. Let $v, u \in PSH(S, \xi, \theta)$ such that $0 \leq u \leq 1$. Then

$$0 \leq \int_S |v| \theta_u^n \wedge \eta \leq \int_S |v| \theta^n \wedge \eta + n(1 + 2 \sup v) \text{vol}_\theta(S)$$

Proof. We first suppose that $v \leq 0$. It is enough to establish the equality for $v_k := \max\{v, -k\}$. Indeed, the sequence $-v_k$ increases to $-v$, which allows us to conclude by monotone convergence theorem. Now let us prove the desired result for v_k . It is clear that v_k is θ -psh. We then have the following chain of inequalities:

$$\begin{aligned} \int_S (-v_k) \theta_u^n \wedge \eta &= \int_S (-v_k) \theta_u^{n-1} \wedge (\theta + \sqrt{-1} \partial_B \bar{\partial}_B u) \wedge \eta \\ &= \int_S (-v_k) \theta_u^{n-1} \wedge \theta \wedge \eta + \int_S (-v_k) \theta_u^{n-1} \wedge \sqrt{-1} \partial_B \bar{\partial}_B u \wedge \eta \\ &= \int_S (-v_k) \theta_u^{n-1} \wedge \theta \wedge \eta + \int_S u \theta_u^{n-1} \wedge (-\sqrt{-1} \partial_B \bar{\partial}_B v_k) \wedge \eta \\ &\leq \int_S (-v_k) \theta_u^{n-1} \wedge \theta \wedge \eta + \int_S \theta_u^{n-1} \wedge \theta \wedge \eta \end{aligned}$$

A simple induction allows us to conclude for the case $v \leq 0$. The general case follows by considering $v' := v - \sup_S v$. \square

Definition 2.13. *The capacity of a Borel set $E \subset S$ is defined as:*

$$Cap_\theta(E) := \sup \left\{ \int_E MA_\theta(u), u \in PSH(S, \xi, \theta), 0 \leq u \leq 1 \right\}$$

This definition makes sense since θ is supposed to be big (otherwise Cap would be identically zero). It is clear by definition that $Cap_\theta(\cdot) \geq 0$.

Now let $PSH^-(S, \xi, \theta)$ be the set of negative, basic θ -psh functions.

Proposition 2.14.

- 1) If $\theta_1 \leq \theta_2$ are two basic semipositive $(1, 1)$ -forms on S , then $Cap_{\theta_1}(\cdot) \leq Cap_{\theta_2}(\cdot)$. Moreover, for all $\delta \geq 1$,

$$Cap_\theta(\cdot) \leq Cap_{\delta\theta}(\cdot) \leq \delta^n Cap_\theta(\cdot)$$

For every Borel set $K \subset E$, we have

$$0 \leq Cap_\theta(K) \leq Cap_\theta(E) \leq Cap_\theta(X) = \text{vol}_\theta(X)$$

- 2) For all $v \in PSH^-(S, \xi, \theta)$, there exists a constant $C = C(S, \theta) > 0$ such that :

$$Cap_\theta(v < -t) \leq \frac{C}{t}$$

for all $t > 0$. In particular, $\lim_{t \rightarrow +\infty} Cap_\theta(v < -t) = 0$.

Proof. 1) It is clear that if $\theta_1 \leq \theta_2$ then $MA_{\theta_1}(\cdot) \leq MA_{\theta_2}(\cdot)$ by a property of the complex Hessian in local coordinates. Moreover, if $\theta_1 \leq \theta_2$, then $PSH(S, \xi, \theta_1) \subset PSH(S, \xi, \theta_2)$, so $Cap_{\theta_1} \leq Cap_{\theta_2}$. For all $\delta \geq 1$ and $u \in PSH(S, \xi, \delta\theta)$, $0 \leq u \leq 1$, we have $u \in PSH(S, \xi, \theta)$ and:

$$0 \leq (u/\delta) \leq (1/\delta) \leq 1, \quad (\delta\theta + d_B d_B^c u)^n = \delta^n \left(\theta + \frac{d_B d_B^c u}{\delta} \right)^n$$

Therefore $Cap_{\delta\theta}(\cdot) \leq \delta^n Cap_\theta(\cdot)$ by definition.

For all $K \subset E$ and all candidate function u in the definition of Cap , $\int_K MA_\theta(u) \leq \int_E MA_\theta(u)$, hence $Cap_\theta(K) \leq Cap_\theta(E) \leq Cap_\theta(X)$. Finally, $Cap_\theta(X) = \text{vol}_\theta(X)$ since a locally bounded function has full mass.

2) By the Chern-Levine-Nirenberg inequality in Lem. 2.12, for a θ -psh function u such that $0 \leq u \leq 1$ and $v \in PSH(S, \xi, \theta)$, $v \leq 0$, we have:

$$(2) \quad \int_S (-v)\theta_u^n \wedge \eta \leq \int_S (-v)\theta^n \wedge \eta + n\text{vol}_\theta(S)$$

This inequality allows us to complete the proof. Indeed, for all $u \in PSH(S, \xi, \theta)$ such that $0 \leq u \leq 1$:

$$\begin{aligned} \int_{\{v < -t\}} \theta_u^n \wedge \eta &\leq \frac{1}{t} \int_S (-v)\theta_u^n \wedge \eta \\ &\leq \frac{1}{t} \left(\int_S (-v)\theta^n \wedge \eta + n\text{vol}_\theta(S) \right) \\ &\leq \frac{1}{t} (C(S, \theta) + n\text{vol}_\theta(S)) \text{ (by Prop. 2.11)} \end{aligned}$$

We conclude then by the definition of capacity. \square

The following uniqueness result still holds in the context of degenerate Sasakian manifolds.

Proposition 2.15. Let $u, v \in PSH(S, \xi, \theta) \cap L^\infty(S)$. If

$$MA_\theta(u) = MA_\theta(v)$$

then $u = v + cst$.

Proof. We borrow the proof from [GZ07, Theorem 3.3] (see also [HL21, Theorem 6.4]), which still applies when θ is only semipositive. Let $f = (u - v)/2$ and $h = (u + v)/2$. We can assume that $u, v \geq -C_\theta$ so that $\int_S (-h)\theta_h^{n-1} \wedge \eta \geq 1$. The key idea is to obtain the following inequalities:

$$(3) \quad \int_S d_B f \wedge d_B^c f \wedge \theta_h^{n-1} \wedge \eta \leq \int_S \frac{f}{2} (\theta_u^n - \theta_v^n) \wedge \eta$$

$$(4) \quad \frac{\int_S d_B f \wedge d_B^c f \wedge \theta_h^{n-1} \wedge \eta}{\int_S (-h)\theta_h^n \wedge \eta} \leq 3^n \left(\int_S d_B f \wedge d_B^c f \wedge \theta_h^{n-1} \wedge \eta \right)^{1/2^{n-1}}$$

As a consequence, if $\theta_u^n \wedge \eta = \theta_v^n \wedge \eta$, then combining (3) and (4) yields $\nabla f = 0$, hence $u = v + cst$ as desired. We give a quick proof of (3).

Note that the current under integration on the lhs of (3) is well-defined since u and v are supposed to be bounded. A direct calculation yields

$$\begin{aligned} \int_S d_B f \wedge d_B^c f \wedge \theta_h^{n-1} \wedge \eta &\leq \sum_{k=1}^{n-1} \int_S d_B f \wedge d_B^c f \wedge \theta_u^k \wedge \theta_v^{n-1-k} \wedge \eta \\ &= \sum \int_S f(d_B d_B^c f) \wedge \theta_u^k \wedge \theta_v^{n-1-k} \wedge \eta \\ &= \int_S \frac{f}{2} (\theta_u^n - \theta_v^n) \wedge \eta \end{aligned}$$

The first inequality follows from $C_{n-1}^k \leq 2^{n-1}$, the second one from Stokes' theorem, and the third from the fact that $2d_B d_B^c f = \theta_u - \theta_v$.

The proof of (4) still goes through unchanged. It consists of proving inductively that for $T = \theta_h^l \wedge \theta^{n-2-l} \wedge \eta$, $l = n-2, \dots, 0$, we have

$$\frac{\int_S df \wedge d^c f \wedge \theta \wedge T}{(\int_S (-h)\theta_h^2 \wedge T)^{1/2}} \leq 3 \left(\int_S df \wedge d^c f \wedge \theta_h \wedge T \right)^{1/2}$$

using an integration by parts and Cauchy-Schwartz inequality. \square

2.3. Extremal functions. Motivated by extremal functions in pluripotential theory, we introduce the following counterpart in the Sasakian setting.

Definition 2.16. *Let $K \subset S$ be a ξ -invariant Borel subset. The extremal function associated to θ and K is defined as:*

$$V_{K,\theta}(p) := \sup \{ \varphi(p), \varphi \in PSH(S, \xi, \theta), \varphi \leq 0 \text{ on } K \}$$

Let $V_{K,\theta}^*$ be the u.s.c. regularization of $V_{K,\theta}$. We say that a ξ -invariant Borel set $K \subset S$ is $PSH(S, \xi, \theta)$ -pluripolar if K belongs to the $-\infty$ locus of a basic θ -psh function. Clearly $\{u = -\infty\}$ is ξ -invariant if u is basic θ -psh. Here we impose the symmetry by ξ on K so that there is no inherent contradiction in the definition of pluripolarity. The pluripolarity of K is determined by its extremal function, as the following lemma shows.

Lemma 2.17. *Let $K \subset S$ be a ξ -invariant Borel set.*

- 1) K is $PSH(S, \xi, \theta)$ -pluripolar $\iff V_{K,\theta}^* = +\infty \iff \sup V_{K,\theta}^* = +\infty$.
- 2) If K is not $PSH(S, \xi, \theta)$ -pluripolar, then $V_{K,\theta}^* \in PSH(S, \xi, \theta)$ and $V_{K,\theta}^* = 0$ on $\text{Int}(K)$. Moreover,

$$\int_{\bar{K}} \text{MA}_\theta(V_{K,\theta}^*) = \int_{\bar{K}} (V_{K,\theta}^*)^n \wedge \eta = \text{vol}_\theta(S), \quad \int_{S \setminus \bar{K}} \text{MA}_\theta(V_{K,\theta}^*) = 0$$

Proof. 1) Suppose that $\sup_S V_{K,\theta}^* = +\infty$. By Choquet's lemma, there exists an increasing sequence of functions $\varphi_j \in PSH(S, \xi, \theta)$ such that $\varphi_j = 0$ on K and $V_{K,\theta}^* = (\lim \nearrow \varphi_j)^*$. Up to extracting a subsequence, we can assume that $\sup_S \varphi_j \geq 2^j$. Define $\psi_j := \varphi_j - \sup_S \varphi_j$. The sequence

$\{\psi_j\}_{j \in \mathbb{N}} \subset PSH(S, \xi, \theta)$ is compact and satisfies $\int_S \psi_j d\mu_{\omega_B} \geq -C(\mu_{\omega_B})$ (cf. Lem. 2.11). Let

$$\psi := \sum_{j \geq 1} 2^{-j} \psi_j$$

The function ψ is basic θ -psh as a limit of basic θ -psh functions, and satisfies $\int \psi d\mu_{\omega_B} \geq -C(\mu_{\omega_B})$. It is clear that $\psi_j(x) = -\sup_S \varphi_j$, $\forall x \in K$, hence $K \subset \{\psi = -\infty\}$.

Now suppose that $K \subset \{\psi = -\infty\}$ where $\psi \in PSH(S, \xi, \theta)$. For all $c \in \mathbb{R}$, $\psi + c \in PSH(S, \xi, \theta)$ and $\psi + c \leq 0$ on K . It follows that $V_{K, \theta}^* \geq \psi + c$, hence $V_{K, \theta}^* = +\infty$ on $S \setminus \{\psi = -\infty\}$. Finally, $V_{K, \theta}^* = +\infty$ on S since $\{\psi = -\infty\}$ has zero mass with respect to $\mu_{\omega_B} = \omega_B^n \wedge \eta$.

2) Clearly $V_{K, \theta}^* = 0$ in $\text{Int}(K)$ by definition. The function $V_{K, \theta}$ is basic as the sup-envelope of basic functions, hence its u.s.c. regularization $V_{K, \theta}^*$ is also basic. The fact that $V_{K, \theta}^*$ is θ -psh follows from (3) of Prop. 2.11. Since a locally bounded θ -psh function has full mass, we have

$$\int_{\overline{K}} \text{MA}_\theta(V_{K, \theta}^*) = \int_S \text{MA}_\theta(V_{K, \theta}^*) = \int_S (\theta + d_B d_B^c V_{K, \theta}^*)^n \wedge \eta = \text{vol}_\theta(S)$$

It only remains to show that $\text{MA}_\theta(V_{K, \theta}^*) = 0$ on $S \setminus \overline{K}$, which is equivalent to showing

$$\int_{U_\alpha} \text{MA}_\theta(V_{K, \theta}^*) = 0$$

on each foliation chart $U_\alpha = B_1(0) \times]-t, t[\subset S \setminus \overline{K}$. By Choquet's lemma, there exists an increasing sequence of functions $\varphi_j \in PSH(S, \xi, \theta)$ such that $\varphi_j = 0$ on K and $V_{K, \theta}^* = (\lim \nearrow \varphi_j)^*$. Let $\tilde{\varphi}_j$ the unique solution of local Dirichlet problem with initial datum φ_j (which exists by Prop. 2.10). In particular,

$$\text{MA}_\theta(\tilde{\varphi}_j) = 0 \text{ on } U_\alpha$$

Moreover, the sequence $(\tilde{\varphi}_j)$ is increasing and $\tilde{\varphi}_j = \varphi_j$ on $S \setminus U_\alpha$, hence $\tilde{\varphi}_j = 0$ on K . This shows that $\tilde{\varphi}_j \leq V_{K, \theta}^*$, therefore $\tilde{\varphi}_j \nearrow V_{K, \theta}^*$. By continuity of the Monge-Ampère operator along a monotone sequence (cf. Thm 2.7), $\text{MA}_\theta(V_{K, \theta}^*) = 0$ on U_α . \square

Let us now state an important comparison theorem between capacity and extremal functions.

Lemma 2.18. Let $M_{K, \theta} := \sup_S V_{K, \theta}^*$. For all compact non-pluripolar and ξ -invariant $K \subset S$ we have:

$$1 \leq \text{vol}_\theta(S)^{1/n} \text{Cap}_\theta(K)^{-1/n} \leq \max(1, M_{K, \theta})$$

Proof. The inequality on the left is clear by Prop. 2.14. First suppose that $M_{K, \theta} \leq 1$, then $V_{K, \theta}^*$ is bounded. Since K is non-pluripolar, $V_{K, \theta}^* \in$

$PSH(S, \xi, \theta)$. Moreover, $MA_\theta(V_{K,\theta}^*)$ is supported in K (cf. Lem. 2.17), hence

$$Cap_\theta(K) \geq \int_K MA_\theta(V_{K,\theta}^*) = \int_S MA_\theta(V_{K,\theta}^*) = \text{vol}_\theta(S),$$

which completes the proof in the $M_{K,\theta} \leq 1$ case.

Assume now that $M := M_{K,\theta} \geq 1$. Since the function $V_{K,\theta}^*/M$ is a candidate in the definition of Cap_θ , it follows that

$$\begin{aligned} Cap_\theta(K) &\geq \int_K MA_\theta(M^{-1}V_{K,\theta}^*) \\ &= \int_S MA_\theta(M^{-1}V_{K,\theta}^*) \text{ (by Lem. 2.17)} \\ &\geq M^{-n} \int_S MA_\theta(V_{K,\theta}^*) = M^{-n} \text{vol}_\theta(S) \end{aligned}$$

This allows us to conclude. \square

2.4. Lelong number and integrability. We define the *Lelong number* of a basic psh function u on a foliation chart U_α at a point p with coordinates (z, x) by

$$\nu(u, p) := \lim_{r \rightarrow 0^+} \frac{1}{\log(r) \text{vol}(B(z, r))} \int_{B(z, r)} u(z) \omega_B^n$$

This number does not depend on the foliation chart since the transition maps restrict to biholomorphisms on transverse neighborhoods and that the right-hand side is invariant under biholomorphisms by a theorem of Siu.

It is clear by our definition that the Lelong number is ξ -invariant. Moreover, in a foliation chart $B_1(0) \times]-t, t[$, the function $x \in]-t, t[\rightarrow \nu(u, (z, x))$ is constant for all $z \in B_1(0)$. The Lelong number at a point p on a Sasakian manifold therefore equals to its value at the projection of p to the transverse holomorphic ball of a foliation chart. Local properties of Lelong number can be translated word by word to the Sasakian setting.

Proposition 2.19. The number

$$\nu(\{\theta\}) := \sup \{ \nu(\varphi, x), (\varphi, x) \in PSH(S, \xi, \theta) \times S \}$$

is finite and depends only on the basic cohomology class of θ .

Proof. Since S is compact, there exists a basic Kähler form θ' such that $\theta' \geq \theta$, hence $PSH(S, \xi, \theta') \supset PSH(S, \xi, \theta)$, so $\nu(\{\theta'\}) \geq \nu(\{\theta\})$. It is then enough to prove the assertion when θ is transverse Kähler.

For $p \in S$, we define χ to be a smooth function equals to 1 in a neighborhood of p and 0 outside a larger neighborhood. Let

$$g_p(\cdot) := \chi(\cdot) \log d(\cdot, p)$$

where d is the Riemannian distance associated to θ . It is clear that g_p is smooth on $S \setminus \{p\}$ and psh on a neighborhood of p , hence $A\theta$ -psh for $A > 0$.

Since S is compact, we can choose a uniform constant $A = A(\theta)$ such that for all $p \in S$:

$$dd^c g_p \geq -A\theta$$

By taking average with respect to the action of the compact torus generated by ξ , we can suppose that g_p is ξ -invariant, hence $g_p \in PSH(S, \xi, A\theta)$.

A basic psh function φ in a foliation chart $B_1(0) \times]-t, t[$ restricts to a psh function on the ball $B_1(0)$, so we have

$$\nu(\varphi, 0) = \int_{\{0_z\}} d_B d_B^c \varphi \wedge (d_B d_B^c \log |z|)^{n-1}$$

with 0_z being the center of $B_1(0)$ (see e.g. [GZ17, Lemma 2.46] for a proof). It follows from this local result that for any $a = (z, x)$

$$(5) \quad \nu(\varphi, a) = \int_{\{z\}} \theta_\varphi \wedge (A\theta + d_B d_B^c g_a)^{n-1}$$

The right-hand side is bounded by $\int_S A^n \theta^n \wedge \eta = A^n \text{vol}_\theta(S)$. This completes our proof. \square

Theorem 2.20. *Let $\mathcal{F}_0 := \{\varphi \in PSH(S, \xi, \theta), \sup_S \varphi = 0\}$. If*

$$A < 2\nu(\{\theta\})^{-1},$$

then

$$\sup_{\varphi \in \mathcal{F}_0} \left\{ \int_S e^{-A\varphi} \omega_B^n \wedge \eta \right\} \leq C$$

for a constant C depending only on ω_B and θ .

Proof. We will reduce the problem to the classic Skoda's integrability theorem. First remark that there exist two covers of S by a finite number of foliations charts $(V_j)_{1 \leq j \leq N}$ and $(U_j)_{1 \leq j \leq N}$, where $U_j = B_1(0) \times]-t, t[$, such that $\bar{V}_j \subset U_j$. We need to show that on each foliation chart U_j , there exists a constant $C_j = C(V_j, \mathcal{F}_0, A)$ satisfying

$$\int_{U_j} e^{-A\varphi} \omega_B^n \wedge \eta \leq C_j$$

But since on U_j , φ depends only on the z coordinates and η coincides with dx , it is enough to show that

$$\int_{U_j} e^{-A\varphi} \omega_B^n \wedge \eta = 2t \int_{B_1(0)} e^{-A\varphi} \omega_B^n \leq C_j$$

This follows from the local Skoda's integrability theorem since the family \mathcal{F}_0 is compact (cf. [GZ17, Theorem 2.50] for a proof). \square

3. REGULARITY OF THE POTENTIAL

This part is dedicated to the proof of our main theorem. Let us first give some preliminaries and outline the arguments of the proof. Consider a Fano cone Y of complex dimension $n + 1$ with a good action by $T \simeq (\mathbb{C}^*)^k$. Let $T_c \simeq (\mathbb{S}^1)^k$ be the maximal compact subtorus of T .

Consider a T_c -equivariant embedding of Y into \mathbb{C}^N such that T_c corresponds to a diagonal group acting linearly on \mathbb{C}^N . Recall that ξ generates the action of a compact torus $T_\xi \subset T_c$. Now fix a locally bounded conical Calabi-Yau potential r^2 and a Reeb vector ξ on Y , whose action by T_ξ extends to \mathbb{C}^N through the embedding. Let r_ξ^2 be the radial function on \mathbb{C}^N associated to ξ with conical metric $\omega_\xi = dd^c r_\xi^2$. Then r_ξ^2 restricts to a ξ -conical potential on Y . The link of Y is homeomorphic to the set $Y \cap \{r_\xi^2 = 1\}$. Now let

$$\pi : X \rightarrow Y$$

be a T -equivariant resolution of Y (which exists by Lem. 3.1). Let

$$\mathcal{U} := \pi^{-1}(Y_{\text{reg}})$$

be the open Zariski subset of X isomorphic to Y_{reg} . Consider the following submanifold of X :

$$(S = \pi^{-1}(Y \cap \{r_\xi^2 = 1\}), \xi, \eta, \omega_B)$$

where by an abuse of notation ξ still denotes the pullback of the given Reeb field on \mathbb{C}^N , ω_B is a transverse Kähler form on S (cf. Lem. 3.3), and $\eta = 2\pi^* d^c \log r_\xi^2$ the contact form on S , which is pullback of the contact form associated to ξ on \mathbb{C}^N . Since $d\theta$ is only semipositive, S is degenerate Sasakian. One can show (see Prop. 3.5) that the conical Calabi-Yau equation

$$(dd^c r)^{n+1} = dV_Y$$

is in fact equivalent to the following transverse equation on $\mathcal{U} \cap S$:

$$(\theta_X + d_B d_B^c \varphi_X)^n \wedge \eta = e^{-(n+1)\varphi_X} e^{(n+1)(\Psi_+ - \Psi_-)} \omega_B^n \wedge \eta$$

Here

- $\theta_X := d\eta$,
- $\varphi_X := \pi^* \varphi$, $\varphi := \log(r^2/r_\xi^2)$,
- Ψ_\pm are basic $A\omega_B$ -quasi-psh on S for $A > 0$ large enough,

Remark that θ_X is a semipositive and big form on S . By construction, φ_X is invariant under the induced actions of ξ and $-J\xi$ on X . In a foliation chart (z_1, \dots, z_n, x) of S , the equation can be written as:

$$\det \left(\theta_{X, i\bar{j}} + \frac{\partial^2 \varphi_X}{\partial z_i \partial \bar{z}_j} \right) = e^{-(n+1)\varphi_X(z)} e^{(n+1)(\Psi_+(z) - \Psi_-(z))} \det(\omega_{B, i\bar{j}})$$

The smoothness of $r^2 = r_\xi^2 e^\varphi$ on Y_{reg} is then equivalent to the regularity of $\varphi_X := \pi^* \varphi$ on $S \cap \mathcal{U}$. Consider the family of equations:

$$(\theta_X + \varepsilon \omega_B + d_B d_B^c \varphi_{j,\varepsilon})^n = e^{(n+1)(\psi_{+,j} - \psi_{-,j})} \omega_B^n$$

where $\psi_{\pm,j}$ are two sequences of basic $A\omega_B$ -qpsH functions decreasing to $\psi_+ := \Psi_+$ and $\psi_- := \Psi_- + \varphi_X$ for $A > 0$ large enough. The existence of a unique $\varphi_{j,\varepsilon}$ verifying $\sup \varphi_{j,\varepsilon} = 0$ is guaranteed by the transverse Calabi-Yau theorem of [EKA90]. Finally, to obtain the regularity of φ_X , we proceed by the following classic steps:

- 1) *Uniform estimate:* The functions $\varphi_{j,\varepsilon}$ are uniformly bounded, i.e. there exists a constant C independent of j and ε , such that:

$$\|\varphi_{j,\varepsilon}\|_{L^\infty(S)} \leq C$$

- 2) *Laplacian uniform estimate:* Using the uniform estimate of the previous step, one can show that there exists C' such that for all j, ε ,

$$\sup_{S \cap \mathcal{U}} |\Delta_{\omega_B} \varphi_{j,\varepsilon}| \leq C'$$

where

$$\text{Tr}_{\omega_B} f := n \frac{d_B d_B^c f \wedge \omega_B^{n-1}}{\omega_B^n}$$

- 3) By the complex Evans-Krylov theory, we obtain the following uniform estimate:

$$\|\varphi_{j,\varepsilon}\|_{C^{2,\beta}(S)} \leq C'',$$

which implies $C^{k+2,\beta}$ -estimates for all $k > 0$ by Schauder estimate and a bootstrapping argument.

The last step is classic and well-known in the literature (cf. [Blo12]). Our focus will be mostly on the first and second steps (see Prop. 3.10 and Prop. 3.12).

3.1. Transverse Kähler form. Let V be an irreducible projective variety. Following [Kol07, Paragraph 3], by a *strong resolution* we mean a proper morphism $\pi : V' \rightarrow V$ such that

- V' is smooth and π is birational.
- $\pi : \pi^{-1}(V_{\text{reg}}) \rightarrow V_{\text{reg}}$ is a biholomorphism.
- $\pi^{-1}(V_{\text{sing}})$ is a divisor with simple normal crossings (s.n.c).

In the sense of [Kol07, Paragraph 4], we say that a resolution is *functorial* if for any varieties V, W with resolutions $\pi_V : V' \rightarrow V$, $\pi_W : W' \rightarrow W$, every smooth morphism $\varphi : V \rightarrow W$ can be lifted to a smooth morphism $\varphi' : V' \rightarrow W'$ such that $\pi_W \circ \varphi' = \varphi \circ \pi_V$.

Lemma 3.1. There exists a smooth T -equivariant resolution of singularities $\pi : X \rightarrow Y$.

Proof. Let us embed Y in a T -equivariant manner into \mathbb{C}^N such that T is identified with a diagonal group. Let $\bar{Y} \subset \mathbb{P}^N$ be the closure of Y in \mathbb{P}^N . There exists a T -equivariant resolution $\pi : \bar{X} \rightarrow \bar{Y}$. Indeed, it is enough to take a π as a strong and functorial resolution in the sense of Kollar as recalled above (see [Kol07, Theorem 36] for a proof of existence).

The functoriality of the resolution implies that the action of all algebraic group on \bar{Y} lifts on \bar{X} such that π is equivariant (see [Kol07, Paragraph 9]). We conclude that $\pi : X := \bar{X} \cap \mathbb{C}^N \rightarrow Y$ is a T -equivariant resolution of Y . \square

Now let (X, π) be the resolution of Y , constructed in the previous lemma. Let $E_0 := \pi^{-1}(0_Y)$ be the ‘‘vertex exceptional divisor’’. Since π is equivariant, the vector fields ξ and $-J\xi$ induce by pullback the respective actions on X (still denoted by ξ and $-J\xi$). The action generated by $-J\xi$ is an action of \mathbb{R}_+^* .

The pullback by π of the holomorphic vector field $v_\xi := (-J\xi - \sqrt{-1}\xi)/2$ defines a holomorphic foliation \mathcal{F}_{v_ξ} on $X \setminus E_0$. At every point $p \in X \setminus E_0$, there exists a *transverse holomorphic coordinates* (z_1, \dots, z_n, w) such that

$$v_\xi \cdot z_j = 0, \quad \frac{\partial}{\partial \Im w} = \xi, \quad \frac{\partial}{\partial \Re w} = (-J\xi)$$

which restrict to the coordinate (z, x) on S . In other words, $w = \pi^* \log r_\xi + \sqrt{-1}x$. A form α on $X \setminus E_0$ is said to be *basic* if

$$\mathcal{L}_V \alpha = i_V \alpha = 0, \quad \forall V \in \mathbb{R} \{ \xi, -J\xi \}$$

The restriction map allows us to identify basic forms on $X \setminus E_0$ and basic forms on S .

Lemma 3.2. There exists a T_c -invariant Kähler form ω on X and a global smooth function Φ_ω defined on U such that

$$dd^c \Phi_\omega = \omega, \quad \Phi_\omega \rightarrow -\infty \text{ near } \partial U$$

Proof. Let $\pi : \bar{X} \rightarrow \bar{Y} \subset \mathbb{P}^N$ be the resolution as in the previous lemma. Let $\mathcal{O}(1)$ be the T -linearized hyperplane line bundle of \mathbb{P}^N and E the exceptional divisor of (\bar{X}, π) . Since π is relatively ample, there exists an ample line bundle A on \bar{X} such that:

$$\pi^* \mathcal{O}(1) = A + E$$

Now let $\|\cdot\|_E$ a T_c -invariant metric on $E = \{s_E = 0\}$ and $\varphi_E := -\log \|s_E\|_E^2$ its potential. Let h_A be a T_c -invariant metric of strictly positive curvature on A . Let h be the T_c -invariant metric $h := h_A e^{-\varphi_E}$ on $\pi^* \mathcal{O}(1)$ and $\Phi_\omega := -\log h$ the potential of h . Since X is contained in an open affine set $\simeq \mathbb{C}^N$ of \mathbb{P}^N , there exists a global trivializing T_c -invariant section s of the line bundle $\pi^* \mathcal{O}(1)|_X$. The global form

$$\omega := dd^c \Phi_\omega|_X = -dd^c \log h(s)$$

is clearly closed and positive definite. Indeed, since $s \neq 0$ in X , we have

$$-dd^c \log h(s) = -dd^c \log h_A|_X > 0$$

Finally, since $s_E \rightarrow 0$ near $\partial\mathcal{U}$, we have $\Phi_\omega = -\log h_A + \varphi_E \rightarrow -\infty$ near $\partial\mathcal{U}$. \square

Lemma 3.3. [Ber20, Prop. 4.3] There exists a global smooth function Φ_B on \mathcal{U} satisfying

$$\mathcal{L}_\xi \Phi_B = 0, \mathcal{L}_{-J\xi} \Phi_B = 2, \Phi_B \rightarrow -\infty \text{ near } \partial\mathcal{U}$$

and a transverse basic Kähler form ω_B on $X \setminus E_0$ such that $dd^c \Phi_B = \omega_B$ on \mathcal{U} .

Remark. The information on the behavior of Φ_B near the border of \mathcal{U} is crucial in the Laplacian estimate of the potential φ_X .

Proof. The proof in [Ber20] is an adaptation of the construction of reduced Kähler metrics on a symplectic quotient (see e.g. [BG04, Formulae 4.5, 4.6]). We provide here the details for the reader's convenience.

Remark however that in our case, the symplectic quotient is not well defined since the action generated by ξ on the level set of the hamiltonian is not free in general. However, the construction still applies since it is local in nature.

Let ω be the T_c -invariant Kähler form on X , constructed in Lem. 3.2. Remark that the action generated by ξ is hamiltonian with respect to ω (since by the embedding of Y into \mathbb{C}^N , ξ is identified with a hamiltonian action on \mathbb{C}^N). It follows that there exists a smooth function $\mathcal{H} : X \rightarrow \mathbb{R}$ such that:

$$d\mathcal{H}(\cdot) = -\omega(\xi, \cdot) = g_\omega(-J\xi, \cdot)$$

where g_ω is the metric associated to ω . In particular, $d\mathcal{H}(-J\xi) > 0$, so $d_x \mathcal{H}$ is surjective for $x \notin E_0$. It follows that \mathcal{H} is a submersion for $x \notin E_0$; hence for λ positive, sufficiently large,

$$S_\lambda = \{\mathcal{H} = \lambda\}$$

is a compact submanifold of $X \setminus E_0$, diffeomorphic to $(X \setminus E_0)/\mathbb{R}_+^*$. Now let

$$\pi_\lambda : X \setminus E_0 \rightarrow S_\lambda, \quad i_\lambda : S_\lambda \rightarrow X \setminus E_0$$

be the natural projection and inclusion. Let Φ_ω be the global potential on \mathcal{U} constructed in Lemma 3.2. Let V_p be the neighborhood of a point $p \in X \setminus E_0$ with local transverse coordinates (z, w) . Consider the following ξ -invariant function on $S_\lambda \cap \mathcal{U}$:

$$\Phi_\lambda := i_\lambda^*(\Phi_\omega - \lambda \Im w)$$

The function

$$(6) \quad \Psi_B = \pi_\lambda^* \Phi_\lambda + \lambda \Im w = \pi_\lambda^* \Phi_\omega|_{S_\lambda} + \lambda(\Im w - i_\lambda^* \Im w)$$

is then ξ -invariant on V_p and well-defined on V_p . Indeed, let $V_{p'}$ be another local transverse neighborhood of a point $p' \in S_\lambda \cap V_p$. By the definition of w , $v_\xi(w - w') = 0$, so there exists a basic transversely holomorphic function $f(z)$ on $V_p \cap V_{p'}$ such that $w - w' = f(z)$. It follows that:

$$\mathfrak{S}(w - w')|_{V_p \cap V_{p'}} = \mathfrak{S}(w - w')|_{S_\lambda \cap V_p \cap V_{p'}} = i_\lambda^* \mathfrak{S}(w - w')|_{V_p \cap V_{p'}}$$

By construction, we have $\mathcal{L}_{-J\xi}\Psi_B = \lambda$, hence Ψ_B extends uniquely to a smooth function on \mathcal{U} . The function

$$\Phi_B := 2(\Psi_B/\lambda)$$

satisfies $\mathcal{L}_\xi\Phi_B = 0$, $\mathcal{L}_{-J\xi}\Phi_B = 2$. We assert that the following global form on \mathcal{U}

$$\omega_B := dd^c\Phi_B$$

defines a transverse Kähler metric on \mathcal{U} . By a direct computation from the equation (6) as in [BG04, Section 9], $2\lambda^{-1}\omega$ is exactly ω_B on S_λ . After replacing Φ_B with $2\lambda^{-1}\Phi_\omega$ on each V_p , we see that ω_B extends to a transverse Kähler metric on $X \setminus E_0$.

It remains to show that $\Phi_B \rightarrow -\infty$ on $\partial\mathcal{U}$. Indeed, on $\mathcal{U} \cap V_p$, $\Phi_B - 2\lambda^{-1}\Phi_\omega = \mathfrak{S}w - i_\lambda^*(\mathfrak{S}w)$ for all $p \in S_\lambda$. It follows that $\Phi_B - 2\lambda^{-1}\Phi_\omega$ is bounded on $S_\lambda \cap \mathcal{U}$, so $\Phi_B = (\Phi_B - 2\lambda^{-1}\Phi_\omega) + 2\lambda^{-1}\Phi_\omega \rightarrow -\infty$ near $\partial\mathcal{U}$ since $\Phi_\omega \rightarrow -\infty$ near $\partial\mathcal{U}$. \square

Since X is a T_c -invariant resolution of Y and that Y has klt singularities, there exists a T_c -invariant divisor D such that:

$$\pi^*K_Y = K_X + D, \quad D = \sum_{a_j > -1} a_j D_j,$$

We have moreover a decomposition $D = D_+ - D_-$ where:

$$D_+ := \sum_{a_j > 0} D_j, \quad D_- := \sum_{a_j < 0} (-a_j) D_j$$

are two effective T_c -invariant \mathbb{Q} -divisors. There exists then a T_c -invariant volume form dV_X on X , two multivalued sections s_\pm and hermitian T_c -invariant metrics h^\pm on D_\pm , such that:

$$(7) \quad \pi^*dV_Y = \|s_+\|_{h^+}^2 \|s_-\|_{h^-}^{-2} dV_X$$

To be precise, we may choose:

$$\|s_+\|_{h^+}^2 := \prod_{a_j > 0} |s_j|_{h_j}^{2a_j}, \quad \|s_-\|_{h^-}^{-2} := \prod_{a_j < 0} |s_j|_{h_j}^{-2a_j}$$

where h_j are T_c -invariant hermitian metrics of the fiber $\mathcal{O}_X(D_j)$. Up to a positive constant, we have the following volume form on S :

$$dV_X(-J\xi, \cdot) = \omega_B^n \wedge \eta$$

Lemma 3.4. There exist two basic psh T_c -invariant functions Ψ_{\pm} on S , smooth on \mathcal{U} and a constant $A > 0$ such that on S ,

$$\pi^* dV_Y(-J\xi, \cdot) = e^{(n+1)(\Psi_+ - \Psi_-)} \omega_B^n \wedge \eta, \quad \frac{i}{2\pi} \partial_B \bar{\partial}_B \Psi_{\pm} \geq -A\omega_B$$

Moreover, $e^{-\Psi_-} \in L^p(S), p > 1$.

Proof. Assume that there exists a positive constant $C > 0$ satisfying:

$$(8) \quad \frac{i}{2\pi} \partial_B \bar{\partial}_B \log \|s_{\pm}\|_{h_{\pm}|_S}^2 \geq -C\omega_B$$

Then by choosing Ψ_{\pm} such that:

$$(n+1)\Psi_{\pm} = \log \|s_{\pm}\|_{h_{\pm}|_S}^2 \in C_B^{\infty}(S)$$

we obtain the equality between volume forms from (7) and the estimate of $\partial_B \bar{\partial}_B \Psi_{\pm}$ follows immediately.

It remains to prove (8). By definition of s_{\pm} and $\|\cdot\|_{h_{\pm}}$, in a transverse holomorphic chart of $X \setminus E_0$ with coordinates (z, w) , there exist T_c -invariant local potentials φ_{\pm} and holomorphic T_c -semi-invariant local functions f_{\pm} such that:

$$\|s_{\pm}\|_{h_{\pm}} = |f_{\pm}(z, w)| e^{-\varphi_{\pm}(z, w)}$$

In particular, there exist $\lambda_{\pm} \in \mathbb{R}$ satisfying:

$$\frac{\partial}{\partial \Im w} f_{\pm} = i\lambda_{\pm} f_{\pm}$$

After replacing f_{\pm} by $f_{\pm} e^{-\lambda_{\pm} w}$, one can suppose that f_{\pm} are ξ -invariant (hence basic), so $\bar{\partial}_B f_{\pm} = 0$. It follows that f_{\pm} are transversely holomorphic, hence $d_B d_B^c \log |f_{\pm}(z, w)|^2 \geq 0$, so locally:

$$d_B d_B^c \log \|s_{\pm}\|_{h_{\pm}|_S}^2 \geq -C d_B d_B^c \varphi_{\pm}$$

for some constant C depending only on the local open set. Moreover, since ω_B is Kähler, one can find in a transverse neighborhood a constant $A > 0$ (which depends only on the neighborhood) such that

$$d_B d_B^c \varphi_{\pm} \leq A\omega_B$$

The compactness of S then completes the proof of (8). Finally, since Y has klt singularities, D_j are normal crossing divisors, hence there exists $p > 1$ such that $pa_j > -1$ for all j , so $e^{-\Psi_-} \in L^p(S)$ for some $p > 1$. \square

3.2. Transverse Monge-Ampère equation.

Proposition 3.5. The conical potential r is a solution in the pluripotential sense of the equation:

$$(9) \quad (dd^c r^2)^{n+1} = dV_Y$$

on Y_{reg} if and only if φ_X satisfies the following equation on $S \cap \mathcal{U}$:

$$(10) \quad (\theta_X + d_B d_B^c \varphi_X)^n \wedge \eta = e^{-(n+1)\varphi_X} e^{(n+1)(\Psi_+ - \Psi_-)} \omega_B^n \wedge \eta$$

In particular, in a transverse holomorphic neighborhood $S \cap \mathcal{U}$,

$$(\theta_X + d_B d_B^c \varphi_X)^n = e^{-(n+1)\varphi_X} e^{-(n+1)(\Psi_+ - \Psi_-)} \omega_B^n$$

Proof. By definition $\Phi = \log r^2$, hence:

$$dd^c r^2 = e^\Phi (dd^c \Phi + d\Phi \wedge d^c \Phi) = r^2 (dd^c \Phi + d\Phi \wedge d^c \Phi)$$

in the current sense. We have

$$(dd^c \Phi + d\Phi \wedge d^c \Phi)^{n+1} = \sum c_{k,n} (dd^c \Phi)^k \wedge (d\Phi \wedge d^c \Phi)^{n-k} = (dd^c \Phi)^n \wedge d\Phi \wedge d^c \Phi$$

Indeed, in the transverse coordinates (z, w) on $X \setminus E_0$,

$$\frac{\partial \Phi}{\partial w} = \frac{\partial \Phi}{\partial \bar{w}} = 1$$

hence $(dd^c \Phi)^{n+1} = 0$. It follows that

$$(dd^c r^2)^{n+1} = dV_Y \iff r^{2n+2} (dd^c \Phi)^n \wedge d\Phi \wedge d^c \Phi = dV_Y$$

Since

$$\mathcal{L}_\xi \Phi = 0,$$

the restriction of Φ in S is basic. It follows that

$$\begin{aligned} (dd^c \Phi)^n \wedge d\Phi \wedge d^c \Phi &= \det \left(\frac{\partial^2 \Phi}{\partial z_l \partial \bar{z}_m} \right) \bigwedge (i/2) dz_k \wedge d\bar{z}_k \wedge d\Phi \wedge d^c \Phi \\ &= (dd^c \Phi)^n \wedge (dw + d\bar{w}) \wedge (d^c w + d^c \bar{w}) \\ &= (dd^c \Phi)^n \wedge 2d\Re w \wedge 2d^c \Re w \end{aligned}$$

The conical Calabi-Yau equation then becomes

$$r^{2n+2} (dd^c \Phi)^n \wedge 2d\Re w \wedge 2d^c \Re w = dV_Y$$

By contracting the equality with $-J\xi$, and using $2d\Re w(-J\xi) = 1$, we have:

$$r^{2n+2} (dd^c \Phi)^n \wedge 2d^c \Re w = dV_Y(-J\xi) = \omega_B^n \wedge \eta$$

By using $dd^c \Phi = \theta + dd^c \varphi = \theta + d_B d_B^c \varphi$, $2d^c \Re w = \eta$, the previous lemma and the fact that $S = X \cap \pi^{-1}(\{r_\xi^2 = 1\})$, we obtain by pullback the following equation on $S \cap \mathcal{U}$:

$$(\theta_X + d_B d_B^c \varphi_X)^n \wedge \eta = e^{-(n+1)\varphi_X} e^{-(n+1)(\Psi_+ - \Psi_-)} \omega_B^n \wedge \eta$$

Finally by applying i_ξ and using that $\eta(\xi) = 1$, the equation on $S \cap \mathcal{U}$ becomes

$$(\theta_X + d_B d_B^c \varphi_X)^n = e^{-(n+1)\varphi_X} e^{-(n+1)(\Psi_+ - \Psi_-)} \omega_B^n$$

The converse is proved in the same manner. \square

3.3. Uniform estimate. Let $\psi_{\pm,j}$ be two sequences of smooth basic quasi-psh functions which decrease to

$$\psi_+ := \Psi_+, \quad \psi_- := \Psi_- + \varphi_X,$$

such that :

$$(11) \quad d_B d_B^c \psi_{\pm,j} \geq -C\omega_B$$

for a uniform constant C independent of j . Such a sequence exists by virtue of Lem. 2.8.

Let $\varepsilon > 0$. Recall that the form $\theta_X = \pi^* dd^c \log r_\xi^2$ is semi-positive, big and basic, hence $\theta_X + \varepsilon\omega_B$ is a transverse Kähler form. Consider the following equation on S for a smooth basic $(\theta_X + \varepsilon\omega_B)$ -psh function $\varphi_{j,\varepsilon}$:

$$(12) \quad (\theta_X + \varepsilon\omega_B + d_B d_B^c \varphi_{j,\varepsilon})^n \wedge \eta = e^{(n+1)(\psi_{+,j} - \psi_{-,j})} \omega_B^n \wedge \eta$$

By the transverse Calabi-Yau theorem of El-Kacimi Alaoui [EKA90, 3.5.5], for all j, ε , there exists a unique basic solution satisfying:

$$\sup \varphi_{j,\varepsilon} = 0$$

Now let μ_j be the smooth volume form $e^{(n+1)(\psi_{+,j} - \psi_{-,j})} \omega_B^n \wedge \eta$ on S . The following lemma is elementary:

Lemma 3.6. Let μ be an inner-regular positive Borel measure on S . Then for all ξ -invariant Borel set $E \subset S$,

$$\mu(E) = \sup \{ \mu(K), K \subset E \text{ compact, } \xi - \text{invariant} \}$$

In particular, μ_j satisfies this property.

Proof. It is enough to show that for all $j \in \mathbb{N}^*$, there exists a compact ξ -invariant K_j such that:

$$\mu(E) \leq \mu(K_j) + \frac{1}{j}$$

By inner regularity of E , there exists a compact $C_j \subset E$ such that:

$$\mu(E) \leq \mu(C_j) + 1/j$$

The idea is to average C_j by the action of T_ξ . We define

$$K_j := \cup_{g \in T_\xi} g.C_j = T_\xi.C_j$$

For each j , the set K_j is compact and ξ -invariant by construction. Moreover, $K_j \subset E$ since $g.C_j \subset g.E \subset E$. Finally, the fact that $C_j \subset K_j$ implies $\mu(C_j) \leq \mu(K_j)$. This completes our proof. \square

We also have the important *domination by capacity* property of the measures μ_j .

Proposition 3.7. The measures μ_j satisfy the $\mathcal{H}(\alpha, A, \theta)$ condition for all α . Namely, for all $\alpha > 0$, there exists a constant A independent of j such that:

$$\mu_j(E) \leq ACap_\theta(E)^{1+\alpha}$$

for all ξ -invariant Borel subset $E \subset S$.

Proof. By inner regularity of μ_j , it is enough to establish the lemma for a compact ξ -invariant $K \subset S$. Indeed, suppose that the inequality is true for all such K , then for all Borel ξ -invariant set E ,

$$\begin{aligned} \mu_j(E) &= \sup \{ \mu_j(K), K \subset E \text{ compact, } \xi - \text{invariant} \} \\ &\leq A \sup \{ Cap_\theta(K)^{1+\alpha}, K \subset E \text{ compact, } \xi - \text{invariant} \} \\ &\leq ACap_\theta(E)^{1+\alpha} \text{ (by Prop. 2.14(1))} \end{aligned}$$

We can suppose furthermore that K is non-pluripolar (otherwise $\mu_j(K) = 0$ and the inequality is then trivial).

Now let K be a compact ξ -invariant and non-pluripolar. Let $p > 1$ be as in Lemma 3.4. By Hölder inequality, we have:

$$0 \leq \mu_j(K) \leq \|f_j\|_{L^p(\omega_B^n \wedge \eta)} \text{vol}_{\omega_B}(K)^{1/q}$$

where $1/p + 1/q = 1$. Since $\psi_{+,j} \leq \psi_{+,1}$ and that $\psi_j \geq \psi_-$, the function $e^{(n+1)(\psi_{+,j} - \psi_{-,j})}$ is bounded in L^p by $e^{(n+1)(C - \psi_-)}$, where $C := \sup_S \psi_{+,1}$. It follows that the norm $\|f_j\|_{L^p(\omega_B^n \wedge \eta)}$ is uniformly bounded, therefore it is enough to show that

$$\text{vol}_{\omega_B}(K) \leq C \exp\left(-\gamma(Cap_\theta(K))^{-1/n}\right)$$

where $C = C(\theta, \omega_B)$, $\gamma = \gamma(\theta)$ are constants independent of j . The conclusion then follows from the elementary equality $\exp(-x^\beta) \leq A_\alpha x^\alpha$, for all $x \in [0, 1]$, $\alpha > 0$.

By Theorem 2.20, for $\gamma := 2/(\nu(\{\theta\}) + 1)$, there exists a constant $C = C(\theta, \omega_B)$ such that:

$$\sup_{\psi \in \mathcal{F}_0} \int_S \exp(-\gamma\psi) \omega_B^n \wedge \eta \leq C$$

In particular, for $\psi := V_{K,\theta}^* - M_{K,\theta}$ (recall that $M_{K,\theta} = \sup V_{K,\theta}^*$), we obtain

$$\int_S \exp(-\gamma V_{K,\theta}^*) \omega_B^n \wedge \eta \leq C \exp(-\gamma M_{K,\theta})$$

Note that $V_{K,\theta}^*$ is well defined thanks to the ξ -invariance of K . Finally, since $V_{K,\theta}^* \leq 0$ μ_{ω_B} -a.e. on K , we have

$$\text{vol}_{\omega_B}(K) \leq C \exp(-\gamma M_{K,\theta})$$

An application of Lemma 2.18 then completes our proof. \square

Let us first establish some more useful lemmas before proving the uniform estimate.

Lemma 3.8. Let $u \in PSH(S, \xi, \theta) \cap L^\infty(S)$ be a negative function. For all $s \geq 0, 0 \leq t \leq 1$,

$$t^n \text{Cap}_\theta(u < -s - t) \leq \int_{\{u < -s\}} \theta_u^n \wedge \eta$$

Proof. Let $v \in PSH(S, \xi, \theta), 0 \leq v \leq 1$. Then

$$\{u < -s - t\} \subset \{u \leq tv - s - t\} \subset \{u < -s\}$$

By definition of the Monge-Ampère operator

$$\int_{\{u < -s - t\}} \text{MA}_\theta(v) \leq \int_{\{u \leq tv - s - t\}} \text{MA}_\theta(v) \leq t^{-n} \int_{\{u \leq tv - s - t\}} \text{MA}_\theta(tv)$$

Applying the comparison principle 2.9 to the functions $u + s + t$ and tv ,

$$t^{-n} \int_{\{u \leq tv - s - t\}} \text{MA}_\theta(tv) \leq t^{-n} \int_{\{u \leq tv - s - t\}} \text{MA}_\theta(u) \leq \int_{\{u < -s\}} \text{MA}_\theta(u)$$

which terminates our proof. \square

Lemma 3.9. [EGZ09, Lem. 2.4] Let $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a right-continuous decreasing function such that $\lim_{s \rightarrow +\infty} f(s) = 0$. If f satisfies the condition

$$H(\alpha, B), \quad tf(s+t) \leq Bf(s)^{1+\alpha}, \quad \forall s \geq 0, 0 \leq t \leq 1$$

then there exists $s_0 = s_0(\alpha, B)$ such that $f(s) = 0, \forall s \geq s_0$.

Proposition 3.10. There exists a uniform constant C such that:

$$\|\varphi_{j,\varepsilon}\|_{L^\infty(S)} \leq C$$

Proof. Let $f(s) := \text{Cap}_\theta(\varphi_{j,\varepsilon} < -s)^{1/n}$. It is clear that $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is right-continuous, and $\lim_{s \rightarrow +\infty} f(s) = 0$ (cf. Prop. 2.14). Moreover, f is decreasing: for all $t > s$, $\{\varphi < -t\} \subset \{\varphi < -s\}$, $\forall t > s$, hence $f(t) \leq f(s)$. Following Lem. 3.8 and the fact that μ_j satisfy $\mathcal{H}(\alpha, A, \theta)$, f satisfies the condition $H(\alpha, B)$ with $B = A^{1/n}$. Indeed,

$$\begin{aligned} t^n f(s+t)^n &\leq t^n \text{Cap}_{\theta + \varepsilon \omega_B}(\varphi_{j,\varepsilon} < -s - t) \\ &\leq \int_{\{\varphi_{j,\varepsilon} < -s\}} (\theta + \varepsilon \omega_B + d_B d_B^c \varphi_{j,\varepsilon})^n \wedge \eta \\ &= \int_{\{\varphi_{j,\varepsilon} < -s\}} \mu_j \leq A \text{Cap}_\theta(\varphi_{j,\varepsilon} < -s)^{1+\alpha} = A f(s)^{n(1+\alpha)} \end{aligned}$$

The first inequality follows from Lem. 2.14, the second is direct from Lem. 3.8, while the fourth is a consequence of Lem. 3.7. Now let $\omega_\varepsilon := \theta_X + \varepsilon \omega_B$. For ε sufficiently small and δ large enough, there exists $\delta = \delta(S) \geq 1$ such that $\omega_\varepsilon \leq \delta \omega_B$. In particular, $\varphi_{j,\varepsilon} \in PSH^-(S, \xi, \delta \omega_B)$. Again by Lem. 2.14,

$$\begin{aligned} f(s)^n &\leq \text{Cap}_{\delta \omega_B}(\varphi_{j,\varepsilon} < -s) \\ &\leq \frac{\delta^n}{s} \left(\int_S (-\varphi_{j,\varepsilon}) \omega_B^n \wedge \eta + n \text{vol}_{\omega_B}(S) \right) \end{aligned}$$

But by (1) of Lem. 2.11:

$$\int_S -\varphi_{j,\varepsilon} d\mu_{\omega_B} \leq -\sup \varphi_{j,\varepsilon} + C(\omega_B) = C(\omega_B)$$

Therefore, $f(s) \leq (C_1/s^{1/n})$, where $C_1 = C_1(\omega_B, \theta_X)$. We can then apply Lem. 3.9 to select $s_0 = s_0(n, \alpha, A, \omega_B, \theta_X)$ as in [EGZ09, Lemma 2.3, Theorem 2.1] such that:

$$\text{Cap}_{\theta_X}(\varphi_{j,\varepsilon} < -s) = 0, \forall s \geq s_0$$

In particular, $\mu_j(\varphi_{j,\varepsilon} < -s_0) = 0$ by Lem. 3.7. Hence $\varphi_{j,\varepsilon} \geq s_0$ on S , so there exists $C = C(n, \alpha, A, \omega_B, \theta_X)$ such that:

$$\|\varphi_{j,\varepsilon}\|_{L^\infty(S)} \leq C$$

□

3.4. Laplacian estimate. We will need the transverse version of the Yau-Aubin inequality, obtained by Siu for two cohomologous forms [Siu87], but the proof can be generalized to any couple of Kähler forms. Let

$$\Delta_{\omega'_B} := \text{Tr}_{\omega'_B} d_B d_B^c$$

be the Laplacian associated to the transverse Kähler form ω'_B .

Lemma 3.11. For all transverse Kähler form ω'_B , there exists a constant κ depending only on the transverse bisectional curvature of ω_B such that:

$$\Delta_{\omega'_B} \log \text{Tr}_{\omega_B} \omega'_B \geq -\kappa \text{Tr}_{\omega'_B} \omega_B - \frac{\text{Tr}_{\omega_B} \text{Ric}(\omega'_B)}{\text{Tr}_{\omega_B} \omega'_B}$$

where $\text{Ric}(\omega'_B)$ is the transverse Ricci curvature.

Proof. On each foliation chart, the transverse Kähler forms depend only on the z -coordinates. The inequality thus follows from the purely local proof in the compact Kähler case. The reader may consult Appendice 4 for a proof. □

The following proposition gives a *a priori* Laplacian estimate of the solution $\varphi_{j,\varepsilon}$ of equation (12). We follow the arguments of [BBE⁺19, Appendice B]. In this section, by a *uniform constant*, we mean a constant independent of the j, ε parameters.

Proposition 3.12. Let $\psi := \Phi_B - r_\xi^2$ and $\omega_\varepsilon = \theta_X + \varepsilon\omega_B$, $\omega'_\varepsilon := \omega_\varepsilon + d_B d_B^c \varphi_{j,\varepsilon}$. There exist uniform constants C_1, C_2 such that:

$$\sup_{S \cap \mathcal{U}} \text{Tr}_{\omega_\varepsilon} \omega'_\varepsilon \leq C_2 e^{-C_1 \psi - \psi_-} \leq C_2 e^{-C_1 \psi - \psi_-}$$

In particular, there exists a uniform constant C_3 such that

$$\sup_{S \cap \mathcal{U}} |\Delta_{\omega_B} \varphi_{j,\varepsilon}| \leq C_3 e^{-C_1 \psi - \psi_-}$$

Proof. The function ψ is clearly basic θ_X -psh and $\psi \rightarrow -\infty$ near $\partial\mathcal{U}$ by the construction of Φ_B in Prop. 3.3. Moreover, $\omega_B|_{\mathcal{U}} = (\theta_X + dd^c\psi)|_{\mathcal{U}}$ is the restriction into \mathcal{U} of the transverse Kähler form ω_B , constructed on $X \setminus E_0$.

Consider the following smooth function on $S \cap \mathcal{U}$:

$$h := \log(\mathrm{Tr}_{\omega_\varepsilon} \omega'_\varepsilon) + n\psi_{-,j} - A_1(\varphi_{j,\varepsilon} - \psi)$$

where $A_1 := A_1(\kappa)$ is a constant sufficiently large and depends on κ . The compacity of S , the L^∞ -estimate in Prop. 3.10, combined with transverse Yau-Aubin inequality in Lem. 3.11 are all the ingredients we need to repeat the arguments of [BBE⁺19, Appendice B] to conclude.

For the reader's convenience, we provide here some details of the proof. By the transverse Yau-Aubin inequality, we have on $S \cap \mathcal{U}$:

$$\Delta_{\omega'_\varepsilon} h \geq \mathrm{Tr}_{\omega'_\varepsilon}(\omega_\varepsilon) - A_2$$

where A_2 depends only on A_1 and n . Since $\varphi_{j,\varepsilon}$ is uniformly bounded and that $\psi \rightarrow -\infty$ near $\partial(S \cap \mathcal{U})$, h attains its maximum at $x_0 \in S \cap \mathcal{U}$. It follows from the maximum principle that

$$0 \geq \Delta_{\omega'_\varepsilon} h(x_0) \geq \mathrm{Tr}_{\omega'_\varepsilon}(\omega_\varepsilon)(x_0) - A_2$$

By local elementary reasonings as in the compact Kähler case, we obtain the following inequality for two transverse Kähler forms:

$$\mathrm{Tr}_{\omega_\varepsilon}(\omega'_\varepsilon) \leq n \frac{(\omega'_\varepsilon)^n}{\omega_\varepsilon^n} (\mathrm{Tr}_{\omega'_\varepsilon}(\omega_\varepsilon))^n = (n+1)e^{\psi_{+,j} - \psi_{-,j}} (\mathrm{Tr}_{\omega'_\varepsilon}(\omega_\varepsilon))^n$$

Taking log on both sides gives us

$$\log(\mathrm{Tr}_{\omega_\varepsilon} \omega'_\varepsilon) \leq \log(n) + (n+1)(\psi_{+,j} - \psi_{-,j}) + n \log(\mathrm{Tr}_{\omega'_\varepsilon} \omega_\varepsilon)$$

hence by definition of h ,

$$h \leq \log(n) + (n+1)\psi_{+,j} + n \log(\mathrm{Tr}_{\omega'_\varepsilon} \omega_\varepsilon) - A_1(\varphi_{j,\varepsilon} - \psi)$$

Therefore

$$\sup_{S \cap \mathcal{U}} h \leq h(x_0) \leq A_3 - A_1 \inf_{S \cap \mathcal{U}} (\varphi_{j,\varepsilon} - \psi) \leq A_3 - A_1 \inf_{S \cap \mathcal{U}} \varphi_{j,\varepsilon}$$

where A_3 is a uniform constant since $\psi_{+,j}$ and $\mathrm{Tr}_{\omega'_\varepsilon} \omega_\varepsilon(x_0)$ are both uniformly bounded. As a consequence, there exists a uniform constant A_4 such that:

$$h := \log(\mathrm{Tr}_{\omega_\varepsilon} \omega'_\varepsilon) + (n+1)\psi_{-,j} - A_1(\varphi_{j,\varepsilon} - \psi) \leq A_4$$

which leads to:

$$\mathrm{Tr}_{\omega_\varepsilon} \omega'_\varepsilon \leq e^{-(n+1)\psi_{-,j}} e^{A_1(\varphi_{j,\varepsilon} - \psi)} e^{A_4}$$

hence the existence of uniform constants A_1, A_5 , depending only on C in inequality (11), κ , and the bound of the L^∞ -estimate 3.10 such that:

$$\sup_{S \cap \mathcal{U}} \mathrm{Tr}_{\omega_\varepsilon} \omega'_\varepsilon \leq A_5 e^{-A_1\psi_{-,j}} \leq A_4 e^{-A_1\psi_{-,j}}$$

For the estimate of $\Delta_{\omega_B} \varphi_{j,\varepsilon}$, we make the following remark. By compacity of S , there exists a uniform constant δ sufficiently large such that

$$\omega_\varepsilon = \theta + \varepsilon \omega_B \leq \delta \omega_B$$

hence

$$\mathrm{Tr}_{\omega_B}(\cdot) \leq \delta^{-1} \mathrm{Tr}_{\omega_\varepsilon}(\cdot)$$

But since

$$\sup_{S \cap \mathcal{U}} \mathrm{Tr}_{\omega_\varepsilon}(\omega_\varepsilon + dd^c \varphi_{j,\varepsilon}) = n + \sup_{S \cap \mathcal{U}} \Delta_{\omega_\varepsilon} \varphi_{j,\varepsilon} \leq A_4 e^{-A_1 \psi - \psi_-},$$

this completes our proof. \square

3.5. Conclusion.

Proof of the main Theorem. By using the L^∞ -estimate in Lem. 3.10 and the transverse Yau-Aubin inequality 3.11, we obtained in Lem. 3.12 the estimate of $\Delta_{\omega_B} \varphi_{j,\varepsilon}$. As a consequence, $\Delta_{\omega_B} \varphi_{j,\varepsilon}$ is locally uniformly bounded on $S \cap \mathcal{U}$ since $\psi_- := \Psi_- + \varphi_X$ is locally bounded by our assumption. It follows that there exists a subsequence $\varphi_{j,\varepsilon(j)}$ which is C^1 -convergent on $S \cap \mathcal{U}$ to

$$\varphi_0 \in L^\infty(S \cap \mathcal{U}), \Delta_{\omega_B} \varphi_0 \in L_{\mathrm{loc}}^\infty(S \cap \mathcal{U})$$

which is a solution of

$$(13) \quad (\theta + d_B d_B^c \varphi_0)^n \wedge \eta = e^{-(n+1)\varphi_X} \pi^* dV_Y(-J\xi, \cdot)$$

on $S \cap \mathcal{U}$. The equation admits a unique solution up to constant (cf. Prop. 2.15), hence:

$$\varphi_0 = \varphi_X + c$$

which implies that $\Delta_{\omega_B} \varphi_X$ is locally bounded. This allows us to obtain a $C^{2,\alpha}$ -estimate of φ_X , as well as higher order estimates using Schauder's estimate and complex Evans-Krylov theory as in [Blo12, 5.3, p.210], hence the smoothness of φ_X on $S \cap \mathcal{U}$.

By definition, $r^2 = r_\xi^2 e^\varphi$ and $\varphi_X = \pi^* \varphi$. Using symmetry by $\mathbb{R}_{>0}$ -action generated by $-J\xi$, we conclude that $\varphi_X = \varphi \circ \pi$ is actually smooth on \mathcal{U} , hence φ is smooth on Y_{reg} . In particular, r^2 is smooth on Y_{reg} . \square

4. APPENDICE : TRANSVERSE YAU-AUBIN INEQUALITY

In the sequel, we will use the summation convention. Let ω_B, ω'_B be two transverse Kähler forms on S . Let (z, x) be the coordinates on a foliation chart of S such that:

$$\omega_B = g_{j\bar{k}} \sqrt{-1} dz^j \wedge d\bar{z}^k, \quad \omega'_B = g'_{j\bar{k}} \sqrt{-1} dz^j \wedge d\bar{z}^k$$

After choosing a normal transverse holomorphic chart, one can suppose that $g_{j\bar{k}} = \delta_{jk}$ and that ω'_B is diagonal. Let $(g^{j\bar{k}})$ denote the inverse of $(g_{j\bar{k}})$. We have:

$$\mathrm{Tr}_{\omega_B} \omega'_B = g^{j\bar{j}} g'_{j\bar{j}} = \sum_j g'_{j\bar{j}}, \quad \mathrm{Tr}_{\omega'_B} \omega_B = g'^{j\bar{j}} g_{j\bar{j}} = \sum_j g'^{j\bar{j}}$$

Denote

$$\partial_j := \frac{\partial}{\partial z_j}, \quad \bar{\partial}_k := \frac{\partial}{\partial \bar{z}_k}, \quad \partial_j \bar{\partial}_k := \frac{\partial^2}{\partial z_j \partial \bar{z}_k}$$

Lemma 4.1. We have the following inequality:

$$g'^{p\bar{p}}(\partial_p g'_{a\bar{a}})(\bar{\partial}_p g'_{b\bar{b}}) \leq (\mathrm{Tr}_{\omega_B} \omega'_B) \sum_{p,a,j} g'^{p\bar{p}} g'^{a\bar{a}} \left| \partial_p g'_{a\bar{j}} \right|^2$$

Proof. The lemma follows from repeated applications of Cauchy-Schwartz inequality:

$$\begin{aligned} \sum_{p,a,b} g'^{p\bar{p}}(\partial_p g'_{a\bar{a}})(\bar{\partial}_p g'_{b\bar{b}}) &\leq \sum_{a,b} (g'^{p\bar{p}} |\partial_p g'_{a\bar{a}}|^2)^{1/2} (g'^{p\bar{p}} |\bar{\partial}_p g'_{b\bar{b}}|^2)^{1/2} \\ &= \left(\sum_a \left(\sum_p g'^{p\bar{p}} |\partial_p g'_{a\bar{a}}|^2 \right)^{1/2} \right)^2 \\ &= \left(\sum_a \sqrt{g'_{a\bar{a}}} \left(\sum_p g'^{p\bar{p}} g'^{a\bar{a}} |\partial_p g'_{a\bar{a}}|^2 \right)^{1/2} \right)^2 \\ &\leq \left(\sum_a g'_{a\bar{a}} \right) \left(\sum_{p,a} g'^{p\bar{p}} g'^{a\bar{a}} |\partial_p g'_{a\bar{a}}|^2 \right) \\ &\leq (\mathrm{Tr}_{\omega_B} \omega'_B) \left(\sum_{p,a,j} g'^{p\bar{p}} g'^{a\bar{a}} \left| \partial_p g'_{a\bar{j}} \right|^2 \right) \end{aligned}$$

□

Recall the statement of the transverse Yau-Aubin inequality:

Lemma 4.2.

$$\Delta_{\omega'_B} \log \mathrm{Tr}_{\omega_B} \omega'_B \geq -\kappa \mathrm{Tr}_{\omega'_B} \omega_B - \frac{\mathrm{Tr}_{\omega_B} \mathrm{Ric}(\omega'_B)}{\mathrm{Tr}_{\omega_B} \omega'_B}$$

Proof. We have:

$$\begin{aligned} \Delta_{\omega'_B} \log \mathrm{Tr}_{\omega_B} \omega'_B &= \frac{\Delta_{\omega'_B} \mathrm{Tr}_{\omega_B} \omega'_B}{\mathrm{Tr}_{\omega_B} \omega'_B} - g'^{p\bar{q}} \frac{(\bar{\partial}_q \mathrm{Tr}_{\omega_B} \omega'_B)(\partial_p \mathrm{Tr}_{\omega_B} \omega'_B)}{(\mathrm{Tr}_{\omega_B} \omega'_B)^2} \\ &= \frac{\Delta_{\omega'_B} \mathrm{Tr}_{\omega_B} \omega'_B}{\mathrm{Tr}_{\omega_B} \omega'_B} - \frac{g'^{p\bar{p}}(\partial_p g'_{a\bar{a}})(\bar{\partial}_p g'_{b\bar{b}})}{(\mathrm{Tr}_{\omega_B} \omega'_B)^2} \end{aligned}$$

By definition,

$$\begin{aligned} \Delta_{\omega'_B} \mathrm{Tr}_{\omega_B} \omega'_B &= g'^{p\bar{q}}(\partial_p \bar{\partial}_q g'^{j\bar{k}})g'_{j\bar{k}} + g'^{p\bar{q}} g'^{j\bar{k}} \partial_p \bar{\partial}_q g'_{j\bar{k}} \\ &= g'^{p\bar{q}}(\partial_p \bar{\partial}_q g'^{j\bar{k}})g'_{j\bar{k}} - g'^{p\bar{q}} g'^{j\bar{k}} R'_{j\bar{k}p\bar{q}} + g'^{p\bar{q}} g'^{j\bar{k}} g'^{a\bar{b}}(\partial_p g'_{j\bar{b}})(\bar{\partial}_q g'_{a\bar{k}}) \end{aligned}$$

where $R'_{j\bar{k}p\bar{q}}$ is the local expression of the transverse curvature form of ω'_B . Let us estimate the three terms of the expression above.

- Since ω_B and ω'_B are diagonal, we have for the first term:

$$g'^{p\bar{q}}(\partial_p \bar{\partial}_q g^{j\bar{k}})g'_{j\bar{k}} = g'^{p\bar{p}}(\partial_p \bar{\partial}_p g^{j\bar{j}})g'_{j\bar{j}} \geq -\kappa(\mathrm{Tr}_{\omega_B} \omega'_B)(\mathrm{Tr}_{\omega'_B} \omega_B)$$

where κ is the infimum of the transverse sectional curvature (which exists since S is compact).

- In the second term, $g'^{p\bar{q}}R_{j\bar{k}p\bar{q}} = R'_{j\bar{k}}$, where $R'_{j\bar{k}}$ is the local expression of the transverse Ricci-form $\mathrm{Ric}(\omega'_B)$.
- For the third term, we have:

$$g'^{p\bar{q}}g^{j\bar{k}}g'^{a\bar{b}}(\partial_p g'_{j\bar{b}})(\bar{\partial}_q g'_{a\bar{k}}) = g'^{p\bar{p}}g'^{a\bar{a}} \left| \partial_p g'_{a\bar{j}} \right|^2$$

It follows that

$$\Delta_{\omega'_B} \mathrm{Tr}_{\omega_B} \omega'_B \geq -\kappa \mathrm{Tr}_{\omega_B} \omega'_B \mathrm{Tr}_{\omega'_B} \omega_B - g^{j\bar{k}} R'_{j\bar{k}} + \sum_{p,a,j} g'^{p\bar{p}} g'^{a\bar{a}} \left| \partial_p g'_{a\bar{j}} \right|^2$$

hence

$$\begin{aligned} \Delta_{\omega'_B} \log \mathrm{Tr}_{\omega_B} \omega'_B &\geq -\kappa \mathrm{Tr}_{\omega'_B} \omega_B - \frac{\mathrm{Tr}_{\omega_B} \mathrm{Ric}(\omega'_B)}{\mathrm{Tr}_{\omega_B} \omega'_B} \\ &+ \frac{\sum_{p,a,j} g'^{p\bar{p}} g'^{a\bar{a}} \left| \partial_p g'_{a\bar{j}} \right|^2}{\mathrm{Tr}_{\omega_B} \omega'_B} - \frac{g'^{p\bar{p}}(\partial_p g'_{a\bar{a}})(\bar{\partial}_p g'_{b\bar{b}})}{(\mathrm{Tr}_{\omega_B} \omega_B)^2} \\ &\geq -\kappa \mathrm{Tr}_{\omega'_B} \omega_B - \frac{\mathrm{Tr}_{\omega_B}(\mathrm{Ric}(\omega'_B))}{\mathrm{Tr}_{\omega_B} \omega'_B} \end{aligned}$$

by the previous lemma. \square

REFERENCES

- [AH06] Klaus Altmann and Jürgen Hausen. Polyhedral divisors and algebraic torus actions. *Math. Ann.*, 334(3):557–607, 2006.
- [BBE⁺19] Robert J. Berman, Sebastien Boucksom, Philippe Eyssidieux, Vincent Guedj, and Ahmed Zeriahi. Kähler-Einstein metrics and the Kähler-Ricci flow on log Fano varieties. *J. Reine Angew. Math.*, 751:27–89, 2019.
- [BEGZ10] Sébastien Boucksom, Philippe Eyssidieux, Vincent Guedj, and Ahmed Zeriahi. Monge-Ampère equations in big cohomology classes. *Acta Math.*, 205(2):199–262, 2010.
- [Ber19] Robert J. Berman. From Monge-Ampère equations to envelopes and geodesic rays in the zero temperature limit. *Math. Z.*, 291(1-2):365–394, 2019.
- [Ber20] Robert J. Berman. Conical Calabi-Yau metrics on toric affine varieties and convex cones. *arXiv.2005.07053*, page 26, 2020.
- [BG04] D. Burns and V. Guillemin. Potential functions and actions of tori on Kähler manifolds. *Commun. Anal. Geom.*, 12(1-2):281–303, 2004.
- [BG08] Charles P. Boyer and Krzysztof Galicki. *Sasakian geometry*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2008.
- [Blo12] Zbigniew Blocki. *The Calabi-Yau Theorem*. Lecture Notes in Mathematics, No. 52. Springer-Verlag, Berlin-Heidelberg, 2012.
- [BT76] Eric Bedford and B. A. Taylor. The Dirichlet problem for a complex Monge-Ampère equation. *Invent. Math.*, 37(1):1–44, 1976.

- [BT82] Eric Bedford and B. A. Taylor. A new capacity for plurisubharmonic functions. *Acta Math.*, 149(1-2):1–40, 1982.
- [CDS15a] Xiuxiong Chen, Simon Donaldson, and Song Sun. Kähler-Einstein metrics on Fano manifolds. I: Approximation of metrics with cone singularities. *J. Amer. Math. Soc.*, 28(1):183–197, 2015.
- [CDS15b] Xiuxiong Chen, Simon Donaldson, and Song Sun. Kähler-Einstein metrics on Fano manifolds. II: Limits with cone angle less than 2π . *J. Amer. Math. Soc.*, 28(1):199–234, 2015.
- [CDS15c] Xiuxiong Chen, Simon Donaldson, and Song Sun. Kähler-Einstein metrics on Fano manifolds. III: Limits as cone angle approaches 2π and completion of the main proof. *J. Amer. Math. Soc.*, 28(1):235–278, 2015.
- [CS19] Tristan C. Collins and Gábor Székelyhidi. Sasaki-Einstein metrics and K-stability. *Geom. Topol.*, 23(3):1339–1413, 2019.
- [DS17] Simon Donaldson and Song Sun. Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry, II. *J. Differential Geom.*, 107(2):327–371, 2017.
- [EGZ09] Philippe Eyssidieux, Vincent Guedj, and Ahmed Zeriahi. Singular Kähler-Einstein metrics. *J. Amer. Math. Soc.*, 22(3):607–639, 2009.
- [EKA90] Aziz El Kacimi-Alaoui. Opérateurs transversalement elliptiques sur un feuilletage riemannien et applications. In *Functional analytic methods in complex analysis and applications to partial differential equations (Trieste, 1988)*, pages 287–340. World Sci. Publ., River Edge, NJ, 1990.
- [GZ05] Vincent Guedj and Ahmed Zeriahi. Intrinsic capacities on compact Kähler manifolds. *J. Geom. Anal.*, 15(4):607–639, 2005.
- [GZ07] Vincent Guedj and Ahmed Zeriahi. The weighted Monge-Ampère energy of quasiplurisubharmonic functions. *J. Funct. Anal.*, 250(2):442–482, 2007.
- [GZ17] Vincent Guedj and Ahmed Zeriahi. *Degenerate Complex Monge-Ampère Equations*. EMS Tracts in Mathematics. European Mathematical Society, 2017.
- [HL21] Weiyong He and Jun Li. Geometric pluripotential theory on sasaki manifolds. *J. Geom. Anal.*, 31(1):1093–1179, 2021.
- [Kol97] János Kollár. Quotient spaces modulo algebraic groups. *Ann. of Math. (2)*, 145(1):33–79, 1997.
- [Kol04] Janos Kollar. Seifert G_m -bundles. *arXiv.0404386*, 2004.
- [Kol07] Janos Kollar. Resolution of singularities. *arXiv.0508332*, 2007.
- [Li22] Chi Li. G -uniform stability and Kähler-Einstein metrics on Fano varieties. *Invent. Math.*, 227(2):661–744, 2022.
- [LLX20] Chi Li, Yuchen Liu, and Chenyang Xu. A guided tour to normalized volume. In *Geometric analysis—in honor of Gang Tian’s 60th birthday*, volume 333 of *Progr. Math.*, pages 167–219. Birkhäuser/Springer, Cham, [2020] ©2020.
- [MSY08] Dario Martelli, James Sparks, and Shing-Tung Yau. Sasaki-Einstein manifolds and volume minimisation. *Comm. Math. Phys.*, 280(3):611–673, 2008.
- [Siu87] Yum Tong Siu. *Lectures on Hermitian-Einstein metrics for stable bundles and Kähler-Einstein metrics*, volume 8 of *DMV Seminar*. Birkhäuser Verlag, Basel, 1987.
- [Sum74] Hideyasu Sumihiro. Equivariant completion. *J. Math. Kyoto Univ.*, 14:1–28, 1974.
- [vC18] Craig van Coevering. Monge-ampère operators, energy functionals and uniqueness of sasaki-extremal metrics. *arXiv.1511.09167*, 2018.