



HAL
open science

Early delivery following chronic and acute ambient temperature exposure: a comprehensive survival approach

Ian Hough, Matthieu Rolland, Ariane Guilbert, Emie Seyve, Barbara Heude, Rémy Slama, Sarah Lyon-Caen, Isabelle Pin, Cécile Chevrier, Itai Kloog, et al.

► To cite this version:

Ian Hough, Matthieu Rolland, Ariane Guilbert, Emie Seyve, Barbara Heude, et al.. Early delivery following chronic and acute ambient temperature exposure: a comprehensive survival approach. International Journal of Epidemiology, In press, 10.1093/ije/dyac190 . hal-03819527v1

HAL Id: hal-03819527

<https://hal.science/hal-03819527v1>

Submitted on 18 Oct 2022 (v1), last revised 25 Oct 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Early delivery following chronic and acute in utero ambient
2 temperature exposure: a comprehensive survival approach

3 Ian Hough^{a,b,*}, Matthieu Rolland^a, Ariane Guilbert^a, Emie Seyve^a, Barbara Heude^c, Rémy Slama^a,
4 Sarah Lyon-Caen^a, Isabelle Pin^{a,d}, Cécile Chevrier^e, Itai Kloog^{b,†}, Johanna Lepeule^{a,†}

5 ¹ *Université Grenoble Alpes, INSERM, CNRS, Institute for Advanced Biosciences (IAB), Team of Environmental*
6 *Epidemiology Applied to Development and Respiratory Health, F-38700 La Tronche, France*

7 ² *Department of Geography and Environmental Development, Ben-Gurion University of the Negev, Be'er Sheva,*
8 *Israel*

9 ³ *Université de Paris Cité, Inserm, INRAE, Centre of Research in Epidemiology and Statistics (CRESS), F-*
10 *75004 Paris, France*

11 ⁴ *Department of Paediatric Pneumology, Grenoble Teaching Hospital, F-38700, La Tronche, France*

12 ⁵ *Université Rennes, INSERM, EHESP, IRSET (Research Institute for Environmental and Occupational Health),*
13 *F-35000 Rennes, France*

14

15 * *Corresponding author.* Institute for Advanced Biosciences (IAB), Site Santé – Allée des Alpes, 38700 La
16 Tronche, France. E-mail: ian.hough@univ-grenoble-alpes.fr

17

18 † *Joint senior authors*

19

20 This is the author's original (preprint) version of an article accepted for publication in
21 *International Journal of Epidemiology* published by Oxford University Press. The version of
22 record is available online at: <https://doi.org/10.1093/ije/dyac190>.

23

24 Abstract

25 **BACKGROUND:** Ambient temperature, particularly heat, is increasingly acknowledged as a
26 trigger of preterm delivery, but study designs have been limited and results mixed. We aimed
27 to comprehensively evaluate the association between ambient temperature throughout
28 pregnancy and preterm delivery.

29 **METHODS:** We estimated daily temperature throughout pregnancy using a cutting-edge
30 spatiotemporal model for 5,347 live singleton births from three prospective cohorts in France,
31 2002-2018. We performed Cox regression (survival analysis) with forward (weekly from
32 conception) and backward (30 days before delivery) distributed lags to evaluate time-varying
33 associations with preterm birth. We examined weekly mean, daytime, night-time, and
34 variability of temperature, and heatwaves accounting for adaptation to location and season.

35 **RESULTS:** Preterm birth risk was higher following cold (5th vs 50th percentile of mean
36 temperature) 7-9 weeks after conception [relative risk (RR) 1.3, 95% confidence interval (CI)
37 1.0-1.6 for 2°C vs 11.6°C] and 10-4 days before delivery (1.6 [1.1-2.1] for 1.2°C vs 12.1°C).
38 Night-time heat (95th vs 50th percentile of minimum temperature; 15.7°C vs 7.4°C) increased
39 risk when exposure occurred within five weeks of conception (2.0 [1.05-3.8]) or 20-26 weeks
40 after conception (2.9 [1.2-6.8]). Overall and daytime heat (high mean and maximum
41 temperature) showed consistent effects. We found no clear associations with temperature
42 variability or heatwave indicators, suggesting they may be less relevant for preterm birth.

43 **CONCLUSION:** In a temperate climate, night-time heat and chronic and acute cold
44 exposures were associated with increased risk of preterm birth. These results suggest night-
45 time heat as a relevant indicator. In the context of rising temperatures and more frequent
46 weather hazards, these results should inform public health policies to reduce the growing
47 burden of preterm births.

48 Keywords

49 preterm birth, heat, cold, exposure windows

50

51 Key Messages

- 52 • We examined chronic and acute windows of susceptibility to temperature during
53 pregnancy in France
- 54 • Cold and night-time heat increased the risk of preterm birth
- 55 • Cold was harmful during weeks 7-9 after conception and days 10-4 before delivery
- 56 • Night-time heat was harmful during weeks 1-5 and weeks 20-26 after conception

57 Introduction

58 Climate change already affects human health, and continued increases in mean temperature
59 and the frequency and intensity of extreme weather events will magnify morbidity and
60 mortality impacts.¹ A growing body of research suggests that extreme ambient temperatures
61 can increase the risk of adverse birth outcomes such as preterm birth.^{2,3} Preterm birth
62 (delivery at <37 weeks amenorrhea) is the leading cause of under-5 mortality worldwide⁴ and
63 increases the risk of adverse health outcomes throughout childhood and into adulthood.⁵
64 Globally, about 11% of births are preterm, and the rate is increasing in many countries.⁶ In
65 France, the rate of preterm birth increased from 5.4% in 1995 to 7.5% in 2016.⁷

66 The biological pathways underlying associations between temperature and preterm birth
67 remain unclear. Chronic exposure throughout pregnancy could increase the risk of preterm
68 delivery and acute exposure could trigger preterm delivery. Pregnant women have increased
69 fat deposition, decreased surface area to mass ratio, weight gain, and higher metabolic heat
70 production (as the foetus contributes), which could make them more susceptible to heat. Heat
71 may cause the release of cytokines involved in labour induction such as prostaglandin and
72 oxytocin,⁸ and their concentration in the blood might be increased by heat-induced
73 dehydration.⁹ Dehydration and shifting of blood flow to the skin to dissipate heat could limit
74 oxygen supply to the foetus,¹⁰ and heat shock proteins might cause inflammation.¹¹ Heat may
75 also be linked to preeclampsia.¹² Fewer mechanisms have been proposed for cold, but
76 thermoregulatory responses can cause peripheral vasoconstriction and increase blood pressure
77 and viscosity, which might restrict blood flow to the placenta or contribute to gestational
78 hypertension.¹¹

79 Many previous studies of temperature and preterm birth focused only on the last days of
80 pregnancy (acute exposure), and most reported that high temperatures or heatwaves were

81 associated with decreased gestational duration or increased risk of preterm or early term
82 birth.^{10,13–15} Fewer studies have examined chronic exposure earlier in pregnancy, and results
83 have been less consistent: chronic heat and cold may increase the risk of preterm birth in
84 some populations,^{16–20} but may protect or have no effect in others.^{17,18,21,22} These mixed
85 findings could in part be due to misidentification of critical windows in studies that did not
86 fully account for correlation in exposure between trimesters.²³ Also, windows of susceptibility
87 may not align with trimesters,²² but very few studies have examined narrow exposure
88 windows. Previous studies mostly estimated temperature exposure for all inhabitants of a city
89 or region based on one or a few monitoring stations. This can lead to exposure error that
90 biases associations towards the null,²⁴ particularly for rural populations, which may live far
91 from monitors, and for urban residents, who may be affected by urban heat islands. To
92 address this, a few recent studies estimated exposure with spatiotemporally resolved models
93 coupled to home addresses.^{17,20,25,26} Previous studies also considered various exposure
94 indicators such as mean, minimum, maximum, and apparent temperature. Some reported
95 stronger associations for certain indicators^{14,27} while others found that all indicators gave
96 similar results.²⁸ Mortality studies suggest that temperature variability may have health
97 impacts²⁹ and that acclimation to location and season may modify associations,³⁰ but few
98 studies have examined these in relation to preterm birth.

99 A recent review highlighted these methodological disparities and recommended that future
100 studies consider cold as well as heat, focus on identifying windows of susceptibility using
101 more accurate exposure data, and use methods such as survival analysis that account for time-
102 varying associations with birth outcomes.² In this study, we estimated daily temperature
103 exposure throughout pregnancy for three French prospective mother-child cohorts using a
104 cutting-edge spatiotemporal model (spatial unit 200 m to 1 km) coupled to participants' exact
105 home addresses. We performed a survival analysis using Cox proportional hazards models
106 with distributed lags to evaluate the time-varying nonlinear association between temperature
107 and preterm birth. We simultaneously considered chronic and acute exposures by exploring
108 gestational-week specific temperatures and daily temperatures preceding delivery. We further
109 examined several exposure indicators: overall (mean), daytime (maximum), and night-time
110 (minimum) temperature, temperature variability, and a heatwave index that accounts for
111 acclimation to location and season.


112 Methods

113 Study population

114 We obtained data from three French prospective mother-child cohorts that were designed to
115 study the effects of prenatal environmental exposures on child development and health:
116 EDEN (Étude des Déterminants pré et post natals du développement et de la santé de
117 l'Enfant),³¹ PELAGIE (Perturbateurs Endocriniens : étude Longitudinale sur les Anomalies de
118 la Grossesse, l'Infertilité, et l'Enfance),³² and SEPAGES (Suivi de l'Exposition à la Pollution
119 Atmosphérique durant la Grossesse et Effets sur la Santé).³³ Briefly, EDEN included 2002
120 women recruited between 2003 and 2006 at <24 weeks amenorrhea in the metropolitan areas
121 of Poitiers and Nancy; PELAGIE included 3421 women recruited between 2002 and 2006 at
122 <19 weeks amenorrhea in the Brittany region; SEPAGES included 484 women recruited
123 between 2014 and 2017 at <19 weeks amenorrhea in the metropolitan area of Grenoble
124 (Figure S1).

125 All three cohorts collected medical and sociodemographic information via clinical
126 examinations and questionnaires during and after pregnancy. Exact home addresses (including
127 any changes during pregnancy) were geocoded. For 43% of PELAGIE participants, only the
128 municipality or neighbourhood of residence at inclusion (on average 10.4 weeks after
129 conception) was available; we assumed these women did not move during pregnancy.

130 We excluded multiple gestation, non-livebirths, pre-existing diabetes or hypertension, and
131 participants lost to follow-up before delivery. To ensure complete equal-length exposure
132 histories, we further excluded participants missing covariates (described below) or missing
133 exposure for more than one day in any of the 26 weeks following conception (among these
134 were five extremely preterm births at <28 weeks amenorrhea; all remaining participants had
135 exposure for the 30 days ending at delivery). This left 5,347 mother-child pairs (Figure 1).



136
137 Figure 1. Flow chart of the study population.

138 Outcome definition

139 Duration of pregnancy (conception to birth) was assessed using both the reported date of the
140 last menstrual period (LMP) and the estimate from the first trimester ultrasound (when LMP
141 was not reported or when the two differed by more than 30 days). When neither of these were
142 available ($n = 3$ in EDEN; $n = 84$ in PELAGIE), we used the obstetrician's estimate of
143 gestational age at delivery. We defined preterm birth as delivery at <37 weeks amenorrhea
144 (<35 weeks since conception).

145 Exposure assessment

146 We estimated daily ambient temperature at women's home address using a multi-resolution
147 hybrid spatiotemporal model.³⁴ The model estimates daily minimum, maximum, and mean air
148 temperature from 2000 to 2018 at a 1 km spatial resolution across France and at a 200 m
149 spatial resolution over urban areas with $>50,000$ inhabitants. Briefly, it uses a multi-stage
150 ensemble approach combining linear mixed models, random forests, and gradient boosting to
151 calibrate air temperature measured at meteorological stations with satellite-derived land
152 surface temperature, elevation, and other spatiotemporal predictors. The model performs well,
153 with cross-validated R^2 better than 0.9 and mean absolute error of about 1°C . We used 200 m
154 temperature for women in urban areas covered by the model ($n = 1,741$; 33%) and 1 km
155 temperature otherwise.

156 We calculated five indicators of exposure based on each woman's daily temperature profile
157 from conception to delivery: 1) weekly mean temperature (Tmean), a marker of overall

158 exposure; 2) weekly average of daily maximum temperature (T_{\max}), a marker of daytime
159 exposure because temperature is usually highest in the afternoon; 3) weekly average of daily
160 minimum temperature (T_{\min}), a marker of night-time exposure because temperature is
161 typically lowest before sunrise; 4) weekly temperature variability (T_{SD} , the standard deviation
162 of daily mean temperature), a marker of exposure to temperature swings; and 5) daily Excess
163 Heat Factor (EHF), a marker of exposure to extreme heat that accounts for both spatial and
164 seasonal acclimation.³⁵ Supplemental appendix A details the EHF calculation.

165 Main analysis

166 We used Cox proportional hazards models with duration of pregnancy (weeks since
167 conception) as the time variable and birth as the outcome (censored at 35 weeks after
168 conception). We fit a separate model for each of the five exposure indicators (T_{mean} , T_{\min} ,
169 T_{\max} , T_{SD} , EHF). We accounted for the time-varying effects of exposure using a distributed
170 lag nonlinear model (DLNM)³⁶ with two exposure matrices: the 26 weeks following
171 conception (weekly chronic exposure) and the 30 days ending at delivery (daily acute
172 exposure). We censored chronic exposure at 26 weeks to avoid excluding a substantial
173 fraction of preterm births. We modelled both the exposure-response and the lag-response
174 relationship using natural cubic splines with equally spaced knots and three degrees of
175 freedom (chosen by testing 3-6 degrees of freedom for the lowest value that minimized the
176 Akaike information criterion).

177 We adjusted all models for possible confounders or predictors of the outcome selected *a*
178 *priori* based on the literature and our reasoning: cohort area, season of conception, urbanicity,
179 normalized difference vegetation index (NDVI), child sex, and maternal characteristics (age at
180 conception, height, pre-pregnancy BMI, parity, education, and smoking during pregnancy).
181 We did not adjust for gestational hypertension or preeclampsia as these may mediate the
182 association between temperature and preterm birth. Urbanicity was based on data from the
183 French National Institute of Statistics and Economics for the home address at birth. NDVI
184 was the mean Landsat satellite NDVI³⁷ in a 500 m buffer around the home address at birth
185 during June-August of the year of birth. We considered NDVI missing if it was unavailable
186 over more than 25% of the buffer.

187 To better understand the associations with temperature and evaluate the robustness of our
188 findings, we repeated our analyses 1) including the 309 participants (of which 16 preterm

189 births) that were missing covariates, which we imputed using the cohort area-specific median
 190 or mode; and 2) using temperature estimated at a 1 km spatial resolution for all participants
 191 (rather than using 200 m temperature for the 33% of participants that lived in large urban
 192 areas).

193 We calculated the relative risk (RR) and 95% confidence interval (CI) of preterm birth
 194 associated with moderate (10th, 90th percentiles), severe (5th, 95th percentiles), and extreme
 195 (1st, 99th percentiles) exposure compared to the median exposure (50th percentile) during the
 196 chronic and acute periods. We conducted all statistical analyses using R version 4.1.0³⁸ with
 197 the packages survival v3.2-11³⁹ and dlnm v2.4.6.⁴⁰

198 Results

199 Over half (58%) of the women lived in Brittany; 18% lived in Nancy, 16% in Poitiers, and
 200 8% in Grenoble (Table 1). Almost half (45%) lived in a rural area. Most women (72%) were
 201 25 to 34 years old at conception, had completed at least one year of post-secondary education
 202 (63%), and were multiparous (56%). A quarter (25%) of women smoked during pregnancy.

203 Table 1. Characteristics of the study population by cohort ($n = 5347$).

	Consortium <i>n (%) or mean (sd)</i>	EDEN <i>n (%) or mean (sd)</i>	PELAGIE <i>n (%) or mean (sd)</i>	SEPAGES <i>n (%) or mean (sd)</i>
Participants	5347 (100%)	1806 (33.8%)	3116 (58.3%)	425 (7.8%)
Preterm births	232 (4.3%)	103 (5.7%)	110 (3.5%)	19 (4.5%)
Duration of pregnancy ^a (weeks)	37.9 (1.7)	37.7 (1.8)	38.1 (1.6)	37.7 (1.5)
Temperature ^b (°C)	11.8 (3.6)	11.3 (4.1)	12.1 (3.2)	12 (4.4)
Temperature variability ^c (°C)	2.0 (0.3)	2.2 (0.2)	1.9 (0.3)	2.1 (0.2)
Excess Heat Factor ^d	0.7 (1.1)	0.8 (1.3)	0.6 (0.9)	0.7 (1.0)
Child sex				
Boy	2737 (51.2%)	940 (52%)	1572 (50.4%)	225 (52.9%)
Girl	2610 (48.8%)	866 (48%)	1544 (49.6%)	200 (47.1%)
Parity				
0	2374 (44.4%)	794 (44.0%)	1386 (44.5%)	194 (45.6%)
1	2021 (37.8%)	670 (37.1%)	1166 (37.4%)	185 (43.5%)
>=2	952 (17.8%)	342 (18.9%)	564 (18.1%)	46 (10.8%)
Maternal age at conception (years)	30 (4.5)	29.4 (4.9)	29.9 (4.3)	32.5 (3.9)
Maternal height				
135-175 cm	4612 (86.3%)	1561 (86.4%)	2719 (87.3%)	332 (78.1%)
170-190 cm	735 (13.7%)	245 (13.6%)	397 (12.7%)	93 (21.9%)
Maternal pre-pregnancy BMI				
<18.5 kg/m ²	412 (7.7%)	157 (8.7%)	229 (7.3%)	26 (6.1%)
18.5 – 25 kg/m ²	3844 (71.9%)	1177 (65.2%)	2346 (75.3%)	321 (75.5%)
>25 kg/m ²	1091 (20.4%)	472 (26.1%)	541 (17.4%)	78 (18.4%)
Maternal education				
Baccalaureate or less	1987 (37.2%)	840 (46.5%)	1125 (36.1%)	22 (5.2%)
Baccalaureate +1 or +2 years	1373 (25.7%)	415 (23.0%)	906 (29.1%)	52 (12.2%)

>= Baccalaureate +3 years	1987 (37.2%)	551 (30.5%)	1085 (34.8%)	351 (82.6%)
Smoking status during pregnancy				
None	4006 (74.9%)	1347 (74.6%)	2257 (72.4%)	402 (94.6%)
Active smoker	1341 (25.1%)	459 (25.4%)	859 (27.6%)	23 (5.4%)
NDVI	0.50 (0.11)	0.46 (0.10)	0.52 (0.10)	0.49 (0.16)
Urbanicity				
City-centre	896 (16.8%)	191 (10.6%)	547 (17.6%)	158 (37.2%)
Small city-centre or suburban	2021 (37.8%)	879 (48.7%)	918 (29.5%)	224 (52.7%)
Rural	2430 (45.4%)	736 (40.8%)	1651 (53%)	43 (10.1%)
Season of conception				
Winter	1321 (24.7%)	431 (23.9%)	762 (24.5%)	128 (30.1%)
Spring	1211 (22.6%)	371 (20.5%)	751 (24.1%)	89 (20.9%)
Summer	1477 (27.6%)	527 (29.2%)	860 (27.6%)	90 (21.2%)
Autumn	1338 (25.0%)	477 (26.4%)	743 (23.8%)	118 (27.8%)
Area				
Nancy	940 (17.6%)	940 (52%)	0 (0%)	0 (0%)
Poitiers	866 (16.2%)	866 (48%)	0 (0%)	0 (0%)
Côtes-d'Armor	898 (16.8%)	0 (0%)	898 (28.8%)	0 (0%)
Finistère	157 (2.9%)	0 (0%)	157 (5.0%)	0 (0%)
Ille-et-Vilaine	2061 (38.5%)	0 (0%)	2061 (66.1%)	0 (0%)
Grenoble	425 (7.9%)	0 (0%)	0 (0%)	425 (100%)

204 Note: BMI, body mass index; NDVI, normalized difference vegetation index.

205 ^a From conception to delivery.

206 ^b Mean during the 26 weeks following conception.

207 ^c Mean weekly standard deviation during the 26 weeks following conception.


208 ^d Mean during days 30-181 after conception (up to 26 weeks following conception).

209 Mean duration of pregnancy (conception to delivery) was 37.9 weeks and 4.3% of births were
210 preterm. Mean temperature \pm standard deviation during the 26 weeks following conception
211 was $11.8 \pm 3.6^\circ\text{C}$; the standard deviation of weekly temperature averaged $2.0 \pm 0.3^\circ\text{C}$. Mean
212 EHF was 0.7 ± 1.1 over days 30 to 181 since conception. Figure S2 shows the distribution of
213 mean temperature and EHF over time for each cohort and Table S2 summarizes the
214 distribution of the exposure indicators.

215 Cold

216 Severe cold (5th vs 50th percentile of Tmean) during weeks 7-9 after conception and days 10-4
217 before delivery increased the risk of preterm birth (Figure 2). A mean temperature of 2°C
218 throughout weeks 7-9 after conception was associated with RR for preterm birth of 1.29 (95%
219 CI: 1.02-1.64) compared to the reference temperature of 11.6°C (Figure 4). A mean
220 temperature of 1.2°C throughout days 10-4 before delivery had RR of 1.55 (95% CI: 1.14-
221 2.11) compared to 12.1°C . Considering a single week or day during each critical window, the
222 mean RR for preterm birth was 1.09 (95% CI: 1.01-1.18) following severe cold on one of
223 weeks 7-9 after conception; the mean RR for preterm birth was 1.06 (95% CI: 1.01-1.12)
224 following severe cold on one of days 10-4 before delivery (Figure S3). For extreme cold (1st


225 percentile of Tmean), the critical windows were 4-9 weeks after conception and 10-4 days
 226 before delivery; the only critical window for moderate cold (10th percentile of Tmean) was
 227 10-5 days before delivery (Figure 4). Chronic daytime cold (low Tmax) showed a longer
 228 critical window than chronic overall cold (low Tmean) (Figure 2), while night-time cold (low
 229 Tmin) was only significant when it was extreme (1st percentile) 7-6 days before delivery
 230 (Figure 4). Imputing missing covariates did not substantially change the associations. Using
 231 only 1 km temperature made the critical windows shorter or widened the confidence intervals.
 232 Maternal education was the only variable that did not satisfy the proportional hazards
 233 assumption; stratifying the Cox models on education did not substantially alter our results.



234
 235 Figure 2. Adjusted relative risk (solid line) and 95% confidence interval (shaded area) for
 236 preterm birth associated with severe cold during the 26 weeks following conception (left) and
 237 the 30 days ending at delivery (right). Top: overall cold (5th percentile of Tmean); middle:
 238 daytime cold (5th percentile of Tmax); bottom: night-time cold (5th percentile of Tmin).
 239 Reference is 50th percentile of temperature. Models included exposure throughout pregnancy,
 240 area, season of conception, urbanicity, NDVI, child sex, and maternal characteristics (age at
 241 conception, height, pre-pregnancy BMI, parity, education, smoking during pregnancy).

242 Heat

243 Heat (high Tmean) during the first weeks following conception, the second half of the second
 244 trimester, and the last days before delivery seemed to correspond with an increased risk of
 245 preterm birth (Figure 3). We identified critical windows for severe night-time heat (95th vs
 246 50th percentile of Tmin; 15.7°C vs 7.4°C) during weeks 1-5 after conception (RR: 2.00; 95%
 247 CI:1.05-3.84) and 20-26 weeks after conception (RR: 2.87; 95% CI: 1.21-6.79) (Figure 3,
 248 Figure 4). Focusing on a single week during each critical window, an average minimum
 249 temperature of 15.7°C during any one of weeks 1-5 or weeks 21-26 after conception was
 250 associated with mean RR of about 1.16 (95% CI: 1.02-1.34) (Figure S3). Moderate (90th
 251 percentile) night-time heat showed a somewhat smaller effect during the same windows,
 252 while extreme (99th percentile) night-time heat showed a larger effect but only 21-26 weeks
 253 after conception (Figure 4). Sensitivity analyses did not substantially alter the results.




254 Figure 3. Adjusted relative risk (solid line) and 95% confidence interval (shaded area) for
 255 preterm birth associated with severe heat during the 26 weeks following conception (left) and
 256 the 30 days ending at delivery (right). Top: overall heat (95th percentile of Tmean); middle:
 257 daytime heat (95th percentile of Tmax); bottom: night-time heat (95th percentile of Tmin).
 258 Reference is 50th percentile of temperature. Models included exposure throughout pregnancy,
 259 area, season of conception, urbanicity, NDVI, child sex, and maternal characteristics (age at
 260 conception, height, pre-pregnancy BMI, parity, education, smoking during pregnancy).
 261

262 Temperature variability and Excess Heat Factor

263 There was no clear association between temperature variability and risk of preterm birth.

264 Moderately variable temperature (90th vs 50th percentile of T_{SD}; 3.3°C vs 1.8°C) 7-8 weeks
265 after conception may have had a protective effect (Figure S4), but the association did not
266 clearly differ from null (RR: 0.85; 95% CI: 0.73-1.00). There were no critical windows for
267 other exposure levels, and sensitivity analyses showed similar results.

268 There was no association between EHF and preterm birth (Figure S5).



1
2 Figure 4. Cumulative adjusted relative risk (RR) for preterm birth associated with ambient temperature exposure throughout an entire critical
3 window. Bars show timing of critical windows; bar labels show cumulative RR (95% confidence interval). Models included exposure throughout
4 pregnancy, area, season of conception, urbanicity, NDVI, child sex, and maternal characteristics (age at conception, height, pre-pregnancy BMI,
5 parity, education, smoking during pregnancy).

6 Discussion

7 Our results are based on a state-of-the-art approach combining survival analysis with daily
8 residence-based temperature exposure, controlling for both chronic (weekly for the 26 weeks
9 following conception) and acute (daily for the 30 days ending at delivery) exposure, and
10 accounting for lagged nonlinear effects. Pregnant women were susceptible to cold from the
11 middle of the first to the middle of the second trimester and about one week before delivery.
12 Women were susceptible to heat during the five weeks following conception and the end of
13 the second trimester. Night-time heat (high T_{min}) seemed to increase preterm birth risk more
14 than daytime (high T_{max}) or overall (high T_{mean}) heat, while night-time cold seemed less
15 harmful than daytime or overall cold.

16 Cold

17 We found a critical window for chronic cold 4-9 weeks after conception. Women were more
18 sensitive to daytime cold: the critical window continued until 18 weeks after conception. We
19 found no critical window for chronic night-time cold. Women may have been less exposed to
20 night-time cold as they were likely inside a heated home.

21 Few previous studies examined cold early during pregnancy in temperate climates, with most
22 finding protective effects.^{17,18,21,22} Consistent with our results, a study in the USA found that
23 cold during the two weeks before or five weeks after conception increased the risk of preterm
24 birth, and studies in China reported increased risk from entire-pregnancy cold in temperate¹⁷
25 and cold areas.¹⁸ Compared to previous studies, we examined narrower windows (single
26 weeks during the first two trimesters), estimated residence-based temperature (rather than
27 using city-wide or regional temperature), and adjusted for temperature later in pregnancy,
28 which may have improved our ability to detect critical windows.

29 We also found a critical window for acute cold 10-4 days before delivery. Previous studies of
30 acute cold in temperate climates have reported conflicting results of no association with
31 preterm birth,^{18,20,41,42} a decreased risk,²² or oscillating positive and negative associations.^{10,14}
32 Most of these studies considered only the last seven days or four weeks of pregnancy, which
33 may have limited their ability to detect a critical window starting about 10 days before
34 delivery, and most did not adjust for temperature earlier in pregnancy as we did. Click or tap
35 here to enter text.

36 Heat

37 We found critical windows for chronic heat during the five weeks after conception and 20-26
38 weeks after conception. The association was clearest for night-time heat, but the shape of the
39 lag-response curve was similar for overall and daytime heat. Heatwave mortality studies have
40 suggested that hot nights following hot days may be particularly dangerous because they limit
41 the ability to recover from daytime exposure,^{43,44} and recent studies in California and Belgium
42 found a clearer association with preterm birth for night-time than daytime heat.^{14,27} Our study
43 may have been particularly suited to examining night-time heat because we estimated
44 residence-based exposure using a model that captures the higher night-time temperatures of
45 urban heat islands.³⁴ Taken together, these results suggest that future studies should consider
46 night-time heat indicators in order to clarify the effects of heat during pregnancy, particularly
47 in countries such as France where only about 13% of homes have air conditioning.⁴⁵

48 Many studies have reported that heat in the last days of pregnancy may trigger preterm
49 delivery,^{2,3} although a few have reported no effect in cold or cool climates.^{10,18,20,41} In our
50 study, heat during the last five days of pregnancy may have increased risk of preterm birth,
51 but the association was unclear. This might be because we adjusted for temperatures earlier in
52 pregnancy whereas most previous studies did not. Our residence-based exposure estimates
53 may also be less accurate during the final days of pregnancy for some women who may have
54 been admitted to maternity units before the day of delivery.

55 Temperature variability and acclimation

56 Temperature variability and acclimation to location and season have been shown to affect the
57 risk of mortality,^{29,30} but few studies have examined them in relation to birth outcomes. A
58 recent study in the Andes associated more variable temperature with lower birth weight,⁴⁶ and
59 a study in France using a subset of our study population associated more variable temperature
60 during weeks 4-18 after conception with lower term birth weight.⁴⁷

61 We did not find an association between preterm birth and the variability of temperature. Nor
62 did we find an association with EHF, a heatwave index that accounts for acclimation to both
63 location and season. This might be related to the fact that we only found critical windows for
64 night-time heat (high T_{min}) whereas EHF is based on T_{mean}. Our sample size may also have
65 limited our ability to examine an infrequent acute exposure such as EHF. We also adjusted for
66 EHF earlier in pregnancy, which may have reduced the effect of EHF shortly before delivery;

67 previous heatwave studies only examined exposure during the last week of pregnancy.^{28,48}
68 Overall, our findings suggest that daily mean, minimum, and maximum temperature may be
69 more relevant for preterm birth than temperature variability or heatwaves.

70 [Strengths and limitations](#)

71 We estimated outdoor temperature at women's home address, but women likely spent time
72 indoors (particularly at night) and at other locations (particularly early in pregnancy). We also
73 lacked complete address history for a quarter of participants (all from the PELAGIE cohort);
74 for these women we used municipality or neighbourhood of residence at inclusion (mean 10.4
75 weeks after conception), which may have increased exposure measurement error and biased
76 our associations towards null.

77 DLNM requires complete equal-length exposure histories, so we only considered exposure
78 during the 26 weeks following conception and the 30 days ending at delivery. This could have
79 led us to miss critical windows early in the third trimester (e.g. weeks 27-30 after conception).
80 A recent study in Rome and Barcelona found that acute heat was more harmful earlier in the
81 third trimester,¹⁵ so future studies should investigate critical windows in the first half of the
82 third trimester. We also did not account for humidity, which may modify the physiological
83 effects of heat.⁴⁹ However, the evidence for humidity is mixed: some studies found it did not
84 substantially modify the association between temperature and mortality^{50,51} or birth
85 outcomes²⁸ while others suggested a significant role⁴⁷ or that humidity's importance varies
86 between locations.⁵²

87 Despite these limitations, our study has several strengths. Pooling three cohorts from different
88 regions of France allowed us to increase the study population and capture greater climatic
89 variability while maintaining detailed health data and similar lifestyles across participants. We
90 estimated exposure at participants' home address with a spatiotemporally resolved
91 temperature model. This likely reduced exposure error for the 45% of women that lived in
92 rural areas, which often have few weather monitors, and the 33% of women that lived in large
93 urban areas, where we were able to use 200 m temperature estimates that better capture urban
94 heat islands and fine-scale spatial patterns. One previous study of temperature and birth
95 weight found that associations disappeared when using coarser monitor-based exposure rather
96 than residence-based exposure.²⁰ Consistently, our sensitivity analyses using 1 km rather than

97 200 m temperature for urban women showed similar trends as the main analysis but the
98 associations were weakened and critical windows shortened.

99 We performed a survival analysis with pregnancy duration as the time variable, which is the
100 most effective method to study time-varying exposures in cohorts.^{53,54} It also avoids possible
101 confounding by temporal trends in conception rates and accounts for the fact that the risk of
102 preterm birth increases exponentially later in pregnancy.⁵⁵ To avoid underestimating
103 gestational age in the case that temperature affects foetal growth during the first trimester, we
104 preferred gestational duration calculated from the last menstrual period rather than from
105 measurements performed at the first ultrasound. We accounted for lags in the effect of
106 exposure, examined narrow windows, and adjusted our estimates of chronic effects (during
107 the 26 weeks following conception) for acute effects (during the 30 days preceding delivery)
108 and vice versa. We further adjusted for potential confounders such as season of conception,
109 maternal age, education, and smoking, but did not adjust for air pollution because it may be
110 on the causal pathway from temperature to preterm birth.⁵⁶ Future research may investigate
111 the possible synergistic effects of air pollution and temperature.^{48,57}

112 Conclusion

113 Our results indicate that, in a temperate climate, cold between the middle of the first and
114 second trimesters may increase the risk of preterm birth and cold late in pregnancy may
115 trigger preterm birth with a lag time of about one week. Night-time heat may be harmful
116 during the five weeks following conception and the 6th month of pregnancy. We found
117 inconclusive evidence for heat as a short-term trigger of preterm birth. In the context of the
118 demonstrated and increasing risks of climate change and preterm birth's association with
119 poorer health in childhood and adulthood, health professionals and policy makers should use
120 these findings to increase awareness of the risks of extreme temperature for pregnant women.

121 Ethics approval

122 EDEN, PELAGIE and SEPAGES were approved by the relevant ethical committees: la
123 Commission Nationale de l'Informatique et des Libertés, le Comité Consultatif pour la
124 Protection des Personnes dans la Recherche Biomédicale du Kremlin Bicêtre, le Comité
125 Consultatif sur le Traitement de l'Information en Matière de Recherche dans le Domaine de la
126 Santé, le Comité de Protection des Personnes Sud-Est V, and le Comité d'Éthique de

127 l'Inserm. All participating women gave informed written consent for themselves and their
128 children.

129 Author contributions

130 IH helped design the study, prepared data, performed analyses, interpreted the results, and
131 wrote the manuscript. MR helped design the study, prepared data, performed analyses, helped
132 interpret the results, and edited the manuscript. AG helped design the study, prepared data,
133 helped interpret the results, and edited the manuscript. ES prepared data and edited the
134 manuscript. BH prepared data and edited the manuscript. RS edited the manuscript. SLC
135 prepared data and edited the manuscript. IP edited the manuscript. CC prepared data and
136 edited the manuscript. IK helped direct the study and edited the manuscript. JL designed and
137 directed the study, helped interpret the results, and edited the manuscript.

138 Data availability

139 The data underlying this article will be shared on reasonable request to the corresponding
140 author with permission of the EDEN, PELAGIE, and SEPAGES steering committees.

141 Supplementary data

142 Supplementary data are available at *IJE* online.

143 Funding

144 This work was supported by the Fondation de France (CLIMATHES grant 00081169), the
145 French Centre National de la Recherche Scientifique and Israel Ministry of Science and
146 Technology (PRC 2018-2020). Ian Hough is supported by a grant from the French National
147 Agency for Research in the framework of the “Investissements d'avenir” program (ANR-15-
148 IDEX-02) and Ben Gurion University of the Negev.

149 The EDEN cohort was supported by the Foundation for Medical Research, the National
150 Agency for Research, the National Institute for Research in Public Health (IRESP: TGIR
151 cohorte sant. 2008 program), the French Ministry of Health, the French Ministry of Research,
152 the Inserm Bone and Joint Diseases National Research and Human Nutrition National
153 Research Programs, Paris–Sud University, Nestlé, the French National Institute for
154 Population Health Surveillance, the French National Institute for Health Education, the

155 European Union FP7 programmes (FP7/2007-2013, HELIX, ESCAPE, ENRIECO, Medall
156 projects), the Diabetes National Research Program (through a collaboration with the French
157 Association of Diabetic Patients), the French Agency for Environmental Health Safety,
158 Mutuelle Générale de l'Éducation Nationale, the French National Agency for Food Security,
159 and the French-speaking Association for the Study of Diabetes and Metabolism.

160 The PELAGIE cohort was supported by Inserm, the French Ministry of Health, the French
161 Ministry of Labor, the French National Agency for Research, the French Agency for
162 Environmental Health Safety, the Fondation de France, the French National Institute for
163 Population Health Surveillance, and the French Ministry of Ecology.

164 The SEPAGES cohort was supported by the European Research Council (N°311765-E-
165 DOHaD), the European Community's Seventh Framework Programme (FP7/2007-2006 –
166 N°308333-892 HELIX), the European Union's Horizon 2020 research and innovation
167 programme (N°874583 ATHLETE Project, N°825712 OBERON Project), the French
168 National Agency for Research (PAPER project ANR-12-PDOC-0029-01, SHALCOH project
169 ANR-14-CE21-0007, ANR-15-IDEX-02 and ANR-15-IDEX5, GUMME project ANR-18-
170 CE36-005, ETAPE project ANR-18-CE36-0005, EDeN project ANR-19-CE36-0003-01), the
171 French Agency for Food, Environmental and Occupational Health & Safety (CNAP project
172 EST-2016-121, PENDORE project EST-2016-121, HyPaxE project EST-2019/1/039), the
173 Plan Cancer (Canc'Air project), the French Cancer Research Foundation, the French
174 Endowment Fund AGIR for chronic diseases (projects PRENAPAR and LCI-FOT), the
175 French Endowment Fund for Respiratory Health, the Fondation de France (CLIMATHES-
176 00081169, SEPAGES 5-00099903).

177 Acknowledgements

178 We acknowledge the commitment of the EDEN mother-child cohort study group: I. Annesi-Maesano,
179 JY. Bernard, J. Botton, MA. Charles, P. Dargent-Molina, B. de Lauzon-Guillain, P. Ducimetière, M. de
180 Agostini, B. Foliguet, A. Forhan, X. Fritel, A. Germa, V. Goua, R. Hankard, B. Heude, M. Kaminski,
181 B. Larroque, N. Lelong, J. Lepeule, G. Magnin, L. Marchand, C. Nabet, F. Pierre, R. Slama, MJ.
182 Saurel-Cubizolles, M. Schweitzer, O. Thiebaugeorges. We also acknowledge the commitment of the
183 SEPAGES study group: E. Eyriey, A. Licinia, A. Vellement (Groupe Hospitalier Mutualiste,
184 Grenoble), I. Pin, P. Hoffmann, E. Hullo, C. Llerena (Grenoble Alpes University Hospital, La
185 Tronche), X. Morin (Clinique des Cèdres, Echirolles), A. Morlot (Clinique Belledonne, Saint-Martin
186 d'Hères), J. Lepeule, S. Lyon-Caen, C. Philippat, I. Pin, J. Quentin, V. Siroux, R. Slama (Grenoble

187 Alpes University, Inserm, CNRS, IAB). We thank the many people that assisted with the cohorts,
188 especially the participants.

189 Conflict of interest

190 None declared.

191 References

- 192 1. Atwoli L, Baqui AH, Benfield T, et al. Call for emergency action to limit global
193 temperature increases, restore biodiversity and protect health. *International Journal of*
194 *Epidemiology*. 2022;**50**(6):1761–1764.
- 195 2. Zhang Y, Yu C, Wang L. Temperature exposure during pregnancy and birth outcomes:
196 An updated systematic review of epidemiological evidence. *Environmental Pollution*.
197 Elsevier Ltd; 2017;**225**:700–712.
- 198 3. Chersich MF, Pham MD, Area A, et al. Associations between high temperatures in
199 pregnancy and risk of preterm birth, low birth weight, and stillbirths: Systematic
200 review and meta-analysis. *The BMJ*. 2020;**371**:1–13.
- 201 4. Liu L, Oza S, Hogan D, et al. Global, regional, and national causes of under-5 mortality
202 in 2000–15: an updated systematic analysis with implications for the Sustainable
203 Development Goals. *The Lancet*. 2016;**388**(10063):3027–3035.
- 204 5. McCormick MC, Litt JS, Smith VC, Zupancic JAF. Prematurity: An Overview and
205 Public Health Implications. *Annual Review of Public Health*. 2011;**32**:367–379.
- 206 6. Chawanpaiboon S, Vogel JP, Moller AB, et al. Global, regional, and national estimates
207 of levels of preterm birth in 2014: a systematic review and modelling analysis. *The*
208 *Lancet Global Health*. 2019;**7**(1):e37–e46.
- 209 7. Blondel B, Coulm B, Bonnet C, Goffinet F, Ray C Le. Trends in perinatal health in
210 metropolitan France from 1995 to 2016: Results from the French National Perinatal
211 Surveys. *Journal of Gynecology Obstetrics and Human Reproduction*.
212 2017;**46**(10):701–713.

- 213 8. Dadvand P, Basagaña X, Sartini C, et al. Climate extremes and the length of gestation.
214 *Environmental Health Perspectives*. 2011;**119**(10):1449–1453.
- 215 9. Schifano P, Lallo A, Asta F, Sario M De, Davoli M, Michelozzi P. Effect of ambient
216 temperature and air pollutants on the risk of preterm birth, Rome 2001-2010.
217 *Environment International*. Elsevier B.V.; 2013;**61**:77–87.
- 218 10. Sun S, Weinberger KR, Spangler KR, Eliot MN, Braun JM, Wellenius GA. Ambient
219 temperature and preterm birth: A retrospective study of 32 million US singleton births.
220 *Environment International*. Elsevier; 2019;**126**(February):7–13.
- 221 11. Basu R, Chen H, Li DK, Avalos LA. The impact of maternal factors on the association
222 between temperature and preterm delivery. *Environmental Research*. Elsevier;
223 2017;**154**(October 2016):109–114.
- 224 12. Shashar S, Kloog I, Erez O, et al. Temperature and preeclampsia: Epidemiological
225 evidence that perturbation in maternal heat homeostasis affects pregnancy outcome.
226 *PLoS ONE*. 2020;**15**(5):1–14.
- 227 13. Vicedo-Cabrera AM, Iñiguez C, Barona C, Ballester F. Exposure to elevated
228 temperatures and risk of preterm birth in Valencia, Spain. *Environmental Research*.
229 Elsevier; 2014;**134**:210–217.
- 230 14. Cox B, Vicedo-Cabrera AM, Gasparini A, et al. Ambient temperature as a trigger of
231 preterm delivery in a temperate climate. *Journal of Epidemiology and Community
232 Health*. 2016;**70**(12):1191–1199.
- 233 15. Schifano P, Asta F, Dadvand P, Davoli M, Basagana X, Michelozzi P. Heat and air
234 pollution exposure as triggers of delivery: A survival analysis of population-based
235 pregnancy cohorts in Rome and Barcelona. *Environment International*. Elsevier B.V.;
236 2016;**88**:153–159.
- 237 16. He J-R, Liu Y, Xia X-Y, et al. Ambient Temperature and the Risk of Preterm Birth in
238 Guangzhou, China (2001–2011). *Environmental Health Perspectives*.
239 2016;**124**(7):1100–1106.

- 240 17. Wang Y-Y, Li Q, Guo Y, et al. Ambient temperature and the risk of preterm birth: A
241 national birth cohort study in the mainland China. *Environment International*.
242 2020;**142**(November 2019).
- 243 18. Guo T, Wang Y-Y, Zhang H, et al. The association between ambient temperature and
244 the risk of preterm birth in China. *Science of the Total Environment*. 2018;**613**–
245 **614**(12):439–446.
- 246 19. Li S, Chen G, Jaakkola JJK, Williams G, Guo Y. Temporal change in the impacts of
247 ambient temperature on preterm birth and stillbirth: Brisbane, 1994–2013. *Science of*
248 *the Total Environment*. Elsevier B.V.; 2018;**634**:579–585.
- 249 20. Kloog I, Melly SJ, Coull BA, Nordio F, Schwartz J. Using Satellite-Based
250 Spatiotemporal Resolved Air Temperature Exposure to Study the Association between
251 Ambient Air Temperature and Birth Outcomes in Massachusetts. *Environmental*
252 *Health Perspectives*. 2015;**123**(10):1053–1058.
- 253 21. Giorgis-Allemand L, Pedersen M, Bernard C, et al. The influence of meteorological
254 factors and atmospheric pollutants on the risk of preterm birth. *American Journal of*
255 *Epidemiology*. 2017;**185**(4):247–258.
- 256 22. Ha S, Liu D, Zhu Y, Kim SS, Sherman S, Mendola P. Ambient temperature and early
257 delivery of Singleton Pregnancies. *Environmental Health Perspectives*.
258 2017;**125**(3):453–459.
- 259 23. Wilson A, Chiu YHM, Hsu HHL, Wright RO, Wright RJ, Coull BA. Potential for Bias
260 When Estimating Critical Windows for Air Pollution in Children’s Health. *American*
261 *Journal of Epidemiology*. 2017;**186**(11):1281–1289.
- 262 24. Zeger SL, Thomas D, Dominici F, et al. Exposure measurement error in time-series
263 studies of air pollution: Concepts and consequences. *Environmental Health*
264 *Perspectives*. 2000;**108**(5):419–426.
- 265 25. Spolter F, Kloog I, Dorman M, Novack L, Erez O, Raz R. Prenatal exposure to ambient
266 air temperature and risk of early delivery. *Environment International*.
267 2020;**142**(February):1–5.

- 268 26. Liu X, Xiao J, Sun X, et al. Associations of maternal ambient temperature exposures
269 during pregnancy with the risk of preterm birth and the effect modification of birth
270 order during the new baby boom: A birth cohort study in Guangzhou, China.
271 *International Journal of Hygiene and Environmental Health*. Elsevier;
272 2020;**225**(160):113481.
- 273 27. Avalos LA, Chen H, Li DK, Basu R. The impact of high apparent temperature on
274 spontaneous preterm delivery: a case-crossover study. *Environmental Health*.
275 *Environmental Health*; 2017;**16**(1):1–13.
- 276 28. Huang M, Strickland MJ, Richards M, et al. Acute associations between heatwaves and
277 preterm and early-term birth in 50 US metropolitan areas: a matched case-control
278 study. *Environmental Health*. *Environmental Health*; 2021;**20**(1):1–14.
- 279 29. Shi L, Kloog I, Zanobetti A, Liu P, Schwartz J. Impacts of temperature and its
280 variability on mortality in New England. *Nature Climate Change*. 2015;**5**:988–991.
- 281 30. Nordio F, Zanobetti A, Colicino E, Kloog I, Schwartz J. Changing patterns of the
282 temperature-mortality association by time and location in the US, and implications for
283 climate change. *Environment International*. Elsevier Ltd; 2015;**81**:80–86.
- 284 31. Heude B, Forhan A, Slama R, et al. Cohort Profile: The EDEN mother-child cohort on
285 the prenatal and early postnatal determinants of child health and development.
286 *International Journal of Epidemiology*. 2016;353–363.
- 287 32. Petit C, Chevrier C, Durand G, et al. Impact on fetal growth of prenatal exposure to
288 pesticides due to agricultural activities: a prospective cohort study in Brittany, France.
289 *Environmental Health*. 2010;**9**(71):1–12.
- 290 33. Lyon-Caen S, Siroux V, Lepeule J, et al. Deciphering the impact of early-life exposures
291 to highly variable environmental factors on foetal and child health: Design of
292 SEPAGES couple-child cohort. *International Journal of Environmental Research and
293 Public Health*. 2019;**16**(20).
- 294 34. Hough I, Just AC, Zhou B, Dorman M, Lepeule J, Kloog I. A multi-resolution air
295 temperature model for France from MODIS and Landsat thermal data. *Environmental
296 Research*. Elsevier Inc.; 2020;**183**(February):109244.

- 297 35. Nairn JR, Fawcett RJB. The excess heat factor: A metric for heatwave intensity and its
298 use in classifying heatwave severity. *International Journal of Environmental Research
299 and Public Health*. 2014;**12**(1):227–253.
- 300 36. Gasparrini A, Armstrong BG, Kenward MG. Distributed lag non-linear models.
301 *Statistics in Medicine*. 2010;**29**(21):2224–2234.
- 302 37. Robinson NP, Allred BW, Jones MO, et al. A dynamic landsat derived normalized
303 difference vegetation index (NDVI) product for the conterminous United States.
304 *Remote Sensing*. 2017;**9**(8):1–14.
- 305 38. R Core Team. R: A language and environment for statistical computing. Vienna,
306 Austria: R Foundation for Statistical Computing; 2021.
- 307 39. Therneau TM. A Package for Survival Analysis in R. 2021.
- 308 40. Gasparrini A. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm.
309 *Journal of statistical software*. 2011;**43**(8):1–20.
- 310 41. Vicedo-Cabrera AM, Olsson D, Forsberg B. Exposure to seasonal temperatures during
311 the last month of gestation and the risk of preterm birth in stockholm. *International
312 Journal of Environmental Research and Public Health*. 2015;**12**(4):3962–3978.
- 313 42. Lee W, Bell ML, Gasparrini A, et al. Mortality burden of diurnal temperature range
314 and its temporal changes: A multi-country study. *Environment International*. Elsevier;
315 2018;**110**(July 2017):123–130.
- 316 43. Laaidi K, Zeghnoun A, Dousset B, et al. The impact of heat islands on mortality in
317 Paris during the August 2003 heat wave. *Environmental Health Perspectives*.
318 2012;**120**(2):254–259.
- 319 44. Murage P, Hajat S, Kovats RS. Effect of night-time temperatures on cause and age-
320 specific mortality in London. *Environmental Epidemiology*. 2017;1.
- 321 45. Randazzo T, Cian E De, Mistry MN. Air conditioning and electricity expenditure: The
322 role of climate in temperate countries. *Economic Modelling*. Elsevier B.V.;
323 2020;**90**(June 2019):273–287.

- 324 46. Molina O, Saldarriaga V. The perils of climate change: In utero exposure to
325 temperature variability and birth outcomes in the Andean region. *Economics and*
326 *Human Biology*. Elsevier B.V.; 2017;**24**:111–124.
- 327 47. Jakpor O, Chevrier C, Kloog I, et al. Term birthweight and critical windows of prenatal
328 exposure to average meteorological conditions and meteorological variability.
329 *Environment International*. Elsevier; 2020;**142**(December 2019):105847.
- 330 48. Sun Y, Ilango SD, Schwarz L, et al. Examining the joint effects of heatwaves, air
331 pollution, and green space on the risk of preterm birth in California. *Environmental*
332 *Research Letters*. 2020;**15**(10).
- 333 49. Davis RE, McGregor GR, Enfield KB. Humidity: A review and primer on atmospheric
334 moisture and human health. *Environmental Research*. 2016;**144**:106–116.
- 335 50. Armstrong BG, Sera F, Vicedo-Cabrera AM, et al. The Role of Humidity in
336 Associations of High Temperature with Mortality: A Multicountry, Multicity Study.
337 *Environmental Health Perspectives*. 2019;**127**(9):1–8.
- 338 51. Barnett AG, Tong S, Clements ACA. What measure of temperature is the best
339 predictor of mortality? *Environmental Research*. Elsevier; 2010;**110**(6):604–611.
- 340 52. Bobb JF, Dominici F, Peng RD. A Bayesian model averaging approach for estimating
341 the relative risk of mortality associated with heat waves in 105 U.S. cities. *Biometrics*.
342 2011;**67**(4):1605–1616.
- 343 53. Lepeule J, Rondeau V, Filleul L, Dartigues JF. Survival analysis to estimate association
344 between short-term mortality and air pollution. *Environmental Health Perspectives*.
345 2006;**114**(2):242–247.
- 346 54. Suh YJ, Kim H, Seo JH, et al. Different effects of PM10 exposure on preterm birth by
347 gestational period estimated from time-dependent survival analyses. *International*
348 *Archives of Occupational and Environmental Health*. 2009;**82**(5):613–621.
- 349 55. Strand LB, Barnett AG, Tong S. Methodological challenges when estimating the
350 effects of season and seasonal exposures on birth outcomes. *BMC Medical Research*
351 *Methodology*. 2011;**11**.

- 352 56. Buckley JP, Samet JM, Richardson DB. Does air pollution confound studies of
353 temperature? *Epidemiology*. 2014;**25**(2):242–245.
- 354 57. Wang Q, Li B, Benmarhnia T, et al. Independent and combined effects of heatwaves
355 and PM2.5 on preterm birth in Guangzhou, China: A survival analysis. *Environmental*
356 *Health Perspectives*. 2020;**128**(1):1–10.