

Conformational variability in proteins bound to single-stranded DNA: a new benchmark for new docking perspectives

Protein-ssDNA benchmark and its analysis

MIAS-LUCQUIN Dominique^{1*}, CHAUVOT DE BEAUCHENE Isaure¹

¹ Universite de Lorraine, CNRS, Inria, LORIA, Nancy, France

* *dominique.mias-lucquin@loria.fr*

ABSTRACT

We explored the Protein Data-Bank (PDB) to collect protein-ssDNA structures and create a multi-conformational docking benchmark including both bound and unbound protein structures. Due to ssDNA high flexibility when not bound, no ssDNA unbound structure is included in the benchmark. For the 91 sequence-identity groups identified as bound-unbound structures of the same protein, we studied the conformational changes in the protein induced by the ssDNA binding. Moreover, based on several bound or unbound protein structures in some groups, we also assessed the intrinsic conformational variability in either bound or unbound conditions, and compared it to the supposedly binding-induced modifications. To illustrate a use case of this benchmark, we performed docking experiments using ATTRACT docking software. This benchmark is, to our knowledge, the first one made to peruse available structures of ssDNA-protein interactions to such an extent, aiming to improve computational docking tools dedicated to this kind of molecular interactions.

KEYWORDS: Single-Stranded DNA, Single-Stranded DNA-Binding Protein, Molecular Docking Analysis, Benchmark,

Introduction

While originally described by Watson and Crick¹ as a double helix, composed of two strands bonded together by hydrogen-bonds, DNA is often found in a transient single-stranded state (ssDNA) during its processing, such as genome replication², or horizontal gene transfer³, and bound to proteins. These complexes (ribosomes⁴, ICE-relaxase⁵, replication fork complex⁶, ...) are potential therapeutic targets in diseases^{7,8}.

The structural analysis of these complexes can help to understand how they achieve their function⁹. For example it can reveal the conformational changes undergone by the protein during nucleic acids (NA) binding, by comparing protein structures with and without bound NA¹⁰.

While very informative, high resolution experimental structures of ssNA-protein complexes are expensive and may be difficult, or even impossible, to obtain, due to the inherent poor ordering of NA, especially ssNA^{11,12}. Several software systems have tried to implement accurate ssRNA-protein docking, including:

- ATTRACT¹³ uses a fragment-based approach, with the need of some knowledge about some protein-RNA contacts;

- RNP-denovo¹⁴, based on Rosetta¹⁵, performs folding and docking of the RNA on the protein simultaneously, but requiring the exact coordinates of few nucleotides;
- RNA-lim¹⁶, models a rough coarse-grained RNA structure (one non-oriented bead per nucleotide) restrained by a set of known binding-sites on the protein surface;

While all these methods advertise a prediction precision from 2 to 10 Å of RMSD between predicted and experimental ssRNA location, none of them was tested yet on ssDNA-protein docking. To our knowledge, no benchmark is available for ssDNA-protein docking. Moreover, while it is possible to query ssDNA-protein complexes with the Nucleic Acid Database¹⁷ (NDB), it seems to find none after 2013, thus limiting the scope of a NDB-derived benchmark. In turn, docking algorithms need experimental ground truth to validate and compare methods. Thus, docking benchmarks based on experimentally resolved structures of complexes are needed. Such docking benchmarks exist for protein-protein¹⁸, membrane protein-protein¹⁹, protein-RNA^{20,21}, and dsDNA-protein²² complexes. And while some works studied ssDNA-protein interactions from

few structures in the PDB²³, none seems to be as exhaustive as possible, with a primary goal to improve ssDNA-Protein docking.

Here, we present a ssDNA-protein docking benchmark based on structures extracted from the PDB, that contains 91 sequence-identity groups of bound-unbound protein chains, created to evaluate ssDNA-protein docking. Due to the high flexibility of unstructured ssNA, it is not relevant to use their unbound forms in the context of macromolecular docking. This is also the reason why the docking programs presented earlier do not require a known unbound ssRNA structure. In consequence, the main aim of this dataset is to provide bound and unbound structures of the proteins but only bound structures of ssDNA, from ssDNA-protein complexes. When possible, we provide several structures for both bound and unbound states, allowing to differentiate binding-specific from binding-independent conformational changes.




Fig.1: Protocol to create the benchmark

Docking experiments were performed to show a use case for this benchmark. It underlines the relevance of using several bound structures as ground truth and to tolerate a minimum conformational deviation from ground truth when evaluating docking results.

Material and methods

All analysis were performed using Python 3.7. Databases were queried on August 18th 2021. Processing steps are summarized in Figure 1.

RCSB Protein Data Bank (PDB) query

We identified the structures containing simultaneously DNA (non hybrid) and proteins by querying the RCSB PDB²⁴ using their search (<https://search.rcsb.org>) and data (<https://data.rcsb.org>) APIs. Another query was performed to extract the PDB ID of all structures containing only proteins without DNA.

These two PDB ID lists were compared to the weekly 100% sequence identity clustering (“clusterNumber100”) of protein chains in the PDB (<ftp://resources.rcsb.org/sequence/clusters/>) to extract identity groups containing chains being part of both DNA-protein and protein-only structures.

Structure alignment, processing and identification of Interacting ssDNA

For each structure containing DNA and protein, the asymmetric units and all biological assemblies (if any) were downloaded from the PDB²⁴. We only kept “ATOM” records describing atoms belonging to proteins or nucleic acids. Nucleic residues involved in double strands are located with 3DNA (script `find_pair`²⁵) in the asymmetric unit and in each biological assembly. A DNA residue is considered single-stranded only if it is not found as double-stranded in any of these structures. Biological assemblies allow the identification of cases where a double strand is formed by the repetition of the asymmetric unit (such as PDB ID:3HZI), while the asymmetric unit eases the processing if the assembly is constituted by the repetition of chains having the same identifier (also like in PDB ID:3HZI). Then VMD-python (Humphrey et al., 1996, <https://github.com/Eigenstate/vmd-python>) was used to compute distances between ssDNA nucleotides and protein residues; ssDNA nucleotides are bound if they are found at less than 5 Å from any protein residue. This bound ssDNA list is processed to only keep the protein chains interacting with a DNA chain of at least 4 consecutive bound and single-stranded nucleotides. For multi-framed asymmetric units (often encountered with NMR models), only the first frame is used. In this case, we assumed a limited variation of conformation between frames, with no impact on DNA 2D state.

Bound protein chains were then sub-grouped by interaction interface: interaction interfaces with ssDNA are computed with VMD-python; two bound chains in a sequence identity cluster are grouped if they share at least one interacting residues. This can lead to one sequence cluster being split into several if distinct interfaces are found. B

From the RCSB PDB sequence clustering, we retrieved protein unbound chains belonging to structures without DNA that have 100% of sequence identity with previously identified bound chains. Unbound chains were compared to their bound counterpart to identify any structure with missing residues at the interacting interface. If one is found, it is removed from the unbound chains list.

In each sequence identity cluster, bound and unbound chains were rigidly superimposed and global RMSD computed with FATCAT 2.0 standalone software²⁵. FATCAT was used because of its ability to superimpose structures with some minor differences between sequences (like missing loop or mutation).

Superposition, RMSD calculation and clustering

The final benchmark is reported in supplementary Table S1 (and RMSD tables in supplementary Table S2), in which chains with a RMSD lower than 0.2 Å are grouped under a single representative chain. This was done to limit the bias from structures containing several times the same chains. These non-redundant dataset is further analyzed. The redundant benchmark and RMSD tables can be found as supplementary Tables 3 and 4 respectively. The sequence identity between clusters is reported in supplementary Table 5.

Interacting ssDNA fragments are reported in supplementary Table 6. We called “fragment” each interacting ssDNA region, while “sequence” refers to unique ssDNA sequence among all clusters. Thus, a DNA chain can have several ssDNA fragments if

separated by non-interacting or double-stranded DNA, and several fragments can have the same sequence.

All docking files are available on <https://github.com/DomML/ssDNAbenchmark>. Because structures can easily be retrieved from the PDB, they were not included in the repository.

Docking and docking evaluation

Docking experiments were performed using ATTRACT²⁶ without explicit restraints. ATTRACT docking being rigid, docking was performed using a library of multiple tri-nucleotides (reported in Tables 2 and 3) built using protNAff (<https://github.com/isaureCdB/ProtNAff>). Tri-nucleotides libraries were used as an exhaustive (at 1Å of heavy-atoms RMSD) ensemble of ligand conformations. For each docking, a random selection of 1000000 initial poses was performed, a pose corresponding to one fragment conformation from the library at one random starting position on sphere of 35Å radius around the protein with one random orientation. The fragments position was optimized by gradient descent minimisation of the protein-DNA energy in ATTRACT coarse-grained force field²⁷.

The best docking solutions according to ATTRACT scoring were analyzed against the bound reference, by measuring the interface RMSD (irmsd), ligand RMSD (lrmsd) and the fraction of native contacts (fnat).

Results & Discussion

Composition of the benchmark

Here, we present a dataset composed of 284 bound and 669 unbound protein chains, distributed in 91 sequence-identity groups and 98 groups with distinct interfaces: 8 identity clusters contain two distinct interfaces, none have more. It covers a wide range of protein structural families, and should be very useful both for a better understanding of the mechanisms




Fig.2: (A, left) Ratio between number of bound and unbound chains in each cluster, (B, right) Mean-RMSD for the clusters made of at least 3 chains

involved in ssDNA-protein binding and as a benchmark to evaluate ssDNA-protein docking software systems.

The most populated group (group 0, endoribonuclease hydrolase) is composed of 104 chains (1 bound, 103 unbound), and 46 groups have at least two chains for both bound and unbound. These multiple conformations allow the study of conformations variability in the bound or unbound state. Several bound structures also allows the definition of not a unique ground-truth structure, but a range of deviations around one bound structure in which a docking solution is considered as good. Moreover, 8 groups have more than one bound chain but only one unbound chain, 31 groups are in the opposite situation, and 13 contain only one chain for both bound and unbound.

Similarly to group 0, most of the groups are unbalanced: they have a difference of at least 20% between the number of bound and unbound chains (Figure 2A).

The presence of several bound or unbound chains for a given group is often purposely avoided in other benchmarks^{20–22} by excluding structures with similar sequences. However, offering alternative conformations and alternative docking solutions for a given structure provides essential piece of information to discriminate between conformational changes occurring independently of the presence or absence of bound DNA and changes induced by ssDNA binding. Moreover, as shown by He et al.²⁸, taking into account pre-binding conformation variability can increase docking performances.

Proteins conformational variability

In each group, the RMSD was computed between each pair of chains to assess the aforementioned conformation variations.

Among the 54 groups with at least 2 bound non-redundant chains ($\text{RMSD} > 0.2\text{\AA}$), 42 have a mean bound-RMSD of 2 Å or below, marking a general stability of the bound forms. Among the 12 other groups, only 2 have a mean-bound RMSD over 10 Å (#61 and 84).

For the 76 groups with at least 2 unbound non-redundant chains ($\text{RMSD} > 0.2\text{\AA}$), 48 have a mean unbound-RMSD under 2 Å, 27 between 2 and 10 Å, and 3 over 10 Å: #3, 61 and 84.

We notice a general stiffening of conformations in bound proteins, with a trend towards lower bound-RMSDs than unbound-RMSDs, probably due to ssDNA-binding. This is consistent with what was reported earlier for dsDNA³³.

Overall, in the 85 groups that contain at least three chains, the median mean RMSD (bound + unbound) is 1.69 Å (Figure 2B). This signals a low general variation in conformations between the chains of a group.

Two groups (#61 and 84, Figure S1 and S2) have a mean-RMSD higher than 10 Å. These high RMSD values are due to different relative positions of sub-domains of the protein. Moreover, RMSD clustering shows the presence of several clusters, corresponding to as many major conformational states, but none of them contains all bound or unbound structures. This is a clear indication towards changes that are not driven by DNA binding. While many related works focus on conformational changes upon RNA²⁹ or DNA^{30,31} binding, ligand binding without conformational change has been modeled³², but poorly studied.

To go further on this track, we dug into the 54 groups containing 2 or more non-redundant bound chains, to test whether there exists a pair of bound-unbound chains closer than a pair of bound-bound chains. If all bound chains are closer to each other than to any unbound chains, this can be a mark of a specific binding-induced rearrangement, including local or low amplitude changes. This is the case for group #3, with a maximum RMSD between bound chains of 3.1 Å and a minimum RMSD between bound and unbound chains of 10.7 Å. Overall, 35 groups are in such a situation, in agreement with what has been observed for RNA-protein binding³³. This marks an ssDNA-induced fit, and the use of unbound conformations in a docking procedure without taking into account such an induced change can only lead to results of limited accuracy. In the remaining 19 groups, some bound chains were closer to unbound chains than to other bound chains, indicating that conformational changes induced by DNA binding, if any, are of lower amplitude than the intrinsic variability of the protein, like in group #2.2. This points to a predominant conformational selection effect in ssDNA-binding. In these cases, using all unbound conformations, if possible with some additional conformational sampling, for docking should increase the chances to reach a close-to-bound conformation, and improve docking results.

Interestingly, our benchmark could provide the necessary data to investigate which features, other than bound-unbound comparison, could indicate in which category (binding-specific or binding-independent conformational changes) a protein lies. Moreover, the fact that most conformational changes are induced by the ssDNA binding is in agreement with the better results obtained in general (not specifically with ssDNA) by flexible docking methods³⁴.

Length and sequence of bound ssDNA

In our benchmark, 325 ssDNA fragments are found bound to proteins, with 148 unique sequences (later called just “sequences”). Among these fragments, 34% (110) are homopolymers: 98 contain only “T”, among which 36 and 24 are respectively 4 and 5 nucleotide

long; 2 contains only A, 9 only C and 1 only G (Table 1). The 4- and 5-mers poly-T are also the most represented fragments (supplementary Table 7).

Besides, 30 fragments (corresponding to 24 different sequences) contain non-canonical nucleotides. While such fragments may not be suitable for general purpose docking methods, we keep them to leave the choice to the end user, as some non-canonical residues have a canonical counterpart.

Docking

Table 1: Occurences of DNA sequences

in Table 2 and 3. Cluster #4 and 8.1 were chosen, with four bound references used. ATTRACT was chosen for its common use in literature and its ability to process DNA and to perform ensemble docking.

The relevance of redundancy in the benchmark is also clearly shown in our docking experiments. Indeed, in both experiments, bound structures were selected

because the same ssDNA sequence was found interacting with the protein, and protein chains were clustered because of the common ssDNA-interface. However, we can observe small variations between the comparison of one docking result with its four bound references, like in the cluster #4, GCT docking on 5tmf_c exhibits irmsd between 4,2Å and 4,4Å depending on the reference. This can be used to set a minimal rmsd clustering criterion of the resulting poses afterdocking experiment, as here 0.2Å in cluster #4.

mechanisms involved in ssDNA-protein interactions and develop mature ssDNA docking protocols.

Conclusion

Unlike ssRNA, the single stranded form of DNA is almost only found as an intermediate state in the DNA processing mechanisms (like DNA replication³⁶), which may be a reason for the lack of study of ssDNA-containing complexes, and ssDNA-protein interactions, from a structural point of view. Yet, those intermediate states play a crucial role in DNA metabolism, and their interactions with proteins are potential targets for therapeutic inhibitors. For instance, the transmission of anti-microbial resistance genes among bacteria could be fought by targeting the excision or transportation of ICE (Integrative Conjugative Elements) ssDNA⁵. Such projects require the knowledge of the 3D structure of such assemblies for rational drug design. To develop computational methods to model their spatial conformation, it is necessary studying the existing experimental structures of such ssDNA-protein complexes.

In this work, we systematically extracted from the PDB all such structures with at least 4 bound single-stranded DNA nucleotides, together with the corresponding unbound structures of the protein, to evaluate and predict the requirements and potential effectiveness of ssDNA-protein docking. We identified groups of proteins with and without ssDNA-induced fit. An application for this benchmark would be as a training set to develop a machine learning tool identifying *a priori* in which category a protein falls. A docking experiment was also performed to illustrate a usecase of this benchmark.


From a biological point of view, we found that the ssDNA composition is biased towards short fragments and homopolymers, with the last feature representing a third of all protein-bound ssDNA sequences. The low number of retrieved sequences (148) may reflect the general low interest for ssDNA-bound protein structures, and a dataset like the one presented here is an important step in their studies.

To our knowledge, this work is the first attempt to aggregate as exhaustively as possible the ssDNA-protein structures available in the PDB. With 98 groups of multi-conformational bound/unbound proteins, this benchmark is an essential first step to understand the

References

- Watson JD, Crick FH. The structure of DNA. *Cold Spring Harb Symp Quant Biol.* 1953;18:123-131. doi:10.1101/sqb.1953.018.01.020
- Seo YS, Lee SH, Hurwitz J. Isolation of a DNA helicase from HeLa cells requiring the multisubunit human single-stranded DNA-binding protein for activity. *J Biol Chem.* 1991;266(20):13161-13170.
- De La Cruz F, Frost LS, Meyer RJ, Zechner EL. Conjugative DNA metabolism in Gram-negative bacteria. *FEMS Microbiol Rev.* 2010;34(1):18-40. doi:10.1111/j.1574-6976.2009.00195.x
- Nikolay R, Schmidt S, Schlömer R, Deuerling E, Nierhaus KH. Ribosome Assembly as Antimicrobial Target. *Antibiotics.* 2016;5(2). doi:10.3391/antibiotics5020018
- Soler N, Robert E, Chauvot de Beauchêne I, et al. Characterization of a relaxase belonging to the MOBT family, a widespread family in Firmicutes mediating the transfer of ICEs. *Mob DNA.* 2019;10:18. doi:10.1186/s13100-019-0160-9
- Leman AR, Noguchi E. The Replication Fork: Understanding the Eukaryotic Replication Machinery and the Challenges to Genome Duplication. *Genes.* 2013;4(1):1-32. doi:10.3391/genes4010001
- Gardner AF, Kelman Z. Editorial: The DNA Replication Machinery as Therapeutic Targets. *Front Mol Biosci.* 2019;6. doi:10.3389/fmolb.2019.00035
- Li X, Du Y, Du P, et al. SXT/R391 integrative and conjugative elements in Proteus species reveal abundant genetic diversity and multidrug resistance. *Sci Rep.* 2016;6. doi:10.1038/srep37372
- Nishimasu H, Ran FA, Hsu PD, et al. Crystal Structure of Cas9 in Complex with Guide RNA and Target DNA. *Cell.* 2014;156(5):935-949. doi:10.1016/j.cell.2014.02.001
- Büttner K, Nehring S, Hopfner K-P. Structural basis for DNA duplex separation by a superfamily-2 helicase. *Nat Struct Mol Biol.* 2007;14(7):647-652. doi:10.1038/nsmb1246
- Ke A, Doudna JA. Crystallization of RNA and RNA-protein complexes. *Methods.* 2004;34(3):408-414. doi:10.1016/j.ymeth.2004.03.027
- Ravindran PP, Héroux A, Ye J-D. Improvement of the crystallizability and expression of an RNA crystallization chaperone. *J Biochem (Tokyo).* 2011;150(5):535-543. doi:10.1093/jb/mvr093
- Chauvot de Beauchêne I, de Vries SJ, Zacharias M. Binding Site Identification and Flexible Docking of Single Stranded RNA to Proteins Using a Fragment-Based Approach. *PLoS Comput Biol.* 2016;12(1). doi:10.1371/journal.pcbi.1004697
- Kappel K, Das R. Sampling Native-like Structures of RNA-Protein Complexes through Rosetta Folding and Docking. *Structure.* 2019;27(1):140-151.e5. doi:10.1016/j.str.2018.10.001
- Chaudhury S, Lyskov S, Gray JJ. PyRosetta: a script-based interface for implementing molecular modeling algorithms using Rosetta. *Bioinformatics.* 2010;26(5):689-691. doi:10.1093/bioinformatics/btq007
- Hall D, Li S, Yamashita K, Azuma R, Carver JA, Standley DM. RNA-LIM: a novel procedure for analyzing protein/single-stranded RNA propensity data with concomitant estimation of interface structure. *Anal Biochem.* 2015;472:52-61. doi:10.1016/j.ab.2014.11.004
- Coimbatore Narayanan B, Westbrook J, Ghosh S, et al. The Nucleic Acid Database: new features and capabilities. *Nucleic Acids Res.* 2014;42(D1):D114-D122. doi:10.1093/nar/gkt980
- Hwang H, Vreven T, Janin J, Weng Z. Protein-Protein Docking Benchmark Version 4.0. *Proteins.* 2010;78(15):3111-3114. doi:10.1002/prot.22830
- Koukos PI, Faro I, van Noort CW, Bonvin AMJJ. A Membrane Protein Complex Docking Benchmark. *J Mol Biol.* 2018;430(24):5246-5256. doi:10.1016/j.jmb.2018.11.005
- Huang S-Y, Zou X. A nonredundant structure dataset for benchmarking protein-RNA computational docking. *J Comput Chem.* 2013;34(4):311-318. doi:10.1002/jcc.23149
- Nithin C, Mukherjee S, Bahadur RP. A non-redundant protein-RNA docking benchmark version 2.0. *Proteins.* 2017;85(2):256-267. doi:10.1002/prot.25211
- van Dijk M, Bonvin AMJJ. A protein-DNA docking benchmark. *Nucleic Acids Res.* 2008;36(14):e88. doi:10.1093/nar/gkn386
- Pal A, Levy Y. Structure, stability and specificity of the binding of ssDNA and ssRNA with proteins. *PLOS Comput Biol.* 2019;15(4):e1006768. doi:10.1371/journal.pcbi.1006768
- Berman H, Henrick K, Nakamura H, Markley JL. The worldwide Protein Data Bank (wwPDB): ensuring a single, uniform archive of PDB data. *Nucleic Acids Res.* 2007;35(Database issue):D301-D303. doi:10.1093/nar/gkl971
- Analyzing and building nucleic acid structures with 3DNA - PubMed. Accessed September 2, 2021. <https://pubmed.ncbi.nlm.nih.gov/23644419/>
- de Vries SJ, Schindler CEM, Chauvot de Beauchêne I, Zacharias M. A Web Interface for Easy Flexible Protein-Protein Docking with ATTRACT. *Biophys J.* 2015;108(3):462-465. doi:10.1016/j.bpj.2014.12.015
- Setny P, Zacharias M. A coarse-grained force field for Protein-RNA docking. *Nucleic Acids Res.* 2011;39(21):9118-9129. doi:10.1093/nar/gkr636
- He J, Tao H, Huang S-Y. Protein-ensemble-RNA docking by efficient consideration of protein flexibility through homology models. *Bioinformatics.* 2019;35(23):4994-5002. doi:10.1093/bioinformatics/btz388
- Wang H, Zeng F, Liu Q, et al. The structure of the ARE-binding domains of Hu antigen R (HuR) undergoes conformational changes during RNA binding. *Acta Crystallogr D Biol Crystallogr.* 2013;69(3):373-380. doi:10.1107/S0917444912047828
- Epstein J, Cai J, Glaser T, Jepeal L, Maas R. Identification of a Pax paired domain recognition sequence and evidence for DNA-dependent conformational changes. *J Biol Chem.* 1994;269(11):8355-8361.31. Poddar S, Chakravarty D, Chakrabarti P. Structural changes in DNA-binding proteins on complexation. *Nucleic Acids Res.* 2018;46(7):3298-3308. doi:10.1093/nar/gky170
- Cooper A, Dryden DTF. Allostery without conformational change. *Eur Biophys J.* 1984;11(2):103-109. doi:10.1007/BF00276625
- Sankar K, Walia RR, Mann CM, Jernigan RL, Honavar VG, Dobbs D. An analysis of conformational changes upon RNA-protein binding. In: *Proceedings of the 5th ACM Conference on Bioinformatics, Computational Biology, and Health Informatics.* BCB '14. Association for Computing Machinery; 2014:592-593. doi:10.1145/2649387.2660791
- Lexa KW, Carlson HA. Protein Flexibility in Docking and Surface Mapping. *Q Rev Biophys.* 2012;45(3):301-343. doi:10.1017/S0033583512000066
- Marmur J, Doty P. Determination of the base composition of deoxyribonucleic acid from its thermal denaturation temperature. *J Mol Biol.* 1962;5(1):109-118. doi:10.1016/S0022-2836(62)80066-7
- Vos SM, Tretter EM, Schmidt BH, Berger JM. All tangled up: how cells direct, manage and exploit topoisomerase function. *Nat Rev Mol Cell Biol.* 2011;12(12):827-841. doi:10.1038/nrm3228

Supplementary



Supplementary figure 1 : A) Superimposition of (blue) 6qem_a, (green) 6qem_f and (red) 6bbm_d by aligning their N-terminal ends ; B) Hierarchical clustering of cluster 61 non-redundant members from RMSD distance matrix ;