

### GABA-A rhythmic inhibition in the claustrum drives network oscillation

Gilles Ouanounou, Ambre Ledoux, Thierry Bal

#### ▶ To cite this version:

Gilles Ouanounou, Ambre Ledoux, Thierry Bal. GABA-A rhythmic inhibition in the claustrum drives network oscillation. FENS forum 2022, Jul 2022, Paris, France. . hal-03824516

### HAL Id: hal-03824516 https://hal.science/hal-03824516

Submitted on 21 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



## GABA-A rhythmic inhibition in the claustrum drives network oscillation Gilles Ouanounou, Ambre Ledoux and Thierry Bal NeuroPsi, CNRS/Paris-Saclay University, Integrative and Computational Neuroscience department (ICN), Saclay, France

## Introduction

The claustrum is phylogenetically highly preserved in amniotes, from reptiles and birds to mammals, including humans (Norimoto et al., 2020). It is a thin structure that lays deep between the insular cortex and the striatum. It contains a small proportion of interneurons of various types, and a majority of far-projecting excitatory neurons. It is a high-level structure holding the most central position in the brain hierachy, being reciprocally connected to all cerebral cortical areas, to the amygdala, hippocampus, and thalamus. Its functional role is a recent focus of research in neurosciences as it remains largely unknown during both sleep and wakefulness. Due to its high connectivity it has been proposed to be a "seat of consciousness" and involved in multi-sensory binding (Crick & Koch, 2005), a hub for attention (Goll et al., 2015; Atlan et al., 2018; Fodoulian et al., 2020). Most importantly for the present work, the claustrum might orchestrate and integrate cortical activity by coordinating synchronized oscillations in cortical areas during sleep and quiet wakefulness (Narikiyo et al., 2020). The claustrum could act as a synchronizing relay between areas, activating their local GABAergic interneurons (Narikiyo et al., 2020). It could also act during sleep as a "conductor" (Crick & Koch, 2005), driving slow-wave sleep activity on its own rhythm.

Our recordings in the claustrum in vitro show the emergence of a highly regular oscillation in the 0,3-1 Hz range in the presence of low doses of the cholinergic agonist carbachol (CCH). Release of acetylcholine from channelrhodopsin-containing cholinergic afferents also boosts the oscillation (not shown). We found that the oscillation is a synchronized network phenomenon that relies on the occurrence of a rhythmic GABA-A inhibition in the excitatory claustral cells, suggesting the involvement of inhibitory interneurons in the claustrum. We also recorded a subset of claustral neurons with endogenous rhythmic behavior. Whether these (inter)neurons play a pacemaker role in the network oscillation, and what kinds of synaptic loops there are within the claustrum and/or between the claustrum and the adjacent insular cortex, remains to be examined.



different types in the claustrum (purple babana shape) and the entorhinal cortex. Upper trace: Oscillations induced by cholinergic agonist carbachol in the claustrum range between 0,3-1 Hz (see Figs 2). Bottom trace: exemple of typical Up-and-Down oscillation generated through different mechanisms in the entorhinal cortex in the same horizontal slice in absence of Carbachol (0,13 +/- 0,02 Hz; Tahvildari et al., 2012). Mouse brain: Allen Brain Explorer.

## **Putative claustrum-cortex wiring circuit**

Working synaptic diagram of the network present in the horizontal slice derived from Kim et al., 2016, and from our investigations. Present data suggest the existence of highly synchronized GABA-A inhibitions (generated by red interneurons) in claustrum principal cells (green). As suggested in the previous figure, the possible involvement of insular cortex (left) remains to be examined. GABAergic neurons (pink; red); Excitatory neurons (cortex: blue; claustrum: green).





A: Mouse brain left hemisphere from Allen Brain Explorer, with our 2 dissection cuts and the claustrum in purple. **B:** Sketch of a horizontal slice obtained from the horizontal cut in (A). **C:** Horizontal slice pictures under infrared light (IR) and blue laser (Fluo). Yellow rectangles correspond to the green rectangle in (B), within which the mean gray values below are measured (sum of the gray values of all the pixels in the rectangle, divided by the number of pixels). Infrared light shines through the slice, revealing a clear band noticeable as a peak of mean gray value. This area has the same location and shape as the region with enhanced VGLUT2/EGFP-expression cells, which corresponds to the claustrum (Fodoulian, 2020). Cla: Claustrum ; EC: Externa

Capsule (myelinated dark curved line at the internal edge of the Cla); IC: Insular Cortex; St: Striatun ; C: IR: n = 58, Fluo: n = 16.



A. Extracellular unit during application of the cholinergic agonist carbachol. Full-field white light stimulation of the slice (yellow bars) in a emx-creXAi35 P14 transgenic mice expressing Archaerhodopsin kick starts the oscillation. B. Whole-cell recording of a claustrum principal cell. Cholinergic modulation (CCH) makes cla principal cells go from rare activity to rhythmic bursting activity. Autocorrelogram computed on spike counts for 200 ms bins. C. Strongly adapting (SA) claustrum principal cell (wild-type C57bl6; P13; CCH). Hyperpolarized Vm reveals rhythmic synaptic drive.



A. Voltage-clamp (VC) recording at -60 mV with CCH and corresponding autocorrelogram (AC). B. VC recording at 0 mV (CCH). Synaptic excitatory inputs are tonic but there is a periodic change in inhibitory conductances, which are GABA-A-related and could be responsible for the voltage oscillation (same cell as in Fig. 2B) as it occurs at the same frequency. C. IV curves resulting from voltage-clamp ramp protocol. Examples of IV curves obtained at high levels (green) and low levels (red) of inhibitory conductance as seen in (B). On average, the reversal potential was  $-73.1 \text{ mV} \pm 1.7 \text{SD}$ , n=6 cells.



**A.** Recording of a pair of excitatory principal cells in the claustrum of a VGLUT2 mouse (Fodoulian, 2020) using two patch pipettes. B. Picture of the two biocytin-filled cells. Squares are details of the spiny dendrites of each cell shown in (C-D) at higher magnifications. E. Current clamp recording of these two archearhodopsin-containing cells showing hyperpolarizations subsequent to 575 nm light exposition, confirming that those are VGLUT2-expressing cells and thus principal neurons. **F.** Dual voltage-clamp recording of cells 1 and 2 at 0 mV under carbachol. Traces are so similar that the synchronized rhythmic inhibitory synaptic inputs we detect with auto- (G, left) and cross- (G, right) correlograms, must presumably come from one or a few interneurons only.



# 

# A pacemaker hypothesis A subset of neurons with intrinsic oscillatory membrane properties 0 G

A. Composite picture of a neuron presenting no dendritic spines, suggesting that it could be an interneuron subtype. B. Firing response to a current pulse C. Current-clamp recording of the neuron. In absence of carbachol (CCH) the membrane potential is quiescent. **D.** CCH induces membrane oscillation which might be an endogenous membrane property. E. Voltage-clamp at -60 mV (left) with corresponding autocorrelogram (right) showing random tonic excitatory synaptic bombardment, as in principal cells. F. Voltage-clamp at 0 mV showing no rhythmic inhibitions. G. Voltage-clamp at 0 mV in another cell showing strong rythmic inhibitions (notice different time scale).

# Conclusion

In a mouse brain slice containing portions of the claustrum and insular cortex, we previously found that claustral excitatory principal cells undergo a slow oscillation under light cholinergic modulation. Here transgenic mice were used to help identify a simple location criterion for the claustrum: its relative clarity under light shown through the living slices. Then we performed single and dual whole-cell patch-clamp recordings in order to characterize the oscillation as well as its synaptic underlying mechanisms.

We showed that the claustrum oscillation is a network phenomenon involving primarily excitatory principal cells simultaneously driven by a tonic excitatory drive and rhythmic GABA-A inhibitions from presumably local interneurons. In addition, we found a putative interneuron subtype that presents endogenous oscillatory membrane properties revealed only in the presence of the cholinergic agonist CCH and under the influence of a tonic excitatory synaptic drive. We propose that this intrinsic oscillator may play a crucial role in the network oscillation by driving it as a pacemaker. It remains to be examined whether the insular cortex, which is synaptically connected to the claustrum, could be involved in the oscillatory network mechanism via a simple excitatory drive or via a synaptic loop with the claustral network (see "putative claustrum-cortex wiring circuit).

## **References**:

Atlan et al., (Curr. Biol., 2018) The Claustrum Supports Resilience to Distraction. Atilgan, H. et al. (Brain 145, 1610–1623 (2022). Human lesions and animal studies link the claustrum to percep-Crick & Koch (Philos Trans R Soc Lond B Biol Sci., 2005) What is the function of the claustrum? et àl. (bioRxiv. 2020) austrum-medial prefrontal cortex network controls attentional set-shifting Goll et al., (Trends Neurosci., 2015) Attention: the claustrum Kim et al., (J. Neurosc., 2016). Synaptic Organization of the Neuronal Circuits of the Claustrum. Narikiyo et al., (Nature Neuroscience, 2020) The Claustrum Coordinates Cortical Slow-Wave Activity. Norimoto, H. et al. (Nature, 2020). A claustrum in reptiles and its role in slow-wave sleep. Tahvildari, et al., (J. Neurosc., 2012). Selective Functional Interactions between Excit. and Inhib. Cortical Neu-

Acknowledgments: Thanks to M. pananceau, C. Monier, A. Daret, C De Medeiros, V. Lavalée; and J. Bouvier (NeuroPSI) for providing mice. D.A. McCormick and B. Tahvildari (U.Oregon and Yale, USA), A. Foust (Imperial, London) for technical support, and M. Descamps-Bal for valuable insights.