



HAL
open science

On a family of holomorphic self-maps of the unit disk

Jean-Marc Sac-Epee, Myriam Ounaïes

► **To cite this version:**

Jean-Marc Sac-Epee, Myriam Ounaïes. On a family of holomorphic self-maps of the unit disk. Results in mathematics = Resultate der Mathematik, 2022, 77, pp.226. <10.1007/s00025-022-01759-5>. <hal-03825023>

HAL Id: hal-03825023

<https://hal.science/hal-03825023v1>

Submitted on 21 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

On a family of holomorphic self-maps of the unit disk

M. Ounaies and J.-M. Sac-Épée*

August 25, 2022

Abstract

We give a characterization of the sets D_p ($1 < p < 2$) of complex numbers c such that $z \mapsto \frac{1+z}{2} + c \left(\frac{1-z}{2}\right)^p$ is a self-map of the closed unit disk and we show that these sets are increasing with respect to p .

1 Introduction

We will denote by \mathbb{D} the open unit disk, by $\overline{\mathbb{D}}$ the closed unit disk and by $C = \overline{\mathbb{D}} \setminus \mathbb{D}$ the unit circle. For $p \geq 0$ and $c \in \mathbb{C}$, let $f_{p,c}$ be the map, analytic over \mathbb{D} and continuous over $\overline{\mathbb{D}}$ defined by

$$f_{p,c}(z) = \frac{1+z}{2} + c \left(\frac{1-z}{2}\right)^p, \quad (1.1)$$

where the principal branch of the logarithm on $\mathbb{C} \setminus \mathbb{R}^-$ is used to define the p 'th powers. Note that for $z \in \overline{\mathbb{D}} \setminus \{1\}$, $\Re\left(\frac{1-z}{2}\right) > 0$. In [3], Mortini and Rhin raised the question of determining precisely the set

$$D_p = \{c \in \mathbb{C} : f_{p,c} \text{ is a self-map of } \overline{\mathbb{D}}\}$$

for $p > 1$. They proved that $[0, 1] \subseteq D_p$ for $p \geq 1$.

The case $0 \leq p \leq 1$ was settled in [2], where the authors showed the following results:

- $D_0 = \{z \in \mathbb{C} : |z + \frac{1}{2}| \leq \frac{1}{2}\}$,
- $D_1 = [-1, 1]$,
- For all $0 < p < 1$, $D_p = \{-\left(\frac{1-a}{2}\right)^{1-p}, a \in \overline{\mathbb{D}}\}$.
- The regions $\{D_p\}_{0 \leq p < 1}$ are decreasing.

*M. Ounaies, IRMA, France, myriam.ounaies@math.unistra.fr
J.-M. Sac-Épée, IECL, France, jean-marc.sac-epée@univ-lorraine.fr

2000 Mathematics Subject Classification: 30A10; 30C99

Keywords : Holomorphic self-maps of the unit disk; Complex inequalities

In another paper (see [1]), Mortini and Sac-Épée showed that

$$D_2 = \left\{ z \in \mathbb{C} : \left| z - \frac{1}{4} \right| \leq \frac{3}{4} \right\} \quad (1.2)$$

and they made the conjecture, based on numerical results, that the regions $\{D_p\}_{p>1}$ are increasing.

In the present work, we will focus on the case where $1 < p \leq 2$. In section 1, we will prove that the regions $\{D_p\}_{1 < p \leq 2}$ are increasing and in section 2, we will give a description of D_p for $1 < p < 2$.

Throughout this paper, we will use the function $g_{p,c}$ defined as follows for $p \geq 0$, $c \in \mathbb{C}$ and $t \in \mathbb{R}$,

$$g_{p,c}(t) = (1+t^2)^p - |c + it(1+it)^{p-1}|^2. \quad (1.3)$$

We will extensively make use of the next proposition which is contained in [1]. We include here a short proof for the sake of completeness.

Proposition 1.1. *Let $p \geq 0$. Then $c \in D_p$ if and only if for all $t \in \mathbb{R}$, $g_{p,c}(t) \geq 0$.*

Proof. Thanks to the maximum principle applied to the analytic function $f_{p,c}$ and the fact that 1 is a fixed point of $f_{p,c}$, we see that $c \in D_p$ if and only if for all $z \in C \setminus \{1\}$, $|f_{p,c}(z)| \leq 1$.

For all $z \in C \setminus \{1\}$, we write $z = -e^{-2i\theta}$ with $\theta \in]-\frac{\pi}{2}, \frac{\pi}{2}[$ and we put $t = \tan \theta$. Thus we have

$$\begin{aligned} \frac{1-z}{2} &= \frac{1+e^{-2i\theta}}{2} = \cos \theta e^{-i\theta} = (1+it)^{-1}, \\ \frac{1+z}{2} &= \frac{1-e^{-2i\theta}}{2} = i \sin \theta e^{-i\theta} = it(1+it)^{-1}, \\ \frac{1+z}{2} + c \left(\frac{1-z}{2} \right)^p &= (1+it)^{-p} (c + it(1+it)^{p-1}). \end{aligned}$$

With this setting, $c \in D_p$ if and only if for all $t \in \mathbb{R}$,

$$(1+t^2)^{-\frac{p}{2}} |c + it(1+it)^{p-1}| \leq 1$$

which is equivalent to $g_{p,c}(t) \geq 0$. ■

We will denote by

$$D_p^+ = D_p \cap \{z \in \mathbb{C} : \Im z \geq 0\}; \quad D_p^- = D_p \cap \{z \in \mathbb{C} : \Im z \leq 0\}.$$

Remark 1.2. Clearly, for all $c \in \mathbb{C}$ and for all $t \in \mathbb{R}$, $g_{p,c}(t) = g_{p,\bar{c}}(-t)$, thus

$$c \in D_p^+ \iff \bar{c} \in D_p^-.$$

From now on for a complex number $c \neq 0$, we will denote by $\theta_c \in]-\pi, \pi]$ the principal argument of c .

2 The regions $\{D_p\}_{1 < p \leq 2}$ are increasing.

Theorem 2.1. *If $1 < p < q \leq 2$, then $D_p \subset D_q$.*

Before proving this theorem, we need two lemmas.

Lemma 2.2. *Let $1 < p < 2$. Then $c \in D_p^+$ if and only if the two following conditions are satisfied:*

- a) $\theta_c \in [0, (p-1)\frac{\pi}{2}]$;
- b) For all $t \in [0, \tan(\frac{\theta_c}{p-1})]$, $g_{p,c}(t) \geq 0$.

In particular, $D_p \cap \mathbb{R} = [0, 1]$.

Proof. Recall that by definition (1.3), $g_{p,c}(t) = (1+t^2)^p - |c+it(1+it)^{p-1}|^2$. Let us prove the following formula:

$$g_{p,c}(t) = (1+t^2)^{p-1} - |c|^2 - 2t(1+t^2)^{\frac{p-1}{2}} |c| \sin(\theta_c - (p-1) \arctan t). \quad (2.1)$$

Note that

$$|c+it(1+it)^{p-1}|^2 = |c|^2 + t^2(1+t^2)^{p-1} + 2t(1+t^2)^{\frac{p-1}{2}} |c| \cos(\theta_c - (p-1) \arctan t - \frac{\pi}{2}).$$

In the previous line, the expression inside the cosine comes from the fact that $(p-1) \arctan t$ is the principal argument of $(1+it)^{p-1}$, so that $(p-1) \arctan t + \frac{\pi}{2}$ is the principal argument of $i(1+it)^{p-1}$.

Thus

$$\begin{aligned} g_{p,c}(t) &= (1+t^2)(1+t^2)^{p-1} - |c+it(1+it)^{p-1}|^2 \\ &= (1+t^2)^{p-1} - |c|^2 - 2t(1+t^2)^{\frac{p-1}{2}} |c| \cos(\theta_c - (p-1) \arctan t - \frac{\pi}{2}), \end{aligned}$$

and (2.1) follows. Assume that $c \in D_p^+$. Then by definition of D_p^+ , $0 \leq \theta_c \leq \pi$. Since $1 < p < 2$, we deduce that

$$-\frac{\pi}{2} \leq -(p-1)\frac{\pi}{2} \leq \theta_c - (p-1)\frac{\pi}{2} \leq \theta_c \leq \pi.$$

Now we multiply (2.1) by t^{-p} for $t > 0$, and we observe that

$$t^{-p} g_{p,c}(t) = t^{-(2-p)}(t^{-2}+1)^{p-1} - |c|^2 t^{-p} - 2(t^{-2}+1)^{\frac{p-1}{2}} |c| \sin(\theta_c - (p-1) \arctan t).$$

We have

$$\lim_{t \rightarrow +\infty} t^{-p} g_{p,c}(t) = -2|c| \sin\left(\theta_c - (p-1)\frac{\pi}{2}\right). \quad (2.2)$$

By Proposition 1.1, $\sin(\theta_c - (p-1)\frac{\pi}{2}) \leq 0$ which implies condition a).

The necessity of condition b) is obvious by Proposition 1.1.

Reciprocally, assume that conditions a) and b) are satisfied.

Let $t \geq \tan(\frac{\theta_c}{p-1})$. Then $-\frac{\pi}{2} \leq \theta_c - (p-1) \arctan t \leq 0$ which means that

$$g_{p,c}(t) \geq (1+t^2)^{p-1} - |c|^2 \geq 1 - |c|^2 = g_{p,c}(0) \geq 0.$$

Now we only need to verify that for $t < 0$, $g_{p,c}(t) \geq 0$.
If $t < 0$, a simple computation shows that

$$g_{p,c}(t) - g_{p,c}(-t) = -4t(1+t^2)^{\frac{p-1}{2}}|c|\sin\theta_c\cos((p-1)\arctan t).$$

Since $-\frac{\pi}{2} < (p-1)\arctan t < 0$, we have $\cos((p-1)\arctan t) \geq 0$. Thus

$$g_{p,c}(t) > g_{p,c}(-t) \geq 0.$$

Hence $g_{p,c}(t) \geq 0$ for all $t \in \mathbb{R}$ and so, by Proposition 1.1, $c \in D_p^+$. ■

Remark 2.3. In the case where $p = 2$, (2.2) is replaced by

$$\lim_{t \rightarrow +\infty} t^{-p}g_{2,c}(t) = 1 + 2\Re c,$$

and we retrieve the fact that $D_2 \subset \{\Re z \geq -\frac{1}{2}\}$ but we don't get the necessity of condition (a). However, it is easy to deduce from the proof of Lemma 2.2 the following:

Assume that $\theta_c \in [0, \frac{\pi}{2}]$, then $c \in D_2$ if and only if for all $t \in [0, \tan\theta_c]$, $g_{2,c}(t) \geq 0$.

Lemma 2.4. *Let $1 < p < q \leq 2$ and $c \in \mathbb{C}$ such that $\theta_c \in [0, (p-1)\frac{\pi}{2}]$. Then for all $t \in [0, \tan(\frac{\theta_c}{q-1})]$,*

$$g_{q,c}(t) \geq (1+t^2)^{q-p}g_{p,c}(t).$$

Proof. For all $t \in \mathbb{R}$ we have

$$\begin{aligned} g_{q,c}(t) - (1+t^2)^{q-p}g_{p,c}(t) &= ((1+t^2)^{q-p} - 1)|c|^2 + \\ &+ 2t(1+t^2)^{\frac{q-1}{2}}|c|\left((1+t^2)^{\frac{q-p}{2}}\sin(\theta_c - (p-1)\arctan t) - \sin(\theta_c - (q-1)\arctan t)\right). \end{aligned}$$

Now for all $0 \leq t \leq \tan(\frac{\theta_c}{q-1})$,

$$0 \leq \theta_c - (q-1)\arctan t \leq \theta_c - (p-1)\arctan t \leq \theta_c \leq \frac{\pi}{2}$$

and consequently

$$\begin{aligned} 0 \leq \sin(\theta_c - (q-1)\arctan t) &\leq \sin(\theta_c - (p-1)\arctan t) \\ &\leq (1+t^2)^{\frac{q-p}{2}}\sin(\theta_c - (p-1)\arctan t). \end{aligned}$$

The desired result follows immediately. ■

Proof of Theorem 2.1. Consider p and q such that $1 < p < q \leq 2$ and let $c \in D_p$. In view of Remark 1.2, we may assume that $c \in D_p^+$. Then by Lemma 2.2 we have

$$0 \leq \theta_c \leq (p-1)\frac{\pi}{2} \leq (q-1)\frac{\pi}{2}.$$

Let $t \in [0, \tan(\frac{\theta_c}{q-1})]$. Then by Proposition 1.1 and Lemma 2.4, we have

$$g_{q,c}(t) \geq (1+t^2)^{q-p}g_{p,c}(t) \geq 0.$$

Now we use Lemma 2.2 (case $q < 2$) and Remark 2.3 (case $q = 2$) to conclude that $c \in D_q^+$. ■




Figure 1: $h_p(t) = (p-1)t \tan((2-p) \arctan t^{-1} + \arctan(pt))$
 $l_p(t) = 1 + (2-p)t^2$.

3 Description of D_p , $1 < p < 2$.

We know from Lemma 2.2 that $D_p \cap \mathbb{R} = [0, 1]$. Our next theorem gives an explicit expression of the remaining elements of D_p .

Theorem 3.1. *Let $1 < p < 2$. Then $c \in D_p$ if and only if either $c \in [0, 1]$ or c is of the form*

$$c = (1 + it)^{p-2}(1 - ipt)^{-1}(\alpha + i(p-1)t)$$

where $t \neq 0$ and α are such that

$$(p-1)t \tan((2-p) \arctan t^{-1} + \arctan(pt)) \leq \alpha \leq 1 + (2-p)t^2. \quad (3.1)$$

In order to carry out the proof of this theorem, we need two preliminary lemmas.

Lemma 3.2.

Let $1 < p \leq 2$ and assume that $\theta_c \in [0, (p-1)\frac{\pi}{2}]$. Then the following properties are satisfied:

- (i) $g_{p,c}(t) \rightarrow +\infty$ as $|t| \rightarrow +\infty$,
- (ii) $g_{p,c}$ has a unique critical point $t_{p,c}$ on \mathbb{R} ,
- (iii) If $\theta_c = 0$, then $t_{p,c} = 0$ and if $\theta_c > 0$, then $0 < t_{p,c} < \tan\left(\frac{\theta_c}{p-1}\right)$.

Obviously, (i) and (ii) imply that $g_{p,c}$ attains its minimum at $t_{p,c}$.

Proof. Property (i) follows from (2.1) and

$$\sin\left(\theta_c - (p-1)\frac{\pi}{2}\right) \leq 0, \quad \sin\left(\theta_c + (p-1)\frac{\pi}{2}\right) \geq 0.$$

Recall that by definition (1.3), for all $t \in \mathbb{R}$,

$$g_{p,c}(t) = (1 + t^2)^p - |c + it(1 + it)^{p-1}|^2.$$

To prove (ii) and (iii), let us compute the derivative of $g_{p,c}$:

$$\begin{aligned}
g'_{p,c}(t) &= 2(pt(1+t^2)^{p-1} + \Re(i(c+it(1+it)^{p-1})(1-it)^{p-2}(1-ipt))) \\
&= 2(1+t^2)^{p-2} (pt(1+t^2) + \Re(i(c+it(1+it)^{p-1})(1+it)^{2-p}(1-ipt))) \\
&= 2(1+t^2)^{p-2} (pt(1+t^2) - \Im(c(1+it)^{2-p}(1-ipt)) - t\Re((1+it)(1-ipt))) \\
&= 2(1+t^2)^{p-2} ((p-1)t - \Im(c(1+it)^{2-p}(1-ipt))) \\
&= 2(1+t^2)^{p-2} \left((p-1)t + |c|(1+t^2)^{\frac{2-p}{2}} (1+p^2t^2)^{\frac{1}{2}} \sin(\phi_{p,c}(t)) \right),
\end{aligned} \tag{3.2}$$

where we have denoted by

$$\phi_{p,c}(t) = \arctan(pt) - (2-p)\arctan t - \theta_c.$$

Observe that whenever $t < 0$, one has the inequalities

$$-\pi < \arctan(pt) - \theta_c \leq \phi_{p,c}(t) \leq (p-1)\arctan t - \theta_c < 0,$$

thus $g'_{p,c}(t) < 0$.

Now we compute the derivative of $\phi_{p,c}$:

$$\phi'_{p,c}(t) = (p-1) \frac{2 + p(p-1)t^2}{(1+t^2)(1+p^2t^2)} \geq 0.$$

Note that for all $t \geq 0$,

$$-\frac{\pi}{2} \leq -\theta_c \leq \phi_{p,c}(t) \leq \arctan(pt) \leq \frac{\pi}{2}.$$

We conclude that the function $t \mapsto \sin(\phi_{p,c}(t))$ is increasing on $[0, +\infty[$ and so is the function $t \mapsto (1+t^2)^{2-p}g'_{p,c}$. Therefore, $g'_{p,c}$ has at most one zero in $[0, +\infty[$.

Now let us prove the existence of a zero $t_{p,c}$ of $g'_{p,c}$.

If $\theta_c = 0$ then obviously $g'_{p,c}(0) = 0$ and we put $t_{p,c} = 0$.

If $\theta_c \in]0, (p-1)\frac{\pi}{2}[$, then $g'_{p,c}(0) = -2\Im c < 0$. Since $g_{p,c}(t) \rightarrow +\infty$ as $t \rightarrow +\infty$, the derivative $g'_{p,c}$ has to vanish at some point $t_{p,c} > 0$.

Besides, if $\theta_c \in]0, (p-1)\frac{\pi}{2}[$, let us put $T = \tan\left(\frac{\theta_c}{p-1}\right)$. Then by (2.1),

$$g_{p,c}(T) = (1+T^2)^{p-1} - |c|^2 > 1 - |c|^2 = g_{p,c}(0).$$

We deduce that $0 < t_{p,c} < T$. ■

Remark 3.3. In the case where $p = 2$, we may solve explicitly the equation $g'_{2,c}(t) = 0$, which gives

$$t_{2,c} = \frac{\Im c}{1 + 2\Re c}$$

Lemma 3.4. *Let $1 < p \leq 2$.*

(i) *Let $t \in \mathbb{R}$. Then $g'_{p,c}(t) = 0$ if and only if c is of the form*

$$c = (1+it)^{p-2}(1-ipt)^{-1}(\alpha + i(p-1)t), \quad \alpha \in \mathbb{R}. \tag{3.3}$$

(ii) For all $t > 0$, $0 < (2-p) \arctan t^{-1} + \arctan(pt) < \frac{\pi}{2}$.

(iii) Assume that c is of the form (3.3) with $t > 0$ and $\alpha > 0$. Then $\theta_c \geq 0$ and we have the following equivalences:

$$g_{p,c}(t) \geq 0 \iff \alpha \leq 1 + (2-p)t^2; \quad (3.4)$$

$$\theta_c \leq (p-1)\frac{\pi}{2} \iff (p-1)t \tan((2-p) \arctan t^{-1} + \arctan(pt)) \leq \alpha. \quad (3.5)$$

Proof. (i) is an obvious consequence of formula (3.2).

For $t > 0$, define $h_p(t) = (2-p) \arctan t^{-1} + \arctan(pt)$. Then the derivative

$$h'_p(t) = \frac{(p-1)(2+p(p-1)t^2)}{(1+t^2)(1+p^2t^2)}$$

is obviously positive, which implies (ii).

Finally, let us prove (iii). Assume that c is of the form (3.3) with $t > 0$ and $\alpha > 0$. The argument of c is given by

$$\theta_c = \arctan(pt) - (2-p) \arctan t + \arctan\left(\frac{(p-1)t}{\alpha}\right)$$

Clearly

$$\theta_c \geq \arctan(pt) - \arctan t > 0.$$

To prove (3.5), it is sufficient to observe that

$$\theta_c = (p-1)\frac{\pi}{2} + (2-p) \arctan t^{-1} + \arctan(pt) - \arctan\left(\frac{\alpha}{(p-1)t}\right).$$

$$(1+t^2)^{2-p}(1+p^2t^2)g_{p,c}(t) = (1+t^2)^2 - (\alpha + (p-1)t^2)^2.$$

Now since $c = (1+it)^{p-2}(1-ipt)^{-1}(\alpha + i(p-1)t)$, we have

$$\begin{aligned} c + it(1+it)^{p-1} &= (1+it)^{p-2}(1-ipt)^{-1}(\alpha + i(p-1)t + it(1+it)(1-ipt)) \\ &= (1+it)^{p-2}(1-ipt)^{-1}(\alpha + (p-1)t^2 + ipt(1+t^2)) \end{aligned}$$

Thus

$$\begin{aligned} |c + it(1+it)^{p-1}|^2 &= (1+t^2)^{p-2}(1+p^2t^2)^{-1}((\alpha + (p-1)t^2)^2 + p^2t^2(1+t^2)^2) \\ &= (1+t^2)^{p-2}(1+p^2t^2)^{-1}((\alpha + (p-1)t^2)^2 - (1+t^2)^2) + (1+t^2)^p. \end{aligned}$$

By the definition (1.3) of $g_{p,c}$,

$$\begin{aligned} g_{p,c}(t) &= (1+t^2)^p - |c + it(1+it)^{p-1}|^2 \\ &= (1+t^2)^{p-2}(1+p^2t^2)^{-1}((1+t^2)^2 - (\alpha + (p-1)t^2)^2). \end{aligned}$$

and we readily deduce (3.4). ■

Proof of Theorem 3.1.

Necessity:

Thank to Remark 1.2 we may assume that $c \in D_p^+$.

If $\theta_c = 0$, then by Lemma 2.2, $c \in [0, 1]$.

If $\theta_c > 0$, then by Lemma 2.2, $\theta_c \in]0, (p-1)\frac{\pi}{2}]$. Consider $t = t_{p,c} > 0$ given by Lemma 3.2. Lemma 3.4 gives a real number α such that

$$c = (1 + it)^{p-2}(1 - ipt)^{-1}(\alpha + i(p-1)t).$$

Thus we have

$$\begin{aligned} \alpha &= \Re(c(1 + it)^{2-p}(1 - ipt)) \\ &= |c|(1 + t^2)^{\frac{2-p}{2}}(1 + p^2t^2)^{\frac{1}{2}} \cos(\theta_c + (2-p)\arctan t - \arctan(pt)). \end{aligned}$$

Thanks to the inequalities

$$-\frac{\pi}{2} < -\arctan(pt) \leq \theta_c + (2-p)\arctan t - \arctan(pt) \leq \theta_c + (1-p)\arctan t < \theta_c \leq \frac{\pi}{2},$$

we deduce that $\alpha > 0$.

Condition (3.1) follows from Lemma 3.4.

Sufficiency:

We know from Lemma 2.2 that $D_p \cap \mathbb{R} = [0, 1]$.

Assume that $c = (1 + it)^{p-2}(1 - ipt)^{-1}(\alpha + i(p-1)t)$ with $t \neq 0$ and α satisfying (3.1). Observe that $\alpha > 0$. Again thanks to Remark 1.2 we may assume that $t > 0$. Using lemma 3.4 we know that $g'_{p,c}(t) = 0$, $\theta_c \in [0, (p-1)\frac{\pi}{2}]$ and $g_{p,c}(t) \geq 0$.

By lemma 3.2, we necessarily have $t = t_{p,c}$ which means that the minimum of $g_{p,c}$ on \mathbb{R} is attained at t . Using Proposition 1.1, we deduce that $c \in D_p$. ■

Remark 3.5. In the case where $p = 2$, we may repeat the proof of Theorem 3.1 and use Remark 2.3 to obtain the following:

$$\begin{aligned} D_2 \cap \{\Re z \geq 0\} &= \{(1 - 2it)^{-1}(\alpha + it), (t, \alpha) \in \mathbb{R}^2 : 2t^2 \leq \alpha \leq 1\} \\ &= \left\{ z \in \mathbb{C} : \left| z - \frac{1}{4} \right| \leq \frac{3}{4} \right\} \cap \{\Re z \geq 0\}. \end{aligned}$$

The second equality comes from (1.2) but may also be proved directly by the following formal equivalences for $\alpha, t, a, b \in \mathbb{R}$ and $c = a + ib$:

$$\begin{aligned} c = \frac{\alpha + it}{1 - 2it} &\iff \frac{1}{2c + 1} = \frac{1 - 2it}{2\alpha + 1} \\ &\text{by taking real and imaginary parts, we obtain} \\ \frac{1}{2\alpha + 1} &= \frac{2a + 1}{|2c + 1|^2} \text{ and } \frac{t}{2\alpha + 1} = \frac{b}{|2c + 1|^2} \\ \iff \alpha &= \frac{2|c|^2 + a}{2a + 1} = 1 + 2\frac{|c - \frac{1}{4}|^2 - \frac{9}{16}}{2a + 1} \text{ and } t = \frac{b}{1 + 2a}. \end{aligned}$$

Note that

$$c = \frac{\alpha - 2t^2}{1 + 4t^2} + i\frac{t(1 + 2\alpha)}{1 + 4t^2}.$$




Figure 2: D_p for three particular values of p

4 Numerical illustration

For illustrating purposes, we use the characterization given by Theorem 3.1 to draw the sets D_p for three particular values of p (Figure 2). The circle with center $\frac{1}{4}$ and radius $\frac{3}{4}$, which is the boundary of D_2 , appears in red. We also provide a hyperlink to an animated image showing the evolution of the D_p 's for p varying from 1.01 to 1.99 with a step of 0.007 ([4]).

We may point out that our Figure 2 and Figure 2 in [1] show a discrepancy when p is close to 2. This is due to the fact that, while our graphs are based on theoretical results, those in [1] are obtained experimentally in the following way: many complex numbers c have been randomly drawn, and each of them has been tested and kept only if it survives a test consisting in verifying that $|f_{p,c}(z)| \leq 1$ for numerous random values of z (thanks to the maximum principle, one takes random z such that $|z| = 1$). The experimental results obtained in [1] for p close to 2 are not validated by the present theoretical study.

5 Acknowledgement

The authors are very grateful to the anonymous referee for her/his valuable advice, which greatly improved the structure and clarity of this work.

References

- [1] R. Mortini, J.-M. Sac-Épée, Complex inequalities involving sums of holomorphic selfmaps of the unit disk and some experimental conjectures. *Complex Anal. Synerg.* 5 (2019), no. 2-4, Paper No. 12, 4 pp. <https://doi.org/10.1007/s40627-019-0037-1>
- [2] R. Mortini, R. Rupp, Sums of holomorphic selfmaps of the unit disk, *Annales Univ. Mariae Curie-Skłodowska* 61 (2007), 107-115.
- [3] R. Mortini, G. Rhin, Sums of holomorphic selfmaps of the unit disk II. *Comput. Methods Funct. Theory* 11 (2011), no. 1, 135-142
- [4] M. Ounaies, J.-M. Sac-Épée <https://plmbox.math.cnrs.fr/f/e68f07608bfd464c8825/>