



# Ocean monitoring indicator (OMI) of the Copernicus marine environment monitoring service global yearly ocean CO<sub>2</sub> sink

Thi Tuyet Trang Chau, Marion Gehlen, Frederic Chevallier

## ► To cite this version:

Thi Tuyet Trang Chau, Marion Gehlen, Frederic Chevallier. Ocean monitoring indicator (OMI) of the Copernicus marine environment monitoring service global yearly ocean CO<sub>2</sub> sink. [Research Report] LSCE. 2021. hal-03826303v2

HAL Id: hal-03826303

<https://hal.science/hal-03826303v2>

Submitted on 10 Aug 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# Quality Information Document

Ocean Monitoring Indicator (OMI) of the  
Copernicus Marine Environment Monitoring Service

## Global yearly ocean CO<sub>2</sub> sink

Issue: 5.0

Contributors : Thi Tuyet Trang Chau, Marion Gehlen, Frédéric Chevallier

Approval date by the CMEMS product quality coordination team : January 2023



MERCATOR OCEAN  
INTERNATIONAL

2 avenue de l'Aérodrome de Montaudran, 31400 Toulouse, FRANCE

Tél : +33 5 61 39 38 02 - Fax : +33 5 61 39 38 99

Société civile de droit français au capital de 2 000 000 € - 522 911 577 RCS Toulouse - SIRET 522 911 577 00016

[marine.copernicus.eu](http://marine.copernicus.eu)

[mercator-ocean.eu](http://mercator-ocean.eu)

CHANGE RECORD

Issue	Date	§	Description of Change	Author	Validated By
1.0	2019/10	all	Creation of the document	Marion Gehlen, Thi Tuyet Trang Chau	
2.0	2019/12	/	Update of Figure, references Modification of method description and validation	Thi Tuyet Trang Chau, Marion Gehlen, Frédéric Chevallier	
3.0	2020/09	/	Update of Figure, references Modification of evaluation results	Thi Tuyet Trang Chau, Marion Gehlen, Frédéric Chevallier	
4.0	2021/09	/	Update of Figure, references Modification of evaluation results	Thi Tuyet Trang Chau, Marion Gehlen, Frédéric Chevallier	Stéphanie Guinehut
4.1	2021/11		Correction following MOI review	Thi Tuyet Trang Chau, Marion Gehlen, Frédéric Chevallier	Sandrine Mulet
5.0	2022/09	/	Update of Figure, references Modification of evaluation results	Thi Tuyet Trang Chau, Marion Gehlen, Frédéric Chevallier	M. Drevillon



<b>I</b>	<b>INTRODUCTION .....</b>	<b>4</b>
<b>I.1</b>	<b>Product name .....</b>	<b>4</b>
<b>I.2</b>	<b>Indicator evaluation method .....</b>	<b>4</b>
<b>I.3</b>	<b>CMEWS products used as input for the computations .....</b>	<b>4</b>
<b>II</b>	<b>SCIENTIFIC EVALUATION.....</b>	<b>5</b>
<b>III</b>	<b>REFERENCES .....</b>	<b>7</b>



## I INTRODUCTION

---

### I.1 Product name

GLOBAL\_OMI\_HEALTH\_carbon\_co2\_flux\_integrated


### I.2 Indicator evaluation method

The OMI global yearly ocean CO<sub>2</sub> sink represents the ocean uptake of CO<sub>2</sub> from the atmosphere computed over the whole ocean. It is computed by multiplying the 12-month running mean of the surface ocean downward CO<sub>2</sub> mass flux (fgco2) [molC m<sup>-2</sup> yr<sup>-1</sup>], a variable from the MULTIOBS\_GLO\_BIO\_CARBON\_SURFACE\_REP\_015\_008 product, by the area of each grid cell, and then taking the sum over all cells. The unit is finally converted to PgC per year (Figure 1). The yearly estimate of the ocean CO<sub>2</sub> sink corresponds to the mean of a 100-member ensemble (see the method described in Chau et al. (2022) and in the QUID for GLOBAL\_OMI\_HEALTH\_carbon\_ph\_area\_averaged). In line with Friedlingstein et al. (2022), yearly uncertainties are estimated as their corresponding ensemble standard deviation (1σ).

### I.3 CMEMS products used as input for the computations

Product name (and Type)	Documentation
MULTIOBS_GLO_BIO_CARBON_SURFACE_REP_015_008	QUID: <a href="https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MOB-QUID-015-008.pdf">https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MOB-QUID-015-008.pdf</a> PUM: <a href="https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MOB-PUM-015-008.pdf">https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MOB-PUM-015-008.pdf</a>

## II SCIENTIFIC EVALUATION



**Figure 1:** Area-integrated yearly surface downward flux of total CO<sub>2</sub> for 1985-2021. Plain line: mean of 100 samples; range: empirical 68 % confidence interval (mean estimate  $\pm$  standard deviation) computed from 100 samples. Trend and uncertainty are defined as the slope and its residual standard deviation estimated with a linear least-squares regression.

The global ocean CO<sub>2</sub> sink corresponds to the global integral of fgco<sub>2</sub> over space and time. The rate of uptake increases by  $0.04 \pm 0.03e-1 \text{ PgC/yr}^2$  over the full period. The yearly flux time series presents a plateau in the 90s followed by an increasing growth rate of  $0.06 \pm 0.04e-1 \text{ PgC/yr}^2$  since 2000. In 2021 [resp. 2020], the global ocean CO<sub>2</sub> sink was  $2.41 \pm 0.13$  [resp.  $2.50 \pm 0.12$ ] PgC/yr (Figure 1). The average of yearly contemporary fluxes over the full period 1985-2020 is  $1.61 \pm 0.10 \text{ PgC/yr}$  with an interannual variability (temporal standard deviation) of  $0.46 \text{ PgC/yr}$ .

The scientific evaluation relies on the comparison with estimates of the anthropogenic CO<sub>2</sub> uptake by the ocean that are published by the [Global Carbon Budget 2021](#) (GCB2021, Friedlingstein et al. 2022). The reconstructed field covers approximately 96.15% of the total ocean area ( $361.9e6 \text{ km}^2$ ). In order to compare the present estimates to Friedlingstein et al. (2022), contemporary fluxes are corrected for the total area and the preindustrial outgassing of riverine carbon by adding  $0.61 \text{ PgC/yr}$  which is in between the estimate by Jacobson et al. (2007) ( $0.45 \pm 0.18 \text{ PgC/yr}$ ) and the one by Resplandy et al. (2018) ( $0.78 \pm 0.41 \text{ PgC/yr}$ ). This



adjustment yields an anthropogenic flux estimate of  $3.15 \pm 0.13$  [resp.  $3.25 \pm 0.12$ ] PgC/yr for 2021 [resp. 2020]. This is in line with the estimate by Friedlingstein et al. (2022) of  $2.9 \pm 0.4$  [resp.  $3.0 \pm 0.4$ ] PgC/yr.

The latter estimate corresponds to the mean of flux estimates from an ensemble of global ocean biogeochemical general circulation models (GOBMs) and an ensemble of data-based reconstruction methods. The GCB2021 uncertainty estimate combines random uncertainty (ensemble standard deviation of model- and data-based flux estimates) and systematic uncertainty (bias of model-based estimates in anthropogenic carbon cumulation). The new ensemble-based approach (Chau et al. 2022) yields spatio-temporal varying uncertainties which are lower than the previous published total estimate of  $\pm 0.4$  PgC/yr. The uncertainty estimates of the annual CO<sub>2</sub> uptake for the years in the 2000s are smaller than the uncertainties estimated for the years in the 1980s or recent years (Figure 1), as the sea surface spco2 observation network densifies. As shown in the following table, the somewhat higher anthropogenic flux estimates of the ensemble approach fall within the full range ( $\pm 3\sigma$ ) of those by Friedlingstein et al. (2022). Further refinement of ocean C sink estimates relies on improved and sustained ocean observation efforts.

Periods	OMI_HEALTH_carbon_co2_flux_integrated [PgC/yr] (Mean $\pm$ uncertainty)		Friedlingstein et al. (2022) - anthropogenic flux estimate [PgC/yr]
	Contemporary flux estimate	Anthropogenic flux estimate (see text for definition)	
2021	$2.41 \pm 0.13$	$3.15 \pm 0.13$	$2.9 \pm 0.4$
2020	$2.50 \pm 0.12$	$3.25 \pm 0.12$	$3.0 \pm 0.4$
2011-2020	$2.18 \pm 0.09$	$2.91 \pm 0.10$	$2.8 \pm 0.4$
2000-2009	$1.53 \pm 0.08$	$2.22 \pm 0.08$	$2.2 \pm 0.4$
1990-1999	$1.30 \pm 0.10$	$1.98 \pm 0.10$	$2.0 \pm 0.4$
1985-1989	$1.03 \pm 0.15$	$1.69 \pm 0.16$	-

A full discussion regarding this OMI can be found in section 2.10 of the Ocean State Report 4 (Gehlen et al., 2020) and in Chau et al. (2022).



### III REFERENCES

---

Chau, T. T. T., Gehlen, M., and Chevallier, F.: A seamless ensemble-based reconstruction of surface ocean pCO<sub>2</sub> and air-sea CO<sub>2</sub> fluxes over the global coastal and open oceans, *Biogeosciences*, 19, 1087–1109, <https://doi.org/10.5194/bg-19-1087-2022>, 2022.

Gehlen M., Thi Tuyet Trang Chau, Anna Conchon, Anna Denvil-Sommer, Frédéric Chevallier, Mathieu Vrac, Carlos Mejia (2020). Ocean acidification. In: Copernicus Marine Service Ocean State Report, Issue 4, *Journal of Operational Oceanography*, 13:sup1, s88–s91; DOI: 10.1080/1755876X.2020.1785097

Friedlingstein, P., Jones, M. W., O'Sullivan, M., Andrew, R. M., Bakker, D. C. E., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Bates, N. R., Becker, M., Bellouin, N., Bopp, L., Chau, T. T. T., Chevallier, F., Chini, L. P., Cronin, M., Currie, K. I., Decharme, B., Djeutchouang, L. M., Dou, X., Evans, W., Feely, R. A., Feng, L., Gasser, T., Gilfillan, D., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Luijkx, I. T., Jain, A., Jones, S. D., Kato, E., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lienert, S., Liu, J., Marland, G., McGuire, P. C., Melton, J. R., Munro, D. R., Nabel, J. E. M. S., Nakao, S.-I., Niwa, Y., Ono, T., Pierrot, D., Poulter, B., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Schwingershackl, C., Séférian, R., Sutton, A. J., Sweeney, C., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Tubiello, F., van der Werf, G. R., Vuichard, N., Wada, C., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, C., Yue, X., Zaehle, S., and Zeng, J.: Global Carbon Budget 2021, *Earth Syst. Sci. Data*, 14, 1917–2005, <https://doi.org/10.5194/essd-14-1917-2022>, 2022.

Jacobson, A. R., Mikaloff Fletcher, S. E., Gruber, N., Sarmiento, J. L., and Gloor, M. (2007), A joint atmosphere-ocean inversion for surface fluxes of carbon dioxide: 1. Methods and global-scale fluxes, *Global Biogeochem. Cycles*, 21, GB1019, doi:[10.1029/2005GB002556](https://doi.org/10.1029/2005GB002556).

Resplandy, L., Keeling, R. F., Rödenbeck, C., Stephens, B. B., Khatiwala, S., Rodgers, K. B., Long, M. C., Bopp, L. and Tans, P. P.: Revision of global carbon fluxes based on a reassessment of oceanic and riverine carbon transport. *Nature Geoscience*, 11(7), p.504, 2018.