



Global ocean acidification - mean sea water pH trend map from Multi-Observations Reprocessing

Thi Tuyet Trang Chau, Marion Gehlen, Frederic Chevallier

► To cite this version:

Thi Tuyet Trang Chau, Marion Gehlen, Frederic Chevallier. Global ocean acidification - mean sea water pH trend map from Multi-Observations Reprocessing. [Research Report] LSCE; EU. 2022. hal-03826462

HAL Id: hal-03826462

<https://hal.science/hal-03826462>

Submitted on 24 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Quality Information Document

Ocean Monitoring Indicator (OMI) of the
Copernicus Marine Environment Monitoring
Service

Global surface ocean pH trends

Issue: 1.1

Contributors : **T. T. Trang Chau, Marion Gehlen, Frédéric Chevallier**

Approval date by the CMEMS product quality coordination team : **January 2023**



**MERCATOR OCEAN
INTERNATIONAL**

2 avenue de l'Aérodrome de Montaudran, 31400 Toulouse, FRANCE

Tél : +33 5 61 39 38 02 - Fax : +33 5 61 39 38 99

Société civile de droit français au capital de 2 000 000 € - 522 911 577 RCS Toulouse - SIRET 522 911 577 00016

marine.copernicus.eu

mercator-ocean.eu



CHANGE RECORD

Issue	Date	§	Description of Change	Author	Validated By
1.0	January 2022	All	Creation of the document	Thi Tuyet Trang Chau, Marion Gehlen, Frédéric Chevallier	S. Mulet
1.1	2022/09	/	Update results with respect to the new release of input data (see I.3)	Thi Tuyet Trang Chau, Marion Gehlen, Frédéric Chevallier	M. Drevillon



I	INTRODUCTION	3
I.1	Product name.....	3
I.2	Indicator evaluation method.....	3
I.3	CMEWS products used as input for the computations.....	5
II	SCIENTIFIC EVALUATION	6
III	REFERENCES.....	9



I INTRODUCTION

I.1 Product name

GLOBAL_OMI_HEALTH_carbon_ph_trend

I.2 Indicator evaluation method

Secular trends and the associated uncertainties ($\pm 1\sigma$) of seawater pH over the global surface ocean are computed from the 100-member ensemble of pH time series. This ensemble covering the period since 1985 was randomly regenerated with a Gaussian distribution where the mean and the standard deviation are, respectively, the best estimate and the total uncertainty of pH available in the Surface Carbon field **MULTIOBS_GLO_BIO_CARBON_SURFACE REP_015_008** (Chau et al., 2022). Surface ocean pH is reported on total scale and the units of its trend and uncertainty are yr^{-1} .

Denote μ_{ij} and σ_{ij} as a linear trend and its uncertainty computed from a batch of 100 yearly mean pH time series ($pH_{t,ij,n}$) at each $1^\circ \times 1^\circ$ -grid cell ij , where t stands for time and n indicates a member in the 100-member ensemble. A linear least-squares regression was used to fit a linear function between two sets of data (i.e., $pH_{t,ij,n}$ against t values) and derive estimates of its slope (μ_{ij}) and residual standard deviation (σ_{ij}). An illustration of the method of linear fits on the pH data at some stations is shown in Figure 1.

The global maps of pH trend and uncertainty are presented in Figure 2 (a,b). A threshold on $\frac{\sigma}{|\mu|}$ of 10% is empirically defined to mask the regions with uncertain estimates of pH local trend (Figures 2c, 2d). The global pH trend map and this

mask are reproduced in Figure 3 - a representative map for this new OMI for surface ocean pH trend.




Figure 1. Linear fits on the 100-member ensemble of yearly mean pH at different stations (see also in Table 1). Grey points represent pH data in the ensemble and black lines stand for their linear fits. $\mu_{ij} \pm \sigma_{ij}$ denotes estimates of a linear slope and its uncertainty over 1985-2021.






Figure 2. (a) Global surface seawater pH trend and (b) uncertainty computed over the period 1985-2021. (c) Ratio ($r_{ij}[\%]$) between uncertainty estimates (σ_{ij}) and the best trend estimates (μ_{ij}). (d) Mask of the regions where ratios (r_{ij}) between uncertainty estimates and local trends exceed 10%.

I.3 CMEMS products used as input for the computations

Product name (and Type)	Documentation
MULTIOBS_GLO_BIO_CARBON_SURFACE_REP_015_008	QUID: https://catalogue.marine.copernicus.eu/documents/QUID/CMEMS-MOB-QUID-015-008.pdf PUM: https://catalogue.marine.copernicus.eu/documents/PUM/CMEMS-MOB-PUM-015-008.pdf



II SCIENTIFIC EVALUATION

For the period since 1985 pH is decreasing at a rate between -0.0008 yr^{-1} and -0.0022 yr^{-1} over most of the global ocean basins (Figure 3). The spatial pattern of pH secular trends is generally consistent with that presented in Lauvset et al (2015). Nearly all values of the trend exceed the associated uncertainty estimate (Figure 2b) but we conservatively exclude the ones associated with the largest uncertainty estimates, i.e., ratio between the uncertainty and trend estimates larger than 10% (Figure 2c), knowing that they may be underestimated. Only a portion of data over some sectors of the Arctic, the Antarctic, and the coastal ocean has been masked indicating that there might be no certain estimates of pH trend (Figures 2 and 3).

Table 1 shows a comparison between the new OMI pH trends and those derived from pH speciation or measurements at 7 stations presented in Bates et al (2014). The location of these time series stations is plotted in Figure 4. The CMEMS OMI trend at each location is computed from monthly pH time series in the **MULTIOBS_GLO_BIO_CARBON_SURFACE REP_015_008** dataset at the four grid boxes nearby. As seen in Table 1, the CMEMS OMI trend identified by for HOTS ($-0.0018 \pm 0.0001 \text{ yr}^{-1}$), BATS ($-0.0014 \pm 0.0001 \text{ yr}^{-1}$), and ESTOC ($-0.0019 \pm 0.0001 \text{ yr}^{-1}$) are in close agreement with those computed from observations in the previous studies (Dore et al., 2009; Gonzalez-Davila et al., 2010; Bates et al., 2014). Our estimate of pH trend at Irminger for 1985-2012 is about $-0.0014 \pm 0.0001 \text{ yr}^{-1}$ which is nearly twice as small as that by Bates et al 2014 but falls within the full range ($\pm 3\sigma$) of the estimate by Pérez et al (2021) for 1983-2013 ($-0.0017 \pm 0.00019 \text{ yr}^{-1}$). The magnitude of pH trend computed here for Iceland Sea [Mudina] is smaller [larger] than reported in Bates et al (2014). However, because of the low sampling frequency the comparison should be viewed with caution. CARIACO is located on the east coast of South America, its four nearest model grid boxes with available CMEMS pH data result in an average latitude with an offset of approximately 2° from this station and thus a mismatch from the estimate by Bates et al 2014 is expected (Table 1).

Overall, secular trends of seawater pH over the global surface ocean have a complex spatial pattern and their estimates are sensitive to physical and carbonate variables fed in pH speciation (Lauvset et al., 2015; Pérez et al., 2021).

Up to present, direct measurements of surface ocean pH are still sparse (Bates et al., 2014; Lauvset et al., 2021), an increase in the observational networks for pH and other carbonate variables is important for understanding and analyzing changes in seawater pH as well as for a better evaluation of their estimates (Lauvset et al., 2015).




Figure 3. Global map of pH trend over the period 1985-2021. The white mask indicates the regions where pH trends with the highest uncertainty estimates are excluded (see indicator evaluation method).






Figure 4. Location of seven time-series stations recording pH and carbonate variables (Bates et al., 2014).

Table 1: Comparison in surface ocean pH trend estimates for the seven time-series stations (Bates et al., 2014). Uncertainty is reported as $\pm 1\sigma$. The CMEMS OMI trend at each station is computed by learning on monthly pH time series in the MULTIOBS_GLO_BIO_CARBON_SURFACE_REP_015_008 dataset at the four grid boxes nearby.

Data	Stations	Iceland Sea	Irminger Sea	BATS	ESTOC	HOT	CARIAC O	Munida
Bates et al. 2014	Years	1983 – 2012	1983 – 2012	1983 – 2012	1995 – 2012	1988 – 2012	1995 – 2012	1998 – 2012
	Latitude, Longitude	68°N, 12.66°W	64.3°N, 28°W	32°N, 64°W	29.04°N, 15.50°W	22.75°N, 158°W	10.5°N, 64.66°W	45.7°S, 171.5°E
	Sampling frequency (per year)	3–4	3–4	14–16	4–6	10	12	~ 4–6
	pH trend (pH data type)	-0.0014 ± 0.0005 (calculated)	-0.0026 ± 0.0006 (calculated)	-0.0017 ± 0.0001 (calculated)	-0.0018 ± 0.0002 (calculated)	-0.0016 ± 0.0001 (measured)	-0.0025 ± 0.0004 (measured)	-0.0013 ± 0.0003 (calculated)
CMEMS-OMI-QUID-GLO-HEALTH-carbon-ph-trend	Years	1985 – 2012	1985 – 2012	1985 – 2012	1995 – 2012	1988 – 2012	1995 – 2012	1998 – 2012
	Latitude, Longitude	68°N, 13°W	64°N, 28°W	32°N, 64°W	29°N, 15°W	23°N, 158°W	12.5°N, 65°W	46.5°S, 172°E
	pH trend	-0.0009 ± 0.0001	-0.0014 ± 0.0001	-0.0014 ± 0.0001	-0.0019 ± 0.0001	-0.0018 ± 0.0001	-0.0015 ± 0.0001	-0.0018 ± 0.0002



III REFERENCES

Bates, N. R., Astor, Y. M., Church, M. J., Currie, K., Dore, J. E., Gonzalez-Davila, M., Lorenzoni, L., Muller-Karger, F., Olafsson, J., and Magdalena Santana-Casiano, J.: A Time-Series View of Changing Surface Ocean Chemistry Due to Ocean Uptake of Anthropogenic CO₂ and Ocean Acidification, *Oceanography*, 27, 126–141, 2014.

Chau, T. T. T., Gehlen, M., Chevallier, F.: Global Ocean Surface Carbon : MULTIOBS_GLO_BIO_CARBON_SURFACE REP_015_008, E.U. Copernicus Marine Service Information, DOI:[10.48670/moi-00047](https://doi.org/10.48670/moi-00047), 2022.

Dore, J.E., Lukas, R., Sadler, D.W., Church, M.J., and Karl, D.M. (2009). Physical and biogeochemical modulation of ocean acidification in the central North Pacific. *Proceedings of the National Academy of Sciences USA*, 106, 12235–40.

González-Dávila, M., Santana-Casiano, J. M., Rueda, M. J., and Llinás, O. (2010). The water column distribution of carbonate system variables at the ESTOC site from 1995 to 2004. *Biogeosciences*, 7, 3067–3081.

Lauvset, S. K., Gruber, N., Landschützer, P., Olsen, A., and Tjiputra, J.: Trends and drivers in global surface ocean pH over the past 3 decades, *Biogeosciences*, 12, 1285–1298, DOI:10.5194/bg-12-1285-2015, 2015.

Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Álvarez, M., Becker, S., Brown, P. J., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E., Jutterström, S., Jones, S. D., Karlsen, M. K., Lo Monaco, C., Michaelis, P., Murata, A., Pérez, F. F., Pfeil, B., Schirnick, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Velo, A., Wanninkhof, R., Woosley, R. J., and Key, R. M.: An updated version of the global interior ocean biogeochemical data product, GLODAPv2.2021, *Earth Syst. Sci. Data*, 13, 5565–5589, DOI:10.5194/essd-13-5565-2021, 2021.

Pérez FF, Olafsson J, Ólafsdóttir SR, Fontela M, Takahashi T. Contrasting drivers and trends of ocean acidification in the subarctic Atlantic. *Sci Rep* 11, 13991, DOI:10.1038/s41598-021-93324-3, 2021.