



HAL
open science

Annealed limit for a diffusive disordered mean-field model with random jumps

Xavier Erny

► **To cite this version:**

Xavier Erny. Annealed limit for a diffusive disordered mean-field model with random jumps. 2022.
hal-03826883v1

HAL Id: hal-03826883

<https://hal.science/hal-03826883v1>

Preprint submitted on 24 Oct 2022 (v1), last revised 14 Sep 2023 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Annealed limit for a diffusive disordered mean-field model with random jumps

Xavier Erny

Ecole Polytechnique, Centre de mathématiques appliquées (CMAP), 91128 Palaiseau

Abstract: We study a sequence of N -particle mean-field systems, each driven by N simple point processes $Z^{N,i}$ in a random environment. Each $Z^{N,i}$ has the same intensity $(f(X_{t-}^N))_t$ and at every jump time of $Z^{N,i}$, the process X^N does a jump of height U_i/\sqrt{N} where the U_i are disordered centered random variables attached to each particle. We prove the convergence in distribution of X^N to some limit process \bar{X} that is solution to an SDE with a random environment given by a Gaussian variable, with a convergence speed for the finite-dimensional distributions. This Gaussian variable is created by a CLT as the limit of the partial sums of the U_i . To prove this result, we use a coupling for the classical CLT relying on the result of [Komlós, Major and Tusnády \(1976\)](#), that allows to compare the conditional distributions of X^N and \bar{X} given the environment variables, with the same Markovian technics as the ones used in [Erny, Löcherbach and Loukianova \(2022\)](#).

MSC2020 subject classifications: 60K37, 60J35, 60J25, 60J60; secondary 60F05, 60G50, 60G55.

Keywords and phrases: Annealed limit in random environment, Central limit theorem coupling, Piecewise deterministic Markov processes, Mean-field model.

1. Introduction

The term disordered comes from physics litterature to designate some "assymmetric systems". Particle system in random environment can often be seen as disordered model, since, in this kind of model, it is possible to attach to each particle (or each pair of particles) a random variable, interpreted as a disordered variable, creating asymmetric interactions between the particles. This is for example the case of models with spin glass dynamics, where for each pair of particle (i, j) , there exists a disordered variable of the form $S_i S_j V_{ij}$, where S_i and S_j are typically $\{-1, +1\}$ -valued variables modelling the spins of the particles i and j , and V_{ij} is a random variable modelling an interaction strength between the two particles. This kind of model has been introduced in the seminal paper of [Sherrington and Kirkpatrick \(1975\)](#).

In the model we study in this paper, unlikely with spin glass dynamics model, we attach disordered variables to each particle (and not to each pair of particles) that are i.i.d. and independent of the time, like in the very similar model of [Pfaffelhuber, Rotter and Stiefel \(2022\)](#). More precisely, we study the solutions of stochastic differential equations with a drift term and a jump term that depends on the disordered variables. There is a natural application of this kind of model in neurosciences (as explained in Section 4 of [Pfaffelhuber, Rotter and Stiefel \(2022\)](#)), where the equations model the dynamics of the membrane potentials of neurons, the jump times model the times at which the neurons receive spikes by another neuron of the network, and the jump heights (given by the disordered variables) model the synaptic weights in the network. The drift term models the (deterministic) dynamics of the membrane potential between its jump times.

The property we want to prove is the convergence in distribution as the number of particles of the system goes to infinity. As we work in a random environment, the convergence of the processes

can be understood in two ways: a quenched convergence (i.e. convergence conditionally on the environment) or an annealed convergence (i.e. convergence when the environment variables are averaged). We refer to [Ben Arous and Guionnet \(1995\)](#) and [Guionnet \(1997\)](#) for the definitions of the terms "quenched" and "annealed" that we use in this paper.

The aim of the paper is to prove the convergence in distribution of a process $(X_t^N)_t$ defined in a random environment. Let us define rigorously this process. If the environment $(u_j^{[N]})_{1 \leq j \leq N} \in \mathbb{R}^N$ is fixed, we define $(X_t^N(u^{[N]}))_t$ as the solution of the following SDE:

$$dX_t^N(u^{[N]}) = b(X_t^N(u^{[N]}))dt + \frac{1}{\sqrt{N}} \sum_{j=1}^N u_j^{[N]} \int_{\mathbb{R}_+} 1_{\{z \leq f(X_t^N(u^{[N]}))\}} d\pi_j(t, z), \quad (1)$$

where b and f are deterministic functions and π_j ($1 \leq j \leq N$) are independent Poisson measures of intensity $dt \cdot dz$, that are independent of X_0^N . We note ν_0^N the distribution of X_0^N . By definition, the process $(X_t^N)_t$ is defined as

$$X_t^N := X_t^N(U^{[N]}), \quad (2)$$

where the variables $U_j^{[N]}$ ($1 \leq j \leq N$) are i.i.d. centered random variables with finite variance σ^2 , that are independent of the Poisson measures π_j ($1 \leq j \leq N$) and of the initial condition X_0^N . We note μ the law of these variables. This law is independent of N and is a parameter of the model. For the sake of notation, instead of defining for each N -particle system a family of N variables $U_j^{[N]}$ ($1 \leq j \leq N$), we introduce a countable sequence of random variables U_j ($j \leq 1$). By definition, we set $U^{[N]} := (U_1, \dots, U_N)$ (e.g. the first N variables of $U^{[N+1]}$ are exactly $U^{[N]}$). In particular the $[N]$ in superscript will be dropped.

The limit of $(X_t^N)_t$ is shown to be a process $(\bar{X}_t)_t$ that is also defined in a random environment. For a fixed $w \in \mathbb{R}$, let $(\bar{X}_t(w))_t$ be defined as the solution of

$$d\bar{X}_t(w) = b(\bar{X}_t(w))dt + wf(\bar{X}_t(w))dt + \sigma\sqrt{f(\bar{X}_t(w))}dB_t, \quad (3)$$

where B is a standard one-dimensional Brownian motion that is independent of \bar{X}_0 . We note $\bar{\nu}_0$ the distribution of \bar{X}_0 . The limit process $(\bar{X}_t)_t$ is defined as

$$\bar{X}_t := \bar{X}_t(W), \quad (4)$$

where W is Gaussian variable with parameter $(0, \sigma^2)$ that is independent of B, \bar{X}_0 .

Heuristically, a simple way to obtain the limit equation (3)–(4) from (1)–(2) consists in writing the SDE of $(X_t^N)_t$ as

$$dX_t^N = b(X_t^N)dt + \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \int_{\mathbb{R}_+} 1_{\{z \leq f(X_t^N)\}} d\tilde{\pi}_j(t, z) + \left(\frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right) f(X_t^N)dt,$$

where $\tilde{\pi}_j(dt, dz) := \pi_j(dt, dz) - dt dz$ is the compensated Poisson measure of π_j ($1 \leq j \leq N$). Under this form, the second term of the SDE above is a locale martingale (conditionally on the environment) whose jump heights vanish as N goes to infinity. So, in the limit equation, it creates the Brownian term in (3). And the third term in the SDE above is a drift term that corresponds to the second term of (3) since, according to the CLT, the variable $N^{-1/2} \sum_{j=1}^N U_j$ converges in distribution to $W \sim \mathcal{N}(0, \sigma^2)$.

The main result of the paper is the convergence in distribution of the process $(X_t^N)_t$ to the process $(\bar{X}_t)_t$. Our approach only allows to prove the annealed convergence of the processes, because there is a tricky problem concerning the quenched result: for a convergence to be true in our model, we need to guarantee $N^{-1/2} \sum_{j=1}^N U_j$ to converge. This convergence can hold only if the variables are centered, so, if we work conditionally on the variables U_j , the previous sum will necessary explode as N goes to infinity. This why, intuitively, the quenched convergence seems hard to obtain in our framework. Even if, in our proofs, we establish a quenched control, we are not able to obtain a quenched convergence (i.e. the upper-bound of the control does not vanish for a.e. trajectory of the environment), but only an annealed convergence (i.e. the upper-bound vanishes averaging the environment). This point is explained more formally below in the introduction and precisely in Remark 2.7 in the proofs.

The dynamics (1)–(2) and (3)–(4) are similar to respectively equations (3.9) and (3.10) of Pfaffelhuber, Rotter and Stiefel (2022). Indeed, if we consider $b(x) := -\alpha x$ for the drift function in our model, we obtain exactly the same dynamics as Pfaffelhuber, Rotter and Stiefel (2022) for the convolution kernel $\varphi(t) := e^{-\alpha t}$. Consequently, our main result Theorem 1.2 can be compared to Theorem 2 of Pfaffelhuber, Rotter and Stiefel (2022). However, the proof of Pfaffelhuber, Rotter and Stiefel (2022) cannot be used to prove Theorem 1.2 since it relies on the assumption that the environment variables U_j ($j \geq 1$) follow Radon distribution, whereas we only assume the environment variables to be centered with some exponential moments. Note that it would also not be possible to prove Theorem 2 of Pfaffelhuber, Rotter and Stiefel (2022) with the proof of our Theorem 1.2 since it requires to have some Markovian structure (conditionally to the environment variables), which is not the case for Hawkes processes with general convolution kernel as in Pfaffelhuber, Rotter and Stiefel (2022). Another interesting point of Theorem 1.2 is that we have an explicit convergence speed.

The model of this paper is also close to the one of Erny, Löcherbach and Loukianova (2022), except that in Erny, Löcherbach and Loukianova (2022) the environment variables U_j ($1 \leq j \leq N$) in (1) were replaced by centered marks of the point processes $(Z_t^{N,j})_t$ defined as

$$Z_t^{N,j} := \int_{[0,t] \times \mathbb{R}_+} \mathbf{1}_{\{z \leq f(X_{s-}^N)\}} d\pi_j(s, z).$$

In term of neurosciences, the model of Erny, Löcherbach and Loukianova (2022), contrary to the model (1)–(2) and Pfaffelhuber, Rotter and Stiefel (2022), is not consistent with the following biological property: the role of a synapse cannot change (i.e. it is either always excitatory or always inhibitory). Mathematically, if the rate function f is monotone, it means that every time some particle i interacts with another particle j through a synaptic strength U_i , the sign of U_i is always the same. This property is not guaranteed in Erny, Löcherbach and Loukianova (2022), where the U_i are marks of some point processes (and hence they change at each spiking time of the same neuron), but it holds true in the model of this paper where they are fixed environment variables.

The model of this paper is a priori harder to study because, in Erny, Löcherbach and Loukianova (2022), the processes $(X_t^N)_t$ (and their limit) are Markov processes and semimartingales, which is not the case here because of the random environment. In order to apply results from the theories of Markov processes and semimartingales, we need to work conditionally on the environment and, in a second time, to integrate over the environment.

To work conditionally on the environment in a proper manner, we use a coupling between the variables U_j ($j \geq 1$) of (2) and the Gaussian variable W of (4), corresponding to a coupling result for the classical CLT. To be more precise and formal, we construct a sequence of identically distributed

and non-independent Gaussian variables $(W^{[N]})_N$ such that, for every $N \geq 2$,

$$\left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j - W^{[N]} \right| \leq K \frac{\ln N}{\sqrt{N}}, \quad (5)$$

where $K > 0$ is a random variable independent of N . This coupling and the control that we use on the random variable K rely on Theorem 1 of [Komlós, Major and Tusnády \(1976\)](#). Indeed, if we assume that the distribution μ admits some exponential moments: there exists $\alpha > 0$ such that

$$\int_{\mathbb{R}} e^{\alpha|x|} d\mu(x) < \infty,$$

then, it is possible (following the reasoning of Section 7.5 of [Ethier and Kurtz \(2005\)](#) that relies on Theorem 1 of [Komlós, Major and Tusnády \(1976\)](#)) to construct on the same probability space (possibly enlarged) as $(U_j)_{j \geq 1}$ a standard one-dimensional Brownian motion $(\beta_t)_t$ such that, the random variable K defined as

$$K := \sup_{N \geq 2} \left| \sum_{j=1}^N U_j - \sigma \beta_N \right| / \ln N$$

is finite almost surely and admits exponential moments (this is stated and proved in Lemma 3.1 for self-containedness). Then, defining $W^{[N]}$ in the following way

$$W^{[N]} := \sigma \beta_N / \sqrt{N}$$

gives exactly (5). This construction is recalled at Section 3.1.

The coupling (5) allows us to compare the conditional distributions of the processes $(X_t^N)_t$ and $(\bar{X}_t^N)_t$ given the environment variables (where \bar{X}^N is defined as $\bar{X}(W^{[N]})$ in (4)). This result can be interpreted as a quenched control between the distributions of the processes $(X_t^N)_t$ and $(\bar{X}_t^N)_t$, however this is not a quenched convergence. Roughly speaking, the bound (almost sure w.r.t. the environment) we obtain by comparing the two conditional distributions is of the form

$$e^{CW^{[N]}} K \frac{\ln N}{\sqrt{N}}$$

for some positive deterministic constant C . Recalling that $W^{[N]} := \sigma \beta_N / \sqrt{N}$ for some standard Brownian motion β , it is of common knowledge that the limit superior (resp. inferior) of $W^{[N]}$ is infinity (resp. minus infinity) almost surely. Hence we cannot guarantee the bound above to vanish almost surely. The reason why the annealed convergence can be deduced from this bound is that the distribution of $W^{[N]}$ is $\mathcal{N}(0, \sigma^2)$ and so does not depend on N .

To prove this result, we need introduce formally \mathcal{E} the sigma-field related to the random environment:

$$\mathcal{E} := \sigma(U_j : j \geq 1) \vee \sigma(W^{[N]} : N \in \mathbb{N}^*).$$

And, in order to compare the conditional distributions of $(X_t^N)_t$ and $(\bar{X}_t^N)_t$, we introduce their (conditional) semigroups $(P_{\mathcal{E},t}^N)_t$ and $(\bar{P}_{\mathcal{E},t}^N)_t$, and their infinitesimal generators $A_{\mathcal{E}}^N$ and $\bar{A}_{\mathcal{E}}^N$ w.r.t. \mathcal{E} (see Section 1.1 for the definitions). Using our coupling, we obtain a bound for the difference

of the generators for sufficiently smooth test-functions, and deduce a bound for the semigroups using the following formula (which can be found in Lemma 1.6.2 of [Ethier and Kurtz \(2005\)](#) and in equation (3.1) of [Erny, Löcherbach and Loukianova \(2022\)](#) with different terminologies and hypotheses): for g smooth enough, $t \geq 0$ and $x \in \mathbb{R}$,

$$P_{\mathcal{E},t}^N g(x) - \bar{P}_{\mathcal{E},t}^N g(x) = \int_0^t P_{\mathcal{E},t-s}^N (A_{\mathcal{E}}^N - \bar{A}_{\mathcal{E}}^N) \bar{P}_{\mathcal{E},s}^N g(x) ds. \quad (6)$$

Note that, to use the formula above, one needs to guarantee some regularity properties on the limit semigroup $(\bar{P}_{\mathcal{E},t}^N)_t$. This is proved using the regularity of the stochastic flow of the process $(\bar{X}_t^N)_t$ (with a proof similar as the one of Proposition 3.4 of [Erny, Löcherbach and Loukianova \(2022\)](#)).

Let us finally mention that the same kind of model (i.e. particle systems directed by SDEs where the jump term depends on random environment variables) has already been studied in normalization N^{-1} in [Chevallier et al. \(2019\)](#) and more recently in [Agathe-Nerine \(2022\)](#). In both of these references, the random environment relies on the spatial structure of the particle system. To the best of our knowledge, [Pfaffelhuber, Rotter and Stiefel \(2022\)](#) is the first paper about the convergence of this type of model in normalization $N^{-1/2}$. However this kind of convergence concerning models where a drift term is driven by a random environment in normalization $N^{-1/2}$ have already been proved: e.g. [Ben Arous and Guionnet \(1995\)](#), [Guionnet \(1997\)](#) and [Dembo, Lubetzky and Zeitouni \(2021\)](#). And, in a similar framework, [Luçon \(2011\)](#) has proved a quenched convergence of the fluctuations of a particle system (with a drift term driven by a random environment) in normalization N^{-1} , which is a similar regime. Note that the tricky problem concerning the quenched convergence mentioned in the previous paragraph (also related to Remark 3.2) is also encountered in [Luçon \(2011\)](#).

Organization. In Section 1.1, we introduce the notation that we use throughout the paper. The assumptions and the main result are stated in Section 1.2. Section 2 is dedicated to prove the main result Theorem 1.2. Finally, Section 3 gathers some technical proofs of lemmas used in Section 2. The CLT coupling is formally recalled and its main property is proved in Section 3.1.

1.1. Notation

In the paper, we use the following notation:

- For $T > 0$, we note $D([0, T], \mathbb{R})$ (resp. $D(\mathbb{R}_+, \mathbb{R})$) the set of càdlàg functions defined on $[0, T]$ (resp. \mathbb{R}_+) endowed with Skorohod topology (see for example Section 12 (resp. Section 16) of [Billingsley \(1999\)](#)).
- For $n \in \mathbb{N}^*$, $C_b^n(\mathbb{R})$ denotes the set of real-valued functions defined on \mathbb{R} that are C^n such that all their derivatives (up to order n) are bounded.
- For $n \in \mathbb{N}^*$, and $g \in C_b^n(\mathbb{R})$, we note

$$\|g\|_{n,\infty} := \sum_{k=0}^n \|g^{(k)}\|_{\infty}.$$

- For ν_1, ν_2 distributions on \mathbb{R} with finite first order moments, we use the notation $d_{KR}(\nu_1, \nu_2)$ for the Kantorovich-Rubinstein metric between ν_1 and ν_2 . This quantity is defined as:

$$d_{KR}(\nu_1, \nu_2) := \sup_g \left\{ \int_{\mathbb{R}} g(x) d\nu_1(x) - \int_{\mathbb{R}} g(x) d\nu_2(x) \right\},$$

where the supremum is taken over the Lipschitz continuous functions $g : \mathbb{R} \rightarrow \mathbb{R}$ whose Lipschitz constants are non-greater than one.

- If $(X_t)_{t \geq 0}$ is a real-valued Markov process, $(X_t^{(x)})_{t \geq 0, x \in \mathbb{R}}$ denotes its stochastic flow. In other words, for $x \in \mathbb{R}$, $(X_t^{(x)})_t$ is the process starting at position $x \in \mathbb{R}$ that is defined by the dynamics of $(X_t)_{t \geq 0}$. In addition, if the stochastic flow is n times differentiable w.r.t. the space variable x at time t , we note $\partial_x^n X_t^{(x)}$ the related derivative.
- We note U_j ($j \geq 1$) the environment variables introduced in (1)–(2) (see Assumption 1 for the hypotheses satisfied by these random variables) and σ^2 their variance. The variables $W^{[N]}$ ($N \geq 1$) coupled with the U_j ($j \geq 1$) are introduced in (5) (or alternatively at Section 3.1). Each $W^{[N]}$ is used to define the version of the limit equation (3)–(4) where the environment is coupled with the one of the N -th equation (1)–(2).
- In all the paper, we always note (Ω, \mathbb{P}) the probability spaces on which we work (only the environments of the equations are coupled, so all the processes need not be defined on the same space). They are assumed to be complete and large enough to define what we need. They will not be given explicitly.
- We note \mathcal{E} the sigma-field of the environment variables:

$$\mathcal{E} := \sigma(U_j : j \geq 1) \vee \sigma(W^{[N]} : N \in \mathbb{N}^*).$$

In addition, $\mathbb{E}_{\mathcal{E}}$ designates the conditional expectation given \mathcal{E} , and $\mathbb{P}_{\mathcal{E}}$ the related probability measure (i.e. $\mathbb{P}_{\mathcal{E}}(A) = \mathbb{E}_{\mathcal{E}}[1_A]$). If ν is a distribution on \mathbb{R} , we may note $\mathbb{E}_{\mathcal{E}, \nu}[h(X^N)]$ the expectation conditionally on \mathcal{E} under which X_0^N has law ν , and a similar notation for \bar{X}^N .

- As stated earlier, ν_0^N and $\bar{\nu}_0$ are the respective distributions of X_0^N and \bar{X}_0 . In the following, the quantity $\mathbb{E}_{\mathcal{E}}[g(X_t^N)]$ always denotes $\mathbb{E}_{\mathcal{E}, \nu_0^N}[g(X_t^N)]$ (with similar notation for the process \bar{X}).
- We note $(P_{\mathcal{E}, t}^N)_t$ and $A_{\mathcal{E}}^N$ respectively the semigroup and the (extended) generator of the process $(X_t^N)_t$ defined at (1)–(2), conditionally on the environment \mathcal{E} . Similarly, we use the notation $(\bar{P}_{\mathcal{E}, t}^N)_t$ and $\bar{A}_{\mathcal{E}}^N$ for the limit process \bar{X}_t^N defined at (3)–(4), defined w.r.t. the environment variable $W := W^{[N]}$. Note that it would not be possible to consider the non-conditional semigroups and generators since these processes are only Markovian conditionally on the environment. For the precise notion of semigroups and (extended) generators we use in this paper, we refer to Appendix A of [Erny, Löcherbach and Loukianova \(2022\)](#): for g in $C_b(\mathbb{R})$, $x \in \mathbb{R}$ and $t \geq 0$,

$$P_{\mathcal{E}, t} g(x) := \mathbb{E}_{\mathcal{E}} [g(X_t^{(x)})] = \mathbb{E}_{\mathcal{E}, \delta_x} [g(X_t)],$$

and, $A_{\mathcal{E}} g$ (if it exists) is characterized as the unique function such that for all $x \in \mathbb{R}, t \geq 0$,

$$P_{\mathcal{E}, t} g(x) - g(x) = \int_0^t P_{\mathcal{E}, s} A_{\mathcal{E}} g(x) ds.$$

- The respective laws of X_0^N ($N \in \mathbb{N}^*$) and \bar{X}_0 are denoted by ν_0^N ($N \in \mathbb{N}^*$) and $\bar{\nu}_0$.
- We note C any positive constant. The value of C can change from line to line in an equation. And, if C depends on some parameter θ , we note C_{θ} instead (the dependency w.r.t. the parameters of the model such as b, f, μ, σ will not be written explicitly).

1.2. Assumptions and main result

Let us state the hypotheses we need to prove our main result. The first assumption concerns the environment variables U_j ($j \geq 1$). It allows to use the result of [Komlós, Major and Tusnády \(1976\)](#) to couple the environment variables U_j ($j \geq 1$) with a sequence of Gaussian variables $W^{[N]}$ used to define the random environment of the limit equation (see [Section 3.1](#) for the formal coupling).

Assumption 1. *The variables U_j ($j \geq 1$) are i.i.d. μ -distributed. The law μ is centered and admits exponential moments: there exists some $\alpha > 0$ such that*

$$\int_{\mathbb{R}} e^{\alpha|x|} d\mu(x) < \infty.$$

We note σ^2 the variance of μ .

The next assumption gives the conditions we need on the coefficients of the SDEs [\(1\)](#) and [\(3\)](#). Let us note that the function f is obviously assumed to be non-negative in all the paper, so we will not write this condition in our assumptions.

Assumption 2. *The functions b , f and \sqrt{f} are C^4 , and, for each $k \in \{1, 2, 3, 4\}$, the functions $b^{(k)}$, $f^{(k)}$ and $\sqrt{f}^{(k)}$ are bounded.*

Note that [Assumption 2](#) implies that the functions b , f and \sqrt{f} are Lipschitz continuous. In particular, under this condition, the equations [\(1\)](#) and [\(3\)](#) are (strongly) well-posed (it is a consequence of [Theorem IV.9.1 of Ikeda and Watanabe \(1989\)](#) and of classical priori estimates that can be established on the processes, conditionally on the environment: see for instance [Proposition C.1 of Erny, Löcherbach and Loukianova \(2022\)](#) or [Proposition 2 of Fournier and Löcherbach \(2016\)](#)). Once the strong well-posedness is established given a fixed environment, it is possible to define an explicit probability space containing the random environment as well as what is needed in the SDEs (i.e. Poisson measures, Brownian motion and initial conditions) such that they are independent. Moreover [Assumption 2](#) is a necessary condition of [Theorem 1.4.1 of Kunita \(1986\)](#) that implies that the stochastic flow of the limit process $(\bar{X}_t)_t$ is C^3 w.r.t. its initial condition (see also [Theorem 4.6.5 of Kunita \(1990\)](#)).

Remark 1.1. *In our proof, the only condition from [Assumption 2](#) that we use directly is that the functions b and f are measurable and sublinear. The stronger conditions allow to guarantee the well-posedness of our equations and to apply [Theorem 1.4.1 of Kunita \(1986\)](#).*

Now we give the hypotheses we need on the laws of the initial conditions of the processes $(X_t^N)_t$ ($N \in \mathbb{N}^*$) and $(\bar{X}_t)_t$.

Assumption 3.

1. *The probability measures ν_0^N converge to $\bar{\nu}_0$ for the Kantorovich-Rubinstein metric d_{KR} . Equivalently, it means that ν_0^N converges weakly to $\bar{\nu}_0$ and that $\int |x| d\nu_0^N(x)$ converges to $\int |x| d\bar{\nu}_0(x)$ as N goes to infinity (see [Theorem 6.9 of Villani \(2008\)](#)).*
2. *The probability measures ν_0^N admit fourth order moments uniformly bounded in $N \in \mathbb{N}^*$, and $\bar{\nu}_0$ admits a first order moment:*

$$\sup_{N \in \mathbb{N}^*} \int_{\mathbb{R}} x^4 d\nu_0^N(x) < \infty \text{ and } \int_{\mathbb{R}} |x| d\bar{\nu}_0(x) < \infty.$$

Since our main result is the convergence in distribution of $(X_t^N)_t$ to $(\bar{X}_t)_t$, we obviously need an assumption on the convergence of their initial conditions. Item 1 of Assumption 3 is a convergence a bit stronger than the necessary convergence in distribution, but it seems to be the most appropriate in our proof. Besides, we need to assume that the initial conditions of the processes admit finite first order moments in order to manipulate Kantorovich-Rubinstein metric. And, for technical reasons, we also need some a priori estimates on the process $(X_t^N)_t$ that are uniform in $N \in \mathbb{N}^*$.

Theorem 1.2. *Grant Assumptions 1, 2 and 3. The process $(X_t^N)_t$ converges to $(\bar{X}_t)_t$ in distribution on $D(\mathbb{R}_+, \mathbb{R})$. In addition, we have the following convergence speed for the finite-dimensional distributions: for any $k \in \mathbb{N}^*$, $T > 0$, $t_1 \leq \dots \leq t_k \leq T$, and $g_1, \dots, g_k \in C_b^3(\mathbb{R})$, there exists some $N_T \in \mathbb{N}^*$ such that, for all $N \geq N_T$,*

$$|\mathbb{E}[g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E}[g_1(\bar{X}_{t_1}) \dots g_k(\bar{X}_{t_k})]| \leq C_{T,k,g_1,\dots,g_k} \left(d_{KR}(\nu_0^N, \bar{\nu}_0) + \ln N / \sqrt{N} \right),$$

for some positive constant C_{T,k,g_1,\dots,g_k} .

The proof of the convergence (and the bound for the convergence speed) of the finite-dimensional distributions of Theorem 1.2 is given in Section 2.2. The convergence in distribution of the processes is deduced from it in Section 2.3.

2. Proof of Theorem 1.2

In the Section 2.1, we state a lemma about a priori estimates for the process $(X_t^N)_t$ that are uniform in N . The fact that we work in a random environment and in normalization $N^{-1/2}$ makes the proof more tricky than in more usual frameworks. Then, Section 2.2 is dedicated to prove the finite-dimensional convergence in distribution of $(X_t^N)_t$ to $(\bar{X}_t)_t$. As explained in the introduction, we begin by controlling the difference of the conditional generators of X^N and \bar{X}^N to deduce a bound for the difference of the conditional semigroups using (6). This allows to control the difference of the conditional one-dimensional distributions of X^N and \bar{X} , then the control between the conditional finite-dimensional distributions is deduced using usual Markov processes technics. Afterwards, the (non-conditional) finite-dimensional convergence in distribution is obtained by integrating over the environment in the control previously obtained. Finally in Section 2.3, we deduce the convergence in distribution of the processes from the convergence of their finite-dimensional distributions using Theorem 13.5 of Billingsley (1999). Section 2.4 is a mere summary of the results of the two previous sections.

2.1. A priori estimates

Since we work in a random environment, we need, in a first time, to work conditionally on the environment to obtain some a priori estimates. However, in our framework, we need to take precautions to prevent the problematic term $N^{-1/2} \sum_{j=1}^N |U_j|$ from appearing, and to control instead $|N^{-1/2} \sum_{j=1}^N U_j|$. In addition, we need to apply Grönwall's lemma conditionally on the environment, which we make arise the previous term within an exponential. The next lemma (whose proof is postponed to Section 3.2) provides a suitable control for this exponential.

Lemma 2.1. *Let U_i ($i \geq 1$) be i.i.d. centered random variables having some finite exponential moments. Let S_N be defined as*

$$S_N := \frac{1}{\sqrt{N}} \sum_{i=1}^N U_i.$$

Then, for all $\gamma > 0$, there exists some $N_\gamma \in \mathbb{N}^$ such that,*

$$\sup_{N \geq N_\gamma} \mathbb{E} \left[e^{\gamma |S_N|} \right] < \infty.$$

As a consequence, for any $p \in \mathbb{N}^$,*

$$\sup_{N \geq 1} \mathbb{E} [|S_N|^p] < \infty.$$

Remark 2.2. *The second part of Lemma 2.1 is easy to prove with straightforward computation. In the proof of Lemma 2.1, we prefer to use the first part of the lemma to prove the second one, since it is a straightforward consequence. Note also that it is necessary to consider some N_γ in the first statement since $S_1 := U_1$ is not assumed to admit all exponential moments.*

The next lemma states the a priori estimates we need to prove our main result. Its proof is postponed to Section 3.3.

Lemma 2.3. *Grant Assumption 1. Assume that b and f are measurable and sublinear. Then, any solution $(X_t^N)_t$ of (1)–(2) satisfies a priori the following: for every $p \in \mathbb{N}^*$, if X_0^N admits a finite $2p$ -order moment, then for any $1 \leq \kappa < 2p$, for all $T > 0$, there exists some $N_{T,p} \in \mathbb{N}^*$ such that*

(i)

$$\sup_{N \geq N_{T,p}} \sup_{t \leq T} \mathbb{E} \left[(X_t^N)^{2p} \right] \leq C_{T,p} (1 + \mathbb{E} [(X_0^N)^{2p}]),$$

(ii) and

$$\sup_{N \geq N_{T,p}} \mathbb{E} \left[\sup_{t \leq T} |X_t^N|^\kappa \right] < \infty.$$

2.2. Convergence of the finite-dimensional distributions

The goal of this section is to prove the convergence speed stated in Theorem 1.2, namely: for any $k \in \mathbb{N}^*$, $T > 0$, $t_1 \leq \dots \leq t_k \leq T$, and $g_1, \dots, g_k \in C_b^3(\mathbb{R})$,

$$\left| \mathbb{E} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E} [g_1(\bar{X}_{t_1}) \dots g_k(\bar{X}_{t_k})] \right| \leq C_{T,k,g_1,\dots,g_k} \left(d_{KR}(\nu_0^N, \bar{\nu}_0) + \ln N / \sqrt{N} \right), \quad (7)$$

for some positive constant C_{T,k,g_1,\dots,g_k} .

Before proving (7), let us state two useful Lemmas. The first one allows to prove that the limit conditional semigroup is C^3 and gives a control of its derivatives. Its proof is postponed to Section 3.4.

Lemma 2.4. *Grant Assumption 2. Then, for all $t \geq 0$, the function $x \in \mathbb{R} \mapsto \bar{P}_{\mathcal{E},t}g(x)$ belongs to $C_b^3(\mathbb{R})$ and*

$$\|\bar{P}_{\mathcal{E},t}g\|_{3,\infty} \leq C_t \|g\|_{3,\infty} e^{C_t |W|},$$

where W is the gaussian variable in (3)–(4).

Remark 2.5. In the statement of Lemma 2.4, we use the version of the limit process $(\bar{X}_t)_t$ with a generic Gaussian variable W for the sake of notation. However, in the proof below, we use this lemma with the version of the limit process $(\bar{X}_t^N)_t$ defined w.r.t. the Gaussian variables $W^{[N]}$ ($N \in \mathbb{N}^*$).

The next lemma recalls the formula (6). It permits to deduce the convergence of the (conditional) semigroups from the convergence of their (conditional) generators. Since in the formula we condition by the environment, the proof of this lemma is exactly the same as the one of Proposition B.2 of Erny, Löcherbach and Loukianova (2022). This proof is therefore omitted

Lemma 2.6. *If Assumption 2 holds true, then for all $t \geq 0, x \in \mathbb{R}$ and $g \in C_b^3(\mathbb{R})$,*

$$P_{\mathcal{E},t}^N g(x) - \bar{P}_{\mathcal{E},t}^N g(x) = \int_0^t P_{\mathcal{E},t-s}^N (A_{\mathcal{E}}^N - \bar{A}_{\mathcal{E}}^N) \bar{P}_{\mathcal{E},s}^N g(x) ds.$$

Using the two previous lemmas, we obtain the convergence speed of the finite-dimensional distribution of Theorem 1.2.

Proof of (7). Step 1. In this first step, we study the convergence of the one-dimensional distributions of the process $(X_t^N)_t$ to the limit process $(\bar{X}_t^N)_t$, conditionally on the environment \mathcal{E} . Recall that conditionally on \mathcal{E} , these processes are Markov processes, hence their conditional one-dimensional distributions are characterized by the (conditional) semigroups of the processes. We note

$$P_{\mathcal{E},t}^N g(x) := \mathbb{E}_{\mathcal{E},\delta_x} [g(X_t^N)] \quad \text{and} \quad \bar{P}_{\mathcal{E},t}^N g(x) := \mathbb{E}_{\mathcal{E},\delta_x} [g(\bar{X}_t^N)]$$

these semigroups.

We have for any $g \in C_b^1(\mathbb{R}), x \in \mathbb{R}$,

$$A_{\mathcal{E}}^N g(x) = b(x)g'(x) + f(x) \sum_{j=1}^N \left[g\left(x + \frac{U_j}{\sqrt{N}}\right) - g(x) \right],$$

and for $g \in C_b^2(\mathbb{R}), x \in \mathbb{R}$,

$$\bar{A}_{\mathcal{E}}^N g(x) = b(x)g'(x) + W^{[N]} f(x)g'(x) + \frac{1}{2}\sigma^2 f(x)g''(x).$$

As a consequence, for all $g \in C_b^3(\mathbb{R}), x \in \mathbb{R}$,

$$\begin{aligned} |A_{\mathcal{E}}^N g(x) - \bar{A}_{\mathcal{E}}^N g(x)| &\leq f(x) \sum_{j=1}^N \left| g\left(x + \frac{U_j}{\sqrt{N}}\right) - g(x) - \frac{U_j}{\sqrt{N}}g'(x) - \frac{U_j^2}{2N}g''(x) \right| \\ &\quad + f(x)g'(x) \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j - W^{[N]} \right| + \frac{1}{2}f(x)g''(x) \left| \frac{1}{N} \sum_{j=1}^N U_j^2 - \sigma^2 \right| \\ &\leq f(x) \|g\|_{3,\infty} \left(\frac{1}{6N\sqrt{N}} \sum_{j=1}^N |U_j|^3 + K \frac{\ln N}{\sqrt{N}} + \frac{1}{2} \left| \frac{1}{N} \sum_{j=1}^N U_j^2 - \sigma^2 \right| \right). \end{aligned}$$

By Lemma 2.6, for any $g \in C_b^3(\mathbb{R})$, $x \in \mathbb{R}$, $t \geq 0$,

$$P_{\mathcal{E},t}^N g(x) - \bar{P}_{\mathcal{E},t}^N g(x) = \int_0^t P_{\mathcal{E},t-s}^N (A_{\mathcal{E}}^N - \bar{A}_{\mathcal{E}}^N) \bar{P}_{\mathcal{E},s}^N g(x) ds,$$

whence

$$\begin{aligned} |P_{\mathcal{E},t}^N g(x) - \bar{P}_{\mathcal{E},t}^N g(x)| &\leq \int_0^t \mathbb{E}_{\mathcal{E},x} [|A_{\mathcal{E}}^N \bar{P}_{\mathcal{E},s}^N g(X_{t-s}^N) - \bar{A}_{\mathcal{E}}^N \bar{P}_{\mathcal{E},s}^N g(X_{t-s}^N)|] ds \\ &\leq \int_0^t \|P_{\mathcal{E},s}^N g\|_{3,\infty} \mathbb{E}_{\mathcal{E},x} [f(X_{t-s}^N)] \left(\frac{1}{6N\sqrt{N}} \sum_{j=1}^N |U_j|^3 + K \frac{\ln N}{\sqrt{N}} + \frac{1}{2} \left| \frac{1}{N} \sum_{j=1}^N U_j^2 - \sigma^2 \right| \right) ds. \end{aligned}$$

Using Lemma 2.4,

$$\begin{aligned} |P_{\mathcal{E},t}^N g(x) - \bar{P}_{\mathcal{E},t}^N g(x)| &\leq \\ C_t \left(\int_0^t \mathbb{E}_{\mathcal{E},\delta_x} [f(X_s^N)] ds \right) e^{C_t |W^{[N]}|} \|g\|_{3,\infty} &\left(\frac{1}{N\sqrt{N}} \sum_{j=1}^N |U_j|^3 + K \frac{\ln N}{\sqrt{N}} + \frac{1}{2} \left| \frac{1}{N} \sum_{j=1}^N U_j^2 - \sigma^2 \right| \right) \\ =: \|g\|_{3,\infty} \varepsilon_N(t, x). & \quad (8) \end{aligned}$$

Step 2. Let us now show that the control between the semigroups above can be extended to the any finite-dimensional distributions of the processes, conditionally on the environment.

Remark 2.7. *The result of this second step can be seen as a quenched control between $(X_t^N)_t$ and $(\bar{X}_t)_t$. However this not a quenched convergence since the RHS of (8) is not guaranteed to vanish as N goes to infinity because the a.s. limit superior (resp. inferior) of $W^{[N]}$ is $+\infty$ (resp. $-\infty$).*

To begin with

$$\begin{aligned} &|\mathbb{E}_{\mathcal{E}} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E}_{\mathcal{E}} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)]| \\ &= \left| \mathbb{E}_{\mathcal{E},\nu_0^N} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E}_{\mathcal{E},\bar{\nu}_0} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)] \right| \\ &\leq \left| \mathbb{E}_{\mathcal{E},\nu_0^N} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E}_{\mathcal{E},\nu_0^N} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)] \right| \\ &\quad + \left| \mathbb{E}_{\mathcal{E},\nu_0^N} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)] - \mathbb{E}_{\mathcal{E},\bar{\nu}_0} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)] \right| =: A_1 + A_2. \end{aligned}$$

Recall that any Markov process $(X_t)_t$ with semigroup $(P_t)_t$ satisfies for all $t_1 \leq t_2 \leq \dots \leq t_k$, and for all continuous and bounded functions g_1, \dots, g_k ,

$$\begin{aligned} &\mathbb{E}_{\nu} [g_1(X_{t_1}) \dots g_{k-1}(X_{t_{k-1}}) g_k(X_{t_k})] \\ &= \int \nu(dx) \int P_{t_1}(x, dx_1) g(x_1) \dots \int P_{t_k - t_{k-1}}(x_{k-1}, dx_k) g_k(x_k) \\ &= \int \nu(dx) \int P_{t_1}(x, dx_1) g(x_1) \dots \int P_{t_{k-1} - t_{k-2}}(x_{k-2}, dx_{k-1}) (g_{k-1} \cdot P_{t_k - t_{k-1}} g_k)(x_{k-1}) \\ &= \mathbb{E}_{\nu} [g_1(X_{t_1}) \dots g_{k-1}(X_{t_{k-1}}) P_{t_k - t_{k-1}} g_k(X_{t_{k-1}})], \end{aligned}$$

Let us show by induction on k that

$$A_1 \leq \|g_1\|_{3,\infty} \dots \|g_k\|_{3,\infty} C_{T,k} e^{C_{T,k}|W_{[N]}|} \sum_{l=0}^{k-1} \mathbb{E}_{\mathcal{E}} [\varepsilon_N(T, X_{t_l}^N)], \quad (9)$$

recalling that $T \geq t_k > t_{k-1} > \dots > t_1$.

The bound of (9) above for $k = 1$ is valid by definition of $\varepsilon_N(T, x)$ (see (8)). To show the inductive step, let us write

$$\begin{aligned} A_1 &= \left| \mathbb{E}_{\mathcal{E}, \nu_0^N} \left[g_1(X_{t_1}^N) \dots (g_{k-1} \cdot P_{t_k - t_{k-1}}^N g_k)(X_{t_{k-1}}^N) \right] - \mathbb{E}_{\mathcal{E}, \nu_0^N} \left[g_1(\bar{X}_{t_1}^N) \dots (g_{k-1} \bar{P}_{t_k - t_{k-1}}^N)(\bar{X}_{t_{k-1}}^N) \right] \right| \\ &\leq \left| \mathbb{E}_{\mathcal{E}, \nu_0^N} \left[g_1(X_{t_1}^N) \dots (g_{k-1} \cdot P_{t_k - t_{k-1}}^N g_k)(X_{t_{k-1}}^N) \right] - \mathbb{E}_{\mathcal{E}, \nu_0^N} \left[g_1(X_{t_1}^N) \dots (g_{k-1} \cdot \bar{P}_{t_k - t_{k-1}}^N g_k)(X_{t_{k-1}}^N) \right] \right| \\ &\quad + \left| \mathbb{E}_{\mathcal{E}, \nu_0^N} \left[g_1(X_{t_1}^N) \dots (g_{k-1} \cdot \bar{P}_{t_k - t_{k-1}}^N g_k)(X_{t_{k-1}}^N) \right] - \mathbb{E}_{\mathcal{E}, \nu_0^N} \left[g_1(\bar{X}_{t_1}^N) \dots (g_{k-1} \bar{P}_{t_k - t_{k-1}}^N)(\bar{X}_{t_{k-1}}^N) \right] \right| \\ &=: A_{11} + A_{12}, \end{aligned}$$

Then, thanks to (8),

$$\begin{aligned} A_{11} &\leq \|g_1\|_{\infty} \dots \|g_{k-1}\|_{\infty} \mathbb{E}_{\mathcal{E}, \nu_0^N} \left[\left| P_{t_k - t_{k-1}}^N g_k(X_{t_{k-1}}^N) - \bar{P}_{t_k - t_{k-1}}^N g_k(X_{t_{k-1}}^N) \right| \right] \\ &\leq \|g_1\|_{3,\infty} \dots \|g_k\|_{3,\infty} \mathbb{E}_{\mathcal{E}, \nu_0^N} [\varepsilon_N(T, X_{t_{k-1}}^N)]. \end{aligned}$$

On the other hand, using the inductive hypothesis we have

$$A_{12} \leq \|g_1\|_{3,\infty} \dots \|g_{k-1}\|_{3,\infty} \left\| \bar{P}_{t_k - t_{k-1}}^N g_k \right\|_{3,\infty} C_{T,k-1} e^{C_{T,k-1}|W_{[N]}|} \sum_{l=0}^{k-2} \mathbb{E}_{\mathcal{E}} [\varepsilon_N(T, X_{t_l}^N)],$$

whence, by Lemma 2.4,

$$A_{12} \leq \|g_1\|_{3,\infty} \dots \|g_k\|_{3,\infty} C_{T,k} e^{C_{T,k}|W_{[N]}|} \sum_{l=0}^{k-2} \mathbb{E}_{\mathcal{E}} [\varepsilon_N(T, X_{t_l}^N)].$$

Then recalling that $A_1 = A_{11} + A_{12}$, we have proved (9).

Now let us show that

$$A_2 \leq \|g'_1\|_{\infty} \dots \|g'_k\|_{\infty} C_{T,k} e^{C_{T,k}|W_{[N]}|} d_{KR}(\nu_0^N, \bar{\nu}_0), \quad (10)$$

for some constant $C_k > 0$.

Let us define

$$h_{g_1, \dots, g_k}^k(x) := \int \bar{P}_{\mathcal{E}, t_1}^N(x, dx_1) g_1(x_1) \dots \int \bar{P}_{\mathcal{E}, t_k - t_{k-1}}^N(x_{k-1}, dx_k) g_k(x_k).$$

By definition of Kantorovich-Rubinstein metric d_{KR} , to show that (10) holds true, it is sufficient to prove that

$$\left\| (h_{g_1, \dots, g_k}^k)' \right\|_{\infty} \leq \|g'_1\|_{\infty} \dots \|g'_k\|_{\infty} C_{T,k} e^{C_{T,k}|W_{[N]}|}. \quad (11)$$

Now let us prove (11) by induction on k . Since

$$h_{g_1}^1(x) := \bar{P}_{\mathcal{E}, t_1}^N g_1(x),$$

the case $k = 1$ is a consequence of Lemma 2.4. To show the inductive step, let us write

$$\begin{aligned} h_{g_1, \dots, g_k}^k(x) &= \int \bar{P}_{\mathcal{E}, t_1}^N(x, dx_1) g_1(x_1) \dots \int \bar{P}_{\mathcal{E}, t_k - t_{k-1}}^N(x_{k-1}, dx_k) g_k(x_k) \\ &= \int \bar{P}_{\mathcal{E}, t_1}^N(x, dx_1) g_1(x_1) \dots \int \bar{P}_{\mathcal{E}, t_{k-1} - t_{k-2}}^N(x_{k-2}, dx_{k-1}) \left(g_{k-1} \cdot \bar{P}_{\mathcal{E}, t_k - t_{k-1}}^N g_k \right) (x_{k-1}) \\ &= h_{g_1, \dots, g_{k-2}, g_{k-1} \cdot \bar{P}_{\mathcal{E}, t_k - t_{k-1}}^N g_k}^{k-1}(x). \end{aligned}$$

By induction hypothesis, and thanks to Lemma 2.4,

$$\begin{aligned} \left\| (h_{g_1, \dots, g_k}^k)' \right\|_{\infty} &\leq \|g_1'\|_{\infty} \dots \|g_{k-1}'\|_{\infty} \|(\bar{P}_{\mathcal{E}, t_k - t_{k-1}}^N g_k)'\|_{\infty} C_{T, k-1} e^{C_{T, k-1} |W^{[N]}|} \\ &\leq \|g_1'\|_{\infty} \dots \|g_k'\|_{\infty} C_{T, k} e^{C_{T, k} |W^{[N]}|}. \end{aligned}$$

Consequently (11) holds true.

So we have proved (9) and (10). This implies that

$$\begin{aligned} &|\mathbb{E}_{\mathcal{E}} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E}_{\mathcal{E}} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)]| \\ &\leq \|g_1\|_{3, \infty} \dots \|g_k\|_{3, \infty} C_{T, k} e^{C_{T, k} |W^{[N]}|} \left(d_{KR}(\nu_0^N, \bar{\nu}_0) + \sum_{l=0}^{k-1} \mathbb{E}_{\mathcal{E}} [\varepsilon_N(T, X_{t_l}^N)] \right). \end{aligned}$$

Step 3. Now we prove (7) by integrating over the environment in the inequality above.

$$\begin{aligned} &|\mathbb{E} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)]| \\ &= |\mathbb{E} [\mathbb{E}_{\mathcal{E}} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E}_{\mathcal{E}} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)]]| \\ &\leq \mathbb{E} [|\mathbb{E}_{\mathcal{E}} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E}_{\mathcal{E}} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)]|] \\ &\leq \|g_1\|_{3, \infty} \dots \|g_k\|_{3, \infty} C_{T, k} \mathbb{E} \left[e^{2C_{T, k} |W^{[N]}|} \right]^{1/2} \left(d_{KR}(\nu_0^N, \bar{\nu}_0) + \mathbb{E} \left[\left(\sum_{l=0}^{k-1} \mathbb{E}_{\mathcal{E}} [\varepsilon_N(T, X_{t_l}^N)] \right)^2 \right]^{1/2} \right) \\ &\leq C_{T, k} \|g_1\|_{3, \infty} \dots \|g_k\|_{3, \infty} \left(d_{KR}(\nu_0^N, \bar{\nu}_0) + \sqrt{k} \sum_{l=0}^{k-1} \sqrt{\mathbb{E} [\varepsilon_N(T, X_{t_l}^N)^2]} \right), \end{aligned}$$

where we have used Cauchy-Schwarz' inequality in the line before last above, and in the last line, Jensen's inequality as well as the facts that for all $x, y \geq 0$, $(x + y)^2 \leq x^2/2 + y^2/2$ and $\sqrt{x + y} \leq \sqrt{x} + \sqrt{y}$.

Recalling that the expression of $\varepsilon_N(T, x)$ is given explicitly in (8), applying Cauchy-Schwarz'

inequality twice and recalling that f is sublinear, we have

$$\begin{aligned} \mathbb{E} [\varepsilon_N(T, X_{t_t}^N)^2] &\leq C_T \left(1 + \sup_{s \leq T} \mathbb{E} [(X_s^N)^4]^{1/2} \right) \mathbb{E} [e^{8C_T |W_{[N]}|}]^{1/4} \\ &\quad \left(\mathbb{E} \left[\left(\frac{1}{N\sqrt{N}} \sum_{j=1}^N |U_j|^3 \right)^8 \right]^{1/4} + \mathbb{E} \left[\left(K \frac{\ln N}{\sqrt{N}} \right)^8 \right]^{1/4} + \mathbb{E} \left[\left| \frac{1}{N} \sum_{j=1}^N U_j^2 - \sigma^2 \right|^8 \right]^{1/4} \right). \end{aligned}$$

Then, by Jensen's inequality,

$$\mathbb{E} \left[\left(\frac{1}{N\sqrt{N}} \sum_{j=1}^N |U_j|^3 \right)^8 \right] = \mathbb{E} \left[\frac{1}{N^4} \left(\frac{1}{N} \sum_{j=1}^N |U_j|^3 \right)^8 \right] \leq \mathbb{E} \left[\frac{1}{N^4} \cdot \frac{1}{N} \sum_{j=1}^N |U_j|^{24} \right] = \frac{1}{N^4} \mathbb{E} [|U_1|^{24}].$$

Now, let $V_j = U_j^2 - \sigma^2$ ($j \geq 1$). Since the variables V_j ($j \geq 1$) are i.i.d. and centered,

$$\mathbb{E} \left[\left| \frac{1}{N} \sum_{j=1}^N U_j^2 - \sigma^2 \right|^8 \right] = \mathbb{E} \left[\left(\frac{1}{N} \sum_{j=1}^N V_j \right)^8 \right] = \frac{1}{N^4} \mathbb{E} \left[\left(\frac{1}{\sqrt{N}} \sum_{j=1}^N V_j \right)^8 \right] \leq \frac{1}{N^4} C,$$

by the second statement of Lemma 2.1 applied to the variables V_j ($j \geq 1$). Indeed, even if these variables are not assumed to admit some exponential moments, the second statement of Lemma 2.1 still hold true (see Remark 2.2).

Then, recalling that K and $W^{[N]} \sim \mathcal{N}(0, \sigma^2)$ admit exponential moments and a fortiori fourth-order moment, and using Lemma 2.3,

$$\mathbb{E} [\varepsilon_N(T, X_{t_t}^N)^2] \leq C_T \left(\frac{1}{N} + \left(\frac{\ln N}{\sqrt{N}} \right)^2 + \frac{1}{N} \right) \leq C_T \left(\frac{\ln N}{\sqrt{N}} \right)^2.$$

Consequently,

$$|\mathbb{E} [g_1(X_{t_1}^N) \dots g_k(X_{t_k}^N)] - \mathbb{E} [g_1(\bar{X}_{t_1}^N) \dots g_k(\bar{X}_{t_k}^N)]| \leq C_{T,k} \|g_1\|_{3,\infty} \dots \|g_k\|_{3,\infty} \left(d_{KR}(v_0^N, \bar{v}_0) + \frac{\ln N}{\sqrt{N}} \right),$$

which gives (7). \square

2.3. Convergence of the processes in distribution

Since the finite-dimensional convergence in distribution of $(X_t^N)_t$ to $(\bar{X}_t)_t$ has been established in the previous section, it only remains to prove the tightness of the processes to obtain the convergence in distribution in Skorohod space. Instead of using the classical Aldous' criterion we rather rely on Theorem 13.5 of Billingsley (1999) since it seems more appropriate for technical reasons. Firstly, a conditional tightness would not be sufficient, since only the annealed finite-dimensional convergence has been proved and not a quenched one (i.e. conditionally on the environment, the processes do not converge). So we would need to work on stopping times adapted to the (non-conditional) history

of the processes. In particular, these stopping times would depend on the random environment in a non-trivial way, which seems hard to handle for the computation.

The following convergence criterion of Theorem 13.5 (where condition (13.13) is replaced by the stronger (13.14)) appears to be more suitable as no stopping times are involved: for any $T > 0$,

- (i) $\bar{X}_T - \bar{X}_{T-\delta}$ converges to zero in law as δ vanishes,
- (ii) for all $r \leq s \leq t \leq T, N \in \mathbb{N}^*$,

$$\mathbb{E} \left[(X_s^N - X_r^N)^\beta (X_t^N - X_s^N)^\beta \right] \leq C (t - r)^\beta,$$

for some $C > 0$ and some $\beta > 1$ that can be chosen as $\beta := 3/2$ (or any other real number in $]1, 2[$).

Item (i) is trivially satisfied since the trajectories of \bar{X} are continuous.

The rest of this section is dedicated to prove Item (ii). For this purpose, let us rewrite the dynamics of X^N in the following way

$$\begin{aligned} X_t^N &= X_0^N + \int_0^t b(X_s^N) ds + \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \int_{[0,t] \times \mathbb{R}_+} \mathbf{1}_{\{z \leq f(X_{s-}^N)\}} d\tilde{\pi}^j(s, z) \\ &\quad + \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \int_0^t f(X_s^N) ds, \end{aligned}$$

where $\tilde{\pi}^j(ds, dz) := \pi^j(ds, dz) - ds dz$ is the compensated Poisson measure of π^j . For any $s \leq t \leq T$, we have

$$\begin{aligned} (X_t^N - X_s^N)^\beta &\leq C(t-s)^{\beta-1} \int_s^t b(X_r^N)^\beta dr + C \frac{1}{N^{\beta/2}} \left| \sum_{j=1}^N U_j \int_{]s,t] \times \mathbb{R}_+} \mathbf{1}_{\{z \leq f(X_{r-}^N)\}} d\tilde{\pi}^j(r, z) \right|^\beta \\ &\quad + C \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^\beta (t-s)^{\beta-1} \int_s^t f(X_r^N)^\beta dr \\ &=: B_{s,t}^N + M_{s,t}^N + F_{s,t}^N. \end{aligned}$$

Since f and b are sublinear, we have

$$\begin{aligned} B_{s,t}^N &\leq C(t-s)^\beta \left(1 + \sup_{v \leq T} |X_v^N|^\beta \right), \\ F_{s,t}^N &\leq C(t-s)^\beta \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^\beta \left(1 + \sup_{v \leq T} |X_v^N|^\beta \right). \end{aligned}$$

Hence, for any $r \leq s \leq t \leq T$,

$$\begin{aligned}
(X_t^N - X_s^N)^\beta (X_s^N - X_r^N)^\beta &\leq (B_{s,t}^N + M_{s,t}^N + F_{s,t}^N) (B_{r,s}^N + M_{r,s}^N + F_{r,s}^N) \\
&\leq C(t-r)^{2\beta} \left(1 + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^{2\beta} \right) \left(1 + \sup_{v \leq T} |X_v^N|^{2\beta} \right) \\
&\quad + C(t-r)^\beta \left(1 + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^\beta \right) \left(1 + \sup_{v \leq T} |X_v^N|^\beta \right) (M_{s,t}^N + M_{r,s}^N) + M_{s,t}^N M_{r,s}^N \\
&=: \mathcal{R}_1^{N,r,t} + \mathcal{R}_2^{N,r,s,t} + \mathcal{R}_3^{N,r,s,t}.
\end{aligned}$$

Define $q_1, q_2 > 1$ such that $1/q_1 + 1/q_2 = 1$ and that $2\beta q_1 < 4$ (which is possible since we choose $1 < \beta < 2$). Using Hölder's inequality,

$$\begin{aligned}
\mathbb{E} \left[\mathcal{R}_1^{N,r,t} \right] &\leq C(t-r)^{2\beta} \left(1 + \mathbb{E} \left[\left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^{2\beta q_2} \right]^{1/q_2} \right) \left(1 + \mathbb{E} \left[\sup_{v \leq T} |X_v^N|^{2\beta q_1} \right]^{1/q_1} \right) \\
&\leq C(t-r)^{2\beta},
\end{aligned}$$

where we have used Lemmas 2.1.(ii) and 2.3.(ii) (with $p = 2$ and $\kappa = 2\beta q_1$) to obtain the last line.

Besides, since for all $x, y \geq 0$, $xy \leq x^2/2 + y^2/2$,

$$\begin{aligned}
\mathbb{E} \left[\mathcal{R}_2^{N,r,s,t} \right] &\leq C(t-r)^\beta \left\{ \mathbb{E} \left[\left(1 + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^{2\beta} \right) \left(1 + \sup_{v \leq T} |X_v^N|^{2\beta} \right) \right] + \mathbb{E} \left[(M_{s,t}^N)^2 + (M_{r,s}^N)^2 \right] \right\} \\
&\leq C(t-r)^\beta \left(1 + \mathbb{E} \left[(M_{s,t}^N)^2 \right] + \mathbb{E} \left[(M_{r,s}^N)^2 \right] \right),
\end{aligned}$$

where the last line has been obtained with the exact same reasoning as the one used for $\mathcal{R}_1^{N,r,s,t}$ above. For $\mathcal{R}_3^{N,r,s,t}$, we have the following control.

$$\mathbb{E} \left[\mathcal{R}_3^{N,r,s,t} \right] \leq C \mathbb{E} \left[(M_{s,t}^N)^2 \right] + C \mathbb{E} \left[(M_{r,s}^N)^2 \right].$$

So, in order to prove Item (ii), it is sufficient to prove that

$$\mathbb{E} \left[(M_{s,t}^N)^2 \right] \leq C(t-s)^\beta.$$

Recall that $\mathbb{E}_{\mathcal{E}}$ denotes the conditional expectation given the environment $(U_j)_{j \geq 1}$ and $(W^{[N]})_{N \geq 1}$,

we have by Burkholder-Davis-Gundy's inequality,

$$\begin{aligned}
\mathbb{E}_{\mathcal{E}} [(M_{s,t}^N)^2] &= \frac{1}{N^\beta} \mathbb{E}_{\mathcal{E}} \left[\left| \sum_{j=1}^N U_j \int_{]s,t] \times \mathbb{R}_+} 1_{\{z \leq f(X_{r-}^N)\}} d\tilde{\pi}^j(r, z) \right|^{2\beta} \right] \\
&\leq C \frac{1}{N^\beta} \mathbb{E}_{\mathcal{E}} \left[\left(\sum_{j=1}^N U_j^2 \int_s^t f(X_r^N) dr \right)^\beta \right] \\
&\leq C (t-s)^\beta \left(1 + \mathbb{E}_{\mathcal{E}} \left[\sup_{v \leq T} |X_v^N|^\beta \right] \right) \left(\frac{1}{N} \sum_{j=1}^N U_j^2 \right)^\beta \\
&\leq C (t-s)^\beta \left(1 + \mathbb{E}_{\mathcal{E}} \left[\sup_{v \leq T} |X_v^N|^\beta \right]^2 + \left(\frac{1}{N} \sum_{j=1}^N U_j^2 \right)^{2\beta} \right),
\end{aligned}$$

where we have again used that, for any $x, y \geq 0$, $xy \leq x^2/2 + y^2/2$ to obtain the last line. Then, by Jensen's inequality,

$$\mathbb{E}_{\mathcal{E}} [(M_{s,t}^N)^2] \leq C (t-s)^\beta \left(1 + \mathbb{E}_{\mathcal{E}} \left[\sup_{v \leq T} |X_v^N|^{2\beta} \right] + \frac{1}{N} \sum_{j=1}^N |U_j|^{4\beta} \right)$$

Finally, integrating over the environment and using Lemmas 2.1 and 2.3 allows to prove the result: for all $r \leq s \leq t \leq T$, $N \in \mathbb{N}^*$,

$$\mathbb{E} \left[(X_s^N - X_r^N)^\beta (X_t^N - X_s^N)^\beta \right] \leq C (t-r)^\beta,$$

with for example $\beta = 3/2$.

2.4. Proof of Theorem 1.2

We have just proved in the two previous sections that, for all $T > 0$,

- the finite-dimensional distributions of $(X_t^N)_t$ converge to those of $(\bar{X}_t)_t$ on $[0, T]$,
- $\bar{X}_T - \bar{X}_{T-\delta}$ vanishes in distribution as δ goes to zero,
- for all $r \leq s \leq t \leq T$, $N \in \mathbb{N}^*$,

$$\mathbb{E} \left[(X_s^N - X_r^N)^\beta (X_t^N - X_s^N)^\beta \right] \leq C (t-r)^\beta,$$

for some positive constant C and for a fixed $\beta > 1$ (whose value can be any in $]1, 2[$).

Then, according to Theorem 13.5 of Billingsley (1999), we know that the process $(X_t^N)_t$ converges in distribution to $(\bar{X}_t)_t$ on $D([0, T], \mathbb{R})$. As this holds true for any $T > 0$, Lemma 16.3 of Billingsley (1999) implies that the convergence in distribution holds true on the space $D(\mathbb{R}_+, \mathbb{R})$.

3. Appendix

3.1. CLT coupling

Recall that the variables U_j ($j \geq 1$) are i.i.d. centered with exponential moments: there exists $\alpha > 0$,

$$\mathbb{E} \left[e^{\alpha |U_1|} \right] < \infty.$$

Then, according to Theorem 1 of [Komlós, Major and Tusnády \(1976\)](#), it is possible to construct on the same probability space (possibly enlarged) a standard Brownian motion $(\beta_t)_t$ for which there exist some positive constants Γ, Λ, λ such that, for any $x > 0, N \in \mathbb{N}^*$,

$$\mathbb{P} \left(\max_{k \leq N} \left| \sum_{j=1}^k U_j - \sigma \beta_k \right| > \Gamma \ln N + x \right) \leq \Lambda e^{-\lambda x}, \quad (12)$$

with σ^2 the variance of the variables U_j ($j \geq 1$).

Then using the technics of Section 7.5 of [Ethier and Kurtz \(2005\)](#), we have the following result (whose proof is written at the end of this subsection for self-containedness).

Lemma 3.1. *The random variable K defined as*

$$K := \sup_{N \geq 2} \left| \sum_{j=1}^N U_j - \sigma \beta_N \right| / \ln N$$

admits exponential moments. More precisely, for any $0 < \gamma < \lambda$,

$$\mathbb{E} \left[e^{\gamma K} \right] < \infty.$$

Consequently, by definition of K , we have that, almost surely, for all $N \geq 2$,

$$\left| \sum_{j=1}^N U_j - \sigma \beta_N \right| \leq K \ln N.$$

Then, by defining the (non-independent) identically distributed Gaussian variables $W^{[N]} := \sigma \beta_N / \sqrt{N}$, we obtain exactly that, almost surely, for every $N \geq 2$,

$$\left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j - W^{[N]} \right| \leq K \frac{\ln N}{\sqrt{N}}.$$

Remark 3.2. *Since $W^{[N]} := \sigma \beta_N / \sqrt{N}$ with $(\beta_t)_t$ some standard Brownian motion, it is known that, with probability equals one, the sequence $(W^{[N]})_N$ is not bounded. This property prevents us to prove the quenched convergence with our approach. See [Remark 2.7](#) for the precise problem.*

Proof of Lemma 3.1. Step 1. In this first step, we prove the following control on the tails distribution of K : for all $x > 0$,

$$\mathbb{P}(K > 2(\Gamma + 1) + x) \leq C_{\Lambda, \lambda} e^{-\lambda x}. \quad (13)$$

This step is a mere rewriting of the proof of Corollary 7.5.5 of [Ethier and Kurtz \(2005\)](#). To begin with, in (12), if we replace x by $x + \ln N$, we obtain

$$\mathbb{P}\left(\max_{k \leq N} \left| \sum_{j=1}^k U_j - \sigma \beta_k \right| > (\Gamma + 1) \ln N + x\right) \leq \Lambda N^{-\lambda} e^{-\lambda x}.$$

Then

$$\begin{aligned} \mathbb{P}(K > 2(\Gamma + 1) + x) &:= \mathbb{P}\left(\sup_{N \geq 2} \left| \sum_{j=1}^N U_j - \sigma \beta_N \right| / \ln N > 2(\Gamma + 1) + x\right) \\ &\leq \sum_{k=2}^{+\infty} \mathbb{P}\left(\max_{2^{k-1} \leq N \leq 2^k} \left| \sum_{j=1}^N U_j - \sigma \beta_N \right| / \ln N > 2(\Gamma + 1) + x\right) \\ &\leq \sum_{k=2}^{+\infty} \mathbb{P}\left(\max_{N \leq 2^k} \left| \sum_{j=1}^N U_j - \sigma \beta_N \right| / \ln(2^{k-1}) > 2(\Gamma + 1) + x\right) \\ &= \sum_{k=2}^{+\infty} \mathbb{P}\left(\max_{N \leq 2^k} \left| \sum_{j=1}^N U_j - \sigma \beta_N \right| > 2(\Gamma + 1) \ln(2^{k-1}) + x \ln(2^{k-1})\right) \\ &\leq \sum_{k=2}^{+\infty} \mathbb{P}\left(\max_{N \leq 2^k} \left| \sum_{j=1}^N U_j - \sigma \beta_N \right| > (\Gamma + 1) \ln(2^k) + x\right) \\ &\leq \Lambda e^{-\lambda x} \sum_{k=2}^{+\infty} 2^{-k\lambda}, \end{aligned}$$

which gives (13).

Step 2. Now, we conclude the proof using (13). For $0 < \gamma < \lambda$,

$$\begin{aligned} \mathbb{E}[e^{\gamma K}] &= \int_0^{+\infty} \mathbb{P}(e^{\gamma K} > t) dt = \int_0^{+\infty} \mathbb{P}\left(K > \frac{1}{\gamma} \ln t\right) dt \\ &= \int_0^{\exp[\gamma(2(\Gamma+1)+1)]} \mathbb{P}\left(K > \frac{1}{\gamma} \ln t\right) dt + \int_{\exp[\gamma(2(\Gamma+1)+1)]}^{+\infty} \mathbb{P}\left(K > \frac{1}{\gamma} \ln t\right) dt \\ &\leq e^{\gamma(2(\Gamma+1)+1)} + \int_{\exp[\gamma(2(\Gamma+1)+1)]}^{+\infty} \mathbb{P}\left(K > 2(\Gamma + 1) + \left(\frac{1}{\gamma} \ln t - 2(\Gamma + 1)\right)\right) dt. \end{aligned}$$

Then, we use (13) to bound the second term of the sum above

$$\int_{\exp[\gamma(2(\Gamma+1)+1)]}^{+\infty} \mathbb{P}\left(K > 2(\Gamma+1) + \left(\frac{1}{\gamma} \ln t - 2(\Gamma+1)\right)\right) dt \\ \leq C_{\Lambda, \lambda} e^{2\lambda(\Gamma+1)} \int_{\exp[\gamma(2(\Gamma+1)+1)]}^{+\infty} t^{-\lambda/\gamma} dt,$$

which is finite since $\lambda/\gamma > 1$. \square

3.2. Proof of Lemma 2.1

The proof relies on the coupling between S_N and $W^{[N]}$ defined in Section 3.1.

$$\mathbb{E}\left[e^{\gamma|S_N|}\right] \leq \mathbb{E}\left[e^{\gamma|S_N - W^{[N]}|} e^{\gamma|W^{[N]}|}\right] \leq \frac{1}{2} \mathbb{E}\left[e^{2\gamma K \ln N / \sqrt{N}}\right] + \frac{1}{2} \mathbb{E}\left[e^{2\gamma|W^{[N]}|}\right],$$

where we have used that for all $x, y \geq 0$, $xy \leq x^2/2 + y^2/2$. Since $W^{[N]}$ is a Gaussian variable with parameters independent of N , the second term of the sum above is finite and independent of N . And, choosing $N_\gamma \in \mathbb{N}^*$ large enough such that $2\gamma(\ln N_\gamma)/\sqrt{N_\gamma} < \lambda$ (where λ is defined in (12)), we know that the second term is finite and bounded uniformly in $N \geq N_\gamma$ thanks to Lemma 3.1. So the first statement of the lemma is proved.

And, since for all $x \geq 0, \beta > 0$ and $p \in \mathbb{N}^*$,

$$x^p = \frac{p!}{\beta^p} \cdot \frac{\beta^p x^p}{p!} \leq \frac{p!}{\beta^p} e^{\beta x},$$

the second statement of the lemma is a direct consequence of the first one.

3.3. Proof of Lemma 2.3

Let us prove Item (i). By Ito's formula,

$$(X_t^N)^{2p} = (X_0^N)^{2p} + 2p \int_0^t (X_s^N)^{2p-1} b(X_s^N) ds \\ + \sum_{j=1}^N \int_{[0, t] \times \mathbb{R}_+} \left[\left(X_{s-}^N + \frac{U_j}{\sqrt{N}} \right)^{2p} - (X_{s-}^N)^{2p} \right] 1_{\{z \leq f(X_{s-}^N)\}} d\pi_j(s, z) \\ \mathbb{E}_{\mathcal{E}} \left[(X_t^N)^{2p} \right] = \mathbb{E} \left[(X_0^N)^{2p} \right] + 2p \int_0^t \mathbb{E}_{\mathcal{E}} \left[(X_s^N)^{2p-1} b(X_s^N) \right] ds \\ + \sum_{j=1}^N \int_0^t \mathbb{E}_{\mathcal{E}} \left[f(X_s^N) \sum_{k=0}^{2p-1} \binom{2p}{k} (X_s^N)^k U_j^{2p-k} N^{-(2p-k)/2} \right] ds.$$

Since

$$\begin{aligned}
\sum_{j=1}^N \sum_{k=0}^{2p-1} \binom{2p}{k} (X_s^N)^k \frac{U_j^{2p-k}}{N^{(2p-k)/2}} &= \sum_{j=1}^N \sum_{k=0}^{2p-2} \binom{2p}{k} (X_s^N)^k \frac{U_j^{2p-k}}{N^{(2p-k)/2}} + \sum_{j=1}^N 2p (X_s^N)^{2p-1} \frac{U_j}{\sqrt{N}} \\
&\leq C_p \frac{1}{N} \sum_{j=1}^N \sum_{k=0}^{2p-2} |U_j|^{2p-k} |X_s^N|^k + C_p \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j (X_s^N)^{2p-1} \\
&\leq C_p \left(1 + \frac{1}{N} \sum_{j=1}^N |U_j|^{2p} \right) (1 + |X_s^N|^{2p-2}) + C_p \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right| \cdot |X_s^N|^{2p-1},
\end{aligned}$$

we have (recalling that f and b are sublinear),

$$\begin{aligned}
\mathbb{E}_{\mathcal{E}} \left[(X_t^N)^{2p} \right] &\leq \mathbb{E} [(X_0^N)^{2p}] + C_p t + C_p \int_0^t \mathbb{E}_{\mathcal{E}} [(X_s^N)^{2p}] ds \\
&\quad + \left(1 + \frac{1}{N} \sum_{j=1}^N |U_j|^{2p} \right) \left(\int_0^t (1 + \mathbb{E}_{\mathcal{E}} [(X_s^N)^{2p-1}]) ds \right) \\
&\quad + C_p \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right| \int_0^t (1 + \mathbb{E}_{\mathcal{E}} [(X_s^N)^{2p}]) ds.
\end{aligned}$$

In order to control the term at the second line above, we use the fact that, for any $x, y \geq 0$, $xy^{2p-1} \leq C_p(x^{2p} + y^{2p})$, whence the term at the second line is smaller than

$$\begin{aligned}
C_{T,p} \left(1 + \left(\frac{1}{N} \sum_{j=1}^N |U_j|^{2p} \right)^{2p} \right) + C_{T,p} \int_0^t \mathbb{E}_{\mathcal{E}} [(X_s^N)^{2p}] ds \\
\leq C_{T,p} \left(1 + \frac{1}{N} \sum_{j=1}^N |U_j|^{4p^2} \right) + C_{T,p} \int_0^t \mathbb{E}_{\mathcal{E}} [(X_s^N)^{2p}] ds.
\end{aligned}$$

Consequently

$$\begin{aligned}
\mathbb{E}_{\mathcal{E}} \left[(X_t^N)^{2p} \right] &\leq \mathbb{E} [(X_0^N)^{2p}] + C_{T,p} \left(1 + \frac{1}{N} \sum_{j=1}^N |U_j|^{4p^2} + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right| \right) \\
&\quad + C_p \left(1 + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right| \right) \int_0^t \mathbb{E}_{\mathcal{E}} [(X_s^N)^{2p}] ds.
\end{aligned}$$

By Grönwall's lemma, we obtain, for all $t \leq T$,

$$\mathbb{E}_{\mathcal{E}} \left[(X_t^N)^{2p} \right] \leq C_{T,p} \left(1 + \mathbb{E} \left[(X_0^N)^{2p} \right] + \frac{1}{N} \sum_{j=1}^N |U_j|^{4p^2} + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right| \right) \exp \left[TC_p \left(1 + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right| \right) \right].$$

Then, to prove Item (i), it is sufficient to integrate over the environment

$$\mathbb{E} \left[(X_t^N)^{2p} \right] \leq C_{T,p} \left(1 + \mathbb{E} \left[|U_1|^{4p^2} \right] + \mathbb{E} \left[\left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right| \right] \right) \mathbb{E} \left[\exp \left(C_{T,p} \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right| \right) \right],$$

and to use Lemma 2.1.

Now let us prove Item (ii). Let us rewrite the dynamics of X^N in the following way

$$X_t^N = X_0^N + \int_0^t b(X_s^N) ds + \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \int_{[0,t] \times \mathbb{R}_+} \mathbf{1}_{\{z \leq f(X_{s-}^N)\}} d\tilde{\pi}_j(s, z) + \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \int_0^t f(X_s^N) ds,$$

where $\tilde{\pi}_j(ds, dz) := \pi_j(ds, dz) - ds dz$ is the compensated Poisson measure of π_j . Then,

$$\begin{aligned} |X_t^N|^\kappa &\leq C_\kappa |X_0^N|^\kappa + C_\kappa t^{\kappa-1} \int_0^t |b(X_s^N)|^\kappa ds \\ &\quad + C_\kappa N^{-\kappa/2} \left| \sum_{j=1}^N U_j \int_{[0,t] \times \mathbb{R}_+} \mathbf{1}_{\{z \leq f(X_{s-}^N)\}} d\tilde{\pi}_j(s, z) \right|^\kappa \\ &\quad + C_\kappa t^{\kappa-1} \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^\kappa \int_0^t f(X_s^N)^\kappa ds. \end{aligned}$$

So,

$$\begin{aligned}
\sup_{t \leq T} |X_t^N|^\kappa &\leq C_\kappa |X_0^N|^\kappa + C_{T,\kappa} \left(1 + \int_0^T |X_s^N|^\kappa ds \right) \\
&\quad + C_\kappa N^{-\kappa/2} \sup_{t \leq T} \left| \sum_{j=1}^N U_j \int_{[0,t] \times \mathbb{R}_+} 1_{\{z \leq f(X_{s-}^N)\}} d\tilde{\pi}_j(s, z) \right|^\kappa \\
&\quad + C_{T,\kappa} \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^\kappa \left(1 + \int_0^T |X_s^N|^\kappa ds \right) \\
\mathbb{E}_\mathcal{E} \left[\sup_{t \leq T} |X_t^N|^\kappa \right] &\leq C_{T,\kappa} \left(1 + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^\kappa \right) \left(1 + \int_0^T \mathbb{E}_\mathcal{E} [|X_s^N|^\kappa] ds \right) \\
&\quad + C_\kappa N^{-\kappa/2} \mathbb{E}_\mathcal{E} \left[\sup_{t \leq T} \left| \sum_{j=1}^N U_j \int_{[0,t] \times \mathbb{R}_+} 1_{\{z \leq f(X_{s-}^N)\}} d\tilde{\pi}_j(s, z) \right|^\kappa \right].
\end{aligned}$$

Then, by Burkholder-Davis-Gundy's inequality,

$$\begin{aligned}
\mathbb{E}_\mathcal{E} \left[\sup_{t \leq T} |X_t^N|^\kappa \right] &\leq C_{T,\kappa} \left(1 + \left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^\kappa \right) \left(1 + \int_0^T \mathbb{E}_\mathcal{E} [|X_s^N|^\kappa] ds \right) \\
&\quad + C_\kappa N^{-\kappa/2} \mathbb{E}_\mathcal{E} \left[\left(\sum_{j=1}^N U_j^2 \int_0^T f(X_s^N) ds \right)^{\kappa/2} \right].
\end{aligned}$$

Set $q_1, q_2 > 1$ such that $1/q_1 + 1/q_2 = 1$ and that $\kappa q_1 < 2p$ (such a q_1 exists since $\kappa < 2p$). Thanks to Hölder's inequality,

$$\begin{aligned}
\mathbb{E} \left[\sup_{t \leq T} |X_t^N|^\kappa \right] &\leq C_{T,\kappa} \left(1 + \mathbb{E} \left[\left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^{\kappa q_2} \right] \right)^{1/q_2} \left(1 + \int_0^T \mathbb{E} [|X_s^N|^{\kappa q_1}] ds \right)^{1/q_1} \\
&\quad + C_{T,\kappa} \mathbb{E} \left[\left| \frac{1}{N} \sum_{j=1}^N U_j^2 \right|^{\kappa/2} \int_0^T \mathbb{E}_\mathcal{E} [f(X_s^N)^{\kappa/2}] ds \right] \\
&\leq C_{T,\kappa} \left(1 + \mathbb{E} \left[\left| \frac{1}{\sqrt{N}} \sum_{j=1}^N U_j \right|^{\kappa q_2} \right] \right)^{1/q_2} \left(1 + \int_0^T \mathbb{E} [|X_s^N|^{2p}] ds \right)^{1/q_1} \\
&\quad + C_{T,\kappa} \mathbb{E} \left[\left| \frac{1}{N} \sum_{j=1}^N U_j^2 \right|^\kappa \right] + C_{T,\kappa} (1 + \mathbb{E} [|X_s^N|^\kappa]),
\end{aligned}$$

where we have used that $\kappa q_1 < 2p$ and that for all $x, y \geq 0$, $xy \leq x^2/2 + y^2/2$ to obtain the last line. Lemma 2.1 and Item (i) proves the result.

3.4. Proof of Lemma 2.4

Let us recall that, in all this proof, we work conditionally on the environment variable W . This proof is very similar to that of Proposition 3.4 of [Erny, Löcherbach and Loukianova \(2022\)](#). We still write the proof here to have the dependency of the control w.r.t. W .

Under Assumption 2, the assumptions of Theorem 1.4.1 of [Kunita \(1986\)](#) are satisfied for the flow of the limit process $(\bar{X}_t^{(x)})_t$. Indeed, the local characteristics (\bar{b}, \bar{a}) is given by

$$\bar{b}(x) = b(x) + Wf(x) \text{ and } a(x, y) = \sigma^2 \sqrt{f(x)f(y)}.$$

Assumptions 1 and 2 of ([Kunita 1986](#), p. 8, 9) are obtained directly from a priori estimates on the second order moments of the process (recalling that we work conditionally on the environment, this point is very classical and therefore omitted). And Assumption 3 (p. 15) is guaranteed by the Lipschitz continuity of b, f, \sqrt{f} (which is weaker than Assumption 2). And the other assumption of Theorem 1.4.1 is exactly Assumption 2.

As a consequence, we know, by Theorem 1.4.1 of [Kunita \(1986\)](#), that for all $t \geq 0$, the function $x \mapsto \bar{X}_t^{(x)}$ is C^3 . Then, to be able to deduce the regularity of the function

$$x \mapsto \bar{P}_{\mathcal{E}, t} g(x) = \mathbb{E}_{\mathcal{E}} \left[\bar{X}_t^{(x)} \right],$$

we just have to differentiate under the expectation. To prove the domination condition for the equations below to be true, we prove a priori estimates on the derivatives of the stochastic flow.

$$\begin{aligned} (\bar{P}_{\mathcal{E}, t} g)'(x) &= \mathbb{E}_x \left[\partial_x \bar{X}_t^{(x)} g'(\bar{X}_t^{(x)}) \right] \\ (\bar{P}_{\mathcal{E}, t} g)''(x) &= \mathbb{E}_x \left[\partial_x^2 \bar{X}_t^{(x)} g'(\bar{X}_t^{(x)}) + \left(\partial_x \bar{X}_t^{(x)} \right)^2 g''(\bar{X}_t^{(x)}) \right] \\ (\bar{P}_{\mathcal{E}, t} g)'''(x) &= \mathbb{E}_x \left[\partial_x^3 \bar{X}_t^{(x)} g'(\bar{X}_t^{(x)}) + 3 \partial_x^2 \bar{X}_t^{(x)} \partial_x \bar{X}_t^{(x)} g''(\bar{X}_t^{(x)}) + \left(\partial_x \bar{X}_t^{(x)} \right)^3 g'''(\bar{X}_t^{(x)}) \right] \end{aligned} \quad (14)$$

Remark 3.3. *To differentiate under the expectation, instead of proving classical domination conditions like*

$$\mathbb{E} \left[\sup_x \left| \partial_x \bar{X}_t^{(x)} \right| \right] < \infty,$$

we rather verify the following uniformly integrable sufficient kind of conditions

$$\sup_x \mathbb{E} \left[\left| \partial_x \bar{X}_t^{(x)} \right|^2 \right] < \infty.$$

See Lemma 3.1 of [Erny \(2022\)](#) (inspired from Lemma 6.1 of [Eldredge \(2018\)](#)) for a formal statement of this result.

To begin with, we know that the first derivative of the stochastic flow is solution to the following SDE.

$$\partial_x \bar{X}_t^{(x)} = 1 + \int_0^t \partial_x \bar{X}_s^{(x)} \left(b'(\bar{X}_s^{(x)}) + W f'(\bar{X}_s^{(x)}) \right) ds + \sigma \int_0^t \partial_x \bar{X}_s^{(x)} (\sqrt{f})'(\bar{X}_s^{(x)}) dB_s.$$

For any $p \in \mathbb{N}^*$, by Ito's formula,

$$\begin{aligned} \mathbb{E}_{\mathcal{E}} \left[\left(\partial_x \bar{X}_t^{(x)} \right)^{2p} \right] &= 1 + \int_0^t \mathbb{E}_{\mathcal{E}} (\partial_x \bar{X}_s^{(x)})^{2p} \left(2pb'(\bar{X}_s^{(x)}) + 2pWf'(\bar{X}_s^{(x)}) + p(2p-1)\sigma^2(\sqrt{f})'(\bar{X}_s^{(x)}) \right) ds \\ &\leq 1 + C_p(1+|W|) \int_0^t \mathbb{E}_{\mathcal{E}} \left[\left(\partial_x \bar{X}_t^{(x)} \right)^{2p} \right] ds. \end{aligned}$$

By Grönwall's lemma,

$$\mathbb{E}_{\mathcal{E}} \left[\left(\partial_x \bar{X}_t^{(x)} \right)^{2p} \right] \leq e^{C_{p,t}(1+|W|)}. \quad (15)$$

Let us prove a similar result on the second derivative of the stochastic flow.

$$\begin{aligned} \partial_x^2 \bar{X}_t^{(x)} &= \int_0^t \partial_x^2 \bar{X}_s^{(x)} \left(b'(\bar{X}_s^{(x)}) + Wf'(\bar{X}_s^{(x)}) \right) ds + \int_0^t \left(\partial_x \bar{X}_s^{(x)} \right)^2 \left(b''(\bar{X}_s^{(x)}) + Wf''(\bar{X}_s^{(x)}) \right) ds \\ &\quad + \sigma \int_0^t \partial_x^2 \bar{X}_s^{(x)} (\sqrt{f})'(\bar{X}_s^{(x)}) dB_s + \sigma \int_0^t \left(\partial_x \bar{X}_s^{(x)} \right)^2 (\sqrt{f})''(\bar{X}_s^{(x)}) dB_s. \end{aligned}$$

Then, by Ito's formula,

$$\begin{aligned} \mathbb{E}_{\mathcal{E}} \left[\left(\partial_x^2 \bar{X}_t^{(x)} \right)^{2p} \right] &= 2p \int_0^t \mathbb{E}_{\mathcal{E}} \left(\partial_x^2 \bar{X}_s^{(x)} \right)^{2p} \left(b'(\bar{X}_s^{(x)}) + Wf'(\bar{X}_s^{(x)}) \right) ds \\ &\quad + 2p \int_0^t \mathbb{E}_{\mathcal{E}} \left(\partial_x^2 \bar{X}_s^{(x)} \right)^{2p-1} \left(\partial_x \bar{X}_s^{(x)} \right)^2 \left(b''(\bar{X}_s^{(x)}) + Wf''(\bar{X}_s^{(x)}) \right) ds \\ &\quad + p(2p-1)\sigma^2 \int_0^t \mathbb{E}_{\mathcal{E}} \left(\partial_x^2 \bar{X}_s^{(x)} \right)^{2p} \left((\sqrt{f})'(\bar{X}_s^{(x)}) \right)^2 ds \\ &\quad + p(2p-1)\sigma^2 \int_0^t \mathbb{E}_{\mathcal{E}} \left(\partial_x^2 \bar{X}_s^{(x)} \right)^{2p-2} \left(\partial_x \bar{X}_s^{(x)} \right)^4 \left((\sqrt{f})''(\bar{X}_s^{(x)}) \right)^2 ds. \end{aligned}$$

Then using that for all $x, y, z \geq 0$,

$$x^{2p-1}y \leq C_p(x^{2p} + y^{2p}) \text{ and } x^{2p-2}z \leq C_p(x^{2p} + z^{2p}),$$

we obtain from the previous inequality

$$\begin{aligned} \mathbb{E}_{\mathcal{E}} \left[\left(\partial_x^2 \bar{X}_t^{(x)} \right)^{2p} \right] &\leq C_p(1+|W|) \int_0^t \mathbb{E}_{\mathcal{E}} \left(\partial_x^2 \bar{X}_s^{(x)} \right)^{2p} ds + C_p(1+|W|) \int_0^t \mathbb{E}_{\mathcal{E}} \left(\partial_x \bar{X}_s^{(x)} \right)^{4p} ds \\ &\quad + C_p \int_0^t \mathbb{E}_{\mathcal{E}} \left(\partial_x^2 \bar{X}_s^{(x)} \right)^{2p} ds + C_p \int_0^t \mathbb{E}_{\mathcal{E}} \left(\partial_x \bar{X}_s^{(x)} \right)^{8p} ds. \end{aligned}$$

Thanks to (15),

$$\mathbb{E}_{\mathcal{E}} \left[\left(\partial_x^2 \bar{X}_t^{(x)} \right)^{2p} \right] \leq C_{p,t} e^{C_{p,t}|W|} + C_p(1+|W|) \int_0^t \mathbb{E}_{\mathcal{E}} \left[\left(\partial_x^2 \bar{X}_s^{(x)} \right)^{2p} \right] ds,$$

whence, by Grönwall's lemma,

$$\mathbb{E}_{\mathcal{E}} \left[\left(\partial_x^2 \bar{X}_t^{(x)} \right)^{2p} \right] \leq C_{p,t} e^{C_{p,t}|W|}. \quad (16)$$

With the same reasoning, we can also prove that

$$\mathbb{E}_{\mathcal{E}} \left[\left(\partial_x^3 \bar{X}_t^{(x)} \right)^{2p} \right] \leq C_{p,t} e^{C_{p,t}|W|}. \quad (17)$$

Finally, using (15), (16) and (17) in the formulae (14) allows to prove the result.

References

- AGATHE-NERINE, Z. (2022). Multivariate Hawkes processes on inhomogeneous random graphs. *Stochastic Processes and their Applications* **152** 86–148.
- BEN AROUS, G. and GUIONNET, A. (1995). Large deviations for Langevin spin glass dynamics. *Probability Theory and Related Fields* **102** 455–509.
- BILLINGSLEY, P. (1999). *Convergence of Probability Measures*, Second ed. Wiley Series In Probability And Statistics.
- CHEVALLIER, J., DUARTE, A., LÖCHERBACH, E. and OST, G. (2019). Mean field limits for nonlinear spatially extended Hawkes processes with exponential memory kernels. *Stochastic Processes and their Applications* **129** 1–27.
- DEMO, A., LUBETZKY, E. and ZEITOUNI, O. (2021). Universality for Langevin-like spin glass dynamics. *The Annals of Applied Probability* **31** 2864–2880. Publisher: Institute of Mathematical Statistics.
- ELDRIDGE, N. (2018). Strong hypercontractivity and strong logarithmic Sobolev inequalities for log-subharmonic functions on stratified Lie groups. *Nonlinear Analysis* **168** 1–26.
- ERNY, X. (2022). Mean field system of a two-layers neural model in a diffusive regime. *Mathematical Neuroscience and Applications* **Volume 2**. Publisher: Episciences.org.
- ERNY, X., LÖCHERBACH, E. and LOUKIANOVA, D. (2022). Mean field limits for interacting Hawkes processes in a diffusive regime. *Bernoulli* **28** 125–149.
- ETHIER, S. and KURTZ, T. (2005). *Markov Processes. Characterization and Convergence*. Wiley Series In Probability And Statistics.
- FOURNIER, N. and LÖCHERBACH, E. (2016). On a toy model of interacting neurons. *Annales de l'Institut Henri Poincaré - Probabilités et Statistiques* **52** 1844–1876.
- GUIONNET, A. (1997). Averaged and quenched propagation of chaos for spin glass dynamics. *Probability Theory and Related Fields* **109** 183–215.
- IKEDA, N. and WATANABE, S. (1989). *Stochastic Differential Equations and Diffusion Processes*, Second ed. North-Holland Publishing Company.
- KOMLÓS, J., MAJOR, P. and TUSNÁDY, G. (1976). An approximation of partial sums of independent RV's, and the sample DF. II. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete* **34** 33–58.
- KUNITA, H. (1986). *Lectures on Stochastic Flows and Applications for the Indian Institute Of Science Bangalore*. Tata Institute Of Fundamental Research Springer-Verlag.
- KUNITA, H. (1990). *Stochastic flows and stochastic differential equations*. Cambridge University Press.

- LUÇON, E. (2011). Quenched Limits and Fluctuations of the Empirical Measure for Plane Rotators in Random Media. *Electronic Journal of Probability* **16** 792–829. Publisher: Institute of Mathematical Statistics and Bernoulli Society.
- PFAFFELHUBER, P., ROTTER, S. and STIEFEL, J. (2022). Mean-field limits for non-linear Hawkes processes with excitation and inhibition. *Stochastic Processes and their Applications* **153** 57–78.
- SHERRINGTON, D. and KIRKPATRICK, S. (1975). Solvable Model of a Spin-Glass. *Physical Review Letters* **35** 1792–1796. Publisher: American Physical Society.
- VILLANI, C. (2008). *Optimal transport, old and new*. Springer.