

Wolbachia and diet modulation interact and affect the performance of an invasive fruit fly

Gwenaëlle Deconninck, Vincent Foray, Jordy Larges, Laureen Beaugeard, Sylvain Pincebourde

▶ To cite this version:

Gwenaëlle Deconninck, Vincent Foray, Jordy Larges, Laureen Beaugeard, Sylvain Pincebourde. Wolbachia and diet modulation interact and affect the performance of an invasive fruit fly. 34e colloque Biotechnocentre, Oct 2022, Nouan-le-Fuzelier, France. . hal-03827196

HAL Id: hal-03827196

https://hal.science/hal-03827196

Submitted on 24 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Wolbachia and diet modulation interact and AFFECT THE PERFORMANCE OF AN INVASIVE FRUIT FLY



GWENAËLLE DECONNINCK, VINCENT FORAY, JORDY LARGES, LAUREEN BEAUGEARD, SYLVAIN PINCEBOURDE

Institut de Recherche sur la Biologie de l'Insecte • UMR 7261 • Université de Tours

Drosophila suzukii



- Drosophilidae from Asia, invasive worldwide
- Highly polyphagous (cultivated and wild fruits)
- Lay eggs on ripening fruits unlike other Drosophila
- Present year-round
- Multiple generations per year (3 to 16)


wSuz

(D. suzukii's Wolbachia strain)

- Variable prevalence (USA:17% / EU: 46%)
- Imperfect maternal transmission but stably maintained in populations
- No consensus on wSuz effect:
 - No cytoplasmic incompatibility
 - Effect on fecundity and viability
- → Wsuz may provide an advantage to D. suzukii when changing host, mitigating the physiological costs induced by its polyphagy

Wolbachia








- Intracellular endosymbiont of arthropods
- Infect ~50% of insect species
- Maternally transmitted
- Maintained by manipulating reproduction or giving ecological advantage

Question: Could wSuz play a role in Drosophila suzukii performance after a diet shift?



Maintained for > 10 generations

Laying box preparation

1 ♀


3 8


Corn or Grape laying medium


N = 25 ♀

25 females of each population were placed individually in laying boxes with 3 males. Each box contained either Corn or Grape medium for the females to lay on. Thus, the female had to lay eggs either on the medium they originated from or on a different medium, thereby simulating a diet shift.

30 days of egg laying monitoring **EVERY DAY** Egg count Pupa count Adult count Egg to Pupa Pupa to Adult Individual weighing survival survival







Medium change Stable medium ***** → ***** Fecundity \oplus Offspring mass

- Fecundity and mass modulated according to the **new diet →** females would favor a laying medium with suitable nutritional composition, otherwise lay fewer and poorer quality eggs
- The effect of diet shift on fecundity and offspring mass lessen after some generations to reach an intermediate value → niche adaptation
- No effect on survival → both media allow complete development

MEDIUM

EFFECT SUMMARY

wSuz might have an energetic cost for the female, reducing its egg production. However, it increased the offspring viability suggesting a mutualistic role. The mutualism might be linked to a **stressful situation** for the fly, such as the medium change.

WSUZ	Stable medium	Medium change
Fecundity	ns / 😑	Θ
Survival	ns	⊕
Offspring mass	ns / 🖯	⊕

PERSPECTIVES

Broaden our knowledge on Wsuz role:

- How can it help D. suzukii in a natural context with multiple fruits?
- What is its physiological cost?
- Can it change if the abiotic environment (e.g., temperature, humidity) vary?
- → Integrative approach is needed to answer
- Slightly decrease in fecundity → Fitness cost
- Enhances larval survival → Better nutrient assimilation?
 - ► Direct effect? E.g. amino acid or iron supplement
 - ▶ Indirect effect? E.g. microbiome modulation
- Increases offspring mass only when medium **changes** → Relevant in a multitrophic context where D. suzukii females lay eggs on different fruits varying in nutritional composition



Gwenaëlle Deconninck

deconninck@univ-tours.fr

ACKNOWLEDGEMENTS

This work is part of the ANR project "What makes Drosophila suzukii such a successful invader? An integrative analysis of its thermal ecology." ANR-20-CE02-0011









