



HAL
open science

Electron Responsive Viologen-Based Supramolecular Gels

Vivien Andrieux, Christophe Bucher, Floris Chevallier, Denis Frath, Thomas Gibaud, Sébastien Manneville

► **To cite this version:**

Vivien Andrieux, Christophe Bucher, Floris Chevallier, Denis Frath, Thomas Gibaud, et al.. Electron Responsive Viologen-Based Supramolecular Gels. Journées de Chimie Supramoléculaire, May 2022, Lyon, France. . hal-03828172

HAL Id: hal-03828172

<https://hal.science/hal-03828172>

Submitted on 25 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Electron Responsive Viologen-Based Supramolecular Gels

*Vivien Andrieux*¹, *Christophe Bucher*¹, *Floris Chevalier*¹, *Denis Frath*¹,
*Thomas Gibaud*², *Sébastien Manneville*²

¹ Laboratoire de Chimie ENS de Lyon 46, allée d'Italie 69364 LYON CEDEX 07, France,

² Laboratoire de Physique ENS de Lyon 46, allée d'Italie 69364 LYON CEDEX 07, France

vivien.andrieux@ens-lyon.fr

In a nutshell, a gel is a material in which a large amount of solvent is trapped in a molecular network. In supramolecular gels, self-assembly of the network is achieved by non-covalent interactions (π - π stacking, H-bonding, ion pairing, solvophobic forces...) between low molecular weight gelators (LMWGs). Unlike polymeric gelators LMWGs have a well-defined chemical structure, offering the possibility to tune the material properties at the molecular level [1]. Stimuli responsive self-assembled molecular materials are capable of changing their organization in response to various stimuli (pH, T, P, magnetic field, light...). Such responsiveness has been proved useful in numerous fields such as catalysis, smart materials, pollutant removal, drug delivery system, development of sensors and memory devices [2]. However little examples of electro-responsive supramolecular materials are to be found in the literature [3]. The potential of such materials has recently been demonstrated through the development of a gel-based memristor exploiting conductivity changes associated to a sol/gel transition triggered by heating or DC bias [4]. Viologens carry much interest for the development of new functional materials and have already shown promising results for applications in molecular electronics [5]. We have recently reported a supramolecular metallo-gel able to undergo a gel-sol transition triggered by photo-reduction of its viologen moiety [6][7][8]. In the wake of this promising result a previously described viologen based gelator [9] has been synthesized and its gelation conditions have been investigated. Rheological and absorbance spectroscopy measurements have shown that the gel properties were strongly dependent on the ionic composition of the solvent, allowing the design of translucent and conductive gels. First photoreduction trials have shown that the viologen based gelator could be photoreduced in its gel state, giving a blue coloration to the material. A home-made cell allowing multi-characterization of the gel redox-responsiveness is currently under development and has shown promising results for the study of the gel electro-responsiveness.

References:

- [1] Panja S. & Adams D. *Chem. Soc. Rev.*, 2021, 50, 5165
- [2] Wu et. al. *J. Adv. Mater.* 2019, 31, 1806204
- [3] Guan et. al. *Supramolecular Chemistry*, 32:11, 578-596
- [4] Zhang et. al. *J. Am. Chem. Soc.* 2017, 139, 41, 14406–14411
- [5] Van Nguyen et. al. *J. Am. Chem. Soc.* 2018, 140, 10131
- [6] Kahlfuss et. al. *Chem. Eur. J.* 2018, 24, 13009 – 13019
- [7] Chowdhury et. al. submitted, (arXiv:2106.15688)
- [8] Clément Roizard et al 2022 *ECS Adv*
- [9] Xue et. al. *Chem. Mater.* 2004, 16, 3702–3707